The formation of microbial exoskeletons is driven by a controlled calcium-concentrating subcellular niche

Alona Keren-Paz[1*], Malena Cohen-Cymberknoh[2*], Dror Kolodkin-Gal[3*], Shani Peretz[1], Iris Karunker[1], Sharon G. Wolf[4], Tsviya Olender[1], Sergey Kapishnikov[4], Vlad Brumfield[4], Simon Dersch[5], Elena Kartvelishvily[4], Peninnah Green-Zelinger[1], Damilola Isola-Adeyanju[1], Ronit Suissa[1], Michal Shteinberg[6], Daniel McLeod[7], Marianna Patrauchan[7], Gideon Zamir[3], Assaf Gal[8], Peter L. Graumann[5], Eitan Kerem[2**], and Ilana Kolodkin-Gal[1**]

[1] Department of Molecular Genetics, Weizmann Institute of Science, Rehovot, Israel

[2] Pediatric Pulmonary Unit, Hadassah-Hebrew University Medical Center, Jerusalem, Israel

[3] Department of Experimental Surgery, Hadassah-Hebrew University Medical Center, Jerusalem, Israel

[4] Chemical Research Support, Weizmann Institute of Science, Rehovot, Israel

[5] Centre for Synthetic Microbiology (SYNMIKRO), and Fachbereich Chemie, Philipps-Universität Marburg, Marburg, Germany.

[6] Pulmonology Institute and CF Center, Carmel Medical Center, Haifa, Israel

[7] Department of Microbiology and Molecular Genetics, Oklahoma State University, Stillwater, OK, USA

[8] Department of Plant and Environmental Sciences, Weizmann Institute of Science, Rehovot, Israel

* Equal Contribution

** Corresponding authors: Ilana.kolodkin-gal@weizmann.ac.il; EitanK@hadassah.org.il

Abstract

In nature, bacteria reside in biofilms - multicellular differentiated communities held together by extracellular matrix. In this work, we identified a novel subpopulation essential for biofilm formation – mineral-forming cells. This subpopulation contains an intracellular calcium-accumulating niche, in which the formation of a calcium carbonate mineral is initiated. As the biofilm colony develops, this mineral grows in a controlled manner, forming a functional macrostructure that serves the entire community.

The molecular mechanisms promoting calcite scaffold formation were conserved between three distant phyla – the Gram-positive *Bacillus subtilis*, Gram-negative *Pseudomonas aeruginosa* and the actinobacterium *Mycobacterium abscessus*. Biofilm development of all three species was similarly impaired by inhibition of calcium uptake and carbonate accumulation. Moreover, chemical inhibition and mutations targeting mineralization both significantly reduced the attachment of *P. aeruginosa* to the lung, as well as the subsequent damage inflicted by biofilms to lung tissues, and restored their sensitivity to antibiotics.

The evolutionary conserved cellular pathway controlling the fundamental feature of biofilm development uncovered in this work offers novel druggable targets for antibiotics to combat otherwise untreatable biofilm infections.

Introduction

While bacteria have been historically studied in homogeneous monocultures, in natural ecosystems and in clinical settings they are typically found in interface-associated multicellular communities called biofilms (Kolter and Greenberg, 2006). It was recently indicated that in soils, oceans, the deep subsurface, phyloshpere and animal microbiomes, 40−80% of bacteria reside in biofilms (Flemming and Wuertz, 2019). Bacterial biofilms are differentiated communities, in which the cells are held together by extracellular matrix. The formation of a biofilm is frequently favored over free-living planktonic life-style, as it increases the fitness of member cells promoting their survival, such as better attachment to hosts, division of metabolic labor and protection from the environment.

Microbial biofilms are of extreme clinical importance, as they are associated with many persistent and chronic infections (Costerton et al., 1999). A prominent feature of biofilms is their inherent resistance to the immune system and to antibiotics (Bryers, 2008; Bucher et al., 2019; Hill et al., 2005) – biofilm cells are up to 1,000 times more tolerant than planktonic bacteria (Bryers, 2008), making the eradication of biofilm infections with currently available antibiotics extremely challenging.

For example, the commensal/aquatic bacterium *Pseudomonas aeruginosa* can cause devastating chronic biofilm infections in compromised hosts, such as cystic fibrosis (CF) patients and those with burn wounds or implanted medical devices (Costerton et al., 1999). Respiratory infections with *P. aeruginosa* are a leading cause of morbidity and mortality in patients with CF (Yoon and Hassett, 2004). Once a chronic infection with *P. aeruginosa* is established, it is almost impossible to eradicate (Cohen-Cymberknoh et al., 2016). Similarly, the biofilms of *Mycobacterium abscessus* cause infections of respiratory tract, skin and central nervous system in patients with CF and chronic obstructive pulmonary disease (Caimmi et al., 2018).

A hallmark of biofilms is their ability to form robust and complex 3D structures, and the distinctive spatial organization provides the residing bacteria with several

benefits. The biofilm architecture was suggested to relieve metabolic stress. For
example, channels formed below the ridges and wrinkles within the colony may
facilitate diffusion of fluids, nutrients and oxygen (Bloom-Ackermann et al., 2016;
Dietrich et al., 2013; Kolodkin-Gal et al., 2013; Wilking et al., 2013). Furthermore,
cells located in different areas of the colony are exposed to different levels of
oxygen, nutrients and quorum sensing molecules, which affect the genetic
programs they express, promoting differentiation (Asally et al., 2012; Hassanov et
al., 2018; Liu et al., 2015; Monds and O'Toole, 2009; Serra and Hengge, 2014;
Stewart and Franklin, 2008). Thus, community structure allows the genetically
identical biofilm cells to display phenotypic heterogeneity and to endure stressful
environments. Due to physical protection offered by the biofilm biomass,
differentiation and physiological adaptations, cells residing deep within the biofilm
are protected from environmental assaults, such as antibiotics. Biofilm colonies are
a compelling model system for biofilm development, and many basic biological
processes of high relevance to bacterial pathogenicity were first discovered in this
well-controlled and robust system (Colvin et al., 2012; Dietrich et al., 2008; El
Mammeri et al., 2019; Hufnagel et al., 2018; Jo et al., 2017; Richards et al., 2019;
Serra et al., 2013; Steinberg et al., 2020; Wermser and Lopez, 2018).

Until recently, the ability of biofilm-forming bacteria to generate complex
architectures was attributed exclusively to self-produced **organic** extracellular
matrix (ECM) (Dragos and Kovacs, 2017; Reichhardt and Parsek, 2019; Steinberg
and Kolodkin-Gal, 2015), composed of carbohydrate-rich polymers (i.e., lipids or
exopolysaccharides), proteins, and nucleic acids (Branda et al., 2005). However,
we and others have recently shown that microbial biofilms contain an organized
internal mineral structure, composed of crystalline calcium carbonate (calcite), that
also contributes to their 3D morphology (Keren-Paz et al., 2018; Keren-Paz and
Kolodkin-Gal, 2020; Li et al., 2015).

Formation of minerals is known to be induced in bacterial biofilms, through passive
surface-mediated processes promoted by by-products of bacterial metabolism.
Biologically induced minerals associated with bacteria include oxides of Fe, Mn,

and other metals; metal sulfates and sulfites; phosphates and carbonates; and Fe
and Fe-Al silicates (Frankel and Bazylinski, 2003). Of all examples of
biomineralization, microbial induced calcium carbonate precipitation (MICCP) is
most frequently associated with microbial communities (Weiner and Dove, 2003),
(Douglas and Beveridge, 1998). In geological settings, bacterial metabolic
processes, such as urea hydrolysis by urease-positive bacteria, increase local
concentration of bicarbonate and elevate pH. When enough environmental
calcium is present, those changes promote spontaneous precipitation of calcium
carbonate, with bacterial envelopes serving as nucleation sites for the growing
mineral, leading to formation of crystalline calcium carbonate (Dhami et al.,
2013). Similar processes were also suggested to promote biofilm-associated
calcification of *Proteus mirabilis* on catheters (Morris and Stickler, 1998), and
during dual-species biofilm formation (Li et al., 2016b) and to contribute to *P.
aeruginosa* virulence in an insect host (Lotlikar et al., 2019). In both clinical and
environmental scenarios, calcium deposition in biofilms was mostly seen as an
uncontrolled and unintentional byproduct of bacterial metabolic activity.

Recently, intracellular amorphous calcium carbonate (ACC) granules were
detected in some species of Gram-negative autotropic bacteria (Blondeau et al.,
2018). Those intracellular deposits were suggested to contribute to single cell
energy metabolism by promoting photosynthesis (Blondeau et al., 2018), or
chemolithoautotrophy (Monteil et al., 2020). While the precise molecular
mechanisms regulating this process are still unresolved, the presence of a
microcompartment dedicated to mineralization raises the possibility that some
bacteria might be capable of controlling the formation of biogenic mineral.

Our recent discovery of precisely organized mineral macrostructures within
biofilms suggests that biomineralization in biofilms is tightly controlled. These
calcite 'skeletons' contributed to fitness of biofilm colonies in two unrelated soil
bacteria: *Bacillus subtilis* and *Mycobacterium smegmatis* (Keren-Paz et al.,
2018; Oppenheimer-Shaanan et al., 2016). For both, the mineral structure acted
as a structural scaffold supporting the 3D architecture of the colony, and as

diffusion barrier preventing the penetration of solutes into the biofilms. The formation of controlled calcium carbonate macrostructures in phylogenetically distinct heterotrophic biofilms raises the possibility that the function of calcium carbonate is not limited to specific species utilizing it for intracellular chemical energy, but is instead a general phenomenon wide-spread in the bacterial kingdom.

Those reports challenge the current view of biofilm development as a process depending solely on organic ECM production. They raise several fundamental questions regarding the developmental role of mineral skeleton formation within a bacterial community, and the relation between structure and function in microbial biofilms. If indeed mineralization is a controlled process resulting in a structure crucial to biofilm development and function, this is a novel aspect of basic biofilm biology. What is the role of mineral structure in the development of differentiated biofilm community? Are there dedicated 'osteoblast-like' cells initiating mineral production? What are the fitness advantages that the mineral exoskeleton confers to the community? Furthermore, organic matrices and their regulation differ between different organisms (Steinberg and Kolodkin-Gal, 2015). However, the formation of calcium carbonate relies on a highly conserved building blocks, calcium and carbonate, generated from carbon dioxide in most heterotopic organisms. Therefore, the generation of mineral scaffolds could be common and conserved across the bacterial domain, contributing to the phenotypical resistance of diverse microbial biofilms. If so, understanding the underlying molecular mechanisms of biofilm mineralization is a crucial step towards our ability to successfully combat biofilm infections.

In this work, we discovered that the calcium-dependent 3D organization of the biofilm colony leads to transcriptional reprogramming of the bacterial community, and was essential for biofilm development. Calcite formation and the assembly of a functional mineralized macro-skeleton supporting the 3D morphology of a biofilm colony was associated with defined intracellular calcium-rich deposits present in a distinct subpopulation of biofilm cells, differing from flagellin and

organic ECM producers. Calcite formation turns out to be an extremely conserved process, occurring not only in the beneficial model organism *B. subtilis*, but also in two unrelated lung pathogens – *P. aeruginosa* and *M. abscessus*. Inhibition of key biomineralization enzymes or calcium uptake prevented biofilm formation by both pathogens. We were able to identify calcite in sputum samples taken from CF patients, suggesting that this conserved process is of clinical importance. Finally, in an *ex vivo* lung model, chemical and genetic inhibition of calcium uptake and of carbonate accumulation blocked biofilm formation and lung colonization, preventing damage inflicted by *P. aeruginosa* to lung tissues, and sensitized *P. aeruginosa* to antibiotic treatment.

Taken together, our results identify a previously overlooked process essential for bacterial biofilm development – **the tightly regulated formation of mineral skeletons by dedicated cells**. Its conservation across the bacterial kingdom highlights the fundamental role it plays in biofilm biology, and could lead to novel therapeutic approaches for combating a broad range of persistent biofilm infections.

Results

In a previous work, we demonstrated that *B. subtilis* biofilm morphology is calcium-dependent (Oppenheimer-Shaanan et al., 2016). Removing calcium acetate from the undefined biofilm-promoting B4 medium prevented the development of the complex 3D architecture characteristic of a biofilm colony, without affecting bacterial growth. This 3D structure required calcium (as no equivalent cation could replace it and as it was induced by various calcium salts) and metabolically produced carbonate (as it was eliminated under anaerobic conditions and increased in CO_2-enriched environment). The mineral component was identified as calcite by FTIR and XRD (Supplementary Fig. 1, (Oppenheimer-Shaanan et al., 2016). It spanned the entire colony and was spatially co-localized with colony wrinkles, suggesting it acted as a "scaffold" supporting them (Oppenheimer-Shaanan et al., 2016). The mineral structure developed and increased over time

as the colony grew (Keren-Paz et al., 2018). Finally, as the biofilm colonies aged
and their 3D spatial structure diminished, the internal and organized mineral
deteriorated (potentially due to fermentation and acidification of the medium by
oxygen-depleted cells within the inner mass of the aging biofilm), and only large
non-structural crystals were left in the colony periphery due to a passive growth of
calcite crystals (Oppenheimer-Shaanan et al., 2016). During biomineralization, the
formation of exopolysaccharides and the amyloid protein TasA was also essential
to form complex morphology, and interactions between the organic and inorganic
matrix were evident (Oppenheimer-Shaanan et al., 2016).

To better understand the molecular mechanisms underlying the formation of
functional mineral macrostructures, we examined the effect of calcium on gene
expression in a *B. subtilis* biofilm colony by sequencing the transcriptome of
colonies grown either with or without excess calcium (Fig. 1A). The addition of
calcium resulted in a dramatic reprogramming of the transcriptional profile of the
colony, with 20% of the genome (n=875) significantly changing between the two
conditions at all time-points tested (Supplementary File 2). Only in the presence of
added calcium, specific developmental processes defining the biofilm state, such
as induction of ECM production and sporulation, and the repression of motility,
were observed (Fig. 1B). These fundamental changes were mostly sustained over
time (Fig. 1C). We next compared the two transcriptional signatures to a collection
of published data, containing 269 mRNA profiles of *B. subtilis* grown under over
100 different conditions (Nicolas et al., 2012) (Fig. 1D). The transcriptome of cells
grown with excess calcium was similar to that of previously analyzed biofilm cells,
regardless of the biofilm promoting medium used. On the other hand, the
transcription profile of colonies grown without added calcium was similar to that of
planktonic cells experiencing nutrient starvation, but was not identified as biofilm
in this unbiased analysis.

These findings highlight the central role of calcium in biofilm development, and
suggest an intimate connection between biofilm structure and function - raising the
possibility that microbial cells are actively regulating calcium carbonate
biomineralization. Indeed, when a biofilm colony was visualized by scanning

electron microscopy (SEM), we observed a highly mineralized subpopulation of cells, tightly associated with mineral crystals (Fig. 1E, Supplementary Fig. 2). Consistently with our previous characterization of bleached biogenic calcite, the unbleached crystals had rough faces, along with a smooth and flat crystal face and displayed elongated prismatic morphology instead of the rhombohedral morphology of calcite grown in pure solution (Oppenheimer-Shaanan et al., 2016).

The crystals were consistent with biogenic calcite generated by *B. subtilis (Oppenheimer-Shaanan et al., 2016)*, as judged by shape; backscatter mode and energy dispersive X-ray spectroscopy (EDX) (Fig. 1F, Supplementary Fig. 2). In the absence of added calcium, mineral crystals were rarely detected (Supplementary Fig. 3). Mineral producers were not associated with organic ECM, did not carry flagella, and did not sporulate (Figures 1E, Supporting Figures 2 and 3), and therefore morphologically differed from previous subpopulations observed in *B. subtilis* biofilms (Vlamakis et al., 2008). To create the 3D reconstruction, a whole, unfixed bacterial colony was transferred to a plastic slide for a microCT scan, and rotated between the X-ray source and the detector positioned at optimal distances for a voxel size of 0.87 μm. 2D projections were taken at different angles until a full rotation (360°) was completed. The full set of images was then used to reconstruct the whole volume of the sample by back projection algorithm and thus a high-resolution 3D image was generated (see Supplementary Movie 1). This microCT X-ray scan confirmed that calcite is organized in a macro-structure, generating a non-uniform continuous layer throughout the wrinkles (Fig. 1G).

One possible mechanism linking calcium mineralization and colony structure is that, in the presence of calcium, some kind of specialized cells located in specific regions of the colony, would serve as nucleation sites and promote localized mineral formation. The observed induction of sporulation regulon transcription by calcium (Fig. 1B) raised the possibility that spores could serve this function, possibly due to unique membrane properties of the spore. However, a mutant strain lacking sporulation sigma factor SigF (de Hoon et al., 2010), which does not form spores and is blocked early during sporulation, was able to form highly structured colonies, in a calcium-dependent manner (Supplementary Fig. 4),

suggesting that biomineralization promotes biofilm development and sporulation,
and not *vice versa*.

The tight association of calcium carbonate mineral with the bacterial cells led us to
examine the role of intracellular calcium. Disruption of intracellular calcium
homeostasis by a chemical inhibitor of P-type ATPases sodium metavanadate
(SMV) (Clausen et al., 2016; Guragain et al., 2013; Kuhlbrandt, 2004), prevented
the formation of a complex 3D structure even in the presence of excess calcium
(Fig. 1H). This morphological defect was not due to inhibition of planktonic growth,
which was only mildly affected by SMV (Fig. 1I). This need for controlled
intracellular calcium levels raised the possibility that carbonate mineralization
starts intracellularly. Therefore, we next examined single cells from biofilms grown
with excess calcium.

3D reconstructions of cryo-fixed bacteria with scanning transmission electron
tomography (CSTET) (Wolf et al., 2014) (Fig. 2A-B) revealed calcium dense foci
inside the cells, associated with the membrane. CryoSTEM imaging and cryo-EDX
measurements verified excess calcium in these deposits (Supplementary Fig. 5).
The appearance of these deposits was calcium dependent, as we could not
observe them in cells grown without added calcium (Supplementary Fig. 6). While
the mineral producing sub-population was small, it was enriched in the structured
zones of the colony (Fig. 2C). Next, we examined the distribution of cellular calcium
levels within the biofilm community. We stained live biofilm cells with calcein-AM –
a dye that can be used for visualization of intracellular calcium in live cells (Hale et
al., 2000). This lipophilic stain can freely enter the cells, but only binds calcium and
becomes fluorescent after the acetoxymethyl (AM) groups are removed by cellular
esterases. Gated-STED fluorescent microscopy revealed heterogeneity in cellular
calcium levels within the biofilm colony, with most cells containing low levels of
calcium, while a smaller fraction contained very high levels (Fig. 2D). The number
of cells showing a strong calcein signal increased during biofilm development, as
the calcified structures matured. While the cytoplasm of most cells was stained
evenly, a small fraction of cells (2.6% out of 300 cells examined) displayed
concrete, membrane-localized calcium deposits (Fig. 2E), comparable in their

frequency, distribution and localization with calcium deposits identified by CSTET

(Fig. 2A-C). Consistent with our previous observation, sporulation does not seem

to be involved in cellular calcium deposition – as Δ*sigF* cells contained calcium-

rich foci similar to those of wild type (Supplementary Fig. 7). Furthermore, we could

detect such deposits associated with the cell envelope on the cytoplasm side,

embedded within the membrane, and closely associated with the cell from the

outside – suggesting different stages of export (Fig. 2F). Similar calcium deposits

were also observed by CSTET in close association with the outer membrane of

bacterial cells (Fig. 2G). Taken together, those finding support the existence of a

calcium-concentrating compartment in a subpopulation of heterotropic biofilm

cells, similar to that recently discovered in autotrophic bacteria (Blondeau et al.,

2018), and that the mechanism for its formation involves an active and regulated

calcium uptake, as well as the export of the stored calcium (Fig.2F).

In *B. subtilis*, the transcriptional regulation of calcium uptake is poorly

characterized. We therefore examined the potential involvement of several general

transcriptional regulators of cation uptake. Non-ferric metal homeostasis in

bacteria is regulated by the MntR, MerR and ArsR familes of transcriptional

regulators (Moore and Helmann, 2005). Strains lacking *arsR, mntR* and *zur*

developed robust and structured biofilm colonies. In contrast, *cueR* mutant had a

severely altered biofilm phenotype and formed completely flat colonies (Fig. 2H).

CSTET analysis was suggestive of diminished calcium-concentrating population

(a maximum of 1.3% compared with 4% of the wild type, n=149). CueR is a poorly

characterized transcriptional regulator belonging to the metal-responsive MerR

family (Moore and Helmann, 2005). CueR binds the promoter of copper

transporting P-type ATPase CopA, but a deletion of *copA* had no effect of biofilm

morphology (Supplementary Fig. 8). The effect on biofilm morphology was also not

due to reduction in growth of *cueR* mutant (Supplementary Fig. 9). Those

observations raise the possibility that CueR regulates additional, yet undiscovered,

P-type ATPase, involved in calcium uptake. To characterize the regulon of CueR,

we repeated the transcriptome analysis of the wild-type strain and of *cueR* mutant,

both grown in the presence of calcium. When compared to the wild type colony,

the expression of matrix genes was upregulated and motility was inhibited in the
cueR mutant, suggesting a general dis-regulation of biofilm transcriptional profile
(Fig. 2I). Biofilm formation is a tightly regulated developmental process, and both
inhibition and overexpression of key components is known to cause severe defects
in biofilm development (for example, both deletion and over-activation of the biofilm
regulator Spo0A prevent normal biofilm development (Veening et al., 2006). High
levels of ECM gene expression further indicates that the featureless morphology
of the *cueR* mutant is not a result of the lack of organic ECM components. The
subsequent analysis of the expression of divalent cation transporters reveled that
their expression is significantly altered in *cueR* mutant, while other transporters are
less affected (Fig. 2I). Among those, we identified three calcium transporters with
transcriptional levels significantly changed in the mutant: *yloB* (ATP-driven Ca^{2+}
pump) (Gupta et al., 2017) *, chaA* (H^+/Ca^{2+} exchanger) (Fujisawa et al., 2009) and
yetJ (pH-dependent calcium leak channel) (Chang et al., 2014) (Supplementary
File 2). Deletion of *yloB* (but not *chaA*, *yetJ*, or both) was sufficient to prevent 3D
colony morphology formation, indicating the importance of calcium uptake and
homeostasis (Fig. 2J and Supplementary Fig. 10).

Taken together, these observations suggest that the first critical step in the
formation of calcite mineral is initiated in saturated mineral precursor
microenvironments within the cell. After the initial nucleation, the precursor seems
to be exported out of the bacterial cell without hampering the membrane integrity
of its producer, and to further grow on organic ECM templates in the highly complex
biofilm microenvironment, until an organized functional mineral macro-structure is
formed.

The organic ECM components and the molecular mechanism governing
transition from planktonic life style to the biofilm state vary immensely between
phylogenetically distant bacteria, making a development of broad range drugs
specifically targeting biofilms challenging. In this sense, biomineralization is an
extremely appealing target, as it relies on simple and conserved mechanisms
allowing accumulation of calcium and carbonate. We have recently reported that,
like *B. subtilis*, the biofilm colonies of soil actinobacterium *M. smegmatis* contain

functional mineral macro-structures (Oppenheimer-Shaanan et al., 2016). One
additional study provided evidence for some spatial organization of calcium
carbonate mineral in submerged biofilms formed by the Gram-negative
pathogen *P. aeruginosa* (Li et al., 2016a). This led us to explore whether we can
detect regulated formation of functional mineral structures in pathogenic biofilm
colonies.

When grown on biofilm-inducing agar, *P. aeruginosa* PA14 forms complex
colonies with a highly characteristic morphology, which were shown to be
dependent on polysaccharides encoded by the *pel* operon (Colvin et al., 2011)).
However, like in *B. subtilis,* these complex 3D structures were co-localized with
spatially organized mineral, as determined by microCT (Fig. 3A), were abolished
by the cation chelator EDTA and EGTA, and could be restored by addition of
excess calcium (Fig. 3B, C). 3D reconstruction confirmed that the densest mineral
areas formed non-continuous layer correlated with colony wrinkles (Fig. 3D),
further suggesting that the 3D structure of *P. aeruginosa* biofilms likely relies on
the formation of calcite mineral scaffolds. The formation of elaborate calcium-
dependent 3D morphology was conserved in an additional strain of *P. aeruginosa*
PA01 (Supplementary Fig. 11). The mineral and exopolysaccharides were both
essential for structure formation, as a mutant for both *pel* and *psl* operons was flat
and featureless, similarly to biofilms grown in the presence of a calcium chelator
(Supplementary Fig. 12).

To indicate whether the formation of mineral scaffolds is a conserved process of
significant clinical potential, we tested whether it occurs in another,
phylogenetically distant, drug-resistant pathogen. We therefore examined the
biofilms of *M. abscessus,* which are held together by extracellular matrix
consisting of mycolic acids (Halloum et al., 2016), which bear little resemblance
to EPS components of *B. subtilis* and *P. aeruginosa*. Nevertheless, this
bacterium also only formed colonies of complex morphology in the presence of
calcium (Fig. 3E). Fourier-transform infrared spectroscopy (FTIR) detected
calcium carbonate mineral in the colony (Supplementary Fig. 13), and X-Ray
analysis revealed that the densest mineral areas were co-localized with colony

wrinkles (Fig. 3F), just like in *B. subtilis* and *P. aeruginosa* colonies. In addition to the organized macro-scale structures present throughout the colony, over time large crystals also formed beneath the colony and in its periphery, reminiscent of crystals passively precipitated in the vicinity of *B. subtilis* colonies at later stages. All the bacteria we tested formed mineral macrostructures in a slightly basic-neutral macro pH (Supplementary Fig. 14). Under such conditions, passive mineral precipitation is unlikely, supporting a potential biological regulation of mineralization.

The cellular pathways of biomineralization present in all three bacteria are strikingly conserved (Fig. 4A), and can be inhibited at several points. The role of urease and carbonic anhydrase (CA) in biomineralization is well established (Jansson and Northen, 2010; Phoenix and Konhauser, 2008). The consequent action of those enzymes increases local bicarbonate levels, and at the same time increases the pH, creating an environment favorable to mineral precipitation. In the section above, we have shown that intracellular calcium was needed for calcite formation. Consistent with the central role of calcite scaffolds in supporting biofilm morphology, inhibition of urease, CA, and especially P-type ATPases, disrupted *P. aeruginosa* biofilm morphology (Fig. 4B) and abolished the mineral macrostructures as judged by microCT X-Ray (Supplementary Fig. 15). This effect could not be explained by inhibition of planktonic growth, which was mostly unaffected (Fig. 4C, Supplementary Fig. 16). Similarly, reducing self-produced and atmospheric carbon dioxide reduced the mineral macrostructure (Supplementary Fig. 17).

Inhibition of complex calcium-dependent colony formation, but not planktonic growth, was also observed in mutants lacking two or more carbonic anhydrase genes or a calcium transporter CalC (Fig. 4D). Biofilm formation on polystyrene, as quantified by crystal violet staining, was also severely compromised by either chemically or genetically impairing carbonate formation or calcium uptake (Fig. 4E-F). The fact that SMV addition or CalC deletion could prevent 3D morphology development and biofilm formation, but had little effect on planktonic growth, is

consistent with our view that carbonate mineralization in biofilms relies on
regulation of intracellular calcium homeostasis.

Potentially broad clinical implication of these findings was supported by testing
the same chemical inhibitors on *M. abscessus*. Colony morphology could be
compromised by inhibiting biomineralization enzymes (Fig. 4G and
Supplementary Fig. 19). Morphology defects could not be explained by inhibition
of cell growth, as most potent morphology inhibitors AHA and SMV had the
smallest effect on planktonic growth (Fig. 4G). We could therefore conclude that
biomineralization is indeed highly conserved, and is mediated by a previously
untargeted druggable pathway.

Biofilm mineralization was not previously demonstrated in the lung
microenvironment. To test the ability of lung pathogens to form structured
communities in such conditions, we grew *M. abscessus* and *P. aeruginosa* on
synthetic CF sputum medium (SCFM). Both species were capable to form highly
organized 3D colonies on SCFM (Supplementary Fig. 20). For *P. aeruginosa*, this
robust colony architecture was enhanced by addition of calcium to the growth
medium, and diminished with the addition of calcium chelator EGTA (Fig. 4H);
suggesting that biofilms may rely on crystalline calcium carbonate scaffolds to
support the community architecture in the lung. Therefore, we examined whether
calcite formation by lung pathogens could be also observed in clinical samples.
We collected sputum samples from CF patients chronically infected with *P.*
aeruginosa and removed the organic matter by bleach. Insoluble material
indicating the presence of a mineral was found in 31 out of 50 samples examined
(Supplementary Table 3). These putatively positive patients were further examined
by FTIR for residual pure minerals, and spectra typical of calcite, lacking a v1 peak
and containing a v_2 peak at 875 cm^{-1}, v_3 peak at 1425 cm^{-1} , and v_4 peak at 713 cm^{-1}
^{1} (Supplementary Figure 21), were identified in 7 samples.

To further explore the role of biomineralization during lung infection by *P.*
aeruginosa, we set up a lung *ex vivo* model. Organ cultures derived from lung and
colon tissues have been applied previously for the study of biofilm infections and

viral tropism (Kolodkin-Gal et al., 2008; Kolodkin-Gal et al., 2007; Massler et al.,

2011; Wu et al., 2021), and thus can potentially be used to study lung infections.

We cultured mice lungs and then infected them with *P. aeruginosa* constitutively

expressing GFP. Fluorescent microscopy revealed biofilms forming on the tissue

prior to its destruction (Supplementary Fig. 22). Using this system, we then tested

the effect of chemically blocking biomineralization during initial infection stages.

When left untreated, *P. aeruginosa* infected tissue (labeled by DAPI for

visualization) was completely deteriorated. Infection in the presence of urease

inhibitor AHA prevented biofilm formation and rescued the lung tissue (Fig. 5A).

While the tissue was rescued, the bacterial load in the growth media remained

similar (Fig. 5B). This is consistent with the *in vitro* results presented above – as in

all cases inhibition of biomineralization prevented biofilm formation, but had little

effect on planktonic growth. This further strengthens our proposed hypothesis,

suggesting biomineralization is a specific and controlled developmental feature of

bacterial biofilms. To quantify the effects of urease inhibition during lung tissue

infection, the tissue was stained with H&A and the nuclei were counted (Fig. 5C-

D). We observed a significant rescue of the infected tissue by AHA, in a dose-

dependent manner. In agreement with these findings, the growth of bacteria in the

medium was not reduced by AHA addition, but their colonization of the lung tissue

was largely compromised (Fig. 5E and 5F).

Significant reduction of lung cell death during infection by *P. aeruginosa* was also

obtained with CA and calcium uptake inhibitors (Fig. 6A-C), and with mutants

lacking CAs (Fig. 6D). Consistent with our observation that both Diamox and AHA

inhibited lung colonization and subsequent cell death, the combination of both was

additive (Figure 6E). At the concentrations used here, AHA and DTNB were slightly

toxic to the uninfected lung tissue, while SMV and Diamox had no adverse effect

(Supplementary Fig. 23). We can therefore conclude that the entire

biomineralization pathway, rather than a specific enzyme, is of crucial importance

to host infection.

Finally, while the ability to rescue host tissue by inhibition of biomineralization is

encouraging, in clinical settings the eradication of bacteria is essential. We

therefore tested whether inhibition of mineralization during lung infection of *P. aeruginosa* could increase its sensitivity to antibiotics. While neither biomineralization inhibitors nor quinolone antibiotics (a first-line antibiotic for treating *P. aeruginosa* infections) (Hewer and Smyth, 2017) could effectively eradicate established biofilms, the combination of both classes of drugs significantly increased the sensitivity of biofilms to treatment (Fig. 6F) – highlighting the potential clinical applications of biomineralization inhibition.

Discussion

Bacterial biofilms are widespread in nature, forming multicellular colonies on various surfaces. The molecular mechanisms guiding the complex network of events leading to the transition from a free-living planktonic bacterium to the differentiated community are well characterized. Much is known about the production of organic matrix, the division of labor between the different cell types, the metabolic adaptations, the methods of communication between the community members – and the fitness advantages of such a communal life style are evident.

In addition, bacterial biofilms were extensively shown to promote biomineralization of metals (such as gold and iron), phosphate and carbonate (Phoenix and Konhauser, 2008). In this work, we uncover a previously overlooked, but fundamental, connection between the highly regulated biofilm development and the seemingly random biomineralization occurring within biofilms. We propose that bacterial communities actively control the formation of mineral macrostructures and that those "skeletons" play an important role in biofilm development and fitness.

Moreover, the conserved and wide-spread nature of biomineralization offers novel therapeutic approaches to target one of the most challenging global threats to human health – drug-resistant chronic biofilm infections.

The first clues to the presence of a structural mineral component in a biofilm came when we demonstrated that a mature 3D structure of biofilm colonies of *B. subtilis* depends on their ability to precipitate precisely organized patterns of calcium carbonate (Oppenheimer-Shaanan et al., 2016). A controlled deposition of a calcium carbonate scaffold structurally supports the complex architecture of biofilm colonies and protects them from the environment by limiting solute diffusion (Keren-Paz et al., 2018).

In this work, we build on those preliminary observations to show that crystalline calcium carbonate production is a controlled process fundamental to biofilm development. Only when calcification occurs, *B. subtilis* can successfully activate and maintain the characteristic biofilm transcriptional program, activating matrix production and sporulation while repressing motility (Fig. 1).

Mechanistically, the formation of mineral macro-structures is initiated within intracellular compartments of a small subpopulation of cells, which secrete small calcium-rich granules, while remaining intact (Fig. 2). The process is likely actively controlled, as the membrane integrity of the exporting cells is not compromised. Initiating mineral formation within the controlled cellular environment allows the organism to regulate the composition, structure, and physical properties of the mineral – therefore controlling the final function.

The best known example of bacteria producing functional minerals formed in a defined intracellular compartment by a highly regulated cellular pathway is the magnetotactic bacteria, forming invagination of membranes to generate magnetosomes (Komeili et al., 2006). Recently, intracellular calcium carbonate mineralization was reported in cyanobacteria (Blondeau et al., 2018) and in a subgroup of poorly characterized magnetotactic bacteria (Monteil et al., 2020). Unlike the above examples, which occur in a specific group of phylogenetically related planktonic aquatic bacteria, the formation of mineralized macrostructures in biofilms is conserved across the microbial domain, highlighting their potential importance for general microbial fitness, unrelated to specific adaptations of single-cell microbial metabolism.

Regulated calcification was found to mediate complex 3D biofilm structure in three
phyla – the Gram-positive spore former *B. subtilis* and actinobacterium *M.*
abscessus, and the Gram-negative *P. aeruginosa* (Fig. 3). The conserved nature
of cellular pathways promoting biomineralization allowed us to prevent
biomineralization in phylogenetically distinct bacteria by chemically and genetically
inhibiting carbonate accumulation and calcium uptake (Fig. 4). These results
support the notion that mineralization in biofilms is a well-defined developmental
process, initiated by a dramatic change in the structure and function at the single
cell level within a subpopulation of cells in bacterial biofilms. Similar to the
formation of bones initiated by the osteoblast cells within higher animals, the
presence of mineral-producing cells within a biofilm leads to the formation of a
macrostructure serving specific functions and increasing the fitness of the
microbial community.

In this work, we demonstrate that, in a subpopulation of cells, calcium uptake and
functionally conserved pathways of carbonate production alter the intracellular
microenvironment, promoting mineral nucleation and leading to formation of a
mineral macro-structure. Numerous studies of *B. subtilis, P. aeruginosa* and
additional biofilm formers, established that complex colony architecture is
dependent on the organic extracellular matrix (Branda et al., 2005; Colvin et al.,
2012; Jones and Wozniak, 2017; Kobayashi and Iwano, 2012; Romero et al., 2010;
Vlamakis et al., 2013), and thus hypothesized that the presence of the organic
ECM components is sufficient for bacterial biofilm morphogenesis. We found that
non-organic ECM components are also necessary for the development of
structured bacterial communities. While our results indicate that mineral formation
initiates within the cells and requires a controlled intracellular environment, the final
mineral macro-structure assembly has to occur in the extracellular
microenvironment, guided by complex interactions with the extracellular organic
matrix (Keren-Paz and Kolodkin-Gal, 2020), as indicated by the scanning electron
microscopy (Figure 4C, (Oppenheimer-Shaanan et al., 2016)). Therefore, both
organic ECM components and biogenic minerals are essential but not sufficient for
the formation of complex structures.

While in our settings bacteria relied on calcite to generate functional scaffolds,
other minerals may serve similar functions. For example, deposition of calcium
phosphate crystals in *P. mirabilis, P. vulgaris* and *Providencia rettgeri* biofilms
blocks catheters in infected patients (Broomfield et al., 2009; Mathur et al., 2006;
Tan et al., 2018), and urease-dependent calcium phosphate mineralization was
shown to increase *P. mirabilis* resistance to ciprofloxacin (Li et al., 2016c) and
increase its survival in dual-species biofilms (Li et al., 2016b). Importantly, the
physiological role of calcium phosphate deposits within these medical biofilms and
whether dedicated cells facilitate this process remains to be determined.

Intriguingly, we could observe the presence of bacterially produced calcite
minerals in the sputum of some CF patients infected with *P. aeruginosa*. This
observation raises the possibility that lung pathogens can generate functional
minerals within the host. While biofilm colonies formed on top of biofilm-inducing
agar are a convenient model system mimicking many of the environmental and
physical conditions of the host, it lacks the host itself. While several murine models
are available for the study of CF (such as *Cftr-/-* knockout mice), they all fail to
replicate the acute lung disease and chronic bacterial infections characteristic of
CF patients (Semaniakou et al., 2018). *Ex vivo* organ cultures are an experimental
system that recapitulates the three-dimensional structure of the tissue and includes
several cell types and a variety of extracellular matrix components. On the other
hand, organ culture is a controlled microenvironment lacking compounding factors
that play a role in any infection, such as efficiency of drug delivery and the immune
system (Kolodkin-Gal et al., 2007).

Using chemical inhibitors and mutants, we demonstrated that biomineralization by
P. aeruginosa was necessary for lung deterioration, and that preventing it could
rescue the infected tissue. The fact the free-living bacteria do not seem to be
affected by inhibiting biomineralization, and remain in the culture, while the tissue
itself is rescued, highlights the importance of host attachment during infection. It is
also possible that the restrictive properties of crystalline minerals on the diffusivity
and viscoelasticity could be deleterious to the host tissue. Moreover, consistent
with the protective role of the mineral for the bacterial biofilm, chemical or genetic

inhibition of mineralization restored the sensitivity the *P. aeruginosa* to antibiotic treatment (Fig. 5 and Fig. 6).

Our work suggests that it is time to reconsider the old definition of bacterial mineralization solely as a passive, unintended byproduct of environmental bacterial activity. Instead, here we provide evidence that it can also be a regulated developmental process originating from a dedicated, previously uncharacterized microbial compartment, conferring clear benefits to biofilm-residing bacteria. Biomineralization is highly conserved, and thus is of enormous clinical significance, as it can yield completely novel classes of broad-spectrum drugs to combat emerging biofilm infections.

Methods

Strains and media

All strains used in this work were either wild type or derivatives from *Bacillus subtilis* NCIB 3610, *Pseudomonas aeruginosa* PA14, PA01 and *Mycobacterium abscessus* ATCC 19977 (recently renamed *Mycobacteroides abscessus*). A list of strains used in this study can be found in Supplementary Table 1. Deletions of *arsR*, *mntR*, *zur*, *cueR*, *copA* and *chaA* were generated by transforming *B. subtilis* NCIB 3610 with genomic DNA isolated from *B. subtilis* 168 deletion library {Koo, 2017 #6} (Addgene, Cat#1000000115) , and verified by PCR. Deletion of *yetJ* and *yloB* was done as previously described, and double and triple mutants were generated by transformation with genomic DNA, and verified by PCR. *P. aureginosa* PA01 single *psCA1* (CA1), double *psCA1-psCA2* (CA1-2), and triple *psCA1-psCA2-psCA3* (CA1-2-3) mutants were reported earlier (Lotlikar et al., 2019). Deletion of *calC* was generated as described in (Hoang et al., 2000).

Biofilm assays

B. subtilis and *M. abscessus* biofilms were grown on B4 biofilm-promoting solid medium (0.4% yeast extract, 0.5% glucose, and 1.5% agar) (Barabesi et al., 2007)

supplemented with calcium acetate at 0.25% v/v, incubated at 30°C and 37°C respectively, in a sealed box for enriched CO_2 environment achieved by using the candle jar method (Oppenheimer-Shaanan et al., 2016). *P. aeruginosa* biofilms were grown on TB medium, as described previously (Dietrich et al., 2013) or on Brain Heart Infusion Broth (BHI, Sigma-Aldrich) supplemented with 1% agar. SCFM medium was prepared as in (Palmer et al., 2005) and supplemented with 1.5% agar. Crystal-violet biofilm assay was carried out as previously described (O'Toole, 2011).

When indicated, the medium was supplemented with the following inhibitors purchased from Sigma-Aldrich: acetohydroxamic acid (AHA) (Cat. #159034); 5,5'-Dithiobis(2-nitrobenzoic acid) (DTNB) (Cat. #D8130); acetazolamide (Diamox) (Cat. #A6011) and sodium metavanadate (SMV) (Cat. #590088). The concentrations used are indicated in the text.

Phase microscopy

Biofilm colonies were observed using a Nikon D3 camera or a Stereo Discovery V20" microscope (Tochigi, Japan) with objectives Plan Apo S × 0.5 FWD 134 mm or Apo S × 1.0 FWD 60 mm (Zeiss, Goettingen, Germany) attached to Axiocam camera, as required. Data were captured using Axiovision suite software (Zeiss).

Planktonic growth assays

All strains were grown from a single colony isolated over lysogeny broth (LB) plates to a mid-logarithmic phase of growth (4 h at 37°C with shaking). Cells were diluted 1:100 in 150 µl liquid B4 medium in 96-well microplate (Thermo Scientific). Cells were grown at 30°C for 20 h in a microplate reader (Synergy 2, BioTek), and the optical density at 600 nm (OD_{600}) was measured every 15 min. Three independent experiments were conducted, with three technical repeats per plate.

CFU assay was carried out as previously described in detail (Bucher et al., 2016).

RNA extraction and library preparation

Biofilm colonies were grown on biofilm-promoting B4 solid medium with and without calcium for 1, 2, 3 and 6 days. Three independent experiments were conducted, with three colonies from each treatment combined for RNA extraction in each experiment. The samples were frozen in liquid nitrogen and stored until extraction. Frozen bacterial pellets were lysed using the Fastprep homogenizer (MP Biomedicals) and RNA was extracted with the FastRNA PROT blue kit (MP Biomedicals, 116025050) according to the manufacturer's instructions. RNA levels and integrity were determined by Qubit RNA BR Assay Kit (Life Technologies, Q10210) and TapeStation, respectively. All RNA samples were treated with TURBO DNase (Life Technologies, AM2238).

A total of 5 ug RNA from each sample was subjected to rRNA depletion using the Illumina Ribo-Zero rRNA Removal Kit (Bacteria, MRZB12424), according to the manufacturers' protocols. RNA quantity and quality post-depletion was assessed as above. RNA-seq libraries were contracted with NEBNext® Ultra™ Directional RNA Library Prep Kit (NEB, E7420) according to the manufacturer's instructions. Libraries concentrations and sizes were evaluated as above, and were sequenced as multiplex indexes in one lane using the Illumina HighSeq2500 platform.

RNAseq processing

Reads were trimmed from their adapter with cutadapt and aligned to the *B. subtilis* genome (subsp. *subtilis* str. NCIB 3610, NZ_CM000488.1) with Bowtie2 version 2.3.4.1 (Langmead and Salzberg, 2012). The number of uniquely mapped reads per gene was calculated with HT-seq (Anders et al., 2015). Normalization and testing for differential expression was performed with DESeq2 v1.16. A gene was considered to be differentially expressed using the following criteria: normalized mean read count $\geq$ 30, fold change $\geq$ 3, and adjusted p value < 0.05. First, we tested for differential expression between samples grown with and without calcium separately for each time point; however, since the results for days 1, 2 and 3 were very similar, we joined days 1-3. The crude read count, normalized read count,

691 and the result of the differentially expression tests are available in Supplementary

692 File 1.

Comparison between growth conditions

We compared our RNAseq expression data to publically available transcriptomes

representing 269 different growth conditions (Nicolas et al., 2012). Because that

study used microarray platform and not RNAseq, the comparison was performed

using the top 10% genes with the highest expression level of every condition and

every replicate (383 genes per sample). We than used Jaccard index to measure

the overlap between the conditions of the two platforms (i.e. the current study and

(Nicolas et al., 2012)). Prior to the analysis, we removed 152 genes that appear

among the top 10% in more than 80% of the conditions.

Scanning electron microscopy and EDX

Biofilm colonies were grown for 1, 3, 6, 10 and 15 days at 30°C on biofilm-

promoting B4 solid medium, with or without calcium. The colonies were fixed

overnight at 4°C with 2% glutaraldehyde, 3% paraformaldehyde, 0.1 M sodium

cacodylate (pH 7.4) and 5 mM $CaCl_2$, dehydrated and dried as in (Bucher et al.,

2016; Bucher et al., 2015b). Clinical samples were first partially or completely

bleached to remove the organic material by 1 hour of incubation in either 3% or

6% sodium hypochlorite, respectively. The insoluble material was collected,

washed three times in PBS, three times in acetone and air-dried for 16 hours.

Mounted samples were coated with 15 nm thick carbon layer in carbon coater

(EDWARDS). The imaging by secondary electron (SE) or back scattered electron

(BSE) detectors and the Energy Dispersive X-ray Spectroscopy (EDS, Bruker)

were performed using Carl Zeiss Ultra 55 or Supra scanning electron microscopes.

Cryo-STEM analysis

Bacterial colonies grown as described were suspended in PBS buffer. Quantifoil
TEM grids were glow-discharged with an Evactron Combi-Clean glow-discharge
device, and 5 microliters of suspended cells were deposited onto the glow-
discharged grids. Ten nm-diameter gold fiducials (Duchesne et al., 2008) were
applied before blotting and verification using a Leica EM-GP automated plunging
device (Leica). No chemical fixation was used to ovoid artifacts

Vitrified samples were observed with a Tecnai F20 S/TEM instrument (Thermo
Fisher Scientific) at 200 kV, with Gatan 805 brightfield and Fischione HAADF
detectors. Microscope conditions: extraction voltage = 4300 V, gun lens = 3 or 6,
and spot size = 5 or 6 with 10 micrometer condenser apertures, yielding probe
diameters of 1-2 nm and semi-convergence angles of ~ 1.3- 2.7 mrad. Images of
2048 X 2048 pixels were recorded with probe dwell times of 8-18 microseconds.
Spatial sampling was set between 1 and 4 nm/pixel. Electron doses were 1–3
electrons/A^2 per image. Single-axis tilt series were recorded using SerialEM
(Kremer et al., 1996). EDX was performed in STEM mode on vitrified cell samples
with the same electron microscope set-up as used for STEM imaging, using a
liquid N$_2$ cooled Si(Li) detector (EDAX). Vitrified grids contacting 100-200 cells per
grid were automatically imaged on a Talos Arctica (Thermo Fisher Scientific)
microscope in STEM mode, and mapped at intermediate magnifications (10-13k)
using SerialEM software to obtain large fields of view for quantification of data
presented in Fig. 2C. The quantification was done by manually counting the cells
with or without dense deposits.

Tomography reconstructions and visualization

The CSTET tomographic tilt series were aligned using fiducial markers and
reconstructed using weighted back projection (Frangakis and Hegerl, 2001) as
implemented in the IMOD software suite (Kremer et al., 1996). Reconstructions
are displayed after non-linear anisotropic diffusion filtering within IMOD.
Segmentation and volume rendering were performed using Amira 6.3 (FEI
Visualization Sciences Group).

STED image acquisition

Cells were isolated from 3-6 day old biofilm colonies and stained with Calcein-AM
(20 μM) and NileRed (100 μg/ml). Immediately after the staining procedure, the
cells were mounted on a coverslip and immobilized by an agarose pad (1%
agarose, S7$_{50}$ minimal media). STED microscopy was performed using a Leica
SP8 confocal microscope with a 100x oil immersion objective (NA: 1.4). A 552 nm
laser line was used for confocal detection of NileRed, a 494 nm excitation laser
and a 592 nm depletion laser were used for G-STED detection of Calcein-AM.
Channels were acquired with 200 Hz (4 x line averaging) and the appropriately set
HyD hybrid detectors. Images processing was performed using Leica LAS X and
the deconvolution of the respective channels was performed using the Huygens
algorithm.

Micro-CT X-ray analysis

763 Images of indicated magnification were taken using a Zeiss micro XCT 400
764 instrument (Pleasanton, CA, USA). Tomography was carried out using a micro-
765 focused source set at 20 kV and 100 μA. 1200 separate 2D images were taken
766 with a pixel size of 0.87 mm over 1800, exposure time of 30 sec. Image analysis
767 was carried out with Avizo software (VSG, Hillsboro, OR, USA).

FTIR spectrophotometer analysis

Calcite was collected as described by Mahamid *et al*., with some modifications
(Mahamid et al., 2008): agar samples were slightly bleached with 6% sodium
hypochlorite for 1 min to remove organic matter, washed with Milli-Q water twice
and dehydrated in acetone.

FTIR spectra of the bleached samples were acquired in KBr pellets by using a
NICOLET iS5 spectrometer (Thermo Scientific, Pittsburgh, PA, USA). The
samples were homogenized in an agate mortar and pestle with about 40 mg of

KBr, and pressed into a 7 mm pellet using a manual hydraulic press (Specac, Orpington, UK). Infrared spectra were obtained at 4 cm−1 resolution for 32 scans.

The infrared calcite spectrum has three characteristic peaks, designated v_2, v_3, and v_4. Calcium carbonate v_3 peak is expected at 1425 cm^{-1} for calcite, 1490 cm^{-1} for vaterite and 1475 cm-1 for aragonite. The calcium carbonate v_2 peak is expected at 875 cm^{-1} for calcite, 850 cm^{-1} for vaterite and 855 cm^{-1} for aragonite. Finally, the calcium carbonate v_4 peak is expected at 713 cm^{-1} for calcite, 750 cm^{-1} for vaterite and 715 cm^{-1} for aragonite (Politi et al., 2004).

Patient's samples

Sputum was collected from adult patients as published previously (Schiller and Millard, 1983) and stored in 4ºC degrees. All patients positive for calcite were carrying chronic *pseudomonas* infections. Specific patient information can be found in Supplementary Table 2. The samples were collected under Helskinki approval to Prof. Eitan Keren 0456-17-HMO. For FTIR analysis, organic matter was removed as described above.

Ex vivo **lung infection system**

Animal work was carried out under ethical approval MD-16-15035-1. Lungs were harvested from 2 mice (one month old) and placed in petri dishes containing DMEM 5% FCS. We divided the tissue into circular pieces 3 mm in diameter with a biopsy punch and transferred to a 24 well plate (4-5 explants/well) with 450 µl DMEM. DMEM was applied with carbenicillin 100 µg/ml (Sigma-Aldrich) and chemical inhibitors, as indicated. To each respective well, either 50 µl DMEM (control) or *P. aeruginosa* (pretreated with chemical inhibitors when indicated and diluted within DMEM to attain OD_{600} 0.4) was added – with three technical repeats for each condition. The plates were incubated at 37ºC for ~2 days (52 hours), washed twice with PBS, fixed with PFA 4% for 10 minutes, and imbedded in either cryosection (OCT) compound or paraffin. Paraffin samples were cut into 7 micron

slices and stained with H&E; whereas cryosections were cut into 10 micron slices and placed on superfrost plus slides.

Cultivable Cell Quantification (CFU)

CFU quantification was done as previously described (Bucher et al., 2015a). For CFU of bacteria infecting *ex vivo* tissue culture: individual punches or their growth media were collected, Punches were resuspended in 1 ml PBS (Biological Industries), and free-living bacteria were pelleted and resuspended in PBS at the same volume. Samples were thoroughly vortexed (5 min). The samples were then mildly sonicated (BRANSON digital sonifier, Model 250, Microtip, amplitude 30%, pulse 2 × 5 s). In all cases, to determine the number of viable cells, samples were serially diluted in PBS, plated on LB plates, and colonies were counted after incubation at 30°C overnight. For

Acknowledgments

The Kolodkin-Gal lab is supported by the Israel Science Foundation grant number 119/16, Israel Foundation grant number JSPS 184.20 , Kamin grant by Israel Chief Scientist no. 67459, Israel Ministry of Science, Technology & Space (grant no. 713454), Ministry of Health (grant no. 713645) , Angel-Fiavovich fund for ecological research, Dr. Barry Sherman Institute for Medicinal Chemistry, Kekst Family Institute for Medical Genetics and by the Helen and Milton A. Kimmelman Center for Biomolecular Structure & Assembly. IKG is a recipient of Rowland and Sylvia Career Development Chair. The electron microscopy studies were partially supported by the Irving and Cherna Moskowitz Center for Nano and BioNano Imaging (Weizmann Institute of Science). We thank Prof. Ehud Banin and to Itzhak Zander (Bar-Ilan University) for helpful discussions, and for providing PA01 strains.

References

Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq-a Python framework to work with high-throughput sequencing data. Bioinformatics *31*, 166-169.

Asally, M., Kittisopikul, M., Rue, P., Du, Y., Hu, Z., Cagatay, T., Robinson, A.B., Lu, H., Garcia-Ojalvo, J., and Suel, G.M. (2012). Localized cell death focuses mechanical forces during 3D patterning in a biofilm. Proc Natl Acad Sci U S A *109*, 18891-18896.

Barabesi, C., Galizzi, A., Mastromei, G., Rossi, M., Tamburini, E., and Perito, B. (2007). Bacillus subtilis gene cluster involved in calcium carbonate biomineralization. J Bacteriol *189*, 228-235.

Blondeau, M., Sachse, M., Boulogne, C., Gillet, C., Guigner, J.M., Skouri-Panet, F., Poinsot, M., Ferard, C., Miot, J., and Benzerara, K. (2018). Amorphous Calcium Carbonate Granules Form Within an Intracellular Compartment in Calcifying Cyanobacteria. Frontiers in Microbiology *9*.

Bloom-Ackermann, Z., Steinberg, N., Rosenberg, G., Oppenheimer-Shaanan, Y., Pollack, D., Ely, S., Storzi, N., Levy, A., and Kolodkin-Gal, I. (2016). Toxin-Antitoxin systems eliminate defective cells and preserve symmetry in Bacillus subtilis biofilms. Environ Microbiol *18*, 5032-5047.

Branda, S.S., Vik, S., Friedman, L., and Kolter, R. (2005). Biofilms: the matrix revisited. Trends Microbiol *13*, 20-26.

Broomfield, R.J., Morgan, S.D., Khan, A., and Stickler, D.J. (2009). Crystalline bacterial biofilm formation on urinary catheters by urease-producing urinary tract pathogens: a simple method of control. J Med Microbiol *58*, 1367-1375.

Bryers, J.D. (2008). Medical biofilms. Biotechnol Bioeng *100*, 1-18.

Bucher, T., Kartvelishvily, E., and Kolodkin-Gal, I. (2016). Methodologies for Studying B. subtilis Biofilms as a Model for Characterizing Small Molecule Biofilm Inhibitors. J Vis Exp.

Bucher, T., Keren-Paz, A., Hausser, J., Olender, T., Cytryn, E., and Kolodkin-Gal, I. (2019). An active beta-lactamase is a part of an orchestrated cell wall stress resistance network of Bacillus subtilis and related rhizosphere species. Environ Microbiol.

Bucher, T., Oppenheimer-Shaanan, Y., Savidor, A., Bloom-Ackermann, Z., and Kolodkin-Gal, I. (2015a). Disturbance of the bacterial cell wall specifically interferes with biofilm formation. Environ Microbiol Rep *7*, 990-1004.

Bucher, T., Oppenheimer-Shaanan, Y., Savidor, A., Bloom-Ackermann, Z., and Kolodkin-Gal, I. (2015b). Disturbance of the bacterial cell wall specifically interferes with biofilm formation. Environmental microbiology reports.

Caimmi, D., Martocq, N., Trioleyre, D., Guinet, C., Godreuil, S., Daniel, T., and Chiron, R. (2018). Positive Effect of Liposomal Amikacin for Inhalation on Mycobacterium abcessus in Cystic Fibrosis Patients. Open Forum Infect Dis *5*, ofy034.

Chang, Y., Bruni, R., Kloss, B., Assur, Z., Kloppmann, E., Rost, B., Hendrickson, W.A., and Liu, Q. (2014). Structural basis for a pH-sensitive calcium leak across membranes. Science *344*, 1131-1135.

Clausen, J.D., Bublitz, M., Arnou, B., Olesen, C., Andersen, J.P., Moller, J.V., and Nissen, P. (2016). Crystal Structure of the Vanadate-Inhibited Ca2+-ATPase. Structure *24*, 617-623.

Cohen-Cymberknoh, M., Gilead, N., Gartner, S., Rovira, S., Blau, H., Mussaffi, H., Rivlin, J., Gur, M., Shteinberg, M., Bentur, L.*, et al.* (2016). Eradication failure of newly acquired

Pseudomonas aeruginosa isolates in cystic fibrosis. Journal of cystic fibrosis : official journal of the European Cystic Fibrosis Society.

Colvin, K.M., Gordon, V.D., Murakami, K., Borlee, B.R., Wozniak, D.J., Wong, G.C., and Parsek, M.R. (2011). The pel polysaccharide can serve a structural and protective role in the biofilm matrix of Pseudomonas aeruginosa. PLoS Pathog *7*, e1001264.

Colvin, K.M., Irie, Y., Tart, C.S., Urbano, R., Whitney, J.C., Ryder, C., Howell, P.L., Wozniak, D.J., and Parsek, M.R. (2012). The Pel and Psl polysaccharides provide Pseudomonas aeruginosa structural redundancy within the biofilm matrix. Environ Microbiol *14*, 1913-1928.

Costerton, J.W., Stewart, P.S., and Greenberg, E.P. (1999). Bacterial biofilms: a common cause of persistent infections. Science *284*, 1318-1322.

de Hoon, M.J., Eichenberger, P., and Vitkup, D. (2010). Hierarchical evolution of the bacterial sporulation network. Curr Biol *20*, R735-745.

Dhami, N.K., Reddy, M.S., and Mukherjee, A. (2013). Biomineralization of calcium carbonate polymorphs by the bacterial strains isolated from calcareous sites. J Microbiol Biotechnol *23*, 707-714.

Dietrich, L.E., Okegbe, C., Price-Whelan, A., Sakhtah, H., Hunter, R.C., and Newman, D.K. (2013). Bacterial community morphogenesis is intimately linked to the intracellular redox state. J Bacteriol *195*, 1371-1380.

Dietrich, L.E., Teal, T.K., Price-Whelan, A., and Newman, D.K. (2008). Redox-active antibiotics control gene expression and community behavior in divergent bacteria. Science *321*, 1203-1206.

Douglas, S., and Beveridge, T.J. (1998). Mineral formation by bacteria in natural microbial communities. Fems Microbiology Ecology *26*, 79-88.

Dragos, A., and Kovacs, A.T. (2017). The Peculiar Functions of the Bacterial Extracellular Matrix. Trends Microbiol *25*, 257-266.

Duchesne, L., Gentili, D., Comes-Franchini, M., and Fernig, D.G. (2008). Robust ligand shells for biological applications of gold nanoparticles. Langmuir *24*, 13572-13580.

El Mammeri, N., Hierrezuelo, J., Tolchard, J., Camara-Almiron, J., Caro-Astorga, J., Alvarez-Mena, A., Dutour, A., Berbon, M., Shenoy, J., Morvan, E.*, et al.* (2019). Molecular architecture of bacterial amyloids in Bacillus biofilms. FASEB J, fj201900831R.

Flemming, H.C., and Wuertz, S. (2019). Bacteria and archaea on Earth and their abundance in biofilms. Nature Reviews Microbiology *17*, 247-260.

Frangakis, A.S., and Hegerl, R. (2001). Noise reduction in electron tomographic reconstructions using nonlinear anisotropic diffusion. Journal of structural biology *135*, 239-250.

Frankel, R.B., and Bazylinski, D.A. (2003). Biologically induced mineralization by bacteria. Rev Mineral Geochem *54*, 95-114.

Fujisawa, M., Wada, Y., Tsuchiya, T., and Ito, M. (2009). Characterization of Bacillus subtilis YfkE (ChaA): a calcium-specific Ca^{2+}/H^+ antiporter of the CaCA family. Arch Microbiol *191*, 649-657.

Gupta, H.K., Shrivastava, S., and Sharma, R. (2017). A Novel Calcium Uptake Transporter of Uncharacterized P-Type ATPase Family Supplies Calcium for Cell Surface Integrity in Mycobacterium smegmatis. MBio *8*.

Guragain, M., Lenaburg, D.L., Moore, F.S., Reutlinger, I., and Patrauchan, M.A. (2013).
Calcium homeostasis in Pseudomonas aeruginosa requires multiple transporters and
modulates swarming motility. Cell Calcium *54*, 350-361.

Hale, L.V., Ma, Y.F., and Santerre, R.F. (2000). Semi-quantitative fluorescence analysis
of calcein binding as a measurement of in vitro mineralization. Calcified Tissue Int *67*, 80-
84.

Halloum, I., Carrere-Kremer, S., Blaise, M., Viljoen, A., Bernut, A., Le Moigne, V.,
Vilcheze, C., Guerardel, Y., Lutfalla, G., Herrmann, J.L.*, et al.* (2016). Deletion of a
dehydratase important for intracellular growth and cording renders rough Mycobacterium
abscessus avirulent. Proc Natl Acad Sci U S A *113*, E4228-4237.

Hassanov, T., Karunker, I., Steinberg, N., Erez, A., and Kolodkin-Gal, I. (2018). Novel
antibiofilm chemotherapies target nitrogen from glutamate and glutamine. Scientific
reports *8*, 7097.

Hewer, S.C.L., and Smyth, A.R. (2017). Antibiotic strategies for eradicating Pseudomonas
aeruginosa in people with cystic fibrosis. Cochrane Db Syst Rev.

Hill, D., Rose, B., Pajkos, A., Robinson, M., Bye, P., Bell, S., Elkins, M., Thompson, B.,
Macleod, C., Aaron, S.D.*, et al.* (2005). Antibiotic susceptabilities of Pseudomonas
aeruginosa isolates derived from patients with cystic fibrosis under aerobic, anaerobic, and
biofilm conditions. Journal of clinical microbiology *43*, 5085-5090.

Hoang, T.T., Kutchma, A.J., Becher, A., and Schweizer, H.P. (2000). Integration-
proficient plasmids for Pseudomonas aeruginosa: Site-specific integration and use for
engineering of reporter and expression strains. Plasmid *43*, 59-72.

Hufnagel, D.A., Price, J.E., Stephenson, R.E., Kelley, J., Benoit, M.F., and Chapman, M.R.
(2018). Thiol Starvation Induces Redox-Mediated Dysregulation of Escherichia coli
Biofilm Components. J Bacteriol *200*.

Jansson, C., and Northen, T. (2010). Calcifying cyanobacteria--the potential of
biomineralization for carbon capture and storage. Curr Opin Biotechnol *21*, 365-371.

Jo, J., Cortez, K.L., Cornell, W.C., Price-Whelan, A., and Dietrich, L.E. (2017). An orphan
cbb3-type cytochrome oxidase subunit supports Pseudomonas aeruginosa biofilm growth
and virulence. Elife *6*.

Jones, C.J., and Wozniak, D.J. (2017). Psl Produced by Mucoid Pseudomonas aeruginosa
Contributes to the Establishment of Biofilms and Immune Evasion. mBio *8*.

Keren-Paz, A., Brumfeld, V., Oppenheimer-Shaanan, Y., and Kolodkin-Gal, I. (2018).
Micro-CT X-ray imaging exposes structured diffusion barriers within biofilms. NPJ
Biofilms Microbiomes *4*, 8.

Keren-Paz, A., and Kolodkin-Gal, I. (2020). A brick in the wall: Discovering a novel
mineral component of the biofilm extracellular matrix. N Biotechnol *56*, 9-15.

Kobayashi, K., and Iwano, M. (2012). BslA(YuaB) forms a hydrophobic layer on the
surface of Bacillus subtilis biofilms. Molecular microbiology *85*, 51-66.

Kolodkin-Gal, D., Zamir, G., Edden, Y., Pikarsky, E., Pikarsky, A., Haim, H., Haviv, Y.S.,
and Panet, A. (2008). Herpes simplex virus type 1 preferentially targets human colon
carcinoma: role of extracellular matrix. Journal of virology *82*, 999-1010.

Kolodkin-Gal, D., Zamir, G., Pikarski, E., Pikarski, A., Shimony, N., Wu, H., Haviv, Y.S.,
and Panet, A. (2007). A novel system to study adenovirus tropism to normal and malignant
colon tissues. Virology *357*, 91-101.

Kolodkin-Gal, I., Elsholz, A.K., Muth, C., Girguis, P.R., Kolter, R., and Losick, R. (2013). Respiration control of multicellularity in Bacillus subtilis by a complex of the cytochrome chain with a membrane-embedded histidine kinase. Genes Dev *27*, 887-899.

Kolter, R., and Greenberg, E.P. (2006). Microbial sciences: the superficial life of microbes. Nature *441*, 300-302.

Komeili, A., Li, Z., Newman, D.K., and Jensen, G.J. (2006). Magnetosomes are cell membrane invaginations organized by the actin-like protein MamK. Science *311*, 242-245.

Kremer, J.R., Mastronarde, D.N., and McIntosh, J.R. (1996). Computer visualization of three-dimensional image data using IMOD. Journal of structural biology *116*, 71-76.

Kuhlbrandt, W. (2004). Biology, structure and mechanism of P-type ATPases. Nat Rev Mol Cell Biol *5*, 282-295.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat Methods *9*, 357-359.

Li, X., Chopp, D.L., Russin, W.A., Brannon, P.T., Parsek, M.R., and Packman, A.I. (2015). Spatial patterns of carbonate biomineralization in biofilms. Appl Environ Microbiol *81*, 7403-7410.

Li, X., Chopp, D.L., Russin, W.A., Brannon, P.T., Parsek, M.R., and Packman, A.I. (2016a). In Situ Biomineralization and Particle Deposition Distinctively Mediate Biofilm Susceptibility to Chlorine. Appl Environ Microbiol *82*, 2886-2892.

Li, X., Lu, N., Brady, H.R., and Packman, A.I. (2016b). Biomineralization strongly modulates the formation of Proteus mirabilis and Pseudomonas aeruginosa dual-species biofilms. FEMS Microbiol Ecol.

Li, X., Lu, N., Brady, H.R., and Packman, A.I. (2016c). Ureolytic Biomineralization Reduces Proteus mirabilis Biofilm Susceptibility to Ciprofloxacin. Antimicrob Agents Chemother *60*, 2993-3000.

Liu, J., Prindle, A., Humphries, J., Gabalda-Sagarra, M., Asally, M., Lee, D.Y., Ly, S., Garcia-Ojalvo, J., and Suel, G.M. (2015). Metabolic co-dependence gives rise to collective oscillations within biofilms. Nature *523*, 550-554.

Lotlikar, S.R., Kayastha, B.B., Vullo, D., Khanam, S.S., Braga, R.E., Murray, A.B., McKenna, R., Supuran, C.T., and Patrauchan, M.A. (2019). Pseudomonas aeruginosa beta-carbonic anhydrase, psCA1, is required for calcium deposition and contributes to virulence. Cell Calcium *84*, 102080.

Mahamid, J., Sharir, A., Addadi, L., and Weiner, S. (2008). Amorphous calcium phosphate is a major component of the forming fin bones of zebrafish: Indications for an amorphous precursor phase. Proceedings of the National Academy of Sciences of the United States of America *105*, 12748-12753.

Massler, A., Kolodkin-Gal, D., Meir, K., Khalaileh, A., Falk, H., Izhar, U., Shufaro, Y., and Panet, A. (2011). Infant lungs are preferentially infected by adenovirus and herpes simplex virus type 1 vectors: role of the tissue mesenchymal cells. J Gene Med *13*, 101-113.

Mathur, S., Suller, M.T., Stickler, D.J., and Feneley, R.C. (2006). Factors affecting crystal precipitation from urine in individuals with long-term urinary catheters colonized with urease-positive bacterial species. Urol Res *34*, 173-177.

Monds, R.D., and O'Toole, G.A. (2009). The developmental model of microbial biofilms: ten years of a paradigm up for review. Trends in microbiology *17*, 73-87.

Monteil, C.L., Benzerara, K., Menguy, N., Bidaud, C.C., Michot-Achdjian, E., Bolzoni, R., Mathon, F.P., Coutaud, M., Alonso, B., Garau, C.*, et al.* (2020). Intracellular amorphous Ca-carbonate and magnetite biomineralization by a magnetotactic bacterium affiliated to the Alphaproteobacteria. ISME J.

Moore, C.M., and Helmann, J.D. (2005). Metal ion homeostasis in Bacillus subtilis. Curr Opin Microbiol *8*, 188-195.

Morris, N.S., and Stickler, D.J. (1998). Encrustation of indwelling urethral catheters by Proteus mirabilis biofilms growing in human urine. J Hosp Infect *39*, 227-234.

Nicolas, P., Mader, U., Dervyn, E., Rochat, T., Leduc, A., Pigeonneau, N., Bidnenko, E., Marchadier, E., Hoebeke, M., Aymerich, S.*, et al.* (2012). Condition-dependent transcriptome reveals high-level regulatory architecture in Bacillus subtilis. Science *335*, 1103-1106.

O'Toole, G.A. (2011). Microtiter dish biofilm formation assay. J Vis Exp.

Oppenheimer-Shaanan, Y., Sibony-Nevo, O., Bloom-Ackermann, Z., Suissa, R., Steinberg, N., Kartvelishvily, E., Brumfeld, V., and Kolodkin-Gal, I. (2016). Spatio-temporal assembly of functional mineral scaffolds within microbial biofilms. NPJ Biofilms Microbiomes *2*, 15031.

Palmer, K.L., Mashburn, L.M., Singh, P.K., and Whiteley, M. (2005). Cystic fibrosis sputum supports growth and cues key aspects of Pseudomonas aeruginosa physiology. Journal of Bacteriology *187*, 5267-5277.

Phoenix, V.R., and Konhauser, K.O. (2008). Benefits of bacterial biomineralization. Geobiology *6*, 303-308.

Politi, Y., Arad, T., Klein, E., Weiner, S., and Addadi, L. (2004). Sea urchin spine calcite forms via a transient amorphous calcium carbonate phase. Science *306*, 1161-1164.

Reichhardt, C., and Parsek, M.R. (2019). Confocal Laser Scanning Microscopy for Analysis of Pseudomonas aeruginosa Biofilm Architecture and Matrix Localization. Front Microbiol *10*, 677.

Richards, J.P., Cai, W.L., Zill, N.A., Zhang, W.J., and Ojha, A.K. (2019). Adaptation of Mycobacterium tuberculosis to Biofilm Growth Is Genetically Linked to Drug Tolerance. Antimicrob Agents Ch *63*.

Romero, D., Aguilar, C., Losick, R., and Kolter, R. (2010). Amyloid fibers provide structural integrity to Bacillus subtilis biofilms. Proc Natl Acad Sci U S A *107*, 2230-2234.

Schiller, N.L., and Millard, R.L. (1983). Pseudomonas-infected cystic fibrosis patient sputum inhibits the bactericidal activity of normal human serum. Pediatr Res *17*, 747-752.

Semaniakou, A., Croll, R.P., and Chappe, V. (2018). Animal Models in the Pathophysiology of Cystic Fibrosis. Front Pharmacol *9*, 1475.

Serra, D.O., and Hengge, R. (2014). Stress responses go three dimensional - the spatial order of physiological differentiation in bacterial macrocolony biofilms. Environ Microbiol *16*, 1455-1471.

Serra, D.O., Richter, A.M., Klauck, G., Mika, F., and Hengge, R. (2013). Microanatomy at cellular resolution and spatial order of physiological differentiation in a bacterial biofilm. MBio *4*, e00103-00113.

Steinberg, N., Keren-Paz, A., Hou, Q., Doron, S., Yanuka-Golub, K., Olender, T., Hadar, R., Rosenberg, G., Jain, R., Camara-Almiron, J.*, et al.* (2020). The extracellular matrix protein TasA is a developmental cue that maintains a motile subpopulation within Bacillus subtilis biofilms. Sci Signal *13*.

Steinberg, N., and Kolodkin-Gal, I. (2015). The Matrix Reloaded: Probing the Extracellular
Matrix Synchronizes Bacterial Communities. J Bacteriol.
Stewart, P.S., and Franklin, M.J. (2008). Physiological heterogeneity in biofilms. Nat Rev
Microbiol *6*, 199-210.
Tan, Y., Leonhard, M., Moser, D., Ma, S., and Schneider-Stickler, B. (2018). Inhibitory
effect of probiotic lactobacilli supernatants on single and mixed non-albicans Candida
species biofilm. Arch Oral Biol *85*, 40-45.
Veening, J.W., Kuipers, O.P., Brul, S., Hellingwerf, K.J., and Kort, R. (2006). Effects of
phosphorelay perturbations on architecture, sporulation, and spore resistance in biofilms of
Bacillus subtilis. J Bacteriol *188*, 3099-3109.
Vlamakis, H., Aguilar, C., Losick, R., and Kolter, R. (2008). Control of cell fate by the
formation of an architecturally complex bacterial community. Genes Dev *22*, 945-953.
Vlamakis, H., Chai, Y., Beauregard, P., Losick, R., and Kolter, R. (2013). Sticking
together: building a biofilm the Bacillus subtilis way. Nature reviews Microbiology *11*,
157-168.
Weiner, S., and Dove, P.M. (2003). An overview of biomineralization processes and the
problem of the vital effect. Rev Mineral Geochem *54*, 1-29.
Wermser, C., and Lopez, D. (2018). Identification of Staphylococcus aureus genes
involved in the formation of structured macrocolonies. Microbiology (Reading) *164*, 801-
815.
Wilking, J.N., Zaburdaev, V., De Volder, M., Losick, R., Brenner, M.P., and Weitz, D.A.
(2013). Liquid transport facilitated by channels in Bacillus subtilis biofilms. Proc Natl
Acad Sci U S A *110*, 848-852.
Wolf, S.G., Houben, L., and Elbaum, M. (2014). Cryo-scanning transmission electron
tomography of vitrified cells. Nat Methods *11*, 423-428.
Wu, B.C., Haney, E.F., Akhoundsadegh, N., Pletzer, D., Trimble, M.J., Adriaans, A.E.,
Nibbering, P.H., and Hancock, R.E.W. (2021). Human organoid biofilm model for
assessing antibiofilm activity of novel agents. NPJ Biofilms Microbiomes *7*, 8.
Yoon, S.S., and Hassett, D.J. (2004). Chronic Pseudomonas aeruginosa infection in cystic
fibrosis airway disease: metabolic changes that unravel novel drug targets. Expert Rev Anti
Infect Ther *2*, 611-623.

1093

1094

1095

1096

1097

1098

1099

1100

Figure Legends:

Figure 1: Calcium is necessary for normal biofilm development and structure

A. Light microscopy images of 6-day-old *B. subtilis* NCIB 3610 biofilm colonies grown on B4 agar, showing the effect of addition of calcium (0.25% v/v calcium acetate) on colony architecture. Scale bar – 2 mm. A representative image (out of n = 3 experiments) is shown.

B. Volcano-plot depicting calcium-dependent changes in the transcriptome in *B. subtilis* biofilm colonies. Significance (P values) vs difference (fold changes) are plotted. Functional categories were determined by DAVID analysis.

C. A heatmap with all differentially expressed genes (n = 876), scaled to the mean expression level of each gene. The map is ordered by age of biofilm colonies, as shown by the left colored bar, where green are genes that are differentially expressed in days 1-3, and day 6; purple genes that are differentially expressed in days 1-3 and yellow genes that are differentially expressed in day 6.

D. The top 20 conditions (Nicolas et al., 2012), showing the highest similarity (expressed as Jaccard index) to the transcriptome of *B. subtilis* 6-day-old biofilm colonies grown with or without calcium.

E. Biofilm cells closely associated with prismatic mineral structures, as visualized by Scanning Electron Microscopy (SEM) image of 10-day-old *B. subtilis* biofilm colony grown with calcium. Upper panel - secondary mode; lower panel - backscattering mode. Magnification – X25000, scale bar – 1 μm. A representative field (out of n=5 fields, from 6 experiments) is shown.

F. EDX analysis of areas indicated in (E), indicating that mineral structures are rich in calcium. Ca – calcium-rich, Cnt – control.

G. 3D re-construction of microCT X-Ray analysis revealing 3D mineral distribution in a 5-day old *B. subtilis* colony grown with calcium. Color indicates intensity, with green representing the densest mineral. Scale bar – 0.2 mm. A representative image (out of n = 3 experiments) is shown.

H. Light microscopy images of 5-day-old *B. subtilis* biofilm colonies grown with calcium, and supplemented with calcium uptake inhibitor (0.1 mg/ml sodium metavanadate (SMV)), as indicated. NT indicated untreated culture. Scale bar – 2 mm. A representative image (out of n = 3 experiments) is shown.

I. The effect of inhibiting calcium uptake on planktonic growth. *B. subtilis* was grown shaking in liquid LB medium with calcium, supplemented with 0.1 mg/ml SMV, as indicated. Results are averages of nine wells, bars represent standard deviations. A representative experiment (out of n=3 experiments) is shown.

Figure 2: Mineralization is initiated intracellularly

A. A 30-nm thick virtual slice through a CSTET 3D reconstruction of *B. subtilis* cells showing intracellular calcium-rich deposits (arrows). Scale bar – 400 nm. A representative field (out of n=20 fields, from 3 experiments) is shown.

B. Volume rendering (orange) and a single orthoslice (greyscale) through the center of the volume, from a 3D CSTET reconstruction. The calcium-rich deposits are artificially colored (blue).

C. Quantification of different subpopulations within a 6-day old *B. subtilis* biofilm colony, showing enrichment of cells with mineral foci in the structured areas (wrinkles) of the colony.

D. Intracellular calcium levels of bacterial cells (n=150) stained with calcein-AM, a calcium-specific fluorescent dye. Cells were isolated from *B. subtilis* biofilm colonies (day 3 and 6) grown with calcium. Inset – a representative image of stained cells isolated from a 6 day-old biofilm. Green – calcein-AM, red – NileRed membrane stain, gray – DIC. Scale bar – 2 µm.

E. and F. G-STED images *B. subtilis* cells isolated from 6 day-old biofilm colonies grown with calcium. Green – calcein-AM, red – NileRed membrane stain, gray – DIC. Scale bar – 2 µm.

G. 30-nm thick virtual slice through a CSTET 3D reconstruction. Arrows indicate calcium-rich deposits associated with a membrane from the outside of the cell. Scale bar – 400 nm.

H. Light microscopy images of 3-day-old biofilm colonies of wild-type *B. subtilis* and its mutant derivatives (see text for details) grown with calcium. Scale bar – 2 mm. A representative experiment (out of n=3 experiments) is shown.

I. Volcano-plot depicting CueR-dependent changes in expression of genes in 3-day old *B. subtilis* biofilm colony. Significance (P values) vs difference (fold changes) are plotted. Functional categories were determined by DAVID analysis.

J. Light microscopy images of 2-day-old biofilm colonies of wild-type *B. subtilis* and its mutant derivatives (see text for details) grown with calcium. Scale bar – 2 mm. A representative experiment (out of n=3 experiments) is shown.

Figure 3: The role of calcium in the biofilm structure is conserved across bacterial species

A. Light microscopy (left) and microCT-X-ray (right) images of 3-day-old *P. aeruginosa* PA14 biofilm colonies grown on TB agar. In the X-Ray images, the bright contrast appears white, and indicates the location of dense mineral, while the organic matter appears dark. Scale bar – 1 mm. A representative image (out of n = 3 experiments) is shown.

B. Light microscopy images of 3-day-old *P. aeruginosa* PA14 of biofilm colonies grown on TB agar supplemented with 1 mg/ml EDTA (chelating agent), either with or without the addition of 700 mM $CaCl_2$. Scale bar – 1 mm. A representative experiment (out of n=3 experiments) is shown.

C. Light microscopy images of 3-day-old *P. aeruginosa* PA14 biofilm colonies, grown on TB agar either untreated (NT), or supplemented with 700 mM $CaCl_2$ and EGTA (calcium chelator), as indicated. Scale bar – 5 mm. A representative image (out of n = 3 experiments) is shown.

D. 3D re-construction of microCT X-Ray analysis revealing mineral distribution in a 5-day old *P. aeruginosa* PA14 colony. Color indicates intensity, with green representing the densest mineral. Scale bar – 0.2 mm. A representative image (out of n = 3 experiments) is shown.

E. Light microscopy images of 5 day-old *M. abscessus* biofilm colonies, grown on B4 agar, supplemented with either calcium acetate (0.25% v/v), or 1.5 mM EDTA, as indicated. Scale bar – 5 mm. A representative image (out of n = 3 experiments) is shown.

F. 3D re-construction of microCT X-Ray analysis revealing mineral distribution in a 5-day old *M. abscessus* colony grown on B4 agar with calcium. Color indicates intensity, with green representing the densest mineral. Scale bar – 0.1 mm. A representative image (out of n = 3 experiments) is shown.

Figure 4: Inhibition of cellular pathways leading to calcium carbonate production disrupts biofilm development in *P. aeruginosa* and *M. abscessus*.

A. A schematic representation of the chemical reaction leading to calcium carbonate production, with the inhibitors used in this study indicated in parenthesis next to the step they inhibit.

B. Light microscopy images of 3-day-old *P. aeruginosa* PA14 biofilm colonies grown on TB agar containing the inhibitors indicated in (A) as follows: untreated (NT), 1.5 mg/ml DTNB, 1.75 mg/ml AHA, 2.5 mg/ml Diamox and 0.01 mg/ml SMV. Scale bar – 2 mm. A representative image (out of n = 3 experiments) is shown.

C. The effect of inhibiting calcium carbonate production on planktonic growth of *P. aeruginosa* PA14. Bacteria were grown shaking in liquid TB medium, supplemented with: 1.5 mg/ml DTNB, 1.75 mg/ml AHA, 2.5 mg/ml Diamox and 0.01, 0.05 and 0.01 mg/ml SMV, or untreated (NT). Growth was monitored by measuring OD600 in a microplate reader every 30 min. Results are averages

of six wells, bars represent standard deviations. A representative experiment

(out of n= 3 experiments) is shown.

D. Light microscopy images of 4-day-old biofilm colonies of wild type *P.*

aeruginosa PA01 and indicated mutants grown on BHI agar: carbonic

anhydrases (*psCA1* – PA0102, *psCA2* - PA2053, *psCA3* – PA4676) and

calcium channel (Δ*calC*). A representative experiment (out of n= 3

experiments) is shown. Insert – the indicated mutant strains were grown

planktonically in shaking liquid BHI medium. Growth was monitored by

measuring OD600 in a spectrophotometer at indicated times. Results are

averages of 4 wells, bars represent standard deviations. Scale bar – 2 mm. A

representative experiment (out of n= 3 experiments) is shown.

E. Crystal violet assay quantifying biofilm formation of *P. aeruginosa* PA14, grown

in BHI supplemented with the indicated inhibitors. Results are averages and

standard deviation of three independent experiments. P values, as determined

by student's t-test are indicated (* pVal <0.01, ** pVal <0.001, when compared

to the untreated control).

F. Crystal violet assay quantifying biofilm formation of wild type and indicated

mutant derivates of *P. aeruginosa* PA01, grown with in liquid BHI medium.

Results are averages and standard deviation of three independent

experiments. P values, as determined by student's t-test are indicated (* pVal

<0.01, when compared to the untreated control).

G. Light microscopy images of 5 day-old *M. abscessus* biofilm colonies, grown on

B4 agar supplemented with calcium and 2.5 mM of the indicated inhibitors or

untreated (NT). A representative experiment (out of n = 3) experiments is

shown. Insert – *M. abscessus* was grown planktonically in liquid B4 medium.

Growth was monitored by measuring OD_{600} in a spectrophotometer at indicated

times. Scale bar – 2 mm. Results are averages of three independent repeats,

bars represent standard deviations.

H. Light microscopy images of 3-day-old *P. aeruginosa* PA14 biofilm colonies

grown on SCFM agar, either untreated (NT) or suplemented with 700 mM $CaCl_2$

and EGTA, as indicated. Scale bar – 5 mm. A representative experiment (out
of n = 3) experiments is shown.

1245

Figure 5: Inhibition of urease prevents biofilm formation and tissue
deterioration in an *ex-vivo* lunge model

A. and B. Fluorescent microscopy images of 10 micron slices of lung tissues,
infected with *P. aeruginosa* PA14 and treated with AHA, as indicated. (A) Blue
– lung cell nuclei stained by DAPI, (magnification x20). (B) Green - GFP
expressing *P. aeruginosa*, (magnification x40). A representative experiment
(out of n = 3) experiments is shown.

C. Light microscopy images of 7 micron slices of lung tissue. Samples were
treated as in (A), and stained with hematoxylin and eosin (H&E). Blue – nuclei,
red – extracellular matrix and cytoplasm, (magnification x40). A representative
experiment (out of n = 3) experiments is shown.

D. Quantification of lung viability. Lung tissue cultures were infected with *P.*
aeruginosa PA14 and treated with AHA, as indicated. ImageJ 1.51g software
was used to automatically count lung cell nuclei in 4 randomly chosen fields for
each treatment. P values, as determined by student's t-test are indicated (**
pVal <0.001).

E. CFU analysis of *P. aeruginosa* PA14 infecting lung tissue and surrounding
media, with and without treatment. P values, as determined by student's t-test
are indicated (* pVal <0.01, ** pVal <0.001, when compared to the untreated
control).

F. Quantification representing the ratios of tissue to medium CFU in (E).

1267

1268 **Figure 6: Inhibition of calcium carbonate production prevents biofilm**
1269 **formation and tissue deterioration in an *ex-vivo* lunge model**

A. Light microscopy images of 7 micron slices of lung tissues infected with *P. aeruginosa* PA14 and treated with biomineralization inhibitors as indicated. Samples were stained with H&E. Blue – nuclei, red – extracellular matrix and cytoplasm, (magnification x40). A representative experiment (out of n = 3) experiments is shown.

B. CFU of *P. aeruginosa* PA14 infecting lung tissue culture and surrounding media with and without indicated inhibitors (as in A). P values, as determined by student's t-test are indicated (* pVal <0.01, ** pVal <0.001, when compared to the untreated control).

C. Quantification of lung viability. Lung tissue cultures were infected with *P. aeruginosa* PA14 and treated with biomineralization inhibitors as indicated. ImageJ 1.51g software was used to automatically count lung cell nuclei in 4 randomly chosen fields for each treatment. P values, as determined by student's t-test are indicated (** pVal <0.001).

D. CFU analysis of wild type *P. aeruginosa* PA01 and the indicated mutants infecting lung tissue culture and surrounding media. Diamox was applied at 1.1 mg/ml, when indicated. P values, as determined by student's t-test are indicated (* pVal <0.01, ** pVal <0.001, when compared to the untreated control).

E. Quantification of lung viability. Lung tissue cultures were infected with *P. aeruginosa* PA14 and treated with biomineralization inhibitors, as indicated (Diamox and AHA applied at 1.1 mg/m). ImageJ 1.51g software was used to automatically count lung cell nuclei in 4 randomly chosen fields for each treatment. P values, as determined by student's t-test are indicated (** pVal <0.001).

F. *P. aeruginosa* PA14 biofilms were grown for 72 h on TB agar. The colonies were physically divided into equal parts, harvested, and half was treated for 2 h, as indicated. CIP – Ciprofloxacin, 3 mg/ml; GC – gentamycin, 10 mg/ml; AHA – 2.5 mg/ml; DTNB – 2 mg/ml. Viability is expressed as the ratio of CFU between the treated and the untreated parts of a colony. Results are averages

1300   and standard deviation of three independent experiments. P values, as

1301   determined by student's t-test are indicated (** pVal <0.001).

1302

A

B

sporulation ● matrix ● flagellum ●

C

D

Figure 2

A

B

C

	Dead	Spore	Mineral producer
Structured (n=510)	12.5%	12.7%	4%
Total colony (n=308)	7.5%	7.5%	1.6%

D

Mean Fluorescence Intensity (AU×10⁴)

Day 3
Day 6

Cell number

E

calcium | membrane
DIC | merged

F

G

H

WT | ΔarsR | ΔmntR | Δzur | ΔcueR

J

WT | ΔyetJ | ΔchaA | ΔyloB | ΔchaAΔyetJ

I

-log10(pVal)

motility/chemotaxis
biofilm formation

all transporters
divalent cation transporters

Log2 Fold Change

Figure 3

Figure 4

A

$H_2O + CO_2$

Urease (AHA)
CA (Diamox/DTNB)

H_2CO_3

$2H^+ + CO_3^{2-}$

$H^+ + HCO_3^-$

$H^+ + CO_3^{2-}$

Ca^{2+}

$CaCO_3$

P-type ATPases
(SMV)

Calcite

B

NT   AHA   Diamox

DTNB   SMV

C

Growth (OD$_{600}$)

DMSO
AHA
Diamox
DTNB

D

PA01   ΔpsCA1

ΔpsCA1-2   ΔpsCA1-2-3

ΔcalC

Growth (OD600)

PA01
ΔpsCA1-2-3
ΔcalC

Time (Hours)

E

Concentration (mg/mL)

DTNB
AHA
SMV

OD595

F

ΔcalC
ΔpsCA1-2-3
ΔpsCA1-2
ΔpsCA1
PA01

OD595

G

NT   Diamox

DTNB   AHA

SMV

Growth (OD600)

NT
AHA
DTNB
SMV
Diamox

Time (hours)

H

NT   + 0.5 mM EGTA   +1 mM EGTA   +Ca^{2+} + 0.5 mM EGTA +Ca^{2+} +1 mM EGTA + Ca^{2+}

Figure 5

A

- *P. aeruginosa* | + *P. aeruginosa* | + *P. aeruginosa* + AHA 2.5 mg/ml | + *P. aeruginosa* + AHA 5 mg/ml

B

+ *P. aeruginosa* + AHA 5 mg/ml

C

- *P. aeruginosa* | + *P. aeruginosa* | + *P. aeruginosa* + AHA 2.5 mg/ml | + *P. aeruginosa* + AHA 5 mg/ml

D

Number of lung nuclei/Area

- *P. aeruginosa*
+ *P. aeruginosa*
+ *P. aeruginosa* + AHA 2.5 mg/ml
+ *P. aeruginosa* + AHA 5 mg/ml

E

CFU

NT | 2.5 mg/ml | 5 mg/ml

□ Tissue ■ Medium

F

CFU Ratio (Tissue /Medium)

NT | 2.5 mg/ml | 5 mg/ml

Figure 6

The formation of microbial exoskeletons is driven by a controlled calcium-concentrating subcellular niche

Supplementary Information

Content

Supplementary Figures

Supplementary Files and Movies Legends

Supplementary Tables

Supplementary figures

Supplementary Figure 1. FTIR spectra of bleached *B. subtilis* biofilm colonies, displaying vibrations characteristic of calcite (crystalline calcium carbonate).

A

B

C

Supplementary Figure 2. (A). Scanning Electron Microscopy (SEM) image of *B. subtilis* biofilm colony grown for 6 days on a biofilm-inducing B4-Ca^{2+} agar. Shown are the surface of the biofilm and rhombohedral mineral structures in secondary electron mode. Magnification – X25000, scale bar – 1 µm. **(B).** Backscattered mode SEM image of the area shown in (**A**). **(C)** EDX spectra of the area shown in (**A**). CA – calcium rich area; Cnt – control area. A representative field (out of n=4 fields, from 8 independent experiments) is shown.

Supplementary Figure 3. Scanning Electron Microscopy (SEM) image of *B. subtilis* biofilm colony grown for 6 days on a B4 agar (without excess calcium). Shown are the surface of the biofilm in secondary electron and in backscattered mode of a representative field. **(A)** Magnification – X2500, scale bar – 10 μm. **(B).** Magnification – X10000, scale bar – 2 μm. Representative fields (out of n=4 fields, from 8 independent experiments) are shown.

Supplementary Figure 4. Light microscopy images of Δ*sigF* mutant strain. Biofilm colonies were grown on B4-Ca^{2+} agar for the indicated time at 30°C. The experiment was repeated 3 times, in a technical quadruplicate – and representative images are shown.

Supplementary Figure 5. Left panel - Bright-field STEM image of representative *B. subtilis* biofilm cells from a showing cellular calcium deposits, from a colony grown 10 days on B4-Ca^{2+} agar. The black arrow indicates the cell used for EDX analysis in panels (B-C). Right panel – EDX analysis of the calcium deposits. A representative mineralizing cell imaged (out of n=50 cells, from 4 independent experiments performed with EDX is shown.

.

Supplementary Figure 6. Cryo-STEM bright-field images of representative biofilm cells from a *B. subtilis* colony grown for 10 days on B4 agar (without calcium). Scale bar 1 µm. A representative field imaged (out of n=30 cells, from 4 independent experiments is shown.

Supplementary Figure 7. Cryo-STEM bright-field images of representative biofilm cells from a Δ*sigF* colony grown for 10 days on B4 agar. Black arrows point to aggregates with edge-on views (left panel) and top views (right panel). Scale bars 500 nm. A representative field imaged (out of n=30 cells), from 4 independent experiments is shown.

Supplementary Figure 8. Light microscopy images of wild type *B. subtilis* (WT) and Δ*copA* mutant. Biofilm colonies were grown on B4-Ca^{2+} agar for 3 days at 30°C. Scale bar – 2 mm. The experiment was repeated 3 times, in a technical quadruplicate – and representative images are shown.

Supplementary Figure 9. Planktonic growth assay. Wild type *B. subtilis* (ET) and *ΔcuerR* mutant were grown at 30°C with shaking in liquid LB, and growth was monitored by measuring OD_{600} in a microplate reader every 30 min. Results are averages of six wells, bars represent standard deviations. A representative experiment out of at least 3 independent experiments is shown.

Supplementary Figure 10. Light microscopy images of wild type (WT) and triple *ΔchaA ΔyloB ΔyetJ* mutant. Biofilm colonies grown on solid B4-Ca^{2+} agar for 3 days at 30°C. Scale bar – 2 mm. The experiment was repeated 3 times, in a technical quadruplicate – and representative images are shown.

Supplementary Figure 11. Light microscopy of biofilm colonies of *P. aeruginosa* PA01 supplemented with EDTA (0.1mg/ml), and divalent ions (1mM), as indicated. Colonies were grown on BHI solid medium for 4 days at 23ºC. Scale bar – 5 mm. The experiment was repeated 3 times, in a technical 3 repeats – and representative images are shown.

	WT	Δ*pelpsl*
NT		
EDTA		

Supplementary figure 12: Light microscopy of biofilm colonies of *P. aeruginosa* PA01 either untreated or supplemented with EDTA (0.1mg/ml) as indicated. Colonies were grown on BHI solid medium for 4 days at 23ºC. Scale bar = 5 mm. The experiment was repeated 2 times, in a technical 3 repeats – and representative images are shown.

Supplementary Figure 13. FTIR spectra of bleached *M. abscessus* biofilm colonies, displaying vibrations characteristic of varterite (crystalline calcium carbonate).

Bacteria	Medium	pH					Medium Langelier Saturation index (LSI)
		medium control	far from colony	close to colony	below the colony	colony center	
B. subtilis	B4Ca^{2+}	6	7.5	7.75	7.75	8	0.51-0.015
P. aeruginosa	TB	6	8	8	8	8	2.4—1.4
M. abscessus	B4Ca^{2+}	6	8.5	8.5	8	8	0.51-0.015

Supplementary Figure 14. pH of colonies and the agar medium they were grown on was assessed using pH-indicator strips (MColorpHast™). The experiment was conducted at least twice for each medium, with at least 4 technical repeats. LSI calculation refers to the indicated growth media. ND – undetermined.

Supplementary Figure 15. MicroCT-XRay image of a wild type *P. aeruginosa* PAO14 colony grown for 3 days at 23°C on TB agar containing SMV (0.01 mg/ml). The absence of visible bright areas indicates lack contrast, indicating no dense mineral (compare to the control colony shown in Fig. 3A). Scale bar – 2 mm. The experiment was repeated twice with similar results.

Supplementary Figure 16. Planktonic growth assay. *P. aeruginosa* PA14 was grown at 37°C with shaking in liquid TB medium, supplemented with: 1.5 mg/ml DTNB, 1.75 mg/ml AHA, 2.5 mg/ml Diamox and 0.01, 0.05 and 0.01 mg/ml SMV. Growth was monitored by measuring OD600 in a microplate reader every 30 min. Results are averages of six wells, bars represent standard deviations. A representative experiment out of at least 3 independent experiments is shown.

| Aerobic | Anaerobic | Aerobic | Anaerobic |

Supplementary Figure 17. Light microscopy of biofilm colonies formed by *P. aeruginosa* PA14 (left) and *B. subtilis* (right). The biofilms were grown TB and B4-Ca^{2+} agar, respectively, with nitrate source (0.5% KN0^3). Under aerobic conditions, CO_2 is available from the atmosphere and also produces as a byproduct of aerobic respiration. Under anaerobic conditions of nitrate reduction (as used here - anaerobic growth chamber, a mix of H_2 and N_2 (5/95%), CO_2 is absent from the atmosphere, and aerobic respiration (and subsequent CO_2 emission) is not feasible due to the lack of oxygen. The experiment was repeated 4 times, in a technical 3 repeats– and representative images are shown.

Supplementary Figure 18. Light microscopy of 3-day-old biofilm colonies formed by *M. abscessus* on B4 agar supplemented with calcium acetate (0.25% v/v) and EGTA, as indicated. Scale bar = 5 mm. The experiment was repeated 3 times, in a technical 3 repeats– and representative images are shown.

Supplementary Figure 19. Light microscopy images of biofilm colonies formed by *M. abscessus*, supplemented with SMV (0.025mg/ml) as indicated. Colonies were grown on B4-Ca²⁺ medium for 5 days, at 37°C. Scale bar = 5 mm. The experiment was repeated 2 times, in a technical 3 repeats – and representative images are shown.

Supplementary Figure 20.

P. aeruginosa and *M. abscessus* biofilm colonies grown on solid SCFM medium for 4 days at 23ºC (*P. aeruginosa*) and 30ºC (*M. abscessus* and *S. aureus*). Scale bar – 2 mm. The experiment was repeated 3 times, in a technical 3 – and representative images are shown.

Supplementary Figure 21. FTIR spectra of bleached sputum samples of *P. aeruginosa* positive CF patients (I-VI). Arrows indicate vibrations characteristic of calcite. For patient details, see Supplementary table 2.

Supplementary Figure 22. Confocal image of *ex-vivo* lung model. Lungs were harvested from one-month-old mice, cultured in DMEM containing carbenicillin 100 μg/ml, and infected with *P. aeruginosa* PA14 constitutively expressing GFP. Green – bacterial cells. Blue - nuclei of lung cells stained with DAPI. Scale bar – 50 μm. The experiment was repeated 3 times, in a technical 4 repeats – and representative image is shown.

Supporting Figure 23. Quantification of lung viability. Lung tissue cultures were treated with biomineralization inhibitors DTNB (2 mg/ml), SMV (0.01 mg/ml), Diamox (1.1 mg/ml) and AHA, as indicated. ImageJ 1.51g software was used to automatically count lung cell nuclei in 4 randomly chosen fields for each treatment. P values, as determined by student's t-test, are indicated (* pVal <0.01, ** pVal <0.001).

WT　　　　　　　ΔyloA

Supporting Figure 24. Light microscopy images of wild type (WT) and *ΔyloA* mutant. Biofilm colonies grown on solid B4-Ca^{2+} agar for 3 days at 30°C. Scale bar – 2 mm. The experiment was repeated 3 times, in a technical quadruplicate – and representative images are shown.

ΔcalC   *ΔcalC, attB::calC*

Supporting Figure 25. Light microscopy of biofilm colonies of *P. aeruginosa* PA01 indicated strains. Colonies were grown on BHI solid medium for 4 days at 23ºC. Scale bar = 5 mm. The experiment was repeated 3 times, in a technical 2 or more repeats – and representative images are shown.

Supplementary Files:

Supplementary File 1: Differentially expressed genes in *B. subtilis* NCIB 3610, + vs - calcium.

Supplementary File 2: Differentially expressed genes *B. subtilis* NCIB 3610, WT vs *cueR*.

Supplementary File 3: Patients information. Clinical parameters of patients analyzed for presence of calcite in sputum samples.

Supplementary Movies:

Supplementary Movie 1: *B. subtilis* colony

Supplementary Movie 2: *P. aeruginosa* PA14 colony

Supplementary Movie 3: *M. abscessus* colony

The intact colonies were visualized under X-ray, and the obtained 2D images were used to generate high-resolution 3D image (see Keren-Paz *et al.,* 2018 for details). Rotation of 360° was taken around an axis perpendicular to the biofilm surface. Results are of a representative experiment out of three independent repeats.

Supplementary Tables:

Supplementary Table 1: Strains used in this study

Genotype	Reference
Bacillus subtilis NCIB 3610	Branda *et al* (2001)
ΔarsR (kan)	This work
ΔmntR (kan)	This work
Δzur (kan)	This work
ΔcueR (kan)	This work
ΔcopA (kan)	This work
ΔyetJ (spec)	This work
ΔchaA (kan)	This work
ΔyloB (mls)	This work, the lack of polar effect was confirmed by the deletion of *yloA* (Figure S24)
ΔchaA (kan) *ΔyetJ* (spec)	This work
ΔchaA (kan) *ΔyetJ* (spec) *ΔyloB* (mls)	This work
Pseudomonas aeruginosa PA14	Lab collection
Pseudomonas aeruginosa PA14, Ppuc-GFP (carb)	[1]
Pseudomonas aeruginosa PA01	Lab collection
Pseudomonas aeruginosa PA01 *ΔpsCA1*	[1]
Pseudomonas aeruginosa PA01 *ΔpsCA1-2*	[1]
Pseudomonas aeruginosa PA01 *ΔpsCA1-2-3*	[1]

Pseudomonas aeruginosa PA01 Δ*calC*	This work, This work, the lack of polar effect was confirmed by complementation of *calC* from an ectopic chromosomal loci (Figure S25), ΔcalC attB::calC
Mycobacterium abscessus ATCC 19977	A kind gift from Daniel Barkan

1   Lotlikar, S. R. *et al.* Pseudomonas aeruginosa beta-carbonic anhydrase, psCA1, is required for calcium deposition and contributes to virulence. *Cell Calcium* **84**, 102080, doi:10.1016/j.ceca.2019.102080 (2019).

Supplementary Table 2:

Patient	Medical	Gender	CFTR mutations	Infection
I 463116	Carmel	M	W1282X/F508	PA+MA
II A19	Hadassah	M	G542X/F508	PA
III 463125	Carmel	F	T360K, Q359K	PA
IV B9	Hadassah	F	W1282X/W1282X	PA
V A21	Hadassah	F	W1282X/F508	PA
VI A32	Hadassah	M	W1282X/W1282X	PA

The details of patients included Supplementary Figure 15 and their respected panels are indicated.