1

A silent disco: Persistent entrainment of low-frequency neural oscillations underlies

beat-based, but not pattern-based temporal expectations

Short title: Persistent effects of beat- and pattern-based temporal expectations

Fleur L. Bouwer[1,2,3*], Johannes J. Fahrenfort[1,3], Samantha K. Millard[3,4,5], Niels A.

Kloosterman[6,7], & Heleen A. Slagter[1,3]

[1] Department of Experimental and Applied Psychology, Institute Brain and Behavior
Amsterdam (IBBA), Vrije Universiteit Amsterdam
[2] Institute for Logic, Language and Computation (ILLC), University of Amsterdam
[3] Department of Psychology, Amsterdam Brain and Cognition (ABC), University of
Amsterdam
[4] Centre for Pain IMPACT, Neuroscience Research Australia (NeuRA), Sydney, New South
Wales, Australia
[5] Faculty of Medicine, University of New South Wales (UNSW), Sydney, New South Wales,
Australia
[6] Max Planck UCL Centre for Computational Psychiatry and Ageing Research, Berlin,
Germany
[7] Center for Lifespan Psychology, Max Planck Institute for Human Development, Berlin,
Germany

* Correspondence concerning this article should be addressed to Dr. Fleur Bouwer,
Department of Psychology, University of Amsterdam (bouwer@uva.nl)

FLB is supported by an ABC Talent Grant awarded by Amsterdam Brain and Cognition, a Veni
grant awarded by the Dutch Research Council NWO (VI.Veni.201G.066), and the ERC starting
grant awarded to HAS. HAS is supported by a European Research Council (ERC) starting grant
(679399).

Highlights

-   Temporal expectations can be based on both regular beats and predictable patterns

-   Behavioral effects are more persistent for beat-based than pattern-based

expectations

-   EEG power tracks the beat, but not the pattern, outlasting rhythmic stimuli

-   Pattern-based, but not beat-based expectations elicit a CNV-like ERP component

-   Decoding and entropy may index temporal expectations in a time-resolved way

Abstract

The brain uses temporal structure in the environment, like rhythm in music and speech,

to predict the timing of events, thereby optimizing their processing and perception. Temporal

expectations can be grounded in different aspects of the input structure, such as a regular beat

or a predictable pattern. One influential account posits that a generic mechanism underlies beat-

based and pattern-based expectations, namely entrainment of low frequency neural oscillations

to rhythmic input, while other accounts assume different underlying neural mechanisms. Here,

we addressed this outstanding issue by examining EEG activity and behavioral responses

during silent periods following rhythmic auditory sequences. We measured responses

outlasting the rhythms both to avoid confounding the EEG analyses with evoked responses, and

to directly test whether beat-based and pattern-based expectations persist beyond stimulation,

as predicted by entrainment theories. To properly disentangle beat-based and pattern-based

expectations, which often occur simultaneously, we used non-isochronous rhythms with a beat,

a predictable pattern, or random timing. In Experiment 1 (N = 32), beat-based expectations

affected behavioral ratings of probe events for two beat-cycles after the end of the rhythm,

while the effects of pattern-based expectations reflected only one interval. In Experiment 2 (N

= 27), using EEG, we found enhanced spectral power at the beat frequency for beat-based

sequences both during listening and the silence, but for pattern-based sequences, enhanced

power at a pattern-specific frequency was only present during listening, not silence. Moreover,

we found a negative deflection in the evoked signal following pattern-based, but not beat-based

sequences. Finally, we show how multivariate pattern decoding and multi scale entropy –

measures sensitive to non-oscillatory components of the signal – can be used to probe temporal

expectations. Together, our results suggest that different mechanisms implement temporal

expectations, depending on the input structure. We suggest climbing activity may reflect

pattern-based, and persistent low frequency oscillations beat-based expectations specifically.

Keywords: Temporal expectations, EEG, rhythm, entrainment, decoding, entropy

Introduction

Predicting the timing of incoming events optimizes processing in our dynamic environment (Nobre & van Ede, 2018), as it allows the brain to increase sensitivity to events at predicted times (Auksztulewicz, Myers, Schnupp, & Nobre, 2019), without the need for constant vigilance (Breska & Deouell, 2017; Rimmele, Morillon, Poeppel, & Arnal, 2018; Schroeder & Lakatos, 2009b, 2009a). Entrainment models (Large & Jones, 1999) provide a mechanistic explanation for temporal expectations, by assuming that the phase and period of low-frequency neural oscillations synchronize to external rhythmic stimulation, causing optimal neural excitability at expected times (Haegens & Zion Golumbic, 2018; Henry & Herrmann, 2014; Schroeder & Lakatos, 2009a). In line with this, behavioral performance is improved for events in phase with an external rhythm (Bouwer & Honing, 2015; Herbst, Stefanics, & Obleser, 2022; Jones, Moynihan, MacKenzie, & Puente, 2002; Large & Jones, 1999), behavioral responses depend on the phase of delta oscillations (Arnal, Doelling, & Poeppel, 2014; Cravo, Rohenkohl, Wyart, & Nobre, 2013; Henry, Herrmann, & Obleser, 2014; Henry & Obleser, 2012), and low frequency oscillations phase lock to rhythmic input (Doelling, Assaneo, Bevilacqua, Pesaran, & Poeppel, 2019; Nozaradan, Peretz, Missal, & Mouraux, 2011; Stefanics et al., 2010).

Entrainment has mainly been studied in the context of periodic ("beat-based") sensory input, but temporal expectations can also be based on memory for absolute intervals (Breska & Deouell, 2017; Breska & Ivry, 2016; Morillon, Schroeder, Wyart, & Arnal, 2016; Teki, Grube, Kumar, & Griffiths, 2011), either in isolation ("cue-based"), or as part of a predictable pattern of intervals ("pattern-based", see (Nobre & van Ede, 2018)). Predictable temporal patterns may be especially important in speech and non-Western music, which is not necessarily periodic. Expectations based on predictable patterns in aperiodic sequences afford similar behavioral benefits as expectations based on a beat (Bouwer, Honing, & Slagter, 2020; Heideman, van Ede, & Nobre, 2018; O'Reilly, McCarthy, Capizzi, & Nobre, 2008), but pose a possible

challenge for entrainment models, which are arguably better suited to explain temporal expectations for periodic input (Breska & Deouell, 2017; Rimmele et al., 2018). Some have suggested that entrainment models can account for pattern-based expectations by assuming multiple coupled oscillators at different frequencies and with different phases (Tichko & Large, 2019), or by assuming flexible top-down phase resets at expected moments, though this would entail some top-down mechanism, making observed entrainment the consequence, rather than the cause of expectations (Meyer, Sun, & Martin, 2019; Obleser & Kayser, 2019; Rimmele et al., 2018).

Alternatively, however, pattern-based and beat-based expectations could arise from dissociable neural mechanisms. For cue-based expectations, tentative evidence for a different underlying mechanism comes from a series of studies looking at the contingent negative variation (CNV), an event-related potential (ERP) component that peaks at expected moments (Praamstra, Kourtis, Kwok, & Oostenveld, 2006). The CNV resolved faster for beat-based than cue-based expectations (Breska & Deouell, 2017), and cerebellar patients showed selective impairments in forming cue-based, but not beat-based expectations (Breska & Ivry, 2018, 2020). However, in these studies, the intended beat-based sequences were isochronous. Isochronous sequences can, in addition to a beat, elicit temporal expectations through learning the repeated, identical interval (Breska & Ivry, 2016; Keele, Nicoletti, Ivry, & Pokorny, 1989). Thus, the differences in responses may here be explained by more precise cue-based or pattern-based expectations in the isochronous, beat-based condition. Moreover, these studies tested temporal expectations based on the contingency between a cue and an interval (e.g., learning a single interval), and it is unclear whether temporal expectations based on patterns are based on the same mechanism (Nobre & van Ede, 2018). In our own recent work, we specifically compared beat-based and pattern-based expectations, and we found no difference in the effects of these expectations on early auditory ERP responses, suggestive of similar modulation of

sensory processing (Bouwer et al., 2020), but we observed suppression of sensory processing

of unexpected events in beat-based rhythms, even when these events were fully predictable

based on their pattern, suggesting different underlying mechanisms (Bouwer et al., 2020).

However, in this study, we did not directly probe and contrasted the neural mechanisms

underlying beat-based and pattern-based expectations, rendering it unclear whether they are

subserved by shared or separate neural dynamics.

In the current study, we directly examined the role of entrainment in beat-based and

pattern-based expectations, using non-isochronous rhythms designed to properly disentangle

these two types of expectations. When studying entrainment, an important challenge has been

to differentiate between real entrainment ("in the narrow sense", see Obleser & Kayser, 2019)

and regular evoked potentials, or similar phase locked responses that resemble entrainment with

common analysis techniques (Zoefel, ten Oever, & Sack, 2018), and that may not differentiate

between beat-based and memory-based expectations (Breska & Deouell, 2017). Crucially, to

sidestep these issues, here we examined responses in a silent window after cessation of the

rhythmic input, directly testing the prediction of entrainment models that entrainment should

outlast sensory stimulation (Haegens & Zion Golumbic, 2018; Obleser & Kayser, 2019; Pesnot

Lerousseau, Trébuchon, Morillon, & Schön, 2021; Zoefel et al., 2018).

Behaviorally, persistent entrainment has been shown for auditory rhythm (Hickok,

Farahbod, & Saberi, 2015; Jones et al., 2002), though this effect is not always found (Bauer,

Jaeger, Thorne, Bendixen, & Debener, 2015; Lin et al., 2021), possibly due to heterogeneity in

the population and effects of musical training (Assaneo et al., 2019; Cameron & Grahn, 2014;

Sun, Michalareas, & Poeppel, 2021). At a neural level, several studies reported persistent

entrainment in the visual (de Graaf et al., 2013; Mathewson et al., 2012), and auditory (Kösem

et al., 2018; Pesnot Lerousseau et al., 2021; van Bree, Sohoglu, Davis, & Zoefel, 2021; Wilsch,

Mercier, Obleser, Schroeder, & Haegens, 2020) domain. However, in these studies,

isochronous stimuli were used, making it unclear whether the expectations probed were based

on a beat, or were formed based on the single repeating interval (i.e., cue- or pattern-based).

Moreover, persistent entrainment was not specific to the frequency of the input (Wilsch et al.,

2020), or only occurred in the gamma (Pesnot Lerousseau et al., 2021), or alpha ranges (de

Graaf et al., 2013; Mathewson et al., 2012), while humans have a preference for forming

temporal expectations at slower rates (Ding et al., 2017; Merchant, Grahn, Trainor, Rohrmeier,

& Fitch, 2015; Zalta, Petkoski, & Morillon, 2020), as naturally present in speech and music

(i.e., the delta and theta range). Thus, not only is evidence for whether entrainment can account

for pattern-based expectations lacking, evidence for persistent entrainment in response to beat-

based rhythms remains elusive as well.

In the current study, participants listened to non-isochronous auditory sequences

(similar to those used in Bouwer et al., 2020) with either a regular beat (eliciting beat-based

expectations), a predictable pattern (eliciting pattern-based expectations), or random timing (no

expectations). The non-isochronous beat-based sequences had a varying surface structure,

similar to patterns used to probe beat-based processing in many neuroimaging (Grahn & Brett,

2007; Grahn & Rowe, 2009; Leow & Grahn, 2014), behavioral (Bouwer et al., 2018, 2021;

Cameron & Grahn, 2014; Povel & Essens, 1985), and electrophysiological studies (Lenc et al.,

2021). While each beat was marked by a sound, between beats, sounds could occur at different

times, or not at all. Therefore, the beat could not be extracted from the rhythmic signal by

simply learning the transition of temporal intervals – as is possible in isochronous sequences.

Moreover, in our previous study, using the same stimuli, we showed that the behavioral

facilitation caused by introducing a regular beat was smaller than the facilitation caused by

introducing a predictable pattern (Bouwer et al., 2020). Hence, the non-isochronous beat-based

sequences used here did not allow for easy learning of the interval of the beat using a pattern-

based strategy, in contrast to the pattern-based sequences, in which the surface structure of the

rhythm was fully predictable.

Each sequence was followed by a silent period. In Experiment 1, we asked participants

to rate how well probe tones, presented at various time points during the silent period, fitted the

preceding sequence. We expected the ratings to be affected by both the beat-based and pattern-

based expectations elicited by the sequences. In Experiment 2, we recorded EEG activity both

during presentation of the sequences and during the silence. If entrainment underlies temporal

expectations, we should see persistent power at the frequency of the beat or pattern during the

silence. Alternatively, if climbing activity in the form of a CNV underlies temporal

expectations, we should see a CNV peaking at expected time points in the silence.

In addition to examining the spectral power at the frequencies of the beat and the pattern,

and the evoked responses, we explored two new methods to index temporal expectations that

do not rely on the EEG signal being a static oscillation. As recently argued, the oscillatory

dynamics underlying such expectations may be subject to changes in power and frequency,

depending on coupling between sound and brain, and on the properties of the neural dynamics

themselves (e.g., to which extent the system shows damping of an oscillation without input, see

Doelling & Assaneo, 2021). Once the rhythmic sensory input ceases, the oscillatory activity in

the brain may quickly return to an intrinsic resonance frequency (Doelling & Assaneo, 2021).

Also, the presence of non-sinusoidal recurring activity may not be captured by traditional

analyses relying on Fourier transforms (Donoghue, Schaworonkow, & Voytek, 2021), while it

may be important for cognition (Waschke, Kloosterman, Obleser, & Garrett, 2021). Therefore,

we here explore indexing temporal expectations in the silence using multi scale entropy (MSE)

– a measure of signal irregularity (Kosciessa, Kloosterman, & Garrett, 2020) – and multivariate

pattern decoding. These methods may provide useful tools to study the neural dynamics

underlying rhythm processing and temporal expectations, which are often hard to study due to

methodological issues with analyzing the EEG signal (Zoefel et al., 2018). While MSE provides

us with a method to look at possible non-sinusoidal contributions to the EEG signal related to

temporal expectations, decoding allows us to look at how entrainment evolves over time. Note

that we consider these analyses exploratory in nature, and results should be interpreted as such.

Materials and methods

Participants

Thirty-two participants (18 women), aged between 18 and 44 years old (M = 24, SD = 5.6) took part in the behavioral Experiment 1, and 32 participants (26 women), aged between 19 and 28 years old (M = 23, SD = 2.5) took part in the EEG experiment (Experiment 2), in exchange for course credit or monetary compensation. Due to technical problems, the EEG data from five participants was not recorded correctly, hence we report the results for 27 participants (21 women, between 19 and 28 years old, M = 23, SD = 2.4). For the behavioral experiments, we used mixed-effects models, which need both the number of participants and the number of items to be taken into account to assess power (Brysbaert & Stevens, 2018). In two similar experiments in which ratings in response to rhythms of varying complexity were analyzed, small-sized effects were replicated with a total number of around 275 responses per condition (Bouwer, Burgoyne, Odijk, Honing, & Grahn, 2018). For our new experimental paradigm, we here included a multiple of this amount of trials (960 responses per condition and probe position in Experiment 1 – 32 participants with 30 responses each in each cell – and, after loss of data was accounted for, 486 responses per condition and probe position in Experiment 2 – 27 participants and 18 responses per cell). Previous EEG experiments examining persistent entrainment in the auditory domain used sample sizes ranging from fifteen (Pesnot Lerousseau et al., 2021) to twenty-one (van Bree et al., 2021), similar to the sample size used in a study looking at different types of temporal expectations (twenty-one, Breska & Deouell, 2017). To obtain robust power, we here tested thirty-two participants, which, even when loss of data is accounted for, thus exceeded typical sample sizes as used previously. None of the participants reported a history of hearing or neurological problems, and all provided written informed consent prior to the onset of the study. The experiment was approved by the Ethics Review Board of the Faculty of Social and Behavioral Sciences of the University of Amsterdam.

Figure 1. Schematic overview of the rhythmic stimuli used and the task. A) Twelve patterns of five temporal intervals with integer ratio durations and an event at each 600 ms period were created to form beat-based sequences. Equivalent patterns without a regular beat every 600 ms were created by using non-integer ratio durations, while keeping the number of intervals and grouping structure the same (random condition). For the pattern-based sequences, only pattern 1 was used, to allow for learning of the intervals. B) Four semi-randomly chosen patterns were concatenated to form rhythmic sequences. In both the beat-based and random sequences, the last pattern was always pattern 1 or 2, to equate the acoustic context preceding the silent period. C) To measure behavioral effects of expectations, a probe tone could appear at various temporal positions in the silent period (indicated by the dashed red lines), predictable based on a beat (B+, light orange), predictable based on the pattern (P+, light purple), or unpredictable based on the beat (B-) or pattern (P-). Subjects had to indicate how well the probe tone fitted the preceding rhythm. In Experiment 1, all 6 probe tone positions were used. In Experiment 2, only the last 3 probe positions were used.

 Stimuli

We used patterns marked by woodblock sounds of 60 ms length, generated in

GarageBand (Apple Inc.), as previously used in (Bouwer et al., 2020) to elicit beat-based and

pattern-based expectations (Figure 1). Each pattern was 1800 ms long and consisted of five

temporal intervals. The number of tones was chosen to be within the range that was previously

shown to allow for learning of a predictable pattern (Schultz, Stevens, Keller, & Tillmann,

2013), while the length of the pattern was such that the formation of beat-based expectations

with a period of the entire pattern would be unlikely in the pattern-based sequences, given that

this would require hearing a beat at 0.55 Hz, which is very far from the range at which humans

can typically perceive a beat (Honing & Bouwer, 2019; London, 2012). Sequences (beat-based,

pattern-based, or with random timing) were constructed by concatenating four patterns and a

final tone, for a total sequence length of 7260 ms (four patterns of 1800 ms, plus 60 ms for the

final tone).

In the twelve patterns used to create beat-based sequences (Figure 1), temporal intervals

were related to each other with integer-ratio durations. The shortest interval had a length of 150

ms, with the relation between the five intervals used of 1:2:2:3:4 (i.e., 150, 300, 300, 450, and

600 ms). The sounds were grouped such that a perceptually accented sound (Povel &

Okkerman, 1981) occurred every 600 ms (every unit length 4), giving rise to a beat at 100 beats

per minute, or 1.67 Hz, within the range of preferred tempo for humans (London, 2012). All

beat-based patterns were strictly metric, with the beat always marked by a sound (Grahn &

Brett, 2007). Sequences of beat-based patterns were constructed from four semi-randomly

chosen patterns, with the restriction that the last pattern of the sequences was always pattern 1

or 2 (see Figure 1). This way, the final 600 ms preceding the silence epoch was equated in terms

of the acoustic context, to make the bleed of auditory ERPs into the silence as similar between

conditions as possible. Note that in beat-based sequences, a sound could be expected every 600

ms based on the beat, but the surface structure of the pattern was unpredictable, due to the

random concatenation of patterns.

To create patterns that did not allow for beat-based expectations ("aperiodic" patterns,

see Figure 1), the ratios by which the temporal intervals were related were changed to be non-

integer (1:1.4:1.4:3:5.2, or 150, 210, 450, and 780 ms respectively). In these patterns, no marked

beat was present at unit length four, nor at any other subdivision of the 1800 ms pattern (Bouwer

et al., 2020), while the patterns were matched to their periodic counterparts in terms of overall

length, event density, number of sounds, and grouping structure.

From the aperiodic patterns, two types of sequences were created: pattern-based and

random sequences. To create sequences allowing for pattern-based expectations, we

concatenated four identical patterns. To be able to use the data with an EEG-based decoding

analysis (Experiment 2), we needed the timing of expectations in the silence to be identical for

each sequence, hence we restricted the pattern-based sequences to only pattern 1. The use of a

single pattern was not only necessary for decoding, but also optimized the experiment for

pattern-based expectations, since participants only had to memorize one pattern, allowing them

to easily form expectations, even if the single sequences were only four patterns long.

For the random sequences, four semi-randomly chosen aperiodic patterns were

concatenated. Like for the beat-based sequences, the final pattern was always pattern 1 or 2,

equating the final 600 ms of the sequences in terms of acoustics. In the random sequences, the

timing of sounds could not be predicted based on the surface structure of the pattern, nor on the

basis of an underlying beat.

A spectral analysis of the stimuli in the range in which a beat can normally be perceived

(the delta-range, 0.5-4 Hz) confirmed that in the beat-based sequences, a peak was present at

the beat frequency of 1.67 Hz as well as at 3.33 Hz (see Figure 2). The 3.33 Hz peak is a

harmonic of the beat frequency, but also the frequency at which participants may perceive

subdivisions of the beat (e.g., an extra layer of perceived metrical regularity with a period of

300 ms). In the pattern-based and random sequences, peaks were more distributed, in line with

the more irregular nature of these rhythms, and the highest peaks in the delta range were at 2.22

and 3.89 Hz.

Figure 2. Spectral analysis of the sound signal of the different rhythmic sequences. Ten sequences per
condition were generated to base the spectral analysis on. Note that these sequences were identical for the
pattern-based condition, but semi-random for the beat-based and random conditions. The envelope of each
sequence was obtained by performing a Hilbert transform, and subsequently, a fast fourier transform (fft)
was used to obtain the spectral decomposition and power values were averaged over ten sequences. n.a.=
normalized amplitude.

In Experiment 1, to assess the persistence of temporal expectations behaviorally, on

each trial, a probe tone was presented at 600, 780, 885, 990, 1200, or 1485 ms after the onset

of the last tone of the sequence (see Figure 1C), and participants provided ratings for how well

probe tones fitted the preceding rhythm. These positions were carefully chosen to represent

times at which a tone could be expected based on the beat (600, 1200 ms), based on memory

for the pattern (780, 990, 1200 ms), or neither (885, 1485 ms). Note that the latter two probe

tones that were unexpected based on the beat (780, 885, 990, and 1485 ms) did not fall on

subdivisions of the beat.

Experiment 2 contained both trials in which a probe tone was presented (25% of the

trials, using only the last three probe positions), and trials in which a 7260 ms sequence was

followed by a silence period without a probe tone (75% of the trials). The latter were used for

EEG analyses, uncontaminated by a probe presentation. The silent period lasted for 2100 ms

after the onset of the last sound, providing 300 ms immediately following the final sound to allow for ERPs to mostly return to baseline, and 1800 ms (three full beat cycles, or a full cycle of the repeating pattern) of silence for the analysis. After the silence, the onset of the next sequence was jittered between 25 and 75 ms to prevent carryover of the beat from previous sequences (e.g., the next trial started between 325 and 375 ms after the last beat in the silence, which is not on the beat, nor at a subdivision of the beat). On trials that contained a probe tone, we chose to only use the last three probe positions, to 1) limit the time for the EEG experiment to prevent fatigue, 2) still provide participants with the incentive to form expectations well into the silence period, as a probe tone could appear as late as 1485 ms into the trial, and 3) obtain some measure of whether participants formed expectations, by including positions that were expected based on the beat and the pattern (1200 ms), based on the pattern only (990 ms) or neither (1485 ms).

Procedure

Participants were tested individually in a dedicated lab at the University of Amsterdam. Upon arrival, participants were provided with information about the experiment, provided written informed consent, and were allowed to practice the task. On probe tone trials (all trials in Experiment 1, 25% of the trials in Experiment 2), participants were asked to judge on a four-point scale ("very poorly", "poorly", "well", "very well") how well the probe tone fitted the preceding sequence, similar to previous studies investigating the perception of musical meter (Manning, Harris, & Schutz, 2017; Manning & Schutz, 2013; Palmer & Krumhansl, 1990). Participants could respond with four buttons on the armrest of their chair, two on each side. The order of the answer options on the screen in front of the participants was randomized on each trial, to avoid any artefacts of motor preparation in participants that anticipated which answer they would provide. There was no time limit for responses and the next trial started after a response was made.

In Experiment 1, each participant was presented with 18 blocks of 30 trials, amounting to 540 trials in total (30 trials per probe position for each condition). In Experiment 2, participants were presented with 18 blocks of 36 trials, for a total of 648 trials (18 per condition and position). In Experiment 2, for each condition, 162 trials were silence trials, and did not contain a probe tone. Fifty-four trials for each condition contained a probe tone. In both experiments, in each block, only one type of sequence (beat-based, pattern-based, or random) could appear to optimize for the formation of expectations. Blocks were semi-randomized, with each type appearing once in a set of three blocks. In each block of Experiment 2, the number of probe trials was varied between 3 and 11, for an average of 25% probe trials, and 75% silent trials.

Sounds were presented through a single speaker positioned in front of the participant using Presentation® software (version 14.9, www.neurobs.com). After completion of the experiment, participants performed the Beat Alignment Task (Iversen & Patel, 2008; Müllensiefen, Gingras, Musil, & Stewart, 2014) to assess their beat perception abilities, and completed the musical training subscale from the Goldsmith Musical Sophistication Index (GMSI) questionnaire to assess their musical training (Müllensiefen et al., 2014). In total, a behavioral session lasted two hours, and the EEG session lasted between 3.5 and 4 hours.

Behavioral analysis

A total of 17280 responses was included in the analysis of Experiment 1 (32 participants, 3 conditions, 6 probe positions, 30 responses each), and 4374 in the analysis of Experiment 2 (27 participants, 3 conditions, 3 probe positions, 18 responses each). To account for the ordinal nature of the Likert-scale responses (Bouwer et al., 2018; Carifio & Perla, 2008; Jamieson, 2004), we used a mixed ordinal regression model. With this model, the ordinal responses are normalized, to correct for potential unequal distances between rating points. The results can subsequently be interpreted similar to the results from a normal mixed model regression. Two

independent variables and their interaction were included in the model as fixed factors:

Condition (beat-based, pattern-based, or random), and Probe Position (600, 780, 885, 990,

1200, or 1485 ms; only the latter three in Experiment 2). Additionally, the score on the GMSI

musical training questionnaire was included as a continuous variable (Musical Training), as

well as its interactions with the two fixed factors. We used a random intercept for each subject

to account for between-subject variation.

The initial model showed a significant effect of Probe Position in the random condition,

most likely due to recency effects. To assess the effect of Probe Position in the beat-based and

pattern-based conditions while accounting for recency effects, for each participant we

subtracted the mean response in the random condition at each position from the responses in

the beat-based and pattern-based condition. Subsequently, we submitted the random-baseline-

corrected ratings to a second ordinal regression model, with only two levels for the factor

Condition (beat-based and pattern-based) and without the random intercept for each participant

(as the baseline correction already corrected for between-subject variability). For both the

original model, and the baseline corrected model, significant interactions were followed up by

tests of simple effects, corrected for multiple comparisons using a Bonferroni correction.

The statistical analysis was conducted in R (R Development Core Team, 2008). The

ordinal mixed model was implemented using the clmm() function from the ordinal package

(Christensen, 2019). Subsequently, we used the Anova() function from the car package (Fox &

Weisberg, 2019) to look at omnibus effects for the main factors of interest, and the emmeans

package (Lenth, 2019) to assess simple effects and compare slopes between conditions.

EEG recording

EEG was recorded at 1024 Hz using a 64-channel Biosemi Active-Two acquisition

system (Biosemi, Amsterdam, The Netherlands), with a standard 10/20 configuration and

additional electrodes for EOG channels (located under and on the left and right sides of the

eye), on the nose, on both mastoids, and on both earlobes.

EEG analysis

Preprocessing was performed in MATLAB, version 2015a (Mathworks) and EEGLAB,

version 14.1.1 (Delorme & Makeig, 2004). Data were offline down-sampled to 256 Hz, re-

referenced to averaged mastoids, bad channels were manually removed, and eye-blinks were

removed using independent component analysis. Subsequently, bad channels were replaced by

values interpolated from the surrounding channels.

ERPs. For the ERP analysis, the continuous data were filtered using 0.1 Hz high-pass

and 40 Hz low-pass finite impulse response filters (as implemented in the standard EEGLAB

filter function pop_eegfiltnew). Epochs were extracted from the data from -1800 till 2100 ms

relative to the onset of the last sound. Epochs with a voltage change of more than 150 microvolts

in a 200 ms sliding window were rejected from further analysis. For each participant and

condition, epochs were averaged to obtain the ERPs, and ERPs were averaged over participants

to obtain grand average waveforms for plotting. All waveforms were initially baseline corrected

using the average voltage of a 50 ms window preceding the onset of the last sound of the

sequence. This baseline can be regarded as preceding the "cue" (the last event before the onset

of the expectation). Such a baseline is customary in CNV analyses. However, visual inspection

suggested that this baseline was biased, as baseline correction resulted in an overall shift of the

waveform amplitude relative to each other, as also reflected in a significant cluster when

comparing beat-based and pattern-based conditions that spanned the entire analysis epoch (see

Supplementary Figure 1). This was likely caused by the rapid succession of sounds preceding

the onset of the silence, which made it impossible to find a clean, unbiased baseline. Therefore,

we repeated the analysis without baseline correction, to confirm that the results were not caused

by a noisy baseline.

As a control analysis, we repeated the above analysis for the longest intervals during the

presentation of the rhythmic streams, to assess whether in those longer intervals, we could

observe a negative deflection like the one observed in the silence. For this analysis, epochs were

extracted from -200 till 880 ms around the onset of the sound preceding the long interval (600

ms in the beat-based condition, and 780 ms in the other conditions, see Figure 1). All

preprocessing steps were identical to the analysis of the ERPs in the silence. The control

analysis is reported in the Supplementary Materials (Supplementary Figure 1).

Three cluster-based permutation tests (Oostenveld, Fries, Maris, & Schoffelen, 2011)

were used to compare all three conditions against each other (i.e., beat-based vs. random;

pattern-based vs. random; beat-based vs. pattern-based), comparing all timepoints from 300 till

1200 ms after the onset of the final sound for the silence (see Figure 1), and all timepoints from

300 till 600 ms after the onset of the preceding sound for the long intervals during the sequences

(as the next sound came in at 600 ms for the beat-based condition, we could not compare the

conditions beyond this timepoint). This window excluded a large portion of the ERP response

to the previous sound, and included both the first expected moments for beat-based (600 ms)

and pattern-based (780 ms) expectations in the silence window, and additional time to allow

for an evaluation of possible return to baseline of the CNV (Breska & Deouell, 2017). For the

ERP analysis, clusters were formed based on adjacent time-electrode samples.

For all EEG analyses, cluster-based tests were evaluated statistically by forming clusters

of samples based on dependent samples t-tests and a threshold of $p < 0.05$, and using

permutation tests with 2000 permutations of the data. We report corrected p-values to account

for two-sided testing (multiplied by a factor of two).

Frequency-domain analysis. To obtain the spectrum of the EEG signal in the silence,

we used the raw, unfiltered data. Epochs were extracted from the continuous data both from -

1800 till 0 ms relative to the onset of the last sound (control window, see Figure 1), and from

300 till 2100 ms relative to the onset of the last sound (silence window, see Figure 1), the latter

starting at 300 ms to avoid contamination from the final ERPs. Both windows thus had equal

length, both spanning three full cycles of the beat. Epochs with an amplitude change of 250

microvolts or more in a sliding 200 ms window were rejected from further analysis. The more

lenient rejection criterium compared to the ERP analysis was used to account for the fact that

these data were unfiltered, and to avoid rejection of too many trials that showed some slow

drift. All epochs were baseline corrected using the mean of the entire epoch. Subsequently,

epochs were averaged for each condition separately to obtain the evoked signal, phase locked

to the onset of the final sound, and similar to previous studies using frequency tagging to look

at beat-based perception (Lenc et al., 2021; Nozaradan et al., 2011; Nozaradan, Peretz, &

Mouraux, 2012).

For each participant and condition separately, the average waveforms were transformed

into the frequency domain using an FFT, with the data zero-padded to 4608 samples (NFFT) to

obtain a better frequency resolution (0.056 Hz), and importantly, be able to extract data at

exactly the frequencies of interest. Note that the zero-padding can only improve the frequency

resolution, but not the frequency precision, which by definition with the 1800 ms epochs is

limited to 0.56 Hz. While the design of the experiment simply does not allow for a better

resolution, the 0.56 Hz does allow us to differentiate between the frequencies of interest, which

are 0.56 Hz or more apart. The obtained power values at each frequency were normalized to

account for the 1/f distribution of noise (Nozaradan et al., 2011, 2012), by subtracting the

average of neighboring bins four to six on either side for all frequencies (e.g., 1.33 – 1.44 Hz

and 1.89 – 2.00 Hz for the beat frequency, 1.89 – 2.00 Hz and 2.44 – 2.56 Hz for the pattern-

based frequency, and 3.00 – 3.11 Hz and 3.56 – 3.67 Hz for the beat subdivisions). To account

for bleeding into neighboring frequency bins (Nozaradan et al., 2011, 2012), for each

frequency, we averaged over 5 bins centered on that frequency (e.g., for the frequencies of

interest, this was 1.56 – 1.78 Hz for the beat frequency, 2.11 – 2.33 Hz for the pattern-based

frequency, and 3.22 – 3.44 Hz for the beat subdivisions).

To statistically test differences between conditions in the evoked power at the

frequencies of interest, we used cluster-based permutation tests. First, this avoided bias by

selecting only a subset of electrodes, as we used all scalp electrodes, as was done in previous

research (Lenc, Keller, Varlet, & Nozaradan, 2018; Nozaradan et al., 2011; Tal et al., 2017).

Second, the permutation tests accounted for the non-normal distribution of the data. Like for

the ERPs, we ran t-tests comparing the normalized data for all conditions and for each frequency

of interest (e.g., those most prominent in the sound signal). We included the frequencies that

showed the highest peaks in the spectral analysis of the sound (i.e. 1.67 Hz, 2.22 Hz, and 3.33

Hz), except for 3.89 Hz, since a peak at this frequency was absent on visual inspection in the

spectral decomposition of the EEG data. For the frequency-domain analysis, clusters were

formed based on adjacent electrodes.

The cluster-based tests yielded null results for the pattern-based condition in the silence

(e.g., there was no larger power at 2.22 Hz in the pattern-based than in the random condition,

see Results). To quantify the possible absence of persistent entrainment for the pattern-based

condition, we performed a Bayesian t test using JASP (JASP, 2019; Wagenmakers et al., 2018).

We compared the power in the pattern-based and random conditions at 2.22 Hz in the silence,

averaged over electrodes contributing to the significant cluster in the silence for the beat-based

condition at 1.67 Hz, to optimize for finding entrainment effects. We estimated Bayes factors

using a Cauchy prior distribution (r =.71) and performed a robustness check to assess the effect

of a different prior (r = 1; see (Jeffreys, 1961; Wagenmakers et al., 2018)).

As an additional control analysis to assess phase alignment, we computed inter-trial

phase consistency at beat times for each condition. Specifically, for each condition, we

extracted the instantaneous phase in the delta range at times 1200 and 1800 after the start of the

silence window. The rationale here is that while phase consistency may be influenced by

climbing neuronal activity (Breska & Deouell, 2017), if there is indeed phase alignment to the

putative beat, the phase at beat times should be the same. As the ERP analysis showed a CNV-

like component only in the first half of the silence window, here we took the beat times at the

end of the silence window to avoid contamination by possible climbing neuronal activity. Phase

was computed by filtering the continuous signal between 0.5 and 4 Hz, and applying a Hilbert

transform to extract instantaneous phase at beat times in the silence window for each condition

separately. Subsequently, phase coherence was calculated as the length of the mean phase

vector (Cohen, 2014).

Multiscale entropy (MSE). MSE is a measure of signal irregularity. To compute MSE,

the EEG signal is divided into patterns of a certain length, and throughout the signal, the number

of repeating patterns is counted. More repetitions indicate a more regular signal, and yield a

lower entropy value. By calculating entropy for patterns of different lengths ("multiscale"), the

contributions of slower and faster timescales in the signal can be assessed. However, the

mapping between entropy timescales and spectral frequencies is not absolute, especially since

entropy is not per se related to a signal being oscillatory in nature (Kloosterman, Kosciessa,

Lindenberger, Fahrenfort, & Garrett, 2020; Kosciessa et al., 2020). The advantage of using

MSE is that it does not require filtering of the data, and it does not assume stationarity (e.g., it

can pick up on regularities that are asymmetrical, or that do not have a fixed amplitude or

period). Here, we computed MSE on the control and silence epochs separately.

We computed MSE on high-pass filtered data (0.5 Hz). Epochs were extracted identical

to the epochs for the frequency domain analysis. To compute MSE, we used the mMSE toolbox,

a plugin to the Fieldtrip toolbox (Kloosterman et al., 2020), with $m = 2$ and $r = 0.5$, as was

done previously for EEG data (Kloosterman et al., 2020). For details on how MSE is computed,

see the Supplementary Materials. As for the frequency-domain analysis, we used cluster-based

permutation tests to assess statistical significance. For each comparison between conditions, we used paired t-tests comparing each electrode-timescale combination (note that the above computation of entropy yields one value per condition for each electrode and timescale) to form clusters.

Multivariate decoding. Our exploratory decoding approach is based on the assumption that temporal expectations are always coupled with feature or spatial expectations (e.g., we cannot predict "when" without also predicting "what"), as suggested by studies showing that we only use temporal expectations to improve perception if we can also predict the content (Morillon et al., 2016; Wollman & Morillon, 2018) or location (O'Reilly et al., 2008) of an upcoming event. Thus, we expected to be able to decode the representation of the expected sound at expected moments. As the expected moments are different for each condition, this then allows us to decode in the silence window whether participants were previously listening to a beat-based, pattern-based, or random sequence.

The decoding was conducted on data that was preprocessed in a similar way as for the MSE analysis, but with epochs extending from -1800 to 2100 ms relative to the onset of the last sound. Since the decoding is done in a time resolved way (e.g., sample by sample), there is no need to leave out the response to the ERPs in the analysis. Additionally, the data were resampled to 32 Hz to increase signal to noise, using shape-preserving piecewise cubic interpolation, as implemented in the Fieldtrip toolbox (Oostenveld et al., 2011). Using the ADAM toolbox (Fahrenfort, van Driel, van Gaal, & Olivers, 2018), we applied a classification algorithm to the preprocessed data for each participant. Using all electrodes, each dataset was split into 10 equally sized subsets, for 10-fold cross-validation of the decoding. For each subset, a linear discriminant classifier trained on the remaining 9 subsets was tested. Each condition was decoded against both other conditions (e.g., beat-based vs. pattern-based, beat-based vs. random, and pattern-based vs. random), creating a temporal generalization matrix of

classification accuracy at each possible combination of training and testing time points (King

& Dehaene, 2014). Subsequently, we examined whether we could observe a pattern of recurrent

activity (King & Dehaene, 2014). Classification accuracies averaged over the 10 folds for each

comparison of two conditions for the silence window (300-2100 ms after the onset of the last

sound) were submitted to cluster-based permutation tests to assess whether they exceeded the

chance level of 0.5. Clusters were based on T-tests with a threshold of 0.05 for each training-

testing time point combination, comparing the accuracy to 0.5.

The initial decoding analysis yielded large effects that may be task-related (see Results

for an explanation of these results). Therefore, we ran an additional decoding analysis in which

we decoded expected and unexpected positions against each other within each condition.

Details on this additional analysis can be found in the Supplementary Materials.

Musical expertise. In an exploratory analysis, we assessed the relationship between the

EEG data and musical expertise. We divided participants in two groups, based on a median split

on the GMSI questionnaire scores. For each EEG marker, we extracted the relevant values

indexing beat-based and pattern-based expectations, and we performed a t-test, comparing the

two musical expertise groups. First, for the CNV-like component in the ERP, we extracted

average amplitudes for each condition in the two windows in which the cluster-based analysis

yielded a significant cluster (300-450 ms and 925-1005 ms, see also Figure 5), from central

electrodes (FC1, FCz, FC2, C1, Cz, C2, CP1, CPz, CP2). To index the effects of temporal

expectations, we then followed the subtraction logic we also used in the analysis of the

behavioral results: the effects of beat-based expectations were quantified as the amplitude in

the beat-based condition minus that in the random condition, and the effects of pattern-based

expectations were quantified as the amplitude in the pattern-based condition minus that in the

random condition. Likewise, for the frequency-domain analysis, we extracted the difference in

power at 1.67 Hz between the beat-based and random conditions, and the difference in power

at 2.22 Hz between the pattern-based and random conditions. Here, we used only electrodes

that contributed to the significant cluster we had found in the silence for the frequency-domain

analysis. For the MSE, we similarly extracted differences between entropy in the beat-based

and pattern-based conditions when compared to the random condition, using only electrodes

and timescales that contributed to the significant cluster found in the silence for the MSE

analysis. For decoding, we extracted the classification accuracy when decoding expected

against unexpected times for each participant from the beat-based and pattern-based conditions,

and again contrasted those with the random condition (see Supplementary Materials for details

on this decoding analysis).

Code and data availability. All datafiles, and code used for data acquisition, data

analysis, and figure creation, are available through https://osf.io/uwny8/.

 Results

 Behavioral effects of beat-based expectations last multiple beat cycles, while those of

 pattern-based expectations reflect only one interval

Figure 3A shows the average ratings for each condition and probe position from

Experiment 1. Visual inspection of this figure suggests that beat-based expectations were

associated with higher fitness ratings for sounds at expected times than unexpected times for

two beat cycles in the silence window (at 600 and 1200 ms), while the effects of pattern-based

expectations appeared to reflect only the first expected time point in the silence window (780

ms). This was confirmed by our statistical analyses. The ordinal regression showed main effects

of Condition ($\chi^2(2) = 300.72$, $p < 0.001$), Position ($\chi^2(5) = 1067.05$, $p < 0.001$), as well as

Musical Training ($\chi^2(1) = 5.16$, $p = 0.023$). However, crucially, these main effects were

accompanied by a very large two-way interaction between Condition and Position ($\chi^2(10) =$

2478.98, $p < 0.001$), showing that the effects of beat-based and pattern-based expectations on

fitness ratings differed, depending on the position of the probe. We found additional smaller

interactions between Position and Musical Training ($\chi^2(5) = 67.26$, $p < 0.001$), Condition and

Musical Training ($\chi^2(2) = 6.01$, $p = 0.05$), and, interestingly, Condition, Position, and Musical

Training ($\chi^2(10) = 204.88$, $p < 0.001$). Following the interactions, tests of simple main effects

showed that the effect of Position was significant in all conditions (all ps < 0.001), and the

effect of Condition was significant for all probe positions (all ps < 0.001). The main effect of

Position in the random condition showed that even after sequences in which no specific

temporal structure was present, ratings depended on the position of the probe. This likely was

due to recency effects. To account for these effects when comparing the ratings at different

positions in the beat-based and pattern-based conditions, we subtracted the ratings in the

random condition at each position from the ratings in the other conditions. The baseline

corrected model (Figure 3B) showed similar interactions between Position and Condition

$(\chi^2(5) = 2147.55, p < 0.001)$, and between Position, Condition, and Musical Training $(\chi^2(5) = 154.46, p < 0.001)$. The latter indicated that the correlation between Musical Training and the rating score depended on both the Position and the Condition of the probe tone.

Figure 3. The effects of beat-based expectations on fitness ratings outlast those of pattern-based expectations, and are associated with musical training. A) Mean ratings for all conditions and positions. Colored asterixis indicate positions where ratings in the beat-based (orange) and pattern-based (purple) conditions differed from the random condition at $p < 0.05$. B) Single participant data, with the random condition subtracted to account for serial position effects. For the beat-based but not the pattern-based condition, ratings follow the expectedness pattern, indicated by colored lines on the bottom of the plots (orange: expected based on the beat; purple: expected based on pattern; grey: neither). C) Data median split based on scores on the musical training questionnaire. The pattern of results, while present for both groups of participants, is enhanced for the group of participants with most musical training ("experts"). Note: the median split is for visualization purposes only, the models were run with musical training as a covariate. Error bars in panels A-C are 2 standard errors (note: these are computed on the complete dataset, not the participant averages, as the ordinal model is run on trial-level data). D) Association between musical training and rating for each condition and position. Colored asterixis show positions in which the association between musical training and the ratings was significantly correlated ($p < 0.05$). A positive association was observed for the beat-based condition at 600 and 1200 ms (expected positions) and for the pattern-based condition at 780 (expected) and 885 ms (unexpected). Negative associations were observed in the beat-based condition at 780, 885, and 990 ms (all unexpected), and in the pattern-based condition at 990 (unexpected) and 1200 ms (expected).

Beat-based expectations. At both 600 ms and 1200 ms (expected in terms of a beat), probes in the beat-based condition were rated as better fitting than probes in the random condition ($p < 0.001$ and $p = 0.004$) as evident from the full model; all simple effects from the full model can be found in Supplementary Table 1). At 780, 885, 990, and 1485 ms (unexpected in terms of a beat), probes in the beat-condition were rated as worse fitting than probes in the random condition (all ps < 0.001). Moreover, within the beat-based condition, at 600 ms, baseline corrected ratings were higher than at any other probe position (all ps < 0.001), and at 1200 ms baseline corrected ratings were higher than at 780, 885, 990, or 1485 ms (all ps < 0.001). Baseline corrected ratings for probes at 780, 885, and 990 ms (all unexpected in terms of the beat) did not differ from each (all ps > 0.93). Probes at 1485 ms (unexpected in terms of the beat) were rated as better fitting than probes at 780, 885, and 990 ms (all ps < 0.001). All simple effects from the corrected model can be found in Supplementary Table 2.

As can be seen in Figure 3C and 3D, higher scores on the Musical Training questionnaire were associated with higher fitness ratings in the beat-based condition at 600 and 1200 ms (expected in terms of the beat), but lower fitness ratings at 780, 885, 990, and 1485 ms (unexpected in terms of the beat). In other words, musically trained participants were better able to differentiate between probes that were in expected and unexpected positions (Figure 3C). Slopes reach significance at all positions except 1485 ms (all ps < 0.022). Also, the association between Musical Training and ratings differed between beat-based and pattern-based conditions, at 600, 780, 885, and 1200 ms (all ps < 0.004).

To sum up, for the beat-based sequences, we could observe a clear pattern in the results indicating that beat-based expectations were used to rate the probes well into the silence window, affecting ratings up to 1200 ms after the onset of the last sound. Beat-based expectations lead to higher ratings for expected probes (600 and 1200 ms), and lower ratings for unexpected probes (780, 885, 990, and 1485 ms), both when comparing ratings for each

position to the random condition, and when comparing ratings for each position within the beat-

based condition. At 1200 ms, these effects resulted in a classic inverted U-curve, as previously

associated with beat-based processing (Bauer et al., 2015; Jones et al., 2002), with optimal

performance on the beat, and diminished performance on either side (e.g., both earlier and

later). The effects of beat-based expectations did diminish over time, as is apparent from

differences between ratings at 600 and 1200 ms, and at 1485 ms and other unexpected time

points. Both the enhancing and attenuating effects of beat-based expectations were correlated

with musical training. It is worth noting that the longer lasting effects of beat-based expectations

(at 1200 ms) were very heterogenous in our participant pool. Out of 32 participants in

Experiment 1, only 18 showed the inverted U, with higher ratings at 1200 than at 990 and 1485

ms.

Pattern-based expectations. For pattern-based sequences, ratings at 600 ms

(unexpected based on the pattern) were lower than for the random sequences ($p < 0.001$) and

lower than at any other position (all ps < 0.001, baseline corrected model), showing that

participants also formed predictions based on the sequences. In line with this, at 780 ms

(expected in terms of the pattern), ratings were numerically higher in the pattern-based

condition than in the random condition, though this difference did not survive the Bonferroni

correction ($p = 0.058$). After this point, ratings did not differ between pattern-based and random

conditions for probes at 885 (unexpected) and 990 (expected) ms, suggesting that the pattern-

based expectations only exerted an influence on the rating in the beginning of the silent period.

In line with this, the ratings continued to deviate from what would be predicted based on the

pattern, with lower ratings for the pattern-based than random condition at 1200 ms (expected;

$p < 0.001$). Ratings were also lower for the pattern-based than random condition at 1485 ms

(unexpected, $p < 0.001$), and ratings at 780 ms (expected) did not differ from ratings at 885 ms

(unexpected), while being marginally higher than at 990 (expected, $p = 0.080$), and higher than

at 1200 (expected), and 1485 (unexpected) ms (both ps < 0.032). In addition, ratings at 885 and

990 ms were higher than at 1200 and 1485 ms (all ps < 0.002). See Supplementary Tables 1

and 2 for all simple effects.

As for beat-based expectations, for pattern-based expectations, there was a positive

association between ratings and musical training at an expected time point (780 ms; p < 0.001),

and a negative, albeit nonsignificant, association at an unexpected time point (600 ms). Thus,

like for beat-based expectations, at these early time points, musicians were better able than non-

musicians at differentiating between expected and unexpected moments in time. However, at

885 ms (unexpected in terms of the pattern), the results behaved like at 780 ms, with higher

ratings associated with more Musical Training (p = 0.002). At 990 and 1200 ms, Musical

Training was associated with lower ratings (both ps < 0.05), but these results are

counterintuitive, as these are expected positions based on the pattern.

The results for pattern-based expectations suggest that just like for beat-based

expectations, participants were able to predict the timing of probes based on the preceding

sequence. However, the results show that while this was still the case at 780 ms after the onset

of the last tone, at later probe positions, the effects of pattern-based expectations did not reflect

the preceding sequence. At 885 ms, the results, both in terms of the ratings and how they were

associated with musical training, behaved similar to at 780 ms. This may indicate that the

pattern-based expectations at this point were not precise enough to differentiate between

positions, even when the deviation in time is half the learned interval (105 of 210 ms). After

this point, the results suggest that participants did not correctly represent the preceding

sequence anymore.

Figure 4 shows the behavioral results obtained from the EEG experiment. Replicating

Experiment 1, we found main effects of Condition ($\chi^2(2)$ = 92.30, p < 0.001) and Position

($\chi^2(2)$ = 49.36, p < 0.001), accompanied by interactions between Condition and Position ($\chi^2(4)$

= 96.62, $p < 0.001$), and Condition, Position, and Musical Training ($\chi^2(4) = 11.04$, $p = 0.03$).

Following the analysis strategy from Experiment 1, we subtracted the ratings from the Random

condition from the ratings for the other two conditions, yielding a baseline corrected model

with similar interactions (Condition and Position: $\chi^2(2) = 31.46$, $p < 0.001$; Condition, Position,

and Musical Training: $\chi^2(2) = 8.29$, $p = 0.02$).

Figure 4. Behavioral results during the EEG experiment replicate findings from Experiment 1. Even
with only 18 trials per participant per condition and position, we could replicate the inverted U-curve for
beat-based sequences at the end of the silence epoch. Like in Experiment 1, the results for the pattern-based
condition do not follow the pattern, but instead, are consistent with building expectations for the next trial.
A) Mean ratings for all conditions and positions. Colored asterixis indicate positions where ratings in the
beat-based (orange) and pattern-based (purple) conditions differed from the random condition (with $p <$
0.05). B) Single participant data, with the random condition subtracted to account for serial position effects.
C) Data median split based on scores on the musical training questionnaire. Error bars in panels A-C are 2
standard errors. D) Association between musical training and rating for each condition and position. See
Figure 3 for more details.

In line with the preceding beat-based sequences, probes at 990 and 1485 ms (both

unexpected times based on the beat) were rated lower in the beat-based condition than in the

random and pattern-based conditions (all ps < 0.012), and within the beat-based condition,

probes at 1200 ms were rated as better fitting than at 990 ($p < 0.001$) and 1485 ms, though the

latter difference did not reach significance. Thus, like in Experiment 1, we found an inverted U-curve at 1200 ms after the final tone, suggestive of beat-based expectations lasting at least two beat cycles. As in Experiment 1, at 1200 ms (expected based on the beat), higher ratings in the beat-based condition were associated with more musical training ($p = 0.047$), suggesting that the effects of beat-based expectations correlate with musical expertise. Additionally, probes at 990 ms were rated lower than at 1485 ms ($p < 0.001$), possibly because the effects of beat-based expectations diminished over the course of the silence.

In the pattern-based condition, ratings did not follow the pattern of the preceding sequences. At 990 ms (expected based on the pattern), probes were rated as worse fitting than in the random condition ($p < 0.001$), and as worse fitting than at 1200 (expected) and 1485 (unexpected) ms (both $ps < 0.023$), while at 1485 ms (unexpected), probes were rated as better fitting than in the random condition, and as better fitting than at 1200 (expected) ms (both $ps < 0.001$). Also, at 1200 ms (expected), higher ratings were associated with less musical training for the pattern-based condition (though after the Bonferroni correction, only marginally so: $p = 0.06$), contrary to what would be expected if the effects of expectations are enlarged in musical experts. All simple effects can be found in Supplementary Tables 1 and 2.

Thus, the behavioral results from Experiment 2, though based on less trials than Experiment 1, suggest a similar pattern as found in Experiment 1: while beat-based expectations exert their effect well into the silent period, with participant faithfully following the beat in their goodness-of-fit ratings, pattern-based expectations fail to affect ratings in the second half of the silent period in a manner consistent with the learned pattern. Albeit speculatively, the results for the pattern-based expectations may be more in line with expectations for the start of the next trial leading to higher ratings for probe positions closer to the end of the silent period, as in Experiment 2, after non-probe trials, the next trial followed each silent period at a somewhat predictable time.

Negative deflection in the evoked potential elicited by pattern-based, but not beat-based sequences

Figure 5 shows the average ERPs for each condition, without baseline correction (see Materials and Methods), and scalp topographies for windows in which we found significant clusters. In line with previous research, we may expect climbing neuronal activity, or a CNV, flexibly adapting its slope to peak at the moment that participants expect the next event (Breska & Deouell, 2017; Breska & Ivry, 2020; Damsma, Schlichting, & van Rijn, 2021; Mento, 2013). A CNV-like frontocentral negativity was indeed observed early in the silence window in the pattern-based and random conditions, but not the beat-based condition (see Supplementary Figure 1 for the difference waveforms between conditions). Without baseline correction, the random condition elicited a significantly more negative deflection ($p = 0.011$) than the beat-based condition in a frontocentral cluster between 300 and 614 ms after the onset of the last sound (though note that we did not include timepoints preceding 300 ms in the cluster-based tests). Likewise, in the same latency range (300 – 473 ms), there was a trend for the pattern-based condition to elicit a larger negative deflection than the beat-based condition ($p = 0.077$). Thus, in a window between approximately 300 and 450 ms, we found tentative evidence for more negative-going waveforms in both the random and pattern-based condition compared to the beat-based condition, with a central scalp topography, as is commonly found for a CNV (Figure 5B). In a later window, the pattern-based condition elicited a second negative deflection, which showed a trend to be larger than in the beat-based condition ($p = 0.077$, 864 – 1005 ms). While these results were somewhat different depending on the choice of baseline (see Supplementary Figure 1 for the results with a traditional pre-cue baseline), the overall picture is the same, with significant clusters in an early and later window, and negative deflections with a frontocentral scalp distribution reminiscent of a CNV for the pattern-based and random but not the beat-based condition.

While the negative deflection observed could be interpreted as something akin to

climbing neuronal activity or a CNV, the time course of the effect deviated from what we

expected: the negative deflection did not peak at the next expected moment in time, but rather,

peaked much earlier. Also, contrary to previous research, we did not find a CNV-like

component for beat-based expectations. To further explore and confirm these results, as a

control analysis, we performed the same ERP analysis on the longest time intervals during the

sound presentation. The waveforms showed a negative deflection very similar in morphology

and scalp distribution to the one we found in the silence window (see Supplementary Figure 1).

During the sound presentation, like in the silence, this negative deflection was largest for the

pattern-based condition, though the difference was only significant when comparing the

pattern-based condition to the random ($p = 0.039$) but not the beat-based condition ($p = 0.1$).

The latter non-significant result may be due to a noisy baseline, as during sound presentation,

the succession of intervals in the beat-based and random sequences was (semi-)randomly

chosen, and therefore, the baseline was not consistent over conditions. Overall, however, this

control analysis yielded very similar results to the analysis of ERPs in the silence window.

As an exploratory analysis, since musical training was related to larger effects of

entrainment in behavior, we also examined whether there was a relationship between the

amplitude of the CNV-like negative deflection in the silence window (extracted from the time-

electrode clusters depicted in Figure 5, see Materials and Methods for details) and musical

expertise. However, we found no difference in the effects of expectations on the amplitude of

the CNV-like component between musically trained and untrained subjects (all ps > 0.3).

Figure 5. A CNV-like component is present for pattern-based and random conditions, but not for the
beat-based condition. A) Left panel show the grand average waveforms for the silence window for a central
electrode cluster (FC1, FCz, FC2, C1, Cz, C2, CP1, CPz, CP2). Time 0 is the onset of the last tone of the
sequence. Colored bars on the bottom of the plots, and vertical orange and purple lines, indicate at which
times a tone would be expected based on the beat (light orange) and memory (light purple). Note that these
are expected times, but no sounds were played during the window shown after time 0. Sounds during the
sequence are depicted in dark orange and purple. B) The scalp distributions for windows in which a
significant cluster was observed.

Frequency-domain analysis shows persistent power at the beat frequency following beat-

based sequences

A strong prediction of entrainment theories is that the entrainment outlasts stimulation

with the entraining stimulus. Therefore, we next looked at the frequency content of the EEG

signal in the silent period. Specifically, we predicted that if entrainment occurs, we would find

enhanced power at the frequencies associated with the rhythmic sounds to be found in the

silence. In Figure 6, the average power for all electrodes, separated for each condition in the

control (i.e. during the auditory sequence) and silence windows is depicted as a function of

frequency. In the control window (Figure 6, left, and Figure 7, top), the frequency response

followed the sound input. That is, at the beat frequency (1.67 Hz), significant clusters indicated

higher power in the control window for the beat-based sequences than the pattern-based ($p <$

0.001) and random ($p = 0.006$) sequences. At 2.22 Hz, prominent in the pattern-based and

random sound sequences, higher power was observed in the EEG signal in the pattern-based

than beat-based ($p = 0.02$) and random ($p = 0.023$) sequences, and higher power was observed

in the random than beat-based sequences ($p = 0.035$). Finally, at 3.33 Hz (subdivisions of the

beat), power in the control window was larger for the beat-based than pattern-based ($p = 0.013$)

and random ($p = 0.005$) conditions. Thus, in the control window, the EEG signal reflected the

spectral properties of the sound signal, as can be expected, since each sound will have elicited

an ERP, which are represented in steady-state potentials, and thus picked up by the frequency

analysis (Keitel, Obleser, Jessen, & Henry, 2021).

Importantly, during the silence (Figure 6, right, and Figure 7, bottom), no significant

clusters were found at 2.22 Hz, but at the beat frequency (1.67 Hz), power was significantly

larger for the beat-based than the pattern-based ($p = 0.01$) and random ($p = 0.012$) conditions.

In addition, at 3.33 Hz, power for the beat-based condition was larger than for the pattern-based

condition ($p = 0.038$), with a trend when comparing the beat-based with the random condition

($p = 0.077$). Thus, while the pattern-based and random conditions showed tracking of the sound

during stimulation (which is sometimes considered entrainment "in the broad sense" (Obleser

& Kayser, 2019)), in the silence, entrainment was only present for the beat-based condition.

This finding fits our behavioral observations that beat-based expectations persisted longer in

the silence than pattern-based expectations. To further substantiate the absence of persistent

entrainment at the pattern frequency in the silence, we performed a Bayesian T-test comparing

the normalized power at 2.22 Hz between the pattern-based and random conditions. We found

moderate evidence in favor of the null hypothesis (no difference between conditions) ($BF_{01} =$

4.5). The results did not change as a function of the prior used (with a more traditional prior of

$r = 1$, $BF_{01} = 6.17$).

The follow-up analysis to look at the actual phase angles at beat times showed more

phase consistency at beat times (1200 ms and 1800 ms after the onset of the last sound) for the

beat-based condition than for the random condition ($p = 0.015$, see Supplementary Figure 2).

The difference between the beat-based and the pattern-based condition, albeit numerically in

the expected direction, did not reach significance, nor did the difference between the pattern-

based and random condition (both *p*s > 0.15). The absence of a difference between the beat-

based and pattern-based condition here may simply be a lack of power. For this additional

analysis of the phase angles, we only considered data points in the last half of the silence

window, by which time, behaviorally, participants showed weaker effects of the beat than

earlier in the silence window. Hence, the effects may also at the neural level be less pronounced.

Figure 6. Oscillatory power at the beat frequency persists during the silence window. In the control
window, during auditory stimulation, peaks can be observed at all frequencies of interest, and for the relevant
conditions (1.67 and 3.33 Hz in the beat-based condition, and 2.22 Hz in the pattern-based condition). In the
silence window, the peaks at 1.67 and 3.33 Hz are larger in the beat-based condition than in the pattern-based
and random conditions, while peaks at 2.22 Hz do not differ between conditions. This suggests that only
beat-based expectations persisted in the silence window. Note: the raw data is depicted here, before the
normalization procedure.

Note that like for the behavioral results indicative of entrainment, there was large

heterogeneity between participants (see Figure 7). While the power differences were significant

in the overall cluster-based analyses, out of 27 participants, only 16 showed on average (i.e.

over all electrodes) numerically larger power in the beat-based condition at the beat frequency

when compared to both the random and pattern-based condition. As for the CNV-like activity,

musical training did not affect the difference in power between the beat-based and random

condition at 1.67 Hz, nor the difference in power between the pattern-based and random

condition at 2.22 Hz (both *p*s > 0.54).

Figure 7. Individual participant data shows heterogeneity of the effects of temporal expectations on spectral power. All plots depict the spectral power at the frequency of interest, averaged over all electrodes, and normalized to account for the 1/f distribution (see Materials and Methods). A) Single participant data, with all data points in grey. Boxplots show the median, with the lower and upper hinges corresponding to the first and third quartiles (25th and 75th percentiles), and the whiskers corresponding to values no further than 1.5 times the inter-quartile range from the hinges. Error bars depict 2 standard errors around the mean. B) Scalp topographies for all conditions. Electrodes contributing to significant clusters in the cluster-based tests are highlighted on the plots for the conditions from each comparison in which power was largest.

Multiscale entropy as a non-stationary measure of temporal expectations

Figure 8 shows sample entropy for each condition separately, as well as the electrodes

contributing to significant clusters in the analysis. Given that MSE indexes signal irregularity,

we would expect entropy to be higher for the random and pattern-based conditions than for the

beat-based condition in the silence. A cluster-based test on all electrodes and timescales showed

that both in the control window, and in the silence window, entropy was higher for the pattern-

based than for the random condition (control: $p = 0.03$; silence: $p = 0.021$). Note that in the

silence window, all but the two highest timescales were included in the cluster. In the control

window, the cluster spanned all timescales from 35 till 406 ms (see Materials and Methods for

an explanation of the timescales). These results suggest that the signal was more irregular in

the pattern-based than the random condition, over a broad range of timescales. Neither the beat-

based condition compared to pattern-based nor the beat-based compared to the random

condition reached significant differences in entropy, in either control or silence windows (all

ps > 0.24).

Entropy has been related to various other EEG measures, such as spectral power and

overall differences in signal variability (Kosciessa et al., 2020). To account for these, we ran

several additional analyses. First, to check whether the differences in signal variability may

have been caused by differences in low frequency activity, we repeated the MSE analysis on

high-pass filtered data (Kloosterman et al., 2020), using a 5 Hz high-pass filter. This completely

removed the effects (all ps > 0.43), suggesting that the differences between conditions were

caused by low frequency activity in the signal (see Supplementary Figure 3).

Second, we checked whether differences in signal variability caused the differences in

entropy, since entropy is calculated relative to the overall signal standard deviation (e.g., a

pattern is considered a match at a lower threshold when overall signal variability is high). As

can be seen in Supplementary Figure 4, the similarity bounds used to compute entropy (derived

from the time-domain signal standard deviation) differed between conditions, and this difference mirrored the differences in entropy, suggesting that at least some of the variance we observed was due to overall signal variability, and not necessarily signal irregularity. Finally, musical training did not affect the difference in sample entropy between the beat-based and random condition, nor the difference in entropy between the pattern-based and random condition (both ps > 0.66).

Figure 8. Entropy was higher for pattern-based than random sequences in both control and silence windows. Entropy in the Control window (left) and Silence window (right), averaged over frontocentral electrodes, and scalp distributions averaged over all timescales, depicting electrodes contributing to significant clusters.

Multivariate decoding as a time-resolved method for studying entrainment

With multivariate decoding, we expected that training at expected times would yield above chance performance when testing at expected times, regardless of whether these time points were the same (e.g., when training at 600 ms, we expected to be able to accurately distinguish the beat-based from the random condition when testing at not just 600 ms, but also at 1200 and 1800 ms, as all these times were on the beat, or similarly expected). Figure 9A shows the temporal generalization matrices for each comparison, with significant clusters indicated by a black contour. In the silence window we found above chance decoding when decoding the beat-based against the random condition ($p = 0.043$), the pattern-based against the random condition ($p < 0.001$), and the beat-based and pattern-based conditions against each

other (p = 0.009), indicating that based on the EEG signal in the silence, we could classify

which type of rhythm participants had heard just before. However, looking at the temporal

generalization matrices, it becomes apparent that contrary to our expectations, this above-

chance decoding was not due to recurrent activity for expected events. Only clusters on the

diagonal were significant for each comparison. In addition, decoding for all three comparisons

was best in the second halve of the silence window, which is where the probe tones were

presented. This suggests that the decoding mainly picked up on task-related differences. While

the probes were physically identical across conditions, participants may have had different

strategies to perform the task, depending on the type of sequence, and this could have resulted

in above-chance decoding related to the task and probes, even when only analyzing the silence

window.

Figure 9. No recurrent activation in group average decoding, example of recurrency in single participant. A) Temporal generalization matrices for each comparison for the group analysis. The Y-axis shows the training time points, and the X-axis testing time points. The color scale indicates the classification accuracy, with 0.5 being chance level (all analyses are based on decoding two conditions against each other). Significance of classification accuracy was assessed by using cluster-based permutation tests on the silence window (300-2100 ms after the last note onset). The black contour indicates significant clusters. Note that the upper right quadrant, highlighted by the black box, is the entire silence window. B) Temporal generalization matrix for a participant showing recurrent activity following the expected beats. When decoding the beat-based vs. the other two conditions (left- and rightmost plots), peaks in accuracy follow a clear oscillatory pattern, with a phase consistent with the beat-based sequence (600 ms between beats and decoding peaks).

Of note, the decoding results varied considerably between participants. In one

participant in particular (Figure 9B), we observed a pattern in the temporal generalization

matrix that was consistent with the hypothesized result for an oscillatory process (King &

Dehaene, 2014). However, even though some other participants also showed recurrent activity,

the exact times of most accurate decoding, and the exact period of the recurrent process, differed

widely between participants, obscuring these effects in the grand averages. These individual

differences may be caused by individual preferences for a level of regularity in the beat-based

stimuli (Drake, Jones, & Baruch, 2000), with some people attending mostly to the beat level

(1.67 Hz), but others possibly attending to subdivisions (3.33 Hz) or the level of the meter (0.83

Hz). Also, for different people, the optimal phase of delta oscillations (i.e., the phase that aligns

with expected moments) may differ (Breska & Deouell, 2017; Henry & Obleser, 2012),

possibly causing optimal decoding at different time points. To circumvent the large task-related

effects apparent in the decoding results, and the possible individual differences in phase and

metrical level attended to, we ran an additional exploratory decoding analysis looking at

decoding within instead of between conditions. Here, we found above-chance decoding of beat

positions against offbeat positions only following the beat-based condition. However, while

decoding was above chance in the beat-based condition, it was in fact not better than in the

other conditions. As such, these results provide only weak support for persistent effects of beat-

based, but not pattern-based expectations. Detailed results of the within-condition decoding

analysis can be found in the Supplementary Materials (Supplementary Figure 5).

Discussion

In the current study, we aimed to identify and directly compare the neural mechanisms

underlying temporal expectations based on a regular beat, as well as temporal expectations

based on learning a predictable pattern, by examining the development of climbing activity and

the persistence of neural entrainment after cessation of rhythmic input with either a regular beat,

a predictable pattern of temporal intervals, or random timing. Instead of relying on isochrony

to elicit beat-based expectations, which can also elicit different forms of memory-based

expectations, we used varying non-isochronous patterns to clearly separate beat-based and

pattern-based expectations. Moreover, we assessed responses in a silent period after the

auditory input ceased, side stepping the many possible confounds associated with acoustic

differences between conditions (Capilla, Pazo-Alvarez, Darriba, Campo, & Gross, 2011;

Haegens & Zion Golumbic, 2018; Novembre & Iannetti, 2018; Zoefel et al., 2018).

We found several indicators of separate mechanisms for beat-based and pattern-based

expectations. First, behaviorally, we found that while the effects of beat-based expectations

spanned at least two beat cycles, the effects of pattern-based expectations only reflected the

first expected moment in time. Second, while pattern-based expectations elicited a CNV-like

component in the evoked potential, this negative deflection was absent or smaller for beat-based

expectations. Third, we observed significantly more power at the beat frequency in the silence

window when participants were previously listening to a beat-based sequence as compared to

a pattern-based or random sequence. The observed difference between the behavioral effects of

beat-based and pattern-based expectations and increased oscillatory power at the beat frequency

following beat-based but not pattern-based sequences both point towards entrainment

underlying beat-based, but not pattern-based expectations. The time-domain results showing a

CNV-like component for pattern-based, but not beat-based sequences suggests that the former

may rely on a mechanism more like climbing neuronal activity. Together, these findings

provide support for the notion that beat-based and pattern-based expectations rely on different

neural mechanisms, as has previously been suggested for cue-based expectations (Breska &

Deouell, 2017; Breska & Ivry, 2020), but our results do not support models that assume shared

mechanisms, be it entrainment (Tichko & Large, 2019), or a general top-down mechanism for

temporal expectations (Rimmele et al., 2018).

Behaviorally, for the beat-based condition, we observed a pattern clearly in line with

entrainment models, with an inverted U-curve, an indication of entrainment (Bauer et al., 2015;

Jones et al., 2002), present as late as 1200 ms after the final tone of each sequence in both

experiments. Beat-based expectations thus affected the fitness ratings for at least two beat

cycles after the end of the sequences, as predicted by nonlinear oscillator models, that assume

oscillations are self-sustaining (Large, 2008; Large & Palmer, 2002). Expectations in the beat-

based condition not only led to higher ratings for expected events, but also to lower ratings for

unexpected events, when compared to the random and pattern-based conditions. Suppression

of unexpected events may be metabolically beneficial (van Atteveldt et al., 2015), and as such,

has been suggested to be a hallmark of entrainment and the associated "rhythmic" mode of

processing (Schroeder & Lakatos, 2009a; Zoefel & Vanrullen, 2017). Indeed, suppression off

the beat has even been proposed to be a better indication of beat-based expectations than

facilitation on the beat (Bouwer et al., 2020; Breska & Deouell, 2017), in line with the current

results, where the effects of beat-based expectations at unexpected time points exceeded those

at expected time points.

Pattern-based expectations similarly affected fitness ratings, with enhanced ratings at

the first expected time point, and lower ratings at the first unexpected time point, showing that

participants did form expectations based on the predictable pattern. However, importantly, for

the pattern-based condition, the results are not suggestive of sustained oscillations, as the effects

of expectations only reflected the first expected moment, but not the subsequent structure of

the pattern. We did observe lower ratings for the last probe positions after pattern-based than

random sequences in Experiment 1. One speculative explanation for this is that while listeners

are able to form expectations following temporal patterns, they only do so one interval at a time

(e.g., they use each event as a cue for the next interval, but in the absence of an event, no next

interval is predicted), and in a probabilistic way (Cannon, 2021; Damsma et al., 2021; van der

Weij, Pearce, & Honing, 2017). In this case, it is possible that the expectation for an event at

780 ms would lead to an inverted U-shape in responses, similar to the inverted U around a beat,

but with a wider distribution. This could explain the ratings being equal for probes at 780 and

885 ms, as they would still fall within the time window where a tone could be expected, while

probes at 1200 and 1485 ms were considered as unexpected, as they fell far from the expected

time.

Crucial to entrainment models of beat-based expectations (Haegens & Zion Golumbic,

2018; Henry & Herrmann, 2014; Large, 2008; Large & Jones, 1999; Obleser & Kayser, 2019),

we found that power at the beat frequency (1.67 Hz) and its harmonic (3.33 Hz) in the EEG

signal during the silence window was larger following beat-based than pattern-based or random

sequences. Such enhanced power was not found for a frequency inherent to the pattern-based

sequence (2.22 Hz). Methodologically, measuring phase locking during rhythmic stimulation

can lead to confounding contributions from tone-evoked responses (Capilla et al., 2011;

Novembre & Iannetti, 2018; Zoefel et al., 2018). Ongoing oscillations in silence, after sensory

input has stopped, are therefore regarded as strong evidence for entrainment (Breska & Deouell,

2017; Haegens & Zion Golumbic, 2018; Obleser & Kayser, 2019; van Bree et al., 2021; Zoefel

et al., 2018). As such, our observation of enhanced power at the beat frequency during silence

provides important novel support for the notion that entrainment of low-frequency neural

oscillations underlies beat-based perception, and our design, using non-isochronous rhythms,

allows us to separate beat-based aspects from other structure present in natural rhythm (Bouwer,

Nityananda, Rouse, & ten Cate, 2021).

Interestingly, a recent paper that assessed ongoing oscillations following auditory

rhythmic input did not find any evidence for persistent entrainment at the frequency of the

rhythm (Pesnot Lerousseau et al., 2021). Two differences between this study and our work may

provide directions for future work. First, as mentioned before, isochronous rhythm allows for

several different ways of forming temporal expectations, including not only beat-based, but

also cue-based and pattern-based expectations. In the current study, this was the motivation to

design stimuli that allowed for differentiating between these types of expectations. However,

an additional concern could be that when presented with stimuli that do not require beat-based

expectations to perform the task of tracking the temporal structure, the brain may not engage in

forming such expectations. Thus, in addition to variation between individuals (Assaneo et al.,

2019), variation in the input signal may also determine which mechanism is used to form

temporal expectations.

Second, the presence of persistent entrainment may also depend on task demands

(Shalev, Nobre, & van Ede, 2019). In our study, the auditory sequences were task relevant, and

the task itself was rhythm-related. In the study by Pesnot Lerousseau et al. (2021), participants

listened to rhythms passively. This raises the possibility that persistent entrainment is the result

of explicit, top-down guided expectations, rather than being the result of some passive,

automatic process (Bouwer, 2022). Interestingly, in our study, the scalp topographies for power

at 1.67 Hz differed between the silence and control windows, with power in the control window

largest above a frontocentral region, and in the silence window above a parieto-central region.

This raises the question to what extent the phase locking as measured during the sound

presentation has the same source as phase locking during the silence. During sound

presentation, oscillations picked up at the scalp likely contain large contributions from evoked

responses in auditory cortex that are phase locked to the input ("entrainment in the broad sense",

see (Obleser & Kayser, 2019). Possibly, instead of resulting from a sustained automatic

oscillation, persistent entrainment could originate from other sources, such as explicit

predictions made by a motor network (Rimmele et al., 2018). The influence of contextual

factors on entrainment, be it person, stimulus, or task, and the source of phase-locked activity,

be it automatic phase alignment in sensory cortices, or active top-down expectations, are

important topics for future research.

The absence of power at 2.22 Hz following the pattern-based sequences suggests that

entrainment does not underlie expectations based on learning a pattern, contrary to a recently

proposed oscillator model that can capture aspects of pattern-based expectations (Tichko &

Large, 2019). In the time-domain, we found a negative deflection in the silence window

following the pattern-based and random sequences, but not the beat-based sequences. The

observed differences in the ERP suggest that rather than one entrainment mechanism for both

regular and irregular rhythms, an alternative mechanism, possibly based on climbing neuronal

activity, specifically supports formation of pattern-based expectations. Contrary to this notion,

previous studies did find a CNV peaking at expected times not only for cue-based, but also for

beat-based expectations (Breska & Deouell, 2012, 2017; Breska & Ivry, 2020; Praamstra et al.,

2006). Importantly, the latter studies used isochronous rhythmic sequences to probe beat-

expectations, and therefore, memory-based expectations, be it pattern-based or cue-based,

could also have been formed in response to these sequences (Bouwer et al., 2020, 2021;

Bouwer, Werner, Knetemann, & Honing, 2016; Breska & Ivry, 2016; Keele et al., 1989). Here,

using a beat-based sequence that did not allow for expectations based on simply learning

transitional probabilities, we did not observe the same negative deflection as in the pattern-

based sequences. This raises the possibility that a CNV (or CNV-like) component is specific to

pattern-based, and likely cue-based (Mento, 2013, 2017), temporal expectations.

While previous studies showed that the peak of the CNV coincided with an expected

time (Breska & Deouell, 2017; Mento, 2017; Praamstra et al., 2006), here, the peak latency of

the negative deflection in the signal was earlier, peaking around 400 ms for the pattern-based

condition, while the first expected time point in the silence was at 780 ms. The negative

deflection in the current experiment may index temporal expectations in a probabilistic way.

While the pattern was fully predictable in the pattern-based sequences, it contained several

temporal intervals with durations between 150 and 780 ms. The peak at 400 ms may have

indexed the average interval presented in the sequence (~360 ms). This explanation is supported

by the presence of a deflection with a similar time course for the random condition. Like in the

pattern-based condition, in the random condition, participants could not use a beat-based

strategy to perform the task. Thus, they may have attempted to predict the timing of an

upcoming sound based on the distribution of the absolute intervals, which while random in

terms of transitional probabilities, was on average identical to the pattern-based condition.

Interestingly, the average interval in the beat-based condition was also 360 ms, but here, the

same component was less present, suggesting that in the presence of a possible beat-based

strategy, the brain may operate in a rhythmic mode of processing (Rimmele et al., 2018;

Schroeder & Lakatos, 2009a). To what extent the negative component we observed is related

to the CNV as found in previous studies remains a topic for future studies. Of note, the CNV

has indeed been shown to be susceptible to its probabilistic context (Damsma et al., 2021).

Probabilistic models incorporating statistical regularities in inter-onset intervals at different

levels have indeed been used to explain aspects of temporal processing (Cannon, 2021; Elliott,

Wing, & Welchman, 2014; van der Weij et al., 2017). In future work, linking such models

directly to neural markers of pattern-based expectations may provide more insight in the

mechanisms underlying pattern-based expectations.

Various other challenges for future research remain. First, in the beat-based condition,

it could be argued that participants used an interval-based strategy to perform the task, in which

they predicted an event every 600 ms. However, listening to strictly metric patterns, as the ones

used here, is associated with activity in a circuit including the basal ganglia, while listening to

non-metric patterns is associated with activity in a circuit including the cerebellum, making it

unlikely that the same, interval-based mechanism would be used for both types of rhythms

(Leow & Grahn, 2014). Also, predicting the timing of events in non-isochronous strictly metric

sequences would require participants to learn not just the transitional probabilities of single

intervals, but also to combine multiple intervals into groups that together last the length of a

beat. It is currently unclear whether humans, when faced with rhythmic patterns, use such a

hierarchical interval-based strategy. Indeed, future research could examine if beat-based

expectations in general can be explained by such multilevel interval learning, akin to a recent

model for beat-based perception (Cannon & Patel, 2021), as this could provide a general

challenge to oscillator models of beat-based perception. In such a view, the difference between

beat-based and pattern-based timing may be the importance of hierarchical structure in beat-

based rhythms (Fitch, 2013), rather than the presence of oscillations.

A second challenge concerns the result from the frequency-domain analysis. The

Fourier transform assumes stationarity in the oscillating signal, while entrainment models

propose a dampening factor to account for decreasing oscillatory power over time (Large,

Herrera, & Velasco, 2015). To assess power at specific frequencies in a time-resolved way,

wavelet convolution is often used as an alternative. But, in the current study, differentiating

between the specific frequencies of the beat and the pattern would require wavelet parameters

that would result in a temporal resolution too low to disentangle activity during and before the

silence (i.e., many wavelet cycles would be needed). Recently, a promising alternative to assess

oscillatory activity in the time domain was proposed in the form of cycle-by-cycle analysis

(Cole & Voytek, 2019). However, this approach requires filtering the signal at the specific

frequency of interest, again posing problems for disentangling low frequency oscillations due

to the beat, the pattern, and ongoing ERPs. Assessing the time course of low frequency

oscillations thus remains a challenge for future research.

Multi scale entropy and multivariate pattern analysis may provide alternative ways to

examine neural entrainment with high temporal precision. At the group level, entropy was

higher following the pattern-based than random sequences. This could be explained by

assuming that the brain uses a vigilance mode to track the pattern-based regularities, which at

the neural level, translates in more irregular patterns of activity. Attention and arousal, which

are both associated with temporal expectations (Schroeder & Lakatos, 2009b), have indeed

been linked to neural variability as well (Waschke et al., 2021). However, the group level results

can be explained at least to some extent by overall differences in signal variability (e.g., the

signal variance), so this hypothesis remains to be confirmed in future research. Considering the

decoding results, the observed above-chance decoding seemed to primarily reflect general task-

related activity. The best decoding was observed in the second half of the silence window,

where probes could be presented. Also, in the group-average decoding results, above chance

decoding was limited to training and testing on the same time points. In other words, we did

not observe recurrent activity, as reflected in stronger decoding accuracy at expected time

points. Yet, we did show a proof of concept for our approach in at least one participant, who

showed a clear oscillatory pattern in decoding accuracy when decoding the beat-based against

the other two conditions. This pattern of activity is in line with the strength or sharpness of the

neural representation of tones varying over time as a function of temporal expectations

(Auksztulewicz et al., 2019, 2018).

At the group level, the behavioral results, along with the results obtained using

frequency-domain analysis and ERPs provide evidence for differential processing of the beat

and the pattern in rhythm, while the results from the decoding and entropy analyses are less

clear. One reason for this is heterogeneity between individuals. This heterogeneity may be

present at several levels. First, people may attend to different levels of regularity in beat-based

perception (Drake et al., 2000), and the optimal phase of delta oscillations (e.g., the phase that

aligns with expected moments) may differ across individuals (Breska & Deouell, 2017; Henry

& Obleser, 2012; Sun et al., 2021). Second, while it is often assumed that most people

automatically form beat-based expectations (Honing, 2012), recent evidence showed phase

locking to speech in only about half of the population (Assaneo et al., 2019). Indeed, in our

study, only about two-thirds of the participants behaviorally showed evidence for beat-based

expectations in the second half of the silence window and we only observed enhanced power

at the beat frequency following beat-based sequences in about half of the participants.

In the current study, the behavioral effects of beat-based and pattern-based expectations

were associated with musical training, consistent with previous research using beat-based

(Bouwer et al., 2018, 2016; Cameron & Grahn, 2014; Matthews, Thibodeau, Gunther, &

Penhune, 2016; Vuust et al., 2005) and pattern-based (Cameron & Grahn, 2014) rhythms. Some

previous studies have failed to show differences between musicians and non-musicians,

however (Bouwer, Van Zuijen, & Honing, 2014; Geiser et al., 2009; Grahn & Brett, 2007),

possibly due to differences in task design (Bouwer et al., 2018). The current study used an

explicit rating task, for which performance may be particularly improved by musical training,

as musically trained participants may have additional strategies to perform the task. The use of

implicit timing tasks may be a better probe of innate differences in timing abilities, which need

not necessarily be related to musical training (Law & Zentner, 2012), and implicit tasks may

also be less susceptible to task-related effects as we observed in the decoding analysis, which

may stem from individual strategies in performing the explicit task.

We did not find an association between musical expertise and the neural results. This

may be due to neural activity being more reflective of innate abilities, as argued above. Also,

our sample scored relatively low on the musical training scale, with a mean of 20.5 on the GMSI

subscale, which is comparable to the 32^{nd} percentile of the norm scores, which have a mean of

26.5 (Müllensiefen et al., 2014). This may have made it hard to find the effects of musical

training. However, with the current sample size, we may have simply lacked power to detect

associations between neural data and musical expertise. Undoubtedly, given the heterogeneity

we and others (Assaneo et al., 2019; Bauer et al., 2015; Sun et al., 2021) have observed in tasks

probing temporal expectations, understanding individual differences is an important direction

for future research, with significant implications for applications of musical rhythm, such as in

motor rehabilitation (Dalla Bella, Dotov, Bardy, & Cochen De Cock, 2018).

Conclusion

In summary, we have shown that beat-based and pattern-based expectations can be

differentiated in terms of their behavioral and neurophysiological effects once sensory input

has ceased. These findings provide novel, more conclusive evidence for the notion that different

mechanisms implement temporal expectations based on periodic and aperiodic input streams,

with the former based on entrainment of low frequency neural oscillations, and the latter on

climbing neural activity indexing a memorized interval.

Acknowledgements

1202   We would like to thank Babak Chawoush and Charlotte van Roijen for their assistance

1203   with the data collection, and fruitful discussions during the project.

References

Arnal, L. H., Doelling, K. B., & Poeppel, D. (2014). Delta-beta coupled oscillations underlie temporal prediction accuracy. *Cerebral Cortex*, *25*(9), 3077–3085. https://doi.org/10.1093/cercor/bhu103

Assaneo, M. F., Ripollés, P., Orpella, J., Lin, W. M., de Diego-Balaguer, R., & Poeppel, D. (2019). Spontaneous synchronization to speech reveals neural mechanisms facilitating language learning. *Nature Neuroscience*, *22*(4), 627–632. https://doi.org/10.1038/s41593-019-0353-z

Auksztulewicz, R., Myers, N. E., Schnupp, J. W., & Nobre, A. C. (2019). Rhythmic temporal expectation boosts neural activity by increasing neural gain. *The Journal of Neuroscience*, *39*(49), 9806–9817. https://doi.org/10.1523/JNEUROSCI.0925-19.2019

Auksztulewicz, R., Schwiedrzik, C. M., Thesen, T., Doyle, W., Devinsky, O., Nobre, A. C., … Melloni, L. (2018). Not all predictions are equal: "What" and "when" predictions modulate activity in auditory cortex through different mechanisms. *The Journal of Neuroscience*, *38*(40), 8680–8693. https://doi.org/10.1523/JNEUROSCI.0369-18.2018

Bauer, A.-K. R., Jaeger, M., Thorne, J. D., Bendixen, A., & Debener, S. (2015). The auditory dynamic attending theory revisited: A closer look at the pitch comparison task. *Brain Research*, *1626*, 198–210. https://doi.org/10.1016/j.brainres.2015.04.032

Bouwer, F. L. (2022). Neural entrainment to auditory rhythms: Automatic or top-down driven? *Journal of Neuroscience*, *42*(11), 2146–2148. https://doi.org/10.1523/JNEUROSCI.2305-21.2022

Bouwer, F. L., Burgoyne, J. A., Odijk, D., Honing, H., & Grahn, J. A. (2018). What makes a rhythm complex? The influence of musical training and accent type on beat perception. *PLOS ONE*, *13*(1), e0190322. https://doi.org/10.1371/journal.pone.0190322

Bouwer, F. L., & Honing, H. (2015). Temporal attending and prediction influence the perception of metrical rhythm: Evidence from reaction times and ERPs. *Frontiers in Psychology*, *6*(July), 1094. https://doi.org/10.3389/fpsyg.2015.01094

Bouwer, F. L., Honing, H., & Slagter, H. A. (2020). Beat-based and memory-based temporal expectations in rhythm: similar perceptual effects, different underlying mechanisms. *Journal of Cognitive Neuroscience*, *32*(7), 1221–1241. https://doi.org/10.1162/jocn_a_01529

Bouwer, F. L., Nityananda, V., Rouse, A. A., & ten Cate, C. (2021). Rhythmic abilities in humans and non-human animals: A review and recommendations from a methodological perspective. *Philosophical Transactions of the Royal Society B*, *376*, 20200335. https://doi.org/10.1098/rstb.2020.0335

Bouwer, F. L., Van Zuijen, T. L., & Honing, H. (2014). Beat processing is pre-attentive for metrically simple rhythms with clear accents: An ERP study. *PLoS ONE*, *9*(5), e97467. https://doi.org/10.1371/journal.pone.0097467

Bouwer, F. L., Werner, C. M., Knetemann, M., & Honing, H. (2016). Disentangling beat perception from sequential learning and examining the influence of attention and musical abilities on ERP responses to rhythm. *Neuropsychologia*, *85*, 80–90. https://doi.org/10.1016/j.neuropsychologia.2016.02.018

Breska, A., & Deouell, L. Y. (2012). Automatic bias of temporal expectations following temporally regular input independently of high-level temporal expectation. *Journal of Cognitive Neuroscience*, *26*(7), 1555–1571. https://doi.org/10.1162/jocn

Breska, A., & Deouell, L. Y. (2017). Neural mechanisms of rhythm-based temporal prediction: Delta phase-locking reflects temporal predictability but not rhythmic entrainment. *PLOS Biology*, *15*(2), e2001665. https://doi.org/10.1371/journal.pbio.2001665

Breska, A., & Ivry, R. B. (2016). Taxonomies of timing: where does the cerebellum fit in? *Current Opinion in Behavioral Sciences*, *8*, 282–288. https://doi.org/10.1016/J.COBEHA.2016.02.034

Breska, A., & Ivry, R. B. (2018). Double dissociation of single-interval and rhythmic temporal prediction in cerebellar degeneration and Parkinson's disease. *Proceedings of the National Academy of Sciences*, *115*(48), 12283–12288. https://doi.org/10.1073/pnas.1810596115

Breska, A., & Ivry, R. B. (2020). Context-specific control over the neural dynamics of temporal attention by the human cerebellum. *Science Advances*, *6*(49), 1–16. https://doi.org/10.1126/sciadv.abb1141

Brysbaert, M., & Stevens, M. (2018). Power analysis and effect size in mixed effects models: A tutorial. *Journal of Cognition*, *1*(1), 1–20. https://doi.org/10.5334/joc.10

Cameron, D. J., & Grahn, J. A. (2014). Enhanced timing abilities in percussionists generalize to rhythms without a musical beat. *Frontiers in Human Neuroscience*, *8*(December), 1003. https://doi.org/10.3389/fnhum.2014.01003

Cannon, J. J. (2021). Expectancy-based rhythmic entrainment as continuous Bayesian inference. *PLoS Computational Biology*, *17*(6), e1009025. https://doi.org/10.1371/journal.pcbi.1009025

Cannon, J. J., & Patel, A. D. (2021). How Beat Perception Co-opts Motor Neurophysiology. *Trends in Cognitive Sciences*, *25*(2), 137–150. https://doi.org/10.1016/j.tics.2020.11.002

Capilla, A., Pazo-Alvarez, P., Darriba, A., Campo, P., & Gross, J. (2011). Steady-State Visual Evoked Potentials

1264    Can Be Explained by Temporal Superposition of Transient Event-Related Responses. *PLoS ONE*, *6*,
1265    e14543. https://doi.org/10.1371/journal.pone.0014543

1266 Carifio, J., & Perla, R. (2008). Resolving the 50-year debate around using and misusing Likert scales. *Medical*
1267    *Education*, *42*(12), 1150–1152. https://doi.org/10.1111/j.1365-2923.2008.03172.x

1268 Christensen, R. H. B. (2019). ordinal - Regression models for Ordinal Data. R package version 2019.4-25.
1269    Retrieved from http://www.cran.r-project.org/package=ordinal/

1270 Cohen, M. X. (2014). *Analyzing neural time series data: theory and practice*. MIT press.

1271 Cole, S., & Voytek, B. (2019). Cycle-by-cycle analysis of neural oscillations. *Journal of Neurophysiology*, *122*,
1272    849–861. https://doi.org/10.1152/jn.00273.2019

1273 Cravo, A. M., Rohenkohl, G., Wyart, V., & Nobre, A. C. (2013). Temporal expectation enhances contrast
1274    sensitivity by phase entrainment of low-frequency oscillations in visual cortex. *The Journal of*
1275    *Neuroscience*, *33*(9), 4002–4010. https://doi.org/10.1523/JNEUROSCI.4675-12.2013

1276 Dalla Bella, S., Dotov, D., Bardy, B., & Cochen De Cock, V. (2018). Individualization of music-based rhythmic
1277    auditory cueing in Parkinson's disease. *Annals of the New York Academy of Sciences*, *1423*, 308–317.
1278    https://doi.org/10.1111/nyas.13859

1279 Damsma, A., Schlichting, N., & van Rijn, H. (2021). Temporal context actively shapes EEG signatures of time
1280    perception. *The Journal of Neuroscience*. https://doi.org/10.1523/JNEUROSCI.0628-20.2021

1281 de Graaf, T. A., Gross, J., Paterson, G., Rusch, T., Sack, A. T., & Thut, G. (2013). Alpha-Band Rhythms in
1282    Visual Task Performance: Phase-Locking by Rhythmic Sensory Stimulation. *PLoS ONE*, *8*(3), e60035.
1283    https://doi.org/10.1371/journal.pone.0060035

1284 Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics
1285    including independent component analysis. *Journal of Neuroscience Methods*, *134*, 9–21.
1286    https://doi.org/10.1016/j.jneumeth.2003.10.009

1287 Ding, N., Patel, A. D., Chen, L., Butler, H., Luo, C., & Poeppel, D. (2017). Temporal modulations in speech and
1288    music. *Neuroscience and Biobehavioral Reviews*, *81*, 181–187.
1289    https://doi.org/10.1016/j.neubiorev.2017.02.011

1290 Doelling, K. B., & Assaneo, M. F. (2021). Neural oscillations are a start toward understanding brain activity
1291    rather than the end. *PLoS Biology*, *19*(5), 1–12. https://doi.org/10.1371/journal.pbio.3001234

1292 Doelling, K. B., Assaneo, M. F., Bevilacqua, D., Pesaran, B., & Poeppel, D. (2019). An oscillator model better
1293    predicts cortical entrainment to music. *Proceedings of the National Academy of Sciences*, *116*(20), 10113–
1294    10121. https://doi.org/10.1073/pnas.1816414116

1295 Donoghue, T., Schaworonkow, N., & Voytek, B. (2021). Methodological considerations for studying neural
1296    oscillations. *European Journal of Neuroscience*, (January), 1–26. https://doi.org/10.1111/ejn.15361

1297 Drake, C., Jones, M. R., & Baruch, C. (2000). The development of rhythmic attending in auditory sequences:
1298    attunement, referent period, focal attending. *Cognition*, *77*(3), 251–288. https://doi.org/10.1016/S0010-
1299    0277(00)00106-2

1300 Elliott, M. T., Wing, A. M., & Welchman, A. E. (2014). Moving in time: Bayesian causal inference explains
1301    movement coordination to auditory beats. *Proceedings of the Royal Society B*, *281*(May), 20140751.

1302 Fahrenfort, J. J., van Driel, J., van Gaal, S., & Olivers, C. N. L. (2018). From ERPs to MVPA Using the
1303    Amsterdam Decoding and Modeling Toolbox (ADAM). *Frontiers in Neuroscience*, *12*, 368.
1304    https://doi.org/10.3389/fnins.2018.00368

1305 Fitch, W. T. (2013). Rhythmic cognition in humans and animals: distinguishing meter and pulse
1306    perception. *Frontiers in systems neuroscience*, *7*, 68. https://doi.org/10.3389/fnsys.2013.00068

1307 Fox, J., & Weisberg, S. (2019). *An R Companion to Applied Regression* (3rd ed.). Thousand Oaks (CA): Sage.
1308    Retrieved from https://socialsciences.mcmaster.ca/jfox/Books/Companion/

1309 Geiser, E., Ziegler, E., Jancke, L., Meyer, M., Jäncke, L., Meyer, M., … Meyer, M. (2009). Early
1310    electrophysiological correlates of meter and rhythm processing in music perception. *Cortex*, *45*(1), 93–
1311    102. https://doi.org/10.1016/j.cortex.2007.09.010

1312 Grahn, J. A., & Brett, M. (2007). Rhythm and beat perception in motor areas of the brain. *Journal of Cognitive*
1313    *Neuroscience*, *19*(5), 893–906. https://doi.org/10.1162/jocn.2007.19.5.893

1314 Grahn, J. A., & Rowe, J. B. (2009). Feeling the beat: premotor and striatal interactions in musicians and
1315    nonmusicians during beat perception. *The Journal of Neuroscience*, *29*(23), 7540–7548.
1316    https://doi.org/10.1523/JNEUROSCI.2018-08.2009

1317 Haegens, S., & Zion Golumbic, E. (2018). Rhythmic facilitation of sensory processing: A critical review.
1318    *Neuroscience and Biobehavioral Reviews*, *86*(March), 150–165.
1319    https://doi.org/10.1016/j.neubiorev.2017.12.002

1320 Heideman, S. G., van Ede, F., & Nobre, A. C. (2018). Early Behavioural Facilitation by Temporal Expectations
1321    in Complex Visual-motor Sequences. *Neuroscience*, *389*, 74–84.
1322    https://doi.org/10.1016/J.NEUROSCIENCE.2018.05.014

1323 Henry, M. J., & Herrmann, B. (2014). Low-frequency neural oscillations support dynamic attending in temporal

context. *Timing & Time Perception*, *2*(1), 62–86. https://doi.org/10.1163/22134468-00002011

Henry, M. J., Herrmann, B., & Obleser, J. (2014). Entrained neural oscillations in multiple frequency bands
comodulate behavior. *Proceedings of the National Academy of Sciences*, *111*(41), 14935–14940.
https://doi.org/10.1073/pnas.1408741111

Henry, M. J., & Obleser, J. (2012). Frequency modulation entrains slow neural oscillations and optimizes human
listening behavior. *Proceedings of the National Academy of Sciences*, *109*(49), 20095–20100.
https://doi.org/10.1073/pnas.1213390109

Herbst, S. K., Stefanics, G., & Obleser, J. (2022). Endogenous modulation of delta phase by expectation–A
replication of Stefanics et al., 2010. *Cortex*, *149*, 226–245. https://doi.org/10.1016/j.cortex.2022.02.001

Hickok, G., Farahbod, H., & Saberi, K. (2015). The Rhythm of Perception: Entrainment to Acoustic Rhythms
Induces Subsequent Perceptual Oscillation. *Psychological Science*, *26*(7), 1006–1013.
https://doi.org/10.1177/0956797615576533

Honing, H. (2012). Without it no music: beat induction as a fundamental musical trait. *Annals of the New York*
Academy of Sciences, *1252*(1), 85–91. https://doi.org/10.1111/j.1749-6632.2011.06402.x

Honing, H., & Bouwer, F. L. (2019). Rhythm. In P. J. Rentfrow & D. Levitin (Eds.), *Foundations in Music*
Psychology: Theory and Research (pp. 33–69). Cambridge, MA: MIT Press.

Iversen, J. R., & Patel, A. D. (2008). The Beat Alignment Test (BAT): Surveying beat processing abilities in the
general population. In K. Miyazaki, Y. Hiraga, M. Adachi, Y. Nakajima, & M. Tsuzaki (Eds.), *10th Intl.*
Conf. on Music Perception and Cognition (pp. 465–468). Adelaide: Causal Productions.

Jamieson, S. (2004). Likert scales: How to (ab)use them. *Medical Education*, *38*(12), 1217–1218.
https://doi.org/10.1111/j.1365-2929.2004.02012.x

JASP, T. (2019). JASP. Retrieved from https://jasp-stats.org

Jeffreys, H. (1961). *Theory of probability* (3rd ed.). Oxford, UK: Oxford University Press.

Jones, M. R., Moynihan, H., MacKenzie, N., & Puente, J. (2002). Temporal aspects of stimulus-driven attending
in dynamic arrays. *Psychological Science*, *13*(4), 313–319. https://doi.org/10.1111/1467-9280.00458

Keele, S. W., Nicoletti, R., Ivry, R. B., & Pokorny, R. A. (1989). Mechanisms of perceptual timing: Beat-based
or interval-based judgements? *Psychological Research*, *50*, 251–256. https://doi.org/10.1007/BF00309261

Keitel, C., Obleser, J., Jessen, S., & Henry, M. J. (2021). Frequency-Specific Effects in Infant
Electroencephalograms Do Not Require Entrained Neural Oscillations: A Commentary on Köster et al .
(2019). *Psychological Science*, *32*(6), 966–971. https://doi.org/10.1177/09567976211001317

King, J.-R., & Dehaene, S. (2014). Characterizing the dynamics of mental representations: the temporal
generalization method. *Trends in Cognitive Sciences*, *18*(4), 203–210.
https://doi.org/10.1016/J.TICS.2014.01.002

Kloosterman, N. A., Kosciessa, J. Q., Lindenberger, U., Fahrenfort, J. J., & Garrett, D. D. (2020). Boosts in
brain signal variability track liberal shifts in decision bias. *ELife*, *9*, 1–22.
https://doi.org/10.7554/ELIFE.54201

Kosciessa, J. Q., Kloosterman, N. A., & Garrett, D. D. (2020). Standard multiscale entropy reflects neural
dynamics at mismatched temporal scales: What's signal irregularity got to do with it? *PLoS Computational*
Biology, *16*(5), e1007885. https://doi.org/10.1371/journal.pcbi.1007885

Kösem, A., Bosker, H. R., Takashima, A., Meyer, A., Jensen, O., & Hagoort, P. (2018). Neural Entrainment
Determines the Words We Hear. *Current Biology*, *28*(18), 2867-2875.e3.
https://doi.org/10.1016/j.cub.2018.07.023

Large, E. W. (2008). Resonating to musical rhythm: Theory and experiment. In S. Grondin (Ed.), *Psychology of*
time (pp. 189–231). Bingley, UK: Emerald Group Publishing. https://doi.org/10.1016/B978-0-08046-977-
5.00006-5

Large, E. W., Herrera, J. A., & Velasco, M. J. (2015). Neural networks for beat perception in musical rhythm.
Frontiers in Systems Neuroscience, *9*(November), 159. https://doi.org/10.3389/fnsys.2015.00159

Large, E. W., & Jones, M. R. (1999). The dynamics of attending: how people track time-varying events.
Psychological Review, *106*(1), 119–159. https://doi.org/10.1037/0033-295X.106.1.119

Large, E. W., & Palmer, C. (2002). Perceiving temporal regularity in music. *Cognitive Science*, *26*(1), 1–37.
https://doi.org/10.1207/s15516709cog2601_1

Law, L. N. C., & Zentner, M. (2012). Assessing Musical Abilities Objectively: Construction and Validation of
the Profile of Music Perception Skills. *PLoS ONE*, *7*(12), e52508.
https://doi.org/10.1371/journal.pone.0052508

Lenc, T., Keller, P. E., Varlet, M., & Nozaradan, S. (2018). Neural tracking of the musical beat is enhanced by
low-frequency sounds. *Proceedings of the National Academy of Sciences*, *115*(32), 8221–8226.
https://doi.org/10.1073/pnas.1801421115

Lenc, T., Merchant, H., Keller, P. E., Honing, H., Varlet, M., & Nozaradan, S. (2021). Mapping between sound,
brain and behaviour: Four-level framework for understanding rhythm processing in humans and non-
human primates. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *376*, 20200325.

https://doi.org/10.1098/rstb.2020.0325

Lenth, R. (2019). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.4. Retrieved from https://cran.r-project.org/package=emmeans

Leow, L., & Grahn, J. A. (2014). Neural mechanisms of rhythm perception: Present findings and future directions. In H. Merchant & V. de Lafuente (Eds.), *Neurobiology of Interval Timing* (pp. 325–338). New York: Springer. https://doi.org/10.1007/978-1-4939-1782-2

Lin, W. M., Oetringer, D. A., Bakker-Marshall, I., Emmerzaal, J., Wilsch, A., ElShafei, H. A., … Haegens, S. (2021). No behavioural evidence for rhythmic facilitation of perceptual discrimination. *European Journal of Neuroscience*. https://doi.org/10.1111/ejn.15208

London, J. (2012). *Hearing in time: Psychological aspects of musical meter.* (2nd ed.). Oxford: Oxford University Press.

Manning, F. C., Harris, J., & Schutz, M. (2017). Temporal prediction abilities are mediated by motor effector and rhythmic expertise. *Experimental Brain Research*, *235*(3), 861–871. https://doi.org/10.1007/s00221-016-4845-8

Manning, F. C., & Schutz, M. (2013). "Moving to the beat" improves timing perception. *Psychonomic Bulletin & Review*, *20*, 1133–1139. https://doi.org/10.3758/s13423-013-0439-7

Mathewson, K. E., Prudhomme, C., Fabiani, M., Beck, D. M., Lleras, A., & Gratton, G. (2012). Making Waves in the Stream of Consciousness: Entraining Oscillations in EEG Alpha and Fluctuations in Visual Awareness with Rhythmic Visual Stimulation. *Journal of Cognitive Neuroscience*, *24*(12), 2321–2333. https://doi.org/10.1162/jocn_a_00288

Matthews, T. E., Thibodeau, J. N. L., Gunther, B. P., & Penhune, V. B. (2016). The Impact of Instrument-Specific Musical Training on Rhythm Perception and Production. *Frontiers in Psychology*, *7*(February), 69. https://doi.org/10.3389/fpsyg.2016.00069

Mento, G. (2013). The passive CNV: Carving out the contribution of task-related processes to expectancy. *Frontiers in Human Neuroscience*, *7*(December), 827. https://doi.org/10.3389/fnhum.2013.00827

Mento, G. (2017). The role of the P3 and CNV components in voluntary and automatic temporal orienting: A high spatial-resolution ERP study. *Neuropsychologia*, *107*, 31–40. https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2017.10.037

Merchant, H., Grahn, J. A., Trainor, L. J., Rohrmeier, M. A., & Fitch, W. T. (2015). Finding the beat: a neural perspective across humans and non-human primates. *Philosophical Transactions of the Royal Society B*, *370*, 20140093. https://doi.org/10.1098/rstb.2014.0093

Meyer, L., Sun, Y., & Martin, A. E. (2019). Synchronous, but not entrained: exogenous and endogenous cortical rhythms of speech and language processing. *Language, Cognition and Neuroscience*, 1–11. https://doi.org/10.1080/23273798.2019.1693050

Morillon, B., Schroeder, C. E., Wyart, V., & Arnal, L. H. (2016). Temporal prediction in lieu of periodic stimulation. *The Journal of Neuroscience*, *36*(8), 2342–2347. https://doi.org/10.1523/JNEUROSCI.0836-15.2016

Müllensiefen, D., Gingras, B., Musil, J., & Stewart, L. (2014). The musicality of non-musicians: An index for assessing musical sophistication in the general population. *PLoS ONE*, *9*(2), e89642. https://doi.org/10.1371/journal.pone.0089642

Nobre, A. C., & van Ede, F. (2018). Anticipated moments: temporal structure in attention. *Nature Reviews Neuroscience*, *19*, 34–48. https://doi.org/10.1038/nrn.2017.141

Novembre, G., & Iannetti, G. D. (2018). Tagging the musical beat: Neural entrainment or event-related potentials? *Proceedings of the National Academy of Sciences*, *115*(47), E11002–E11003. https://doi.org/10.1073/pnas.1815311115

Nozaradan, S., Peretz, I., Missal, M., & Mouraux, A. (2011). Tagging the neuronal entrainment to beat and meter. *The Journal of Neuroscience*, *31*(28), 10234–10240. https://doi.org/10.1523/JNEUROSCI.0411-11.2011

Nozaradan, S., Peretz, I., & Mouraux, A. (2012). Selective neuronal entrainment to the beat and meter embedded in a musical rhythm. *The Journal of Neuroscience*, *32*(49), 17572–17581. https://doi.org/10.1523/JNEUROSCI.3203-12.2012

O'Reilly, J. X., McCarthy, K. J., Capizzi, M., & Nobre, A. C. (2008). Acquisition of the temporal and ordinal structure of movement sequences in incidental learning. *Journal of Neurophysiology*, *99*, 2731–2735. https://doi.org/10.1152/jn.01141.2007

Obleser, J., & Kayser, C. (2019). Neural Entrainment and Attentional Selection in the Listening Brain. *Trends in Cognitive Sciences*, *23*(11), 913–926. https://doi.org/10.1016/J.TICS.2019.08.004

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, *2011*, 156869. https://doi.org/10.1155/2011/156869

Palmer, C., & Krumhansl, C. L. (1990). Mental representations for musical meter. *Journal of Experimental*

Psychology. Human Perception and Performance, *16*(4), 728–741. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/2148588

Pesnot Lerousseau, J., Trébuchon, A., Morillon, B., & Schön, D. (2021). Frequency selectivity of persistent cortical oscillatory responses to auditory rhythmic stimulation. *The Journal of Neuroscience*, *41*(38), 7991–8006. https://doi.org/10.1523/JNEUROSCI.0213-21.2021

Povel, D.-J., & Essens, P. (1985). Perception of Temporal Patterns. *Music Perception*, *2*(4), 411–440. https://doi.org/10.2307/40285311

Povel, D.-J., & Okkerman, H. (1981). Accents in equitone sequences. *Perception & Psychophysics*, *30*(6), 565–572. https://doi.org/10.3758/BF03202011

Praamstra, P., Kourtis, D., Kwok, H. F., & Oostenveld, R. (2006). Neurophysiology of implicit timing in serial choice reaction-time performance. *The Journal of Neuroscience*, *26*(20), 5448–5455. https://doi.org/10.1523/JNEUROSCI.0440-06.2006

R Development Core Team. (2008). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from www.R-project.org

Rimmele, J. M., Morillon, B., Poeppel, D., & Arnal, L. H. (2018). Proactive sensing of periodic and aperiodic auditory patterns. *Trends in Cognitive Sciences*, *22*(10), 870–882. https://doi.org/10.1016/J.TICS.2018.08.003

Schroeder, C. E., & Lakatos, P. (2009a). Low-frequency neuronal oscillations as instruments of sensory selection. *Trends in Neurosciences*, *32*(1), 9–18. https://doi.org/10.1016/j.tins.2008.09.012

Schroeder, C. E., & Lakatos, P. (2009b). The Gamma Oscillation: Master or Slave? *Brain Topography*, *22*, 24–26. https://doi.org/10.1007/s10548-009-0080-y

Schultz, B. G., Stevens, C. J., Keller, P. E., & Tillmann, B. (2013). The implicit learning of metrical and nonmetrical temporal patterns. *The Quarterly Journal of Experimental Psychology*, *66*(2), 360–380. https://doi.org/10.1080/17470218.2012.712146

Shalev, N., Nobre, A. C., & van Ede, F. (2019). Time for What? Breaking Down Temporal Anticipation. *Trends in Neurosciences*, *42*(6), 373–374. https://doi.org/10.1016/J.TINS.2019.03.002

Stefanics, G., Hangya, B., Hernádi, I., Winkler, I., Lakatos, P., & Ulbert, I. (2010). Phase Entrainment of Human Delta Oscillations Can Mediate the Effects of Expectation on Reaction Speed. *The Journal of Neuroscience*, *30*(41), 13578–13585. https://doi.org/10.1523/JNEUROSCI.0703-10.2010

Sun, Y., Michalareas, G., & Poeppel, D. (2021). The impact of phase entrainment on auditory detection is highly variable: Revisiting a key finding. *European Journal of Neuroscience*, (June), 1–18. https://doi.org/10.1111/ejn.15367

Tal, I., Large, E. W., Rabinovitch, E., Wei, Y., Schroeder, C. E., Poeppel, D., & Golumbic, E. Z. (2017). Neural entrainment to the beat: The "missing-pulse" phenomenon. *The Journal of Neuroscience*, *37*(26), 6331–6341. https://doi.org/10.1523/JNEUROSCI.2500-16.2017

Teki, S., Grube, M., Kumar, S., & Griffiths, T. D. (2011). Distinct neural substrates of duration-based and beat-based auditory timing. *The Journal of Neuroscience*, *31*(10), 3805–3812. https://doi.org/10.1523/JNEUROSCI.5561-10.2011

Tichko, P., & Large, E. W. (2019). Modeling infants' perceptual narrowing to musical rhythms: neural oscillation and Hebbian plasticity. *Annals of the New York Academy of Sciences*, *1453*(April), 125–139. https://doi.org/10.1111/nyas.14050

van Atteveldt, N., Musacchia, G., Zion-Golumbic, E., Sehatpour, P., Javitt, D. C., & Schroeder, C. E. (2015). Complementary fMRI and EEG evidence for more efficient neural processing of rhythmic vs. unpredictably timed sounds. *Frontiers in Psychology*, *6*(OCT), 1–11. https://doi.org/10.3389/fpsyg.2015.01663

van Bree, S., Sohoglu, E., Davis, M. H., & Zoefel, B. (2021). Sustained neural rhythms reveal endogenous oscillations supporting speech perception. *PLOS Biology*, *19*(2), e3001142. https://doi.org/10.1371/journal.pbio.3001142

van der Weij, B., Pearce, M. T., & Honing, H. (2017). A probabilistic model of meter perception: simulating enculturation. *Frontiers in Psychology*, *8*(May), 824. https://doi.org/10.3389/fpsyg.2017.00824

Vuust, P., Pallesen, K. J., Bailey, C., Van Zuijen, T. L., Gjedde, A., Roepstorff, A., & Østergaard, L. (2005). To musicians, the message is in the meter: Pre-attentive neuronal responses to incongruent rhythm are left-lateralized in musicians. *NeuroImage*, *24*(2), 560–564. https://doi.org/10.1016/j.neuroimage.2004.08.039

Wagenmakers, E.-J., Love, J., Marsman, M., Jamil, T., Ly, A., Verhagen, J., … Morey, R. D. (2018). Bayesian inference for psychology. Part II: Example applications with JASP. *Psychonomic Bulletin & Review*, *25*(1), 58–76. https://doi.org/10.3758/s13423-017-1323-7

Waschke, L., Kloosterman, N. A., Obleser, J., & Garrett, D. D. (2021). Behavior needs neural variability. *Neuron*, *109*(5), 751–766. https://doi.org/10.1016/j.neuron.2021.01.023

Wilsch, A., Mercier, M. R., Obleser, J., Schroeder, C. E., & Haegens, S. (2020). Spatial Attention and Temporal Expectation Exert Differential Effects on Visual and Auditory Discrimination. *Journal of Cognitive*

Neuroscience, *32*(8), 1562–1576. https://doi.org/10.1162/jocn_a_01567

Wollman, I., & Morillon, B. (2018). Organizational principles of multidimensional predictions in human
auditory attention. *Scientific Reports*, *8*(1), 13466. https://doi.org/10.1038/s41598-018-31878-5

Zalta, A., Petkoski, S., & Morillon, B. (2020). Natural rhythms of periodic temporal attention. *Nature*
Communications, *11*(1), 1051. https://doi.org/10.1038/s41467-020-14888-8

Zoefel, B., ten Oever, S., & Sack, A. T. (2018). The Involvement of Endogenous Neural Oscillations in the
Processing of Rhythmic Input: More Than a Regular Repetition of Evoked Neural Responses. *Frontiers in*
Neuroscience, *12*, 95. https://doi.org/10.3389/fnins.2018.00095

Zoefel, B., & Vanrullen, R. (2017). Oscillatory mechanisms of stimulus processing and selection in the visual
and auditory systems: State-of-the-art, speculations and suggestions. *Frontiers in Neuroscience*, *11*(May),
296. https://doi.org/10.3389/fnins.2017.00296

1515

1516

Supplementary materials for "A silent disco: Persistent entrainment of low-frequency neural oscillations underlies beat-based, but not pattern-based temporal expectations"

(Bouwer, Fahrenfort, Millard, Kloosterman, & Slagter, 2022)

Supplementary Figure 1. CNV-like component with traditional 50 ms baseline and difference waves.
A) With the more traditional pre-cue baseline, there was a trend for the random condition to elicit a more negative deflection than the beat-based condition ($p = 0.058$, $300 – 446$ ms) and for the pattern-based condition to elicit a more negative deflection than the random condition (early window: $p = 0.083$, $380 – 551$ ms; later window: $p = 0.022$, $774 – 1200$ ms). Note that a significant cluster spanning the entire analysis window of $300 – 1200$ ms was found when comparing between the pattern-based and beat-based condition with baseline correction ($p = 0.011$), indicative of a possible overall shift due to the baseline correction. However, the overall picture stays the same as when not using a baseline, a CNV-like negative deflection can be seen around 300-450 ms after the last sound onset for the pattern-based and random, but not the beat-based condition. B) Scalp distributions for windows with a significant cluster (see also Figure 5). C) Difference waves depicting condition differences and the 95% confidence interval around the mean difference waveforms. The scalp distributions similarly depict condition differences. D) Evoked potential showing the CNV-like component during sound presentation. Time 0 here is the presentation of a sound, with the next sound being presented at 600 (beat-based condition) or 780 ms (pattern-based and random condition). As in the silence window, a CNV-like negative deflection can be observed, which was larger for the pattern-based than random condition ($p = 0.039$), and peaks around 350 ms after the onset of the previous sound. The difference between the pattern-based and beat-based condition here did not reach significance ($p = 0.1$), likely because of the noisy baseline.

Supplementary Figure 2. Phase consistency at beat times (1200 and 1800 ms) in the silence is higher for beat-based than random sequences. All plots depict the inter-trial phase consistency for beat times (1200 and 1800 ms after the last sound onset) in the silence in the delta range (0.5-4 Hz). A) Single participant data, with all data points in grey. Boxplots show the median, with the lower and upper hinges corresponding to the first and third quartiles (25^{th} and 75^{th} percentiles), and the whiskers corresponding to values no further than 1.5 times the inter-quartile range from the hinges. Error bars depict 2 standard errors around the mean. Values are averaged over all electrodes. B) Scalp topographies for all conditions. Electrodes contributing to significant clusters in the cluster-based tests are highlighted on the plots for the conditions from each comparison in which inter-trial phase coherence was largest.

Supplementary Figure 3. MMSE after highpass filtering at 5 Hz. Here, all differences between conditions are eliminated by the highpass filtering.

Supplementary Figure 4. Similarity bounds used to calculate MMSE values mirror the condition differences. This finding suggests that condition differences found in the entropy measure, with higher entropy for the pattern-based than random condition in the silence window, may be due to overall variability differences, with lower variability for the pattern-based than random condition, rather than differences in entropy of the signal.

Supplementary Table 1. Simple effects per position from the full models, comparing three conditions. All *p*-values have been Bonferroni corrected for 18 (Experiment 1) or 9 (Experiment 2) comparisons.

Position	Contrast	Experiment 1				Experiment 2			
		Estimate	SE	Z ratio	*p*	Estimate	SE	Z ratio	*p*
600 ms	Random – Pattern	2.4	0.09	26.2	<.0001				
	Random – Beat	-1.2	0.09	13.3	<.0001				
	Pattern - Beat	-3.5	0.09	-38.3	<.0001				
780 ms	Random – Pattern	-0.3	0.09	-2.9	0.058				
	Random – Beat	1.6	0.09	18.3	<.0001				
	Pattern - Beat	1.8	0.09	20.7	<.0001				
885 ms	Random – Pattern	-0.1	0.09	-1.6	1				
	Random – Beat	1.5	0.09	17.0	<.0001				
	Pattern - Beat	1.6	0.09	18.4	<.0001				
990 ms	Random – Pattern	0.1	0.09	1.1	1	0.6	0.1	4.8	<.0001
	Random – Beat	1.4	0.09	16.2	<.0001	1.4	0.1	11.4	<.0001
	Pattern - Beat	1.3	0.09	15.0	<.0001	0.8	0.1	6.5	<.0001
1200 ms	Random – Pattern	0.6	0.08	7.5	<.0001	0.2	0.1	1.6	0.92
	Random – Beat	-0.3	0.09	-3.7	0.004	0.1	0.1	0.9	1
	Pattern - Beat	-1.0	0.09	-10.9	<.0001	-0.1	0.1	-0.7	1
1485 ms	Random – Pattern	0.5	0.09	5.2	<.0001	-0.5	0.1	-4.4	<.0001
	Random – Beat	0.6	0.09	6.5	<.0001	0.4	0.1	3.2	0.011
	Pattern - Beat	0.1	0.09	1.3	1	0.9	0.1	7.5	<.0001

Supplementary Table 2. Simple effects per condition from the baseline corrected model, using the random condition as baseline, for all positions. All *p*-values have been Bonferroni corrected for 30 (Experiment 1) or 6 (Experiment 2) comparisons.

Condition	Contrast	Experiment 1				Experiment 2			
		Estimate	SE	Z ratio	*p*	Estimate	SE	Z ratio	*p*
Beat-based	600 ms - 780 ms	2.3	0.08	28.7	<.0001				
	600 ms - 885 ms	2.2	0.08	27.8	<.0001				
	600 ms - 990 ms	2.1	0.08	27.0	<.0001				
	600 ms - 1200 ms	0.6	0.08	6.9	<.0001				
	600 ms - 1485 ms	1.5	0.08	18.0	<.0001				
	780 ms - 885 ms	-0.1	0.08	-1.3	1				
	780 ms - 990 ms	-0.2	0.08	-1.9	1				
	780 ms - 1200 ms	-1.8	0.08	-21.3	<.0001				
	780 ms - 1485 ms	-0.9	0.08	-10.5	<.0001				
	885 ms - 990 ms	-0.1	0.08	-0.7	1				
	885 ms - 1200 ms	-1.7	0.08	-20.3	<.0001				
	885 ms - 1485 ms	-0.8	0.08	-9.4	<.0001				
	990 ms - 1200 ms	-1.6	0.08	-19.5	<.0001	-1.1	0.1	-9.6	<.0001
	990 ms - 1485 ms	-0.7	0.08	-8.7	<.0001	-0.9	0.1	-7.9	<.0001
	1200 ms - 1485 ms	0.9	0.08	10.9	<.0001	0.2	0.1	1.8	0.46
Pattern-based	600 ms - 780 ms	-2.3	0.09	-27.0	<.0001				
	600 ms - 885 ms	-2.2	0.08	-26.4	<.0001				
	600 ms - 990 ms	-2.1	0.08	-24.8	<.0001				
	600 ms - 1200 ms	-1.7	0.08	-20.2	<.0001				
	600 ms - 1485 ms	-1.7	0.09	-20.3	<.0001				
	780 ms - 885 ms	0.1	0.08	1.0	1				
	780 ms - 990 ms	0.2	0.08	3.0	0.080				
	780 ms - 1200 ms	0.6	0.08	7.6	<.0001				
	780 ms - 1485 ms	0.6	0.08	6.9	<.0001				
	885 ms - 990 ms	0.2	0.08	2.0	1				
	885 ms - 1200 ms	0.5	0.08	6.7	<.0001				
	885 ms - 1485 ms	0.5	0.08	6.0	<.0001				
	990 ms - 1200 ms	0.4	0.08	4.7	<.0001	-0.3	0.1	-2.9	0.022
	990 ms - 1485 ms	0.3	0.08	4.1	0.001	-0.9	0.1	-8.3	<.0001
	1200 ms - 1485 ms	0.0	0.08	-0.5	1.00	-0.6	0.1	-5.4	<.0001

<center>**Supplementary Methods: MSE computation**</center>

To compute entropy, epochs were concatenated. Sample entropy was then calculated by taking the following steps:

1. In each time series, a template is selected, consisting of m samples. The calculation of entropy is an iterative process, using each sample in the time series as a starting point for the template once.

2. Throughout the time series, the algorithm searches for patterns that match the template pattern. A section of samples is considered a match if it resembles the template pattern enough to fall within a set boundary, which is defined as r x SD (the similarity bound).

3. The number of pattern matches is counted.

4. Subsequently, the same procedure is followed for patterns of $m + 1$ samples long.

5. For the total counts of pattern matches throughout the time series, sample entropy is then calculated as the logarithm of the ratio between pattern matches of length m and pattern matches of length $m + 1$.

Thus, sample entropy reflects the proportion of patterns in the time series that stays similar when an extra sample is added to the pattern. Here, we used $m = 2$ and $r = 0.5$, as was done previously for EEG data (Kloosterman, Kosciessa, Lindenberger, Fahrenfort, & Garrett, 2020).

Sample entropy is then repeated for multiple timescales, to account for contributions of both low and high frequency neural activity. The time series is coarsened step by step, by taking the average of a group of adjacent samples with step-wise increasing group size. This means that long, or coarse, timescales are equivalent to low frequency activity. For example, at a sampling rate of 256 Hz, a timescale of 4 (e.g., averaged over four adjacent samples, or 15.6 ms) is roughly the equivalent of looking at activity at 64 Hz, while a timescale of 153 at that sample rate corresponds to averaging over 598 ms, or the equivalent of activity at roughly 1.67 Hz. Here, we used twenty timescales ranging from 4 till 153, with 153 being the maximum timescale given the length of an epoch of 1800 ms (e.g., one epoch equals 460 samples, but

since entropy is calculated on patterns of 2 and 3 samples, the maximum coarsening to retain the possibility of having a 3-sample pattern is by averaging over 153 samples). Similarity bounds were recomputed for each time scale (Kloosterman et al., 2020; Kosciessa, Kloosterman, & Garrett, 2020). To control for the contribution of delta, we repeated the analysis on high-pass filtered data (5 Hz).

Supplementary methods: Within condition decoding

The initial decoding analysis yielded large effects that may be task-related, and individual differences in phase and metrical level attended to may have additionally hampered finding condition differences at the group level. Therefore, we ran an additional decoding analysis in which we decoded expected and unexpected positions against each other within each condition. For this analysis, we only included frontocentral electrodes (C1, C2, C3, C4, Cz, FC1, FC2, FC3, FC4, FCz, F1, F2, F3, F4, Fz), which were the same electrodes that showed the largest P1 responses to the initial sounds of the sequences, indicative of representing the auditory cortex. Here, we defined time points in the silence as expected or unexpected based on the rhythmic sequences. To equate the choice of expected and unexpected time points as much as possible in terms of number of time points included and distance between time points, we used times that were also used in the probe tone experiment: 600, 780, 885, and 1200 ms. For beat-based sequences, time points 600 and 1200 were expected (on the beat), while time points 780 and 885 were unexpected (offbeat). The same time points take on a different meaning if preceded by the pattern-based sequences. Then, 780 and 1200 ms are considered expected (predictable based on the pattern), while both 600 and 885 ms are unexpected (unpredictable based on the pattern). For all conditions, we examined whether we could classify above chance whether a time window of 100 ms centered on the time point of interest was an expected or unexpected moment in time. As for the first decoding analysis, the data were resampled to improve signal to noise, here to 128 Hz to retain enough data points in the 100 ms window for analysis. After the decoding, we extracted the average classification accuracy for the 100 ms time window to test significance for each condition separately against chance level, using t-tests against 0.5, and between conditions in the silence, using a repeated measures

ANOVA, with condition as an independent factor, a random intercept for participant, and

decoding accuracy as the dependent variable.

References

Kloosterman, N. A., Kosciessa, J. Q., Lindenberger, U., Fahrenfort, J. J., & Garrett, D. D. (2020). Boosts in brain signal variability track liberal shifts in decision bias. *ELife*, *9*, 1–22. https://doi.org/10.7554/ELIFE.54201

Kosciessa, J. Q., Kloosterman, N. A., & Garrett, D. D. (2020). Standard multiscale entropy reflects neural dynamics at mismatched temporal scales: What's signal irregularity got to do with it? *PLoS Computational Biology*, *16*(5), e1007885. https://doi.org/10.1371/journal.pcbi.1007885