1

A silent disco: Differential effects of beat-based and pattern-based temporal

expectations on persistent entrainment of low-frequency neural oscillations

Short title: Persistent effects of beat- and pattern-based temporal expectations

Fleur L. Bouwer[1,2,3*], Johannes J. Fahrenfort[1,3], Samantha K. Millard[3,4,5], Niels A.

Kloosterman[6,7], & Heleen A. Slagter[1,3]

[1] Department of Experimental and Applied Psychology, Institute Brain and Behavior
Amsterdam (IBBA), Vrije Universiteit Amsterdam
[2] Institute for Logic, Language and Computation (ILLC), University of Amsterdam
[3] Department of Psychology, Amsterdam Brain and Cognition (ABC), University of
Amsterdam
[4] Centre for Pain IMPACT, Neuroscience Research Australia (NeuRA), Sydney, New South
Wales, Australia
[5] Faculty of Medicine, University of New South Wales (UNSW), Sydney, New South Wales,
Australia
[6] Max Planck UCL Centre for Computational Psychiatry and Ageing Research, Berlin,
Germany
[7] Center for Lifespan Psychology, Max Planck Institute for Human Development, Berlin,
Germany
27
28   [*] Correspondence concerning this article should be addressed to Dr. Fleur Bouwer,
29   Department of Psychology, University of Amsterdam (bouwer@uva.nl)

30
31

Abstract

The brain uses temporal structure in the environment, like rhythm in music and speech,

to predict the timing of events, thereby optimizing their processing and perception. Temporal

expectations can be grounded in different aspects of the input structure, such as a regular beat

or a predictable pattern. One influential account posits that a generic mechanism underlies beat-

based and pattern-based expectations, namely entrainment of low frequency neural oscillations

to rhythmic input, while other accounts assume different underlying neural mechanisms. Here,

we addressed this outstanding issue by examining EEG activity and behavioral responses

during silent periods following rhythmic auditory sequences. We measured responses

outlasting the rhythms both to avoid confounding the EEG analyses with evoked responses, and

to directly test whether beat-based and pattern-based expectations persist beyond stimulation,

as predicted by entrainment theories. To properly disentangle beat-based and pattern-based

expectations, which often occur simultaneously, we used non-isochronous rhythms with a beat,

a predictable pattern, or random timing. In Experiment 1 (N = 32), beat-based expectations

affected behavioral ratings of probe events for two beat-cycles after the end of the rhythm. The

effects of pattern-based expectations reflected expectations for one interval. In Experiment 2

(N = 27), using EEG, we found enhanced spectral power at the beat frequency for beat-based

sequences both during listening and silence. For pattern-based sequences, enhanced power at a

pattern-specific frequency was present during listening, but not silence. Moreover, we found a

difference in the evoked signal following pattern-based and beat-based sequences. Finally, we

show how multivariate pattern decoding and multi scale entropy – measures sensitive to non-

oscillatory components of the signal – can be used to probe temporal expectations. Together,

our results suggest that the input structure used to form temporal expectations may affect the

associated neural mechanisms. We suggest climbing activity and low-frequency oscillations

may be differentially associated with pattern-based and beat-based expectations.

Keywords: Temporal expectations, EEG, rhythm, entrainment, Perception: Auditory processing

Introduction

Predicting the timing of incoming events optimizes processing in our dynamic environment (Nobre & van Ede, 2018), as it allows the brain to increase sensitivity to events at predicted times (Auksztulewicz et al., 2019), without the need for constant vigilance (Breska & Deouell, 2017; Rimmele et al., 2018; Schroeder & Lakatos, 2009a, 2009b). Entrainment models (Large & Jones, 1999) provide a mechanistic explanation for temporal expectations, by assuming that the phase and period of low-frequency neural oscillations synchronize to external rhythmic stimulation, causing optimal neural excitability at expected times (Haegens & Zion Golumbic, 2018; Henry & Herrmann, 2014; Schroeder & Lakatos, 2009b). In line with this, behavioral performance is improved for events in phase with an external rhythm (Bouwer & Honing, 2015; Herbst et al., 2022; Jones et al., 2002; Large & Jones, 1999), behavioral responses depend on the phase of delta oscillations (Arnal et al., 2014; Cravo et al., 2013; Henry et al., 2014; Henry & Obleser, 2012), and low frequency oscillations phase lock to rhythmic input (Doelling et al., 2019; Nozaradan et al., 2011; Stefanics et al., 2010). Note that in the current study, we focus on such perceptual entrainment – the putative alignment of low-frequency oscillations to an external stimulus and how this is associated with temporal expectations – and not on the broader connotation of entrainment that includes motor synchronization to rhythm (Damm et al., 2019; Repp, 2005).

Entrainment has mainly been studied in the context of periodic ("beat-based") sensory input, but temporal expectations can also be based on memory for absolute intervals (Breska & Deouell, 2017; Breska & Ivry, 2016; Morillon et al., 2016; Teki et al., 2011), either in isolation ("cue-based"), or as part of a predictable pattern of intervals ("pattern-based", see (Nobre & van Ede, 2018)). Predictable temporal patterns may be especially important in speech and non-Western music, which is not necessarily based on a metrical structure with a prominent periodicity in the range of human beat-based perception (London, 2012; London et al., 2017;

Polak et al., 2016). Expectations based on predictable patterns in aperiodic sequences afford

similar behavioral benefits as expectations based on a beat (Bouwer et al., 2020; Heideman et

al., 2018; O'Reilly et al., 2008), but pose a possible challenge for entrainment models, which

are arguably better suited to explain temporal expectations for periodic input (Breska &

Deouell, 2017; Rimmele et al., 2018). Some have suggested that entrainment models can

account for pattern-based expectations by assuming multiple coupled oscillators at different

frequencies and with different phases (Tichko & Large, 2019), or by assuming flexible top-

down phase resets at expected moments, though this would entail some top-down mechanism,

making observed entrainment the consequence, rather than the cause of expectations (Meyer et

al., 2019; Obleser & Kayser, 2019; Rimmele et al., 2018).

Alternatively, however, pattern-based and beat-based expectations could arise from

dissociable neural mechanisms. For cue-based expectations, tentative evidence for a different

underlying mechanism comes from a series of studies looking at the contingent negative

variation (CNV), an event-related potential (ERP) component that peaks at expected moments

(Praamstra et al., 2006). The CNV resolved faster for beat-based than cue-based expectations

(Breska & Deouell, 2017), and cerebellar patients showed selective impairments in forming

cue-based, but not beat-based expectations (Breska & Ivry, 2018, 2020). However, in these

studies, the intended beat-based sequences were isochronous. Isochronous sequences can, in

addition to a beat, elicit temporal expectations through learning the repeated, identical interval

(Breska & Ivry, 2016; Keele et al., 1989). Thus, the differences in responses may here be

explained by more precise cue-based or pattern-based expectations in the isochronous, beat-

based condition. Moreover, these studies tested temporal expectations based on the contingency

between a cue and an interval (e.g., learning a single interval), and it is unclear whether temporal

expectations based on patterns are based on the same mechanism (Nobre & van Ede, 2018). In

our own recent work, we specifically compared beat-based and pattern-based expectations, and

we found no difference in the effects of these expectations on early auditory ERP responses, suggestive of similar modulation of sensory processing (Bouwer et al., 2020), but we observed suppression of sensory processing of unexpected events in beat-based rhythms, even when these events were fully predictable based on their pattern, suggesting different underlying mechanisms (Bouwer et al., 2020). However, in this study, we did not directly probe the neural mechanisms underlying beat-based and pattern-based expectations, rendering it unclear whether they are subserved by shared or separate neural dynamics.

In the current study, we directly examined the role of entrainment in beat-based and pattern-based expectations, using non-isochronous rhythms designed to properly disentangle these two types of expectations. When studying entrainment, an important challenge has been to differentiate between real entrainment ("in the narrow sense", see (Obleser & Kayser, 2019) and regular evoked potentials, or similar phase locked responses that resemble entrainment with common analysis techniques (Zoefel et al., 2018), and that may not differentiate between beat-based and memory-based expectations (Breska & Deouell, 2017). Crucially, to sidestep these issues, here we examined responses in a silent window after cessation of the rhythmic input, directly testing the prediction of entrainment models that entrainment should outlast sensory stimulation (Haegens & Zion Golumbic, 2018; Obleser & Kayser, 2019; Pesnot Lerousseau et al., 2021; Zoefel et al., 2018).

Behaviorally, the effects of persistent entrainment on perception have been shown for auditory rhythm (Hickok et al., 2015; Jones et al., 2002), though this effect is not always found (Bauer et al., 2015; Lin et al., 2021), possibly due to heterogeneity in the population, effects of musical training, and the intricacies of stimulus design (Assaneo et al., 2019; Cameron & Grahn, 2014; Saberi & Hickok, 2022a; Sun et al., 2021). At a neural level, several studies reported persistent entrainment in the visual (de Graaf et al., 2013; Mathewson et al., 2012), and auditory (Kösem et al., 2018; Pesnot Lerousseau et al., 2021; van Bree et al., 2021; Wilsch

et al., 2020) domain (see (Saberi & Hickok, 2022b) for an overview). However, in these studies,

isochronous stimuli were used, making it unclear whether the expectations probed were based

on a beat, or were formed based on the single repeating interval (i.e., cue- or pattern-based).

Moreover, persistent entrainment was not specific to the frequency of the input (Wilsch et al.,

2020), or only occurred in the gamma (Pesnot Lerousseau et al., 2021), or alpha ranges (de

Graaf et al., 2013; Mathewson et al., 2012), while humans have a preference for forming

temporal expectations at slower rates (Ding et al., 2017; Merchant et al., 2015; Zalta et al.,

2020), as naturally present in speech and music (i.e., the delta and theta range). Thus, not only

is evidence for whether entrainment can account for pattern-based expectations lacking,

evidence for persistent neural entrainment in response to beat-based rhythms remains elusive

as well.

In the current study, participants listened to non-isochronous auditory sequences

(similar to those used in (Bouwer et al., 2020)) with either a regular beat (eliciting beat-based

expectations), a predictable pattern (eliciting pattern-based expectations), or random timing (no

expectations). The non-isochronous beat-based sequences had a varying surface structure,

similar to patterns used to probe beat-based processing in many neuroimaging (Grahn & Brett,

2007; Grahn & Rowe, 2009; Leow & Grahn, 2014), behavioral (Bouwer et al., 2018, 2021;

Cameron & Grahn, 2014; Povel & Essens, 1985), and electrophysiological studies (Lenc et al.,

2021). Therefore, the beat could not be extracted from the rhythmic signal by simply learning

the transition of temporal intervals – as is possible in isochronous sequences.

Each sequence was followed by a silent period. In Experiment 1, we asked participants

to rate how well probe tones, presented at various time points during the silent period, fitted the

preceding sequence. We expected the ratings to be affected by both the beat-based and pattern-

based expectations elicited by the sequences. In Experiment 2, we recorded EEG activity both

during presentation of the sequences and during the silence. If entrainment underlies temporal

expectations, we should see persistent power at the frequency of the beat or pattern during the

silence. Alternatively, if climbing activity in the form of a CNV underlies temporal

expectations, we should see a CNV peaking at expected time points in the silence. As a CNV

is generally contingent on a preceding stimulus (Kononowicz & Penney, 2016), we expect the

CNV to only occur once after cessation of the rhythm, with its timing dependent on the one

interval that is expected at that point, and therefore, to be dissociable from persistent power at

frequencies prominent in the overall rhythmic pattern.

In addition to examining spectral power and evoked responses, we explored indexing

temporal expectations in the silence using multi scale entropy (MSE) – a measure of signal

irregularity (Kosciessa et al., 2020) – and multivariate pattern decoding. While MSE provides

us with a method to look at possible non-sinusoidal contributions to the EEG signal, decoding

allows us to look at how entrainment evolves over time. As recently argued, the oscillatory

dynamics underlying temporal expectations may be subject to changes in power and frequency,

depending on coupling between sound and brain, and on the properties of the neural dynamics

themselves (Doelling & Assaneo, 2021). Once the rhythmic sensory input ceases, the

oscillatory activity in the brain may quickly return to an intrinsic resonance frequency (Doelling

& Assaneo, 2021). MSE and decoding may provide useful tools to study these neural dynamics.

Note that we consider these analyses exploratory in nature, and results should be interpreted as

such.

Materials and methods

Participants

Thirty-two participants (18 women), aged between 18 and 44 years old (M = 24, SD = 5.6) took part in the behavioral Experiment 1, and 32 participants (26 women), aged between 19 and 28 years old (M = 23, SD = 2.5) took part in the EEG experiment (Experiment 2), in exchange for course credit or monetary compensation. Due to technical problems, the EEG data from five participants was not recorded correctly, hence we report the results for 27 participants (21 women, between 19 and 28 years old, M = 23, SD = 2.4). For the behavioral experiments, we used mixed-effects models, which need both the number of participants and the number of items to be taken into account to assess power (Brysbaert & Stevens, 2018). In two similar experiments in which ratings in response to rhythms of varying complexity were analyzed, small-sized effects were replicated with a total number of around 275 responses per condition (Bouwer et al., 2018). For our new experimental paradigm, we included a multiple of this number of trials (960 responses per condition and probe position in Experiment 1 – 32 participants with 30 responses each in each cell – and 486 responses per condition and probe position in Experiment 2 – 27 participants with 18 responses). Previous EEG experiments examining persistent entrainment in the auditory domain used sample sizes ranging from fifteen (Pesnot Lerousseau et al., 2021) to twenty-one (van Bree et al., 2021), similar to the sample size used in a study looking at different types of temporal expectations (Breska & Deouell, 2017). Here, we report data from twenty-seven participants, thus exceeded typical sample sizes used previously. None of the participants reported a history of hearing or neurological problems, and all provided written informed consent prior to the onset of the study. The experiment was approved by the Ethics Review Board of the Faculty of Social and Behavioral Sciences of the University of Amsterdam.

Figure 1. Schematic overview of the rhythmic stimuli used and the task. A) Twelve patterns of five temporal intervals with integer ratio durations and an event at each 600 ms period were created to form beat-based sequences. Equivalent patterns without a regular beat every 600 ms were created by using non-integer ratio durations, while keeping the number of intervals and grouping structure the same (random condition). For the pattern-based sequences, only pattern 1 was used, to allow for learning of the intervals. B) Four semi-randomly chosen patterns were concatenated to form rhythmic sequences. In both the beat-based and random sequences, the last pattern was always pattern 1 or 2, to equate the acoustic context preceding the silent period. C) To measure behavioral effects of expectations, a probe tone could appear at various temporal positions in the silent period (indicated by the dashed red lines), predictable based on a beat (B+, light orange), predictable based on the pattern (P+, light purple), or unpredictable based on the beat (B-) or pattern (P-). Subjects had to indicate how well the probe tone fitted the preceding rhythm. In Experiment 1, all 6 probe tone positions were used. In Experiment 2, only the last 3 probe positions were used.

 Stimuli

217   We used patterns marked by woodblock sounds of 60 ms length, generated in

218 GarageBand (Apple Inc.), as previously used in (Bouwer et al., 2020) to elicit beat-based and

219 pattern-based expectations (Figure 1). Each pattern was 1800 ms long and consisted of five

220 temporal intervals. The number of tones was chosen to be within the range that was previously

221 shown to allow for learning of a predictable pattern (Schultz et al., 2013), while the length of

222 the pattern was such that the formation of beat-based expectations with a period of the entire

223 pattern would be unlikely in the pattern-based sequences, given that this would require hearing

224 a beat at 0.55 Hz, which is very far from the range at which humans can typically perceive a

225 beat (Honing & Bouwer, 2019; London, 2012). Sequences (beat-based, pattern-based, or with

226 random timing) were constructed by concatenating four patterns and a final tone, for a total

227 sequence length of 7260 ms (four patterns of 1800 ms, plus 60 ms for the final tone).

228   In the twelve patterns used to create beat-based sequences (Figure 1), temporal intervals

229 were related to each other with integer-ratio durations. The shortest interval had a length of 150

230 ms, with the relation between the five intervals used of 1:2:2:3:4 (i.e., 150, 300, 300, 450, and

231 600 ms). The sounds were grouped such that a perceptually accented sound (Povel &

232 Okkerman, 1981) occurred every 600 ms (every unit length 4), giving rise to a beat at 100 beats

233 per minute, or 1.67 Hz, within the range of preferred tempo for humans (London, 2012). All

234 beat-based patterns were strictly metric, with the beat always marked by a sound (Grahn &

235 Brett, 2007). Sequences of beat-based patterns were constructed from four semi-randomly

236 chosen patterns, with the restriction that the last pattern of the sequences was always pattern 1

237 or 2 (see Figure 1). This way, the final 600 ms preceding the silence epoch was equated in terms

238 of the acoustic context, to make the bleed of auditory ERPs into the silence as similar between

239 conditions as possible. Note that in beat-based sequences, a sound could be expected every 600

ms based on the beat, but the surface structure of the pattern was unpredictable, due to the

random concatenation of patterns.

To create patterns that did not allow for beat-based expectations ("aperiodic" patterns,

see Figure 1), the ratios by which the temporal intervals were related were changed to be non-

integer (1:1.4:1.4:3:5.2, or 150, 210, 450, and 780 ms respectively). In these patterns, no marked

beat was present at unit length four, nor at any other subdivision of the 1800 ms pattern (Bouwer

et al., 2020), while the patterns were matched to their periodic counterparts in terms of overall

length, event density, number of sounds, and grouping structure.

From the aperiodic patterns, two types of sequences were created: pattern-based and

random sequences. To create sequences allowing for pattern-based expectations, we

concatenated four identical patterns. To be able to use the data with an EEG-based decoding

analysis (Experiment 2), we needed the timing of expectations in the silence to be identical for

each sequence, hence we restricted the pattern-based sequences to only pattern 1. The use of a

single pattern was not only necessary for decoding, but also optimized the experiment for

pattern-based expectations, since participants only had to memorize one pattern, allowing them

to easily form expectations, even if the single sequences were only four patterns long.

For the random sequences, four semi-randomly chosen aperiodic patterns were

concatenated. Like for the beat-based sequences, the final pattern was always pattern 1 or 2,

equating the final 600 ms of the sequences in terms of acoustics. In the random sequences, the

timing of sounds could not be predicted based on the surface structure of the pattern, nor on the

basis of an underlying beat.

A spectral analysis of the stimuli confirmed that in the beat-based sequences, a peak

was present at the beat frequency of 1.67 Hz as well as at 3.33 Hz (see Figure 2). The 3.33 Hz

peak is a harmonic of the beat frequency, but also the frequency at which participants may

perceive subdivisions of the beat (e.g., an extra layer of perceived metrical regularity with a

period of 300 ms). In the pattern-based and random sequences, peaks were more distributed, in

line with the more irregular nature of these rhythms, and the highest peaks in the range in which

a beat can normally be perceived (the delta-range, 0.5-4 Hz) were at 2.22 and 3.89 Hz.

Figure 2. Spectral analysis of the sound signal of the different rhythmic sequences. Ten sequences per
condition were generated to base the spectral analysis on. Note that these sequences were identical for the
pattern-based condition, but semi-random for the beat-based and random conditions. The envelope of each
sequence was obtained by performing a Hilbert transform, and subsequently, a fast fourier transform (fft)
was used to obtain the spectral decomposition and power values were averaged over ten sequences. n.a.=
normalized amplitude.
275

In Experiment 1, to assess the persistence of temporal expectations behaviorally, on

each trial, a probe tone was presented at 600, 780, 885, 990, 1200, or 1485 ms after the onset

of the last tone of the sequence (see Figure 1C), and participants provided ratings for how well

probe tones fitted the preceding rhythm. These positions were carefully chosen to represent

times at which a tone could be expected based on the beat (600, 1200 ms), based on memory

for the pattern (780, 990, 1200 ms), or neither (885, 1485 ms). Note that the latter two probe

tones that were unexpected based on the beat (780, 885, 990, and 1485 ms) did not fall on

subdivisions of the beat.

Experiment 2 contained both trials in which a probe tone was presented (25% of the

trials, using only the last three probe positions), and trials in which a 7260 ms sequence was

followed by a silence period without a probe tone (75% of the trials). The latter were used for

EEG analyses, uncontaminated by a probe presentation. The silent period lasted for 2100 ms

after the onset of the last sound, providing 300 ms immediately following the final sound to

allow for ERPs to mostly return to baseline, and 1800 ms (three full beat cycles, or a full cycle

of the repeating pattern) of silence for the analysis. After the silence, the onset of the next

sequence was jittered between 25 and 75 ms to prevent carryover of the beat from previous

sequences (e.g., the next trial started between 325 and 375 ms after the last beat in the silence,

which is not on the beat, nor at a subdivision of the beat). On trials that contained a probe tone,

we chose to only use the last three probe positions, to 1) limit the time for the EEG experiment

to prevent fatigue, 2) still provide participants with the incentive to form expectations well into

the silence period, as a probe tone could appear as late as 1485 ms into the trial, and 3) obtain

some measure of whether participants formed expectations, by including positions that were

expected based on the beat and the pattern (1200 ms), based on the pattern only (990 ms) or

neither (1485 ms).

Procedure

301   Participants were tested individually in a dedicated lab at the University of Amsterdam.

Upon arrival, participants were provided with information about the experiment, provided

written informed consent, and were allowed to practice the task. On probe tone trials (all trials

in Experiment 1, 25% of the trials in Experiment 2), participants were asked to judge on a four-

point scale ("very poorly", "poorly", "well", "very well") how well the probe tone fitted the

preceding sequence, similar to previous studies investigating the perception of musical meter

(Manning et al., 2017; Manning & Schutz, 2013; Palmer & Krumhansl, 1990). Participants

could respond with four buttons on the armrest of their chair, two on each side. The order of

the answer options on the screen in front of the participants was randomized on each trial, to

avoid any artefacts of motor preparation in participants that anticipated which answer they

would provide. There was no time limit for responses and the next trial started after a response

was made.

In Experiment 1, each participant was presented with 18 blocks of 30 trials, amounting

to 540 trials in total (30 trials per probe position for each condition). In Experiment 2,

participants were presented with 18 blocks of 36 trials, for a total of 648 trials (216 per

condition). For each condition, 162 trials were silence trials, and did not contain a probe tone,

and 54 trials contained a probe tone (18 per position). The number of probe trials per block was

varied between 3 and 11. In both experiments, in each block, only one type of sequence (beat-

based, pattern-based, or random) could appear to optimize for the formation of expectations.

Blocks were semi-randomized, with each type appearing once in a set of three blocks.

Sounds were presented through a single speaker positioned in front of the participant

using Presentation® software (version 14.9, www.neurobs.com). After the experiment,

participants filled out the musical training subscale from the Goldsmith Musical Sophistication

Index (GMSI) questionnaire (Müllensiefen et al., 2014). In total, a behavioral session lasted

two hours, and the EEG session lasted between 3.5 and 4 hours.

Behavioral analysis

A total of 17280 responses was included in the analysis of Experiment 1 (32 participants,

3 conditions, 6 probe positions, 30 responses each), and 4374 in the analysis of Experiment 2

(27 participants, 3 conditions, 3 probe positions, 18 responses each). To account for the ordinal

nature of the Likert-scale responses (Bouwer et al., 2018; Carifio & Perla, 2008; Jamieson,

2004), we used a mixed ordinal regression model. With this model, the ordinal responses are

normalized, to correct for potential unequal distances between rating points. The results can

subsequently be interpreted similar to the results from a normal mixed model regression. Two

independent variables were included in the model as fixed factors: Condition (beat-based,

pattern-based, or random), and Probe Position (600, 780, 885, 990, 1200, or 1485 ms; only the

latter three in Experiment 2). Additionally, GMSI score was included as a continuous variable

(Musical Training), as well as interactions between all factors, and a random intercept for each

subject to account for between-subject variation.

The initial model showed a significant effect of Probe Position in the random condition,

most likely due to recency effects. To assess the effect of Probe Position in the beat-based and

pattern-based conditions while accounting for this, for each participant we subtracted the mean

response in the random condition at each position from the responses in the beat-based and

pattern-based condition. Subsequently, we used these baseline-corrected ratings in a second

ordinal regression model, with only two levels for the factor Condition (beat-based and pattern-

based) and without the random intercept for each participant (as the baseline correction already

corrected for between-subject variability). For both the original model, and the baseline

corrected model, significant interactions were followed up by tests of simple effects, corrected

for multiple comparisons using a Bonferroni correction.

The statistical analysis was conducted in R (R Development Core Team, 2008). The

ordinal mixed model was implemented using the clmm() function from the ordinal package

(Christensen, 2019). Subsequently, we used the Anova() function from the car package (Fox &

Weisberg, 2019) to look at omnibus effects for the main factors of interest, and the emmeans

package (Lenth, 2019) to assess simple effects and compare slopes between conditions.

EEG recording

EEG was recorded at 1024 Hz using a 64-channel Biosemi Active-Two acquisition

system (Biosemi, Amsterdam, The Netherlands), with a standard 10/20 configuration and

additional electrodes for EOG channels (located under and on the left and right sides of the

eye), on the nose, on both mastoids, and on both earlobes.

EEG analysis

Preprocessing was performed in MATLAB, version 2015a (Mathworks) and EEGLAB,

version 14.1.1 (Delorme & Makeig, 2004). Data were offline down-sampled to 256 Hz, re-

referenced to averaged mastoids, bad channels were manually removed, and eye-blinks were

removed using independent component analysis. Subsequently, bad channels were replaced by

values interpolated from the surrounding channels.

ERPs. For the ERP analysis, the continuous data were filtered using 0.1 Hz high-pass

and 40 Hz low-pass finite impulse response filters (as implemented in the standard EEGLAB

filter function pop_eegfiltnew). Epochs were extracted from the data from -1800 till 2100 ms

relative to the onset of the last sound. Epochs with a voltage change of more than 150 microvolts

in a 200 ms sliding window were rejected from further analysis. For each participant and

condition, epochs were averaged to obtain the ERPs, and ERPs were averaged over participants

to obtain grand average waveforms for plotting. All waveforms were initially baseline corrected

using the average voltage of a 50 ms window preceding the onset of the last sound of the

sequence. This baseline can be regarded as preceding the "cue" (the last event before the onset

of the expectation). Such a baseline is customary in CNV analyses. However, visual inspection

suggested that this baseline was biased, as baseline correction resulted in an overall shift of the

waveform amplitude relative to each other, as also reflected in a significant cluster when

comparing beat-based and pattern-based conditions that spanned the entire analysis epoch (see

Appendix, Figure A1). This was likely caused by the rapid succession of sounds preceding the

onset of the silence, which made it impossible to find a clean, unbiased baseline. Therefore, we

repeated the analysis without baseline correction, to confirm that the results were not caused by

a noisy baseline.

As a control analysis, we repeated the above analysis for the longest intervals during the

presentation of the rhythmic streams, to assess whether in those longer intervals, we could

observe similar deflections in the evoked potentials as observed in the silence. For this analysis,

epochs were extracted from -200 till 880 ms around the onset of the sound preceding the long

interval (600 ms in the beat-based condition, and 780 ms in the other conditions, see Figure 1).

All preprocessing steps were identical to the analysis of the ERPs in the silence. The control analysis is reported in the Appendix (Figure A1).

Three cluster-based permutation tests (Oostenveld et al., 2011) were used to compare all three conditions against each other (i.e., beat-based vs. random; pattern-based vs. random; beat-based vs. pattern-based), comparing all timepoints from 300 till 1200 ms after the onset of the final sound for the silence (see Figure 1), and all timepoints from 300 till 600 ms after the onset of the preceding sound for the long intervals during the sequences (as the next sound came in at 600 ms for the beat-based condition, we could not compare the conditions beyond this timepoint). This window excluded a large portion of the ERP response to the previous sound, and included both the first expected moments for beat-based (600 ms) and pattern-based (780 ms) expectations in the silence window, and additional time to allow for an evaluation of possible return to baseline of the CNV (Breska & Deouell, 2017). For the ERP analysis, clusters were formed based on adjacent time-electrode samples.

For all EEG analyses, cluster-based tests were evaluated statistically by forming clusters of samples based on dependent samples t-tests and a threshold of $p < 0.05$, and using permutation tests with 2000 permutations of the data. We report corrected p-values to account for two-sided testing (multiplied by a factor of two).

Frequency-domain analysis. To obtain the spectrum of the EEG signal in the silence, we used the raw, unfiltered data. Epochs were extracted from the continuous data both from -1800 till 0 ms relative to the onset of the last sound (control window, see Figure 1), and from 300 till 2100 ms relative to the onset of the last sound (silence window, see Figure 1), the latter starting at 300 ms to avoid contamination from the final ERPs. Both windows thus had equal length, both spanning three full cycles of the beat. Epochs with an amplitude change of 250 microvolts or more in a sliding 200 ms window were rejected from further analysis. The more lenient rejection criterium compared to the ERP analysis was used to account for the fact that

these data were unfiltered, and to avoid rejection of too many trials that showed some slow drift. All epochs were baseline corrected using the mean of the entire epoch. Subsequently, epochs were averaged for each condition separately to obtain the evoked signal, phase locked to the onset of the final sound, and similar to previous studies using frequency tagging to look at beat-based perception (Lenc et al., 2021; Nozaradan et al., 2011, 2012).

For each participant and condition separately, the average waveforms were transformed into the frequency domain using an FFT, with the data zero-padded to 4608 samples (NFFT) to obtain a better frequency resolution (0.056 Hz), and importantly, be able to extract data at exactly the frequencies of interest. Note that the zero-padding can only improve the frequency resolution, but not the frequency precision, which by definition with the 1800 ms epochs is limited to 0.56 Hz. While the design of the experiment simply does not allow for a better resolution, the 0.56 Hz does allow us to differentiate between the frequencies of interest, which are 0.56 Hz or more apart. The obtained power values at each frequency were normalized to account for the 1/f distribution of noise (Nozaradan et al., 2011, 2012), by subtracting the average of neighboring bins four to six on either side for all frequencies (e.g., 1.33 – 1.44 Hz and 1.89 – 2.00 Hz for the beat frequency, 1.89 – 2.00 Hz and 2.44 – 2.56 Hz for the pattern-based frequency, and 3.00 – 3.11 Hz and 3.56 – 3.67 Hz for the beat subdivisions). To account for bleeding into neighboring frequency bins (Nozaradan et al., 2011, 2012), for each frequency, we averaged over 5 bins centered on that frequency (e.g., for the frequencies of interest, this was 1.56 – 1.78 Hz for the beat frequency, 2.11 – 2.33 Hz for the pattern-based frequency, and 3.22 – 3.44 Hz for the beat subdivisions).

To statistically test differences between conditions in the evoked power at the frequencies of interest, we used cluster-based permutation tests. First, this avoided bias by selecting only a subset of electrodes, as we used all scalp electrodes, as was done in previous research (Lenc et al., 2018; Nozaradan et al., 2011; Tal et al., 2017). Second, the permutation

tests accounted for the non-normal distribution of the data. Like for the ERPs, we ran t-tests

comparing the normalized data for all conditions and for each frequency of interest (e.g., those

most prominent in the sound signal). We included the frequencies that showed the highest peaks

in the spectral analysis of the sound (i.e. 1.67 Hz, 2.22 Hz, and 3.33 Hz), except for 3.89 Hz,

since a peak at this frequency was absent on visual inspection in the spectral decomposition of

the EEG data. For the frequency-domain analysis, clusters were formed based on adjacent

electrodes.

The cluster-based tests yielded null results for the pattern-based condition in the silence

(e.g., there was no larger power at 2.22 Hz in the pattern-based than in the random condition,

see Results). To quantify the possible absence of persistent entrainment for the pattern-based

condition, we performed a Bayesian t test using JASP (JASP, 2019; Wagenmakers et al., 2018).

We compared the power in the pattern-based and random conditions at 2.22 Hz in the silence,

averaged over electrodes contributing to the significant cluster in the silence for the beat-based

condition at 1.67 Hz, to optimize for finding entrainment effects. We estimated Bayes factors

using a Cauchy prior distribution (r =.71) and performed a robustness check to assess the effect

of a different prior (r = 1; see (Jeffreys, 1961; Wagenmakers et al., 2018)).

Multiscale entropy (MSE). MSE is a measure of signal irregularity. To compute MSE,

the EEG signal is divided into patterns of a certain length, and throughout the signal, the number

of repeating patterns is counted. More repetitions indicate a more regular signal, and yield a

lower entropy value. By calculating entropy for patterns of different lengths ("multiscale"), the

contributions of slower and faster timescales in the signal can be assessed. However, the

mapping between entropy timescales and spectral frequencies is not absolute, especially since

entropy is not per se related to a signal being oscillatory in nature (Kloosterman et al., 2020;

Kosciessa et al., 2020). The advantage of using MSE is that it does not require filtering of the

data, and it does not assume stationarity (e.g., it can pick up on regularities that are

asymmetrical, or that do not have a fixed amplitude or period). Here, we computed MSE on the

control and silence epochs separately.

We computed MSE on high-pass filtered data (0.5 Hz). Epochs were extracted identical

to the epochs for the frequency domain analysis. To compute MSE, we used the mMSE toolbox,

a plugin to the Fieldtrip toolbox (Kloosterman et al., 2020), with $m = 2$ and $r = 0.5$, as was

done previously for EEG data (Kloosterman et al., 2020). For details on how MSE is computed,

see the Appendix. As for the frequency-domain analysis, we used cluster-based permutation

tests to assess statistical significance. For each comparison between conditions, we used paired

t-tests comparing each electrode-timescale combination (note that the above computation of

entropy yields one value per condition for each electrode and timescale) to form clusters.

Multivariate decoding. Our exploratory decoding approach is based on the assumption

that temporal expectations are always coupled with feature or spatial expectations (e.g., we

cannot predict "when" without also predicting "what"), as suggested by studies showing that

we only use temporal expectations to improve perception if we can also predict the content

(Morillon et al., 2016; Wollman & Morillon, 2018) or location (O'Reilly et al., 2008) of an

upcoming event. Thus, we expected to be able to decode the representation of the expected

sound at expected moments. As the expected moments are different for each condition, this

then allows us to decode in the silence window whether participants were previously listening

to a beat-based, pattern-based, or random sequence.

The decoding was conducted on data that was preprocessed in a similar way as for the

MSE analysis, but with epochs extending from -1800 to 2100 ms relative to the onset of the last

sound. Since the decoding is done in a time resolved way (e.g., sample by sample), there is no

need to leave out the response to the ERPs in the analysis. Additionally, the data were resampled

to 32 Hz to increase signal to noise, using shape-preserving piecewise cubic interpolation, as

implemented in the Fieldtrip toolbox (Oostenveld et al., 2011). Using the ADAM toolbox

(Fahrenfort et al., 2018), we applied a classification algorithm to the preprocessed data for each

participant. Using all electrodes, each dataset was split into 10 equally sized subsets, for 10-

fold cross-validation of the decoding. For each subset, a linear discriminant classifier trained

on the remaining 9 subsets was tested. Each condition was decoded against both other

conditions (e.g., beat-based vs. pattern-based, beat-based vs. random, and pattern-based vs.

random), creating a temporal generalization matrix of classification accuracy at each possible

combination of training and testing time points (King & Dehaene, 2014). Subsequently, we

examined whether we could observe a pattern of recurrent activity (King & Dehaene, 2014).

Classification accuracies averaged over the 10 folds for each comparison of two conditions for

the silence window (300-2100 ms after the onset of the last sound) were submitted to cluster-

based permutation tests to assess whether they exceeded the chance level of 0.5. Clusters were

based on T-tests with a threshold of 0.05 for each training-testing time point combination,

comparing the accuracy to 0.5.

The initial decoding analysis yielded large effects that may be task-related (see Results

for an explanation of these results). Therefore, we ran an additional decoding analysis in which

we decoded expected and unexpected positions against each other within each condition.

Details on this additional analysis can be found in the Appendix.

Results

Behavioral effects of beat-based expectations last multiple beat cycles, while those of pattern-based expectations reflect one interval

Figure 3A shows the average ratings for each condition and probe position from Experiment 1. Visual inspection of this figure suggests that beat-based expectations were associated with higher fitness ratings for sounds at expected times than unexpected times for two beat cycles in the silence window (at 600 and 1200 ms), while the effects of pattern-based expectations appeared to reflect mainly the first expected time point in the silence window (780 ms). This was confirmed by our statistical analyses. The ordinal regression showed main effects of Condition ($\chi^2(2) = 300.72$, $p < 0.001$), Position ($\chi^2(5) = 1067.05$, $p < 0.001$), as well as Musical Training ($\chi^2(1) = 5.16$, $p = 0.023$). However, crucially, these main effects were accompanied by a very large two-way interaction between Condition and Position ($\chi^2(10) = 2478.98$, $p < 0.001$), showing that the effects of beat-based and pattern-based expectations on fitness ratings differed, depending on the position of the probe. We found additional smaller interactions between Position and Musical Training ($\chi^2(5) = 67.26$, $p < 0.001$), Condition and Musical Training ($\chi^2(2) = 6.01$, $p = 0.05$), and, interestingly, Condition, Position, and Musical Training ($\chi^2(10) = 204.88$, $p < 0.001$). Following the interactions, tests of simple main effects showed that the effect of Position was significant in all conditions (all ps < 0.001), and the effect of Condition was significant for all probe positions (all ps < 0.001). The main effect of Position in the random condition showed that even after sequences in which no specific temporal structure was present, ratings depended on the position of the probe. This likely was due to recency effects. To account for these effects when comparing the ratings at different positions in the beat-based and pattern-based conditions, we subtracted the ratings in the random condition at each position from the ratings in the other conditions. The baseline corrected model (Figure 3B) showed similar interactions between Position and Condition

$(\chi^2(5) = 2147.55, p < 0.001)$, and between Position, Condition, and Musical Training $(\chi^2(5) =$

154.46, $p < 0.001$). The latter indicated that the correlation between Musical Training and the

rating score depended on both the Position and the Condition of the probe tone.

Figure 3. The effects of beat-based expectations on fitness ratings can be differentiated from those of pattern-based expectations, and are associated with musical training. A) Mean ratings for all conditions and positions. Colored asterixis indicate positions where ratings in the beat-based (orange) and pattern-based (purple) conditions differed from the random condition at $p < 0.05$ (Bonferroni corrected). B) Single participant data, with the random condition subtracted to account for serial position effects. The expectedness pattern is indicated by colored lines on the bottom of the plots (orange: expected based on the beat; purple: expected based on pattern; grey: neither). For the beat-based condition, ratings followed this pattern for two beat cycles. For the pattern-based condition, ratings followed the pattern for one interval. C) Data median split based on scores on the musical training questionnaire. The pattern of results, while present for both groups of participants, is enhanced for the group of participants with most musical training ("experts"). Note: the median split is for visualization purposes only, the models were run with musical training as a covariate. Error bars in panels A-C are 2 standard errors (note: these are computed on the complete dataset, not the participant averages, as the ordinal model is run on trial-level data). D) Association between musical training and rating for each condition and position. Colored asterixis show positions in which the association between musical training and the ratings was significantly correlated ($p < 0.05$, Bonferroni corrected). A positive association was observed for the beat-based condition at 600 and 1200 ms (expected positions) and for the pattern-based condition at 780 (expected) and 885 ms (unexpected). Negative associations were observed in the beat-based condition at 780, 885, and 990 ms (all unexpected), and in the pattern-based condition at 990 (unexpected) and 1200 ms (expected).

Beat-based expectations. At both 600 ms and 1200 ms (expected in terms of a beat), probes in the beat-based condition were rated as better fitting than probes in the random condition ($p < 0.001$ and $p = 0.004$) as evident from the full model; all simple effects from the full model can be found in the Appendix, Table A1). At 780, 885, 990, and 1485 ms (unexpected in terms of a beat), probes in the beat-condition were rated as worse fitting than probes in the random condition (all ps < 0.001). Moreover, within the beat-based condition, at 600 ms, baseline corrected ratings were higher than at any other probe position (all ps < 0.001), and at 1200 ms baseline corrected ratings were higher than at 780, 885, 990, or 1485 ms (all ps < 0.001). Baseline corrected ratings for probes at 780, 885, and 990 ms (all unexpected in terms of the beat) did not differ from each (all ps > 0.93). Probes at 1485 ms (unexpected in terms of the beat) were rated as better fitting than probes at 780, 885, and 990 ms (all ps < 0.001). All simple effects from the corrected model can be found in the Appendix, Table A2.

As can be seen in Figure 3C and 3D, higher scores on the Musical Training questionnaire were associated with higher fitness ratings in the beat-based condition at 600 and 1200 ms (expected in terms of the beat), but lower fitness ratings at 780, 885, 990, and 1485 ms (unexpected in terms of the beat). In other words, musically trained participants were better able to differentiate between probes that were in expected and unexpected positions (Figure 3C). Slopes reach significance at all positions except 1485 ms (all ps < 0.022). Also, the association between Musical Training and ratings differed between beat-based and pattern-based conditions, at 600, 780, 885, and 1200 ms (all ps < 0.004).

To sum up, for the beat-based sequences, we could observe a clear pattern in the results indicating that beat-based expectations were used to rate the probes well into the silence window, affecting ratings up to 1200 ms after the onset of the last sound. Beat-based expectations lead to higher ratings for expected probes (600 and 1200 ms), and lower ratings for unexpected probes (780, 885, 990, and 1485 ms), both when comparing ratings for each

position to the random condition, and when comparing ratings for each position within the beat-based condition. At 1200 ms, these effects resulted in a classic inverted U-curve, as previously associated with beat-based processing (Bauer et al., 2015; Jones et al., 2002), with optimal performance on the beat, and diminished performance on either side (e.g., both earlier and later). The effects of beat-based expectations did diminish over time, as is apparent from differences between ratings at 600 and 1200 ms, and at 1485 ms and other unexpected time points. Both the enhancing and attenuating effects of beat-based expectations were correlated with musical training. It is worth noting that the longer lasting effects of beat-based expectations (at 1200 ms) were very heterogenous in our participant pool. Out of 32 participants in Experiment 1, only 18 showed the inverted U, with higher ratings at 1200 than at 990 and 1485 ms.

Pattern-based expectations. For pattern-based sequences, ratings at 600 ms (unexpected based on the pattern) were lower than for the random sequences ($p < 0.001$) and lower than at any other position (all ps < 0.001, baseline corrected model), showing that participants also formed predictions based on the sequences. In line with this, at 780 ms (expected in terms of the pattern), ratings were numerically higher in the pattern-based condition than in the random condition, though this difference did not survive the Bonferroni correction ($p = 0.058$). After this point, ratings did not differ between pattern-based and random conditions for probes at 885 (unexpected) and 990 (expected) ms, suggesting that the responses followed the rhythmic pattern mainly in the beginning of the silent period. In line with this, in the remainder of the silence window, the ratings continued to deviate from what would be predicted based on the pattern, with lower ratings for the pattern-based than random condition at 1200 ms (expected; $p < 0.001$). Ratings were also lower for the pattern-based than random condition at 1485 ms (unexpected, $p < 0.001$), and ratings at 780 ms (expected) did not differ from ratings at 885 ms (unexpected), while being marginally higher than at 990 (expected, $p =$

0.080), and higher than at 1200 (expected), and 1485 (unexpected) ms (both ps < 0.032). In

addition, ratings at 885 and 990 ms were higher than at 1200 and 1485 ms (all ps < 0.002). See

the Appendix, Tables A1 and A2, for all simple effects.

As for beat-based expectations, for pattern-based expectations, there was a positive

association between ratings and musical training at an expected time point (780 ms; p < 0.001),

and a negative, albeit nonsignificant, association at an unexpected time point (600 ms). Thus,

like for beat-based expectations, at these early time points, musicians were better able than non-

musicians at differentiating between expected and unexpected moments in time. However, at

885 ms (unexpected in terms of the pattern), the results behaved like at 780 ms, with higher

ratings associated with more Musical Training (p = 0.002). At 990 and 1200 ms, Musical

Training was associated with lower ratings (both ps < 0.05), but these results are

counterintuitive, as these are expected positions based on the pattern.

The results for pattern-based expectations suggest that just like for beat-based

expectations, participants were able to predict the timing of probes based on the preceding

sequence. However, the results show that while this was still the case at 780 ms after the onset

of the last tone, at later probe positions, the effects of pattern-based expectations did not reflect

the preceding sequence. At 885 ms, the results, both in terms of the ratings and how they were

associated with musical training, behaved similar to at 780 ms. After this point, the results

suggest that participants did not use the preceding sequence to guide their responses, but

instead, used a different heuristic.

Figure 4 shows the behavioral results obtained from the EEG experiment. Replicating

Experiment 1, we found main effects of Condition ($\chi^2(2)$ = 92.30, p < 0.001) and Position

($\chi^2(2)$ = 49.36, p < 0.001), accompanied by interactions between Condition and Position ($\chi^2(4)$

= 96.62, p < 0.001), and Condition, Position, and Musical Training ($\chi^2(4)$ = 11.04, p = 0.03).

Following the analysis strategy from Experiment 1, we subtracted the ratings from the Random

condition from the ratings for the other two conditions, yielding a baseline corrected model

with similar interactions (Condition and Position: $\chi^2(2) = 31.46$, $p < 0.001$; Condition, Position,

and Musical Training: $\chi^2(2) = 8.29$, $p = 0.02$).

Figure 4. Behavioral results during the EEG experiment replicate findings from Experiment 1. Even
with only 18 trials per participant per condition and position, we could replicate the inverted U-curve for
beat-based sequences at the end of the silence epoch. Like in Experiment 1, the results for the pattern-based
condition do not follow the pattern, but instead, are consistent with building expectations for the next trial.
A) Mean ratings for all conditions and positions. Colored asterixis indicate positions where ratings in the
beat-based (orange) and pattern-based (purple) conditions differed from the random condition (with $p < 0.05$,
Bonferroni corrected). B) Single participant data, with the random condition subtracted to account for serial
position effects. C) Data median split based on scores on the musical training questionnaire. Error bars in
panels A-C are 2 standard errors. D) Association between musical training and rating for each condition and
position. See Figure 3 for more details.

In line with the preceding beat-based sequences, probes at 990 and 1485 ms (both

unexpected times based on the beat) were rated lower in the beat-based condition than in the

random and pattern-based conditions (all ps < 0.012), and within the beat-based condition,

probes at 1200 ms were rated as better fitting than at 990 ($p < 0.001$) and 1485 ms, though the

latter difference did not reach significance. Thus, like in Experiment 1, we found an inverted

U-curve at 1200 ms after the final tone, suggestive of beat-based expectations lasting at least

two beat cycles. As in Experiment 1, at 1200 ms (expected based on the beat), higher ratings in the beat-based condition were associated with more musical training ($p = 0.047$), suggesting that the effects of beat-based expectations correlate with musical expertise. Additionally, probes at 990 ms were rated lower than at 1485 ms ($p < 0.001$), possibly because the effects of beat-based expectations diminished over the course of the silence.

In the pattern-based condition, ratings did not follow the pattern of the preceding sequences. At 990 ms (expected based on the pattern), probes were rated as worse fitting than in the random condition ($p < 0.001$), and as worse fitting than at 1200 (expected) and 1485 (unexpected) ms (both ps < 0.023), while at 1485 ms (unexpected), probes were rated as better fitting than in the random condition, and as better fitting than at 1200 (expected) ms (both ps < 0.001). Also, at 1200 ms (expected), higher ratings were associated with less musical training for the pattern-based condition (though after the Bonferroni correction, only marginally so: $p = 0.06$), contrary to what would be expected if the effects of expectations are enlarged in musical experts. All simple effects can be found in the Appendix, Tables A1 and A2.

Thus, the behavioral results from Experiment 2, though based on less trials than Experiment 1, suggest a similar pattern as found in Experiment 1: while beat-based expectations exert their effect well into the silent period, with participant faithfully following the beat in their goodness-of-fit ratings, pattern-based expectations do not affect ratings in the second half of the silent period in a manner consistent with the learned pattern. Albeit speculatively, the results for the pattern-based expectations may be more in line with expectations for the start of the next trial leading to higher ratings for probe positions closer to the end of the silent period, as in Experiment 2, after non-probe trials, the next trial followed each silent period at a somewhat predictable time.

Figure 5. Beat-based and pattern-based can be differentiated based on ERPs. A) Left panel show the grand average waveforms for the silence window for a central electrode cluster (FC1, FCz, FC2, C1, Cz, C2, CP1, CPz, CP2). Time 0 is the onset of the last tone of the sequence. Colored bars on the bottom of the plots, and vertical orange and purple lines, indicate at which times a tone would be expected based on the beat (light orange) and the pattern (light purple). Note that these are expected times, but no sounds were played during the window shown after time 0. Sounds during the sequence are depicted in dark orange and purple. B) The scalp distributions for windows in which a significant cluster was observed.

Differences in the evoked potential elicited by beat-based and pattern-based sequences

Figure 5 shows the average ERPs for each condition, without baseline correction (see Materials and Methods), and scalp topographies for windows in which we found significant clusters. In line with previous research, we may expect climbing neuronal activity, or a CNV, flexibly adapting its slope to peak at the moment that participants expect the next event (Breska & Deouell, 2017; Breska & Ivry, 2020; Damsma et al., 2021; Mento, 2013). We did observe differences in the ERPs between conditions, but the peak of the differences was not at the expected time, but rather, fell earlier (see the Appendix, Figure A1, for the difference waveforms between conditions). Without baseline correction, the random condition elicited a significantly more negative ERP ($p = 0.011$) than the beat-based condition in a frontocentral cluster between 300 and 614 ms after the onset of the last sound (though note that we did not include timepoints preceding 300 ms in the cluster-based tests). Likewise, in the same latency range ($300 - 473$ ms), there was a trend for the pattern-based condition to elicit a more negative ERP than the beat-based condition ($p = 0.077$). Thus, in a window between approximately 300 and 450 ms, we found tentative evidence for more negative-going waveforms in both the

random and pattern-based condition compared to the beat-based condition, with a central scalp

topography (Figure 5B). In a later window, the pattern-based condition elicited a second

negative deflection, which showed a trend to be larger than in the beat-based condition ($p =$

0.077, 864 – 1005 ms). While these results were somewhat different depending on the choice

of baseline (see Figure A1 in the Appendix for the results with a traditional pre-cue baseline),

the overall picture is the same, with significant clusters in an early and later window indicating

differences between conditions in the ERPs.

The time course of the effect deviated from what we expected: the negative deflection

did not peak at the next expected moment in time, but rather, peaked much earlier. Also,

contrary to previous research (Breska & Deouell, 2017), the beat-based condition elicited the

most positive-going ERP, instead of a typical CNV. To further explore and confirm these

results, as a control analysis, we performed the same ERP analysis on the longest time intervals

during the sound presentation. The waveforms showed a negative deflection very similar in

morphology and scalp distribution to the one we found in the silence window (see Figure A1

in the Appendix). During the sound presentation, like in the silence, this negative deflection

was largest for the pattern-based condition, though the difference was only significant when

comparing the pattern-based condition to the random ($p = 0.039$) but not the beat-based

condition ($p = 0.1$). The latter non-significant result may be due to a noisy baseline, as during

sound presentation, the succession of intervals in the beat-based and random sequences was

(semi-)randomly chosen, and therefore, the baseline was not consistent over conditions.

Overall, however, this control analysis yielded very similar results to the analysis of ERPs in

the silence window.

Figure 6. Power at the beat frequency persists during the silence window. In the control window, during auditory stimulation, peaks in power can be observed at all frequencies of interest, and for the relevant conditions (1.67 and 3.33 Hz in the beat-based condition, and 2.22 Hz in the pattern-based condition). In the silence window, the peaks in power at 1.67 and 3.33 Hz were larger in the beat-based condition than in the pattern-based and random conditions, while peaks at 2.22 Hz did not differ between conditions. Note: the raw data is depicted here, before the normalization procedure. Data shown is averaged over all electrodes.

Frequency-domain analysis shows persistent power at the beat frequency following beat-based sequences

A strong prediction of entrainment theories is that the entrainment outlasts stimulation with the entraining stimulus. Therefore, we next looked at the frequency content of the EEG signal in the silent period. Specifically, we predicted that if entrainment occurs, we would find enhanced power at the frequencies associated with the rhythmic sounds to be found in the silence. In Figure 6A, the average power for all electrodes, separated for each condition in the control (i.e. during the auditory sequence) and silence windows is depicted as a function of frequency. In the control window (Figure 6, left, and Figure 7, top), the frequency response followed the sound input. That is, at the beat frequency (1.67 Hz), significant clusters indicated higher power in the control window for the beat-based sequences than the pattern-based ($p <$ 0.001) and random ($p =$ 0.006) sequences. At 2.22 Hz, prominent in the pattern-based and random sound sequences, higher power was observed in the EEG signal in the pattern-based than beat-based ($p =$ 0.02) and random ($p =$ 0.023) sequences, and higher power was observed in the random than beat-based sequences ($p =$ 0.035). Finally, at 3.33 Hz (subdivisions of the beat), power in the control window was larger for the beat-based than pattern-based ($p =$ 0.013)

and random ($p = 0.005$) conditions. Thus, in the control window, the EEG signal reflected the

spectral properties of the sound signal, which is likely caused by each sound eliciting an ERP,

which are represented in steady-state potentials, and thus picked up by the frequency analysis

(Keitel et al., 2021), especially since the ERPs elicited by our stimuli extended for hundreds of

milliseconds, and our analysis focused on frequencies >1 Hz (Zhou et al., 2016).

Importantly, during the silence (Figure 6, right, and Figure 7, bottom), no significant

clusters were found at 2.22 Hz, but at the beat frequency (1.67 Hz), power was significantly

larger for the beat-based than the pattern-based ($p = 0.01$) and random ($p = 0.012$) conditions.

In addition, at 3.33 Hz, power for the beat-based condition was larger than for the pattern-based

condition ($p = 0.038$), with a trend when comparing the beat-based with the random condition

($p = 0.077$). Thus, while the pattern-based and random conditions showed tracking of the sound

during stimulation (which is sometimes considered entrainment "in the broad sense" (Obleser

& Kayser, 2019)), in the silence, entrainment was only present for the beat-based condition.

This finding fits our behavioral observations that differentiate qualitatively between the effects

of beat-based expectations and pattern-based expectations in the silence. To further substantiate

the absence of persistent entrainment at the pattern frequency in the silence, we performed a

Bayesian T-test comparing the normalized power at 2.22 Hz between the pattern-based and

random conditions. We found moderate evidence in favor of the null hypothesis (no difference

between conditions) ($BF_{01} = 4.5$). The results did not change as a function of the prior used

(with a more traditional prior of $r = 1$, $BF_{01} = 6.17$).

Note that like for the behavioral results indicative of entrainment, there was large

heterogeneity between participants (see Figure 7). While the power differences were significant

in the overall cluster-based analyses, out of 27 participants, only 16 showed on average (i.e.

over all electrodes) numerically larger power in the beat-based condition at the beat frequency

when compared to both the random and pattern-based condition.

Figure 7. Individual participant data shows heterogeneity of the effects of temporal expectations on spectral power. All plots depict the spectral power at the frequency of interest, averaged over all electrodes, and normalized to account for the 1/f distribution (see Materials and Methods). A) Single participant data, with all data points in grey. Boxplots show the median, with the lower and upper hinges corresponding to the first and third quartiles (25th and 75th percentiles), and the whiskers corresponding to values no further than 1.5 times the inter-quartile range from the hinges. Error bars depict 2 standard errors around the mean. B) Scalp topographies for all conditions. Electrodes contributing to significant clusters in the cluster-based tests are highlighted on the plots for the conditions from each comparison in which power was largest.

Figure 8. Entropy was higher for pattern-based than random sequences in both control and silence windows. Entropy in the Control window (left) and Silence window (right), averaged over frontocentral electrodes, and scalp distributions averaged over all timescales, depicting electrodes contributing to significant clusters.

Multiscale entropy as a non-stationary measure of temporal expectations

Figure 8 shows sample entropy for each condition separately, as well as the electrodes contributing to significant clusters in the analysis. Given that MSE indexes signal irregularity, we would expect entropy to be higher for the random and pattern-based conditions than for the beat-based condition in the silence. A cluster-based test on all electrodes and timescales showed that both in the control window, and in the silence window, entropy was higher for the pattern-based than for the random condition (control: $p = 0.03$; silence: $p = 0.021$). Note that in the silence window, all but the two highest timescales were included in the cluster. In the control window, the cluster spanned all timescales from 35 till 406 ms (see Materials and Methods for an explanation of the timescales). These results suggest that the signal was more irregular in the pattern-based than the random condition, over a broad range of timescales. Neither the difference in entropy between the beat-based and pattern-based condition, nor between the beat-based and random condition was significant, in either control or silence windows (all ps > 0.24).

Entropy has been related to various other EEG measures, such as spectral power and overall differences in signal variability (Kosciessa et al., 2020). To account for these, we ran several additional analyses. First, to check whether the differences in signal variability may

have been caused by differences in low frequency activity, we repeated the MSE analysis on

high-pass filtered data (Kloosterman et al., 2020), using a 5 Hz high-pass filter. This completely

removed the effects (all ps > 0.43), suggesting that the differences between conditions were

caused by low frequency activity in the signal (see Figure A2 in the Appendix).

Second, we checked whether differences in signal variability caused the differences in

entropy, since entropy is calculated relative to the overall signal standard deviation (e.g., a

pattern is considered a match at a lower threshold when overall signal variability is high). As

can be seen in the Appendix (Figure A3), the similarity bounds used to compute entropy

(derived from the time-domain signal standard deviation) differed between conditions, and this

difference mirrored the differences in entropy, suggesting that at least some of the variance we

observed was due to overall signal variability, and not necessarily signal irregularity.

Multivariate decoding as a time-resolved method for studying entrainment

With multivariate decoding, we expected that training at expected times would yield

above chance performance when testing at expected times, regardless of whether these time

points were the same (e.g., when training at 600 ms, we expected to be able to accurately

distinguish the beat-based from the random condition when testing at not just 600 ms, but also

at 1200 and 1800 ms, as all these times were on the beat, or similarly expected). Figure 9A

shows the temporal generalization matrices for each comparison, with significant clusters

indicated by a black contour. In the silence window we found above chance decoding when

decoding the beat-based against the random condition ($p = 0.043$), the pattern-based against the

random condition ($p < 0.001$), and the beat-based and pattern-based conditions against each

other ($p = 0.009$), indicating that based on the EEG signal in the silence, we could classify

which type of rhythm participants had heard just before. However, looking at the temporal

generalization matrices, it becomes apparent that contrary to our expectations, this above-

chance decoding was not due to recurrent activity for expected events. Only clusters on the

diagonal were significant for each comparison. In addition, decoding for all three comparisons

was best in the second halve of the silence window, which is where the probe tones were

presented. This suggests that the decoding mainly picked up on task-related differences. While

the probes were physically identical across conditions, participants may have had different

strategies to perform the task, depending on the type of sequence, and this could have resulted

in above-chance decoding related to the task and probes, even when only analyzing the silence

window.

Figure 9. No recurrent activation in group average decoding, example of recurrency in single
participant. A) Temporal generalization matrices for each comparison for the group analysis. The Y-axis
shows the training time points, and the X-axis testing time points. The color scale indicates the classification
accuracy, with 0.5 being chance level (all analyses are based on decoding two conditions against each other).
Significance of classification accuracy was assessed by using cluster-based permutation tests on the silence
window (300-2100 ms after the last note onset). The black contour indicates significant clusters. Note that
the upper right quadrant, highlighted by the black box, is the entire silence window. B) Temporal
generalization matrix for a participant showing recurrent activity following the expected beats. When
decoding the beat-based vs. the other two conditions (left- and rightmost plots), peaks in accuracy follow a
clear oscillatory pattern, with a phase consistent with the beat-based sequence (600 ms between beats and
decoding peaks).

Of note, the decoding results varied considerably between participants. In one

participant in particular (Figure 9B), we observed a pattern in the temporal generalization

matrix that was consistent with the hypothesized result for an oscillatory process (King &

Dehaene, 2014). However, even though some other participants also showed recurrent activity,

the exact times of most accurate decoding, and the exact period of the recurrent process, differed

widely between participants, obscuring these effects in the grand averages. These individual

differences may be caused by individual preferences for a level of regularity in the beat-based

stimuli (Drake et al., 2000), with some people attending mostly to the beat level (1.67 Hz), but

others possibly attending more to subdivisions (3.33 Hz) or the level of the meter (0.83 Hz).

Also, for different people, the optimal phase of delta oscillations (i.e., the phase that aligns with

expected moments) may differ (Breska & Deouell, 2017; Henry & Obleser, 2012), possibly

causing optimal decoding at different time points. To circumvent the large task-related effects

apparent in the decoding results, and the possible individual differences in phase and metrical

level attended to, we ran an additional exploratory decoding analysis looking at decoding within

instead of between conditions. Here, we found above-chance decoding of beat positions against

offbeat positions only following the beat-based condition. However, while decoding was above

chance in the beat-based condition, it was in fact not better than in the other conditions. As

such, these results provide only weak support for persistent effects of beat-based, but not

pattern-based expectations. Detailed results of the within-condition decoding analysis can be

found in the Appendix (Figure A4).

Discussion

In the current study, we aimed to directly compare the neural mechanisms underlying temporal expectations based on regular beats and predictable patterns, by examining the development of climbing activity and the persistence of neural entrainment after cessation of rhythmic input. Instead of isochronous sequences, which can elicit both beat-based and memory-based expectations, we used varying non-isochronous patterns to separate beat-based and pattern-based expectations. Moreover, by assessing responses in silence, we sidestepped possible confounds associated with acoustic differences between conditions (Capilla et al., 2011; Haegens & Zion Golumbic, 2018; Novembre & Iannetti, 2018; Zoefel et al., 2018). First, behaviorally, we found that while the effects of beat-based expectations spanned two beat cycles, the effects of pattern-based expectations reflected primarily the first expected moment in time. Second, a negative ERP component at around 300 to 450 ms after the onset of the last sound was larger for pattern-based than beat-based sequences. Third, we observed more power at the beat frequency in the silence window when participants were previously listening to a beat-based sequence as compared to a pattern-based or random sequence. Taken together, these findings suggest that beat-based and pattern-based expectations may rely on different neural mechanisms, as previously found for cue-based expectations (Breska & Deouell, 2017), and not shared mechanisms, be it entrainment (Tichko & Large, 2019), or a general top-down mechanism (Rimmele et al., 2018). Tentatively, our results could be interpreted as entrainment underlying beat-based, but not pattern-based expectations, with pattern-based expectations relying on a different mechanism, such as climbing neuronal activity.

Behaviorally, for the beat-based condition, we observed an inverted U-curve as late as 1200 ms after the final tone of each sequence in both experiments, in line with entrainment models (Bauer et al., 2015; Jones et al., 2002) that assume oscillations are self-sustaining (Large, 2008; Large & Palmer, 2002). Expectations in the beat-based condition led to both

higher ratings for expected and lower ratings for unexpected events compared to the random condition. Divesting resources from times at which an event is unlikely may be metabolically beneficial (van Atteveldt et al., 2015), and as such, has been suggested as a hallmark of entrainment and the associated "rhythmic" mode of processing (Schroeder & Lakatos, 2009b; Zoefel & Vanrullen, 2017). Indeed, the unexpectedness of events off the beat has been proposed to be a better indication of beat-based expectations than the expectedness of events on the beat (Bouwer et al., 2020; Breska & Deouell, 2017), in line with the current results, where the effects of beat-based expectations at unexpected time points exceeded those at expected time points.

Pattern-based expectations led to higher ratings at the first expected time point, and lower ratings at the first unexpected time point, showing that participants did learn the pattern. However, the behavioral effects of the predictable pattern only reflected the first expected moment, but not the subsequent structure. One speculative explanation for this is that listeners form expectations following temporal patterns one interval at a time (e.g., in the absence of an event, no next interval is predicted), and in a probabilistic way (Cannon, 2021; van der Weij et al., 2017). Possibly, the expectation for an event at 780 ms led to an inverted U-shape in responses, similar to the inverted U around a beat, but with a wider distribution, and only occurring once. This could explain the ratings being equal for probes at 780 and 885 ms, both within a time window where a tone could be expected, and ratings being lower for probes at 1200 and 1485 ms, which fell further from the expected time. An alternative explanation could be that the repetitiveness of the pattern lead participants to use a different heuristic during the pattern-based sequences to guide their ratings, considering only one interval at a time, while the varying rhythmic pattern of the beat-based condition induced a strategy whereby participants were more inclined to consider positions after the first expected tone.

In line with entrainment models of beat-based expectations (Haegens & Zion Golumbic, 2018; Henry & Herrmann, 2014; Large & Jones, 1999; Obleser & Kayser, 2019), we found that

EEG power at the beat frequency (1.67 Hz) and its harmonic (3.33 Hz) in the silence window was larger following beat-based than pattern-based or random sequences. Such enhanced power was not found for a frequency inherent to the pattern-based sequence (2.22 Hz). Ongoing oscillations in silence, after sensory input has stopped, are regarded as strong evidence for entrainment (Breska & Deouell, 2017; Haegens & Zion Golumbic, 2018; Obleser & Kayser, 2019; van Bree et al., 2021; Zoefel et al., 2018). Our observation of enhanced power at the beat frequency during silence thus provides converging evidence for the notion that entrainment of low-frequency neural oscillations underlies beat-based perception, and our design, using non-isochronous rhythms, allows us to separate beat-based aspects from other structures present in natural rhythm (Bouwer et al., 2021), similar to other studies combining frequency tagging with non-isochronous rhythms (Lenc et al., 2021; Nozaradan et al., 2012; Tal et al., 2017).

Previously, no evidence for persistent entrainment at the frequency of isochronous rhythmic stimulation was found (Pesnot Lerousseau et al., 2021). Possibly, when presented with stimuli that do not require beat-based expectations to track the temporal structure, the brain may not engage in forming such expectations. Also, while in our study, the auditory sequences were task relevant, and the task itself was rhythm-related, in the study by Pesnot Lerousseau et al. (2021), participants listened to rhythms passively. Thus, the presence of persistent entrainment may also depend on task demands (Shalev et al., 2019). The influence of contextual factors on entrainment, like stimulus and task, are important topics for future research, as is the influence of movement. While we instructed participants not to move to the sequences, we cannot rule out that small, unintentional movements may have affected the EEG signal. More generally, movement may affect rhythm processing (Su & Pöppel, 2012), and thus entrainment.

The presence of persistent entrainment only in active listening conditions may suggest that beat-based expectations result from an explicit, top-down guided process, rather than some passive, automatic one (Bouwer, 2022). Technically, at the scalp, phase locking due to

sustained oscillations and phase locking due to recurrent top-down signals may be indistinguishable, just as phase locking during rhythmic stimulation is often confounded with contributions from tone-evoked responses (Capilla et al., 2011; Novembre & Iannetti, 2018; Zoefel et al., 2018). Interestingly, in our study, the scalp topographies for power at 1.67 Hz differed between the silence and control windows, with power in the control window largest above a frontocentral region, and in the silence window above a parieto-central region. This may suggest that instead of resulting from a sustained automatic oscillation that would have a similar source during sound presentation and silence, persistent entrainment could originate from a different source, such as explicit predictions made by a motor network (Rimmele et al., 2018). Our study raises two questions in this regard, notably whether such a top-down signal would function differentially for beat-based and pattern-based expectations, as suggested by our results, and whether such a top-down signal would be related to CNV-like climbing neuronal activity.

The absence of power at 2.22 Hz following the pattern-based sequences suggests that entrainment does not underlie expectations based on learning a pattern, contrary to a recently proposed oscillator model that captures aspects of pattern-based expectations (Tichko & Large, 2019). In the time-domain, we found a negative deflection in the silence window following the pattern-based and random sequences, akin to a CNV. However, while previous studies showed a CNV peaking at an expected time (Breska & Deouell, 2017; Mento, 2017; Praamstra et al., 2006), here, the peak latency of the negative deflection in the signal was earlier, peaking around 400 ms for the pattern-based condition, while the first expected time point in the silence was at 780 ms. This may be explained by assuming that the CNV indexes temporal expectations in a probabilistic, and context-dependent way (Capizzi, Correa, & Sanabria, 2013; Damsma et al., 2021; Los & Heslenfeld, 2005; see also Hassall et al., 2022). The pattern used in the current study contained temporal intervals with durations between 150 and 780 ms. The ERP peaking

at 400 ms may have indexed the average interval presented in the sequence (~360 ms). This explanation is supported by the presence of a similar ERP deflection for the random condition, which while random in terms of transitional probabilities, was identical to the pattern-based condition in terms of the absolute intervals used. Of note, probabilistic models indexing statistical regularities in inter-onset intervals have indeed been used to explain aspects of temporal processing (Cannon, 2021; Elliott et al., 2014; van der Weij et al., 2017). In future work, linking such models directly to neural markers of pattern-based expectations may provide more insight in the mechanisms underlying pattern-based expectations and how they relate to the CNV.

Interestingly, the same ERP component was less present in the beat-based condition, despite an identical average duration of intervals. In previous research, a CNV was found peaking at expected times not only for cue-based, but also for beat-based expectations (Breska & Deouell, 2012, 2017; Breska & Ivry, 2020; Praamstra et al., 2006), though typically, with isochronous stimulation. Therefore, memory-based expectations, be it pattern-based or cue-based, could also have been formed in response to these sequences (Bouwer et al., 2016, 2020, 2021; Breska & Ivry, 2016; Keele et al., 1989), and may have contributed to the elicitation of a CNV. Here, using a beat-based sequence that did not allow for expectations based on simply learning transitional probabilities, we did not observe the same negative deflection as in the pattern-based sequences. This raises the possibility that a mechanism related to climbing neuronal activity (or CNV) may specifically support the formation of pattern-based and cue-based (Mento, 2013, 2017), temporal expectations, and that if possible given the input, the brain operates in a rhythmic mode of processing (Rimmele et al., 2018; Schroeder & Lakatos, 2009b).

It could be argued that the differences we observed in ERPs were caused by differences in the P3 response to the last sound of each sequence rather than the CNV, which would be apparent at a similar latency. The P3 is indeed susceptible to temporal expectations, with larger

amplitude responses for temporally predictable targets (Lange, 2009; Mento, 2017; Schmidt-

Kassow, Schubotz, & Kotz, 2009). However, these effects can be observed for beat-based and

memory-based expectations in a similar direction (Breska & Deouell, 2017; Breska & Ivry,

2020; Mento, 2017; Schmidt-Kassow et al., 2009). Hence the P3, if anything, should have been

larger for the pattern-based than random sequences, which is not the case. Thus, we feel it is

unlikely that the ERP differences are caused by differences in the P3, and we tentatively suggest

that here, the ERP results are more likely to be due to a CNV-like mechanism.

One alternative explanation for our results could be that participants used an interval-

based strategy in the beat-based condition, predicting an event every 600 ms, and that the

differences between conditions were due to easier learning in the beat-based than pattern-based

condition, both because the length of the beat (600 ms) was shorter than the length of the pattern

(1800 ms), and because the perceived beat occurred three times as often as the pattern.

However, first, because of the varying surface structure in the beat-based condition, in absolute

terms the beat interval occurred exactly as often as the pattern intervals. Second, the pattern

was kept consistent throughout the experiment, for a total of 216 presentations, making it

unlikely that participants did not learn the pattern. Indeed, a previous study showed that even

untrained participants could reproduce non-metrical patterns to some extent after only one

presentation (Cameron & Grahn, 2014). Third, in our previous study, using the same stimuli,

we showed that the behavioral facilitation caused by introducing a regular beat was smaller

than the facilitation caused by introducing a predictable pattern (Bouwer et al., 2020),

regardless of differences in length and number of presentations. Finally, all participants except

one were able to judge the first probe after the pattern-based sequences as not fitting the pattern,

as apparent from lower ratings for this probe in the pattern-based than random condition. This

is comparable to the results for probes at unexpected times in the beat-based condition. Hence,

differences in ease of learning are unlikely to explain our results.

In addition, listening to strictly metric patterns, as the ones used here, is associated with

activity in a circuit including the basal ganglia, while listening to non-metric patterns is

associated with activity in a circuit including the cerebellum, making it unlikely that the same,

interval-based mechanism would be used for both types of rhythms (Leow & Grahn, 2014).

Also, predicting the timing of events in non-isochronous strictly metric sequences would

require participants to learn not just the transitional probabilities of single intervals, but also to

combine multiple intervals into groups that together last the length of a beat. It is currently

unclear whether humans, when faced with rhythmic patterns, use such a hierarchical interval-

based strategy. Indeed, future research could examine if beat-based expectations in general can

be explained by such multilevel interval learning, akin to a recent model for beat-based

perception (Cannon & Patel, 2021), as this could provide a general challenge to oscillator

models of beat-based perception. In such a view, the difference between beat-based and pattern-

based timing may be the importance of hierarchical structure in beat-based rhythms (Fitch,

2013), rather than the presence of oscillations.

In addition to behavioral, ERP, and frequency-domain analyses, we looked at multi scale

entropy and multivariate pattern analysis as alternative ways to examine neural entrainment. At

the group level, entropy was higher following the pattern-based than random sequences. This

could be explained by assuming that the brain uses a vigilance mode to track the pattern-based

regularities, which at the neural level, translates in more irregular patterns of activity. Attention

and arousal, which are both associated with temporal expectations (Schroeder & Lakatos,

2009a), have indeed been linked to neural variability as well (Waschke et al., 2021). However,

the entropy results at least to some extent were explained by overall differences in signal

variability (e.g., the signal variance), so this hypothesis remains to be confirmed in future

research. Considering the decoding results, the observed above-chance decoding seemed to

primarily reflect general task-related activity. The best decoding was observed in the second

half of the silence window, where probes could be presented. Also, in the group-average

decoding results, above chance decoding was limited to training and testing on the same time

points, and we did not observe recurrent activity. Yet, we did show a proof of concept for our

approach in at least one participant, who showed a clear oscillatory pattern in decoding accuracy

when decoding the beat-based against the other two conditions. This pattern of activity is in

line with the strength or sharpness of the neural representation of tones varying over time as a

function of temporal expectations (Auksztulewicz et al., 2018, 2019).

The strength of the decoding approach is that neural dynamics can be assessed in real

time, but this may also be its weakness, particularly in the current study. Due to the absence of

sensory input, individuals may differ in how consistently they persist in their beat percept. Also,

people may attend to different levels of regularity in beat-based perception (Drake et al., 2000),

and the optimal phase of delta oscillations (e.g., the phase that aligns with expected moments)

may differ across individuals (Breska & Deouell, 2017; Henry & Obleser, 2012; Sun et al.,

2021). One option to overcome the weaknesses of the decoding approach could be to transform

the EEG signal to allow frequency and phase drift. Recent technical implementations of time

warping algorithms may be fruitful in this regard (Chemin et al., 2018; van Bree et al., 2022).

The more traditional frequency tagging approach allows for an assessment of activity

across multiple frequencies, as present in rhythm in a metrical structure. Phase-based analyses

may not be suitable to answer the questions we ask here, due to differences in preferred phase

between individuals, as well as the fact that phase may not differentiate between beat-based and

memory-based expectations (Breska & Deouell, 2017). However, a frequency tagging approach

is not sensitive to phase differences between individuals. This may explain why the results for

the frequency-based analysis were more robust than those obtained with decoding. A weakness

of the frequency tagging approach may be that the Fourier transform assumes stationarity in the

oscillating signal, while entrainment models propose a dampening factor to account for

decreasing oscillatory power over time (Doelling & Assaneo, 2021; Large et al., 2015). To

assess power at specific frequencies in a time-resolved way, wavelet convolution is often used

as an alternative. But, in the current study, differentiating between the specific frequencies of

the beat and the pattern would require wavelet parameters that would result in a temporal

resolution too low to disentangle activity during and before the silence (i.e., many wavelet

cycles would be needed). Recently, a promising alternative to assess oscillatory activity in the

time domain was proposed in the form of cycle-by-cycle analysis (Cole & Voytek, 2019).

However, this approach requires filtering the signal at the specific frequency of interest, again

posing problems for disentangling low frequency oscillations due to the beat, the pattern, and

ongoing ERPs. Assessing the time course of low frequency oscillations thus remains a

challenge for future research.

In our current study, we observed large heterogeneity between individuals. While it is

often assumed that most people automatically form beat-based expectations (Honing, 2012),

recent evidence showed phase locking to speech in only about half of the population (Assaneo

et al., 2019). Indeed, in our study, only about two-thirds of the participants behaviorally showed

evidence for beat-based expectations in the second half of the silence window and we only

observed enhanced power at the beat frequency following beat-based sequences in about half

of the participants. The behavioral effects of beat-based and pattern-based expectations were

also associated with musical training, consistent with previous research using beat-based

(Bouwer et al., 2016, 2018; Cameron & Grahn, 2014; Matthews et al., 2016; Vuust et al., 2005)

and pattern-based (Cameron & Grahn, 2014) rhythms. Some previous studies have failed to

show differences between musicians and non-musicians, however (Bouwer et al., 2014; Geiser

et al., 2009; Grahn & Brett, 2007), possibly due to differences in task design (Bouwer et al.,

2018). The current study used an explicit rating task, for which performance may be particularly

improved by musical training, as musically trained participants may have additional strategies

to perform the task. The use of implicit timing tasks may be a better probe of innate differences

in timing abilities, which need not necessarily be related to musical training (Law & Zentner,

2012), and implicit tasks may also be less susceptible to task-related effects as we observed in

the decoding analysis, which may stem from individual strategies in performing the explicit

task. Undoubtedly, given the heterogeneity we and others (Assaneo et al., 2019; Bauer et al.,

2015; Sun et al., 2021) have observed in tasks probing temporal expectations, understanding

individual differences is an important direction for future research, with significant implications

for applications of musical rhythm, such as in motor rehabilitation (Dalla Bella et al., 2018).

Conclusion

In summary, we have shown that beat-based and pattern-based expectations can be

differentiated in terms of their behavioral and neurophysiological effects once sensory input

has ceased. These findings provide novel evidence for the notion that different mechanisms

implement temporal expectations based on periodic and aperiodic input streams, with,

tentatively, the former based on entrainment of low frequency neural oscillations, and the latter

on climbing neural activity indexing a memorized interval.

1107

 Appendix

1109 **Multi Scale Entropy computation**

1110   To compute entropy, epochs were concatenated. Sample entropy was then calculated by

1111 taking the following steps:

1. In each time series, a template is selected, consisting of *m* samples. The calculation

1113   of entropy is an iterative process, using each sample in the time series as a starting

1114   point for the template once.

2. Throughout the time series, the algorithm searches for patterns that match the

1116   template pattern. A section of samples is considered a match if it resembles the

1117   template pattern enough to fall within a set boundary, which is defined as *r* x SD

1118   (the similarity bound).

3. The number of pattern matches is counted.

4. Subsequently, the same procedure is followed for patterns of *m* + 1 samples long.

5. For the total counts of pattern matches throughout the time series, sample entropy is

then calculated as the logarithm of the ratio between pattern matches of length *m*

and pattern matches of length *m* + 1.

Thus, sample entropy reflects the proportion of patterns in the time series that stays similar

when an extra sample is added to the pattern. Here, we used *m* = 2 and *r* = 0.5, as was done

previously for EEG data (Kloosterman et al., 2020).

Sample entropy is then repeated for multiple timescales, to account for contributions of

both low and high frequency neural activity. The time series is coarsened step by step, by taking

the average of a group of adjacent samples with step-wise increasing group size. This means

that long, or coarse, timescales are equivalent to low frequency activity. For example, at a

sampling rate of 256 Hz, a timescale of 4 (e.g., averaged over four adjacent samples, or 15.6

ms) is roughly the equivalent of looking at activity at 64 Hz, while a timescale of 153 at that

sample rate corresponds to averaging over 598 ms, or the equivalent of activity at roughly 1.67

Hz. Here, we used twenty timescales ranging from 4 till 153, with 153 being the maximum

timescale given the length of an epoch of 1800 ms (e.g., one epoch equals 460 samples, but

since entropy is calculated on patterns of 2 and 3 samples, the maximum coarsening to retain

the possibility of having a 3-sample pattern is by averaging over 153 samples). Similarity

bounds were recomputed for each time scale (Kloosterman et al., 2020; Kosciessa et al., 2020).

To control for the contribution of delta, we repeated the analysis on high-pass filtered data (5

Hz).

Decoding within condition

The initial decoding analysis yielded large effects that may be task-related, and

individual differences in phase and metrical level attended to may have additionally hampered

finding condition differences at the group level. Therefore, we ran an additional decoding

analysis in which we decoded expected and unexpected positions against each other within each

condition. For this analysis, we only included frontocentral electrodes (C1, C2, C3, C4, Cz,

FC1, FC2, FC3, FC4, FCz, F1, F2, F3, F4, Fz), which were the same electrodes that showed

the largest P1 responses to the initial sounds of the sequences, indicative of representing the

auditory cortex. Here, we defined time points in the silence as expected or unexpected based

on the rhythmic sequences. To equate the choice of expected and unexpected time points as

much as possible in terms of number of time points included and distance between time points,

we used times that were also used in the probe tone experiment: 600, 780, 885, and 1200 ms.

For beat-based sequences, time points 600 and 1200 were expected (on the beat), while time

points 780 and 885 were unexpected (offbeat). The same time points take on a different meaning

if preceded by the pattern-based sequences. Then, 780 and 1200 ms are considered expected

(predictable based on the pattern), while both 600 and 885 ms are unexpected (unpredictable

based on the pattern). For all conditions, we examined whether we could classify above chance

whether a time window of 100 ms centered on the time point of interest was an expected or

unexpected moment in time. As for the first decoding analysis, the data were resampled to

improve signal to noise, here to 128 Hz to retain enough data points in the 100 ms window for

analysis. After the decoding, we extracted the average classification accuracy for the 100 ms

time window to test significance for each condition separately against chance level, using t-

tests against 0.5, and between conditions in the silence, using a repeated measures ANOVA,

with condition as an independent factor, a random intercept for participant, and decoding

accuracy as the dependent variable.

1166

1167

1168

Figure A1. ERPs with traditional 50 ms baseline and difference waves. A) With the more traditional pre-cue baseline, there was a trend for the random condition to elicit a more negative deflection than the beat-based condition ($p = 0.058$, $300 – 446$ ms) and for the pattern-based condition to elicit a more negative deflection than the random condition (early window: $p = 0.083$, $380 – 551$ ms; later window: $p = 0.022$, $774 – 1200$ ms). Note that a significant cluster spanning the entire analysis window of $300 – 1200$ ms was found when comparing between the pattern-based and beat-based condition with baseline correction ($p = 0.011$), indicative of a possible overall shift due to the baseline correction. However, the overall picture stays the same as when not using a baseline, a negative deflection can be seen around 300-450 ms after the last sound onset for the pattern-based and random, but not the beat-based condition. B) Scalp distributions for windows with a significant cluster (see also Figure 5). C) Difference waves depicting condition differences and the 95% confidence interval around the mean difference waveforms. The scalp distributions similarly depict condition differences. D) Evoked potential showing the ERPs during sound presentation. Time 0 here is the presentation of a sound, with the next sound being presented at 600 (beat-based condition) or 780 ms (pattern-based and random condition). As in the silence window, a negative deflection can be observed, which was larger for the pattern-based than random condition ($p = 0.039$), and peaks around 350 ms after the onset of the previous sound. The difference between the pattern-based and beat-based condition here did not reach significance ($p = 0.1$), likely because of the noisy baseline.

1188

Figure A2. MMSE after highpass filtering at 5 Hz. Here, all differences between conditions are eliminated
by the highpass filtering.

1191

1192

Figure A3. Similarity bounds used to calculate MMSE values mirror the condition differences. This
finding suggests that condition differences found in the entropy measure, with higher entropy for the pattern-
based than random condition in the silence window, may be due to overall variability differences, with lower
variability for the pattern-based than random condition, rather than differences in entropy of the signal.

1197

1198

Figure A4. Above chance decoding for beat vs offbeat time points only after listening to beat-based sequences. For the within-condition decoding analysis, we used a classification algorithm to differentiate between times that were expected or unexpected based on the beat (on the beat: 600 and 1200 ms after the onset of the last tone; offbeat: 780 and 885 ms after the onset of the last tone) or pattern (predicted: 780 and 1200 ms after the onset of the last tone; unpredicted: 600 and 880 ms after the onset of the last tone). The graphs depict the average classification accuracy in a 100 ms time window centered on these expected and unexpected time points in the silence window. Here, the beat-offbeat comparison should yield better decoding if preceded by the beat-based sequences, while the predicted-unpredicted comparison should be decoded better if preceded by the pattern-based sequences. Only decoding of beat vs offbeat positions after the beat-based sequences yielded above-chance decoding ($t_{26} = 2.84$, $p = 0.009$). None of the other comparisons lead to decoding above chance (all $ps > 0.18$), but decoding of predicted vs unpredicted positions in the beat-based condition did yield below-chance decoding ($p = 0.026$). However, a subsequent ANOVA comparing decoding of beat against offbeat positions in all three conditions showed that decoding was not significantly better for the beat-based condition than the other conditions (effect of condition in the silence: $\chi^2 = 2.1$, $p = 0.35$). Like for the other analyses, there were larger differences between participants, and decoding, while above chance, did not exceed 0.55 accuracy. As for the other EEG measures, the difference in decoding accuracy between the beat-based and random condition, and the patten-based and random condition did not depend on musical training (both $ps > 0.21$).

1217
1218

1219 **Table A1**

1220 *Simple effects per position from the full models, comparing three conditions.*

		Experiment 1				Experiment 2			
Position	Contrast	Estimate	SE	Z ratio	*p*	Estimate	SE	Z ratio	*p*
600 ms	Random – Pattern	2.4	0.09	26.2	<.0001				
	Random – Beat	-1.2	0.09	13.3	<.0001				
	Pattern – Beat	-3.5	0.09	-38.3	<.0001				
780 ms	Random – Pattern	-0.3	0.09	-2.9	0.058				
	Random – Beat	1.6	0.09	18.3	<.0001				
	Pattern – Beat	1.8	0.09	20.7	<.0001				
885 ms	Random – Pattern	-0.1	0.09	-1.6	1				
	Random – Beat	1.5	0.09	17.0	<.0001				
	Pattern – Beat	1.6	0.09	18.4	<.0001				
990 ms	Random – Pattern	0.1	0.09	1.1	1	0.6	0.1	4.8	<.0001
	Random – Beat	1.4	0.09	16.2	<.0001	1.4	0.1	11.4	<.0001
	Pattern – Beat	1.3	0.09	15.0	<.0001	0.8	0.1	6.5	<.0001
1200 ms	Random – Pattern	0.6	0.08	7.5	<.0001	0.2	0.1	1.6	0.92
	Random – Beat	-0.3	0.09	-3.7	0.004	0.1	0.1	0.9	1
	Pattern – Beat	-1.0	0.09	-10.9	<.0001	-0.1	0.1	-0.7	1
1485 ms	Random – Pattern	0.5	0.09	5.2	<.0001	-0.5	0.1	-4.4	<.0001
	Random – Beat	0.6	0.09	6.5	<.0001	0.4	0.1	3.2	0.011
	Pattern – Beat	0.1	0.09	1.3	1	0.9	0.1	7.5	<.0001

1221 *Note.* All *p*-values have been Bonferroni corrected for 18 (Experiment 1) or 9 (Experiment 2)

1222 comparisons.

1223

 Table A2

 Simple effects per condition from the baseline corrected model, using the random condition as

 baseline, for all positions.

Condition	Contrast	Experiment 1				Experiment 2			
		Estimate	SE	Z ratio	*p*	Estimate	SE	Z ratio	*p*
Beat-based	600 ms - 780 ms	2.3	0.08	28.7	<.0001				
	600 ms - 885 ms	2.2	0.08	27.8	<.0001				
	600 ms - 990 ms	2.1	0.08	27.0	<.0001				
	600 ms - 1200 ms	0.6	0.08	6.9	<.0001				
	600 ms - 1485 ms	1.5	0.08	18.0	<.0001				
	780 ms - 885 ms	-0.1	0.08	-1.3	1				
	780 ms - 990 ms	-0.2	0.08	-1.9	1				
	780 ms - 1200 ms	-1.8	0.08	-21.3	<.0001				
	780 ms - 1485 ms	-0.9	0.08	-10.5	<.0001				
	885 ms - 990 ms	-0.1	0.08	-0.7	1				
	885 ms - 1200 ms	-1.7	0.08	-20.3	<.0001				
	885 ms - 1485 ms	-0.8	0.08	-9.4	<.0001				
	990 ms - 1200 ms	-1.6	0.08	-19.5	<.0001	-1.1	0.1	-9.6	<.0001
	990 ms - 1485 ms	-0.7	0.08	-8.7	<.0001	-0.9	0.1	-7.9	<.0001
	1200 ms - 1485 ms	0.9	0.08	10.9	<.0001	0.2	0.1	1.8	0.46
Pattern-based	600 ms - 780 ms	-2.3	0.09	-27.0	<.0001				
	600 ms - 885 ms	-2.2	0.08	-26.4	<.0001				
	600 ms - 990 ms	-2.1	0.08	-24.8	<.0001				
	600 ms - 1200 ms	-1.7	0.08	-20.2	<.0001				
	600 ms - 1485 ms	-1.7	0.09	-20.3	<.0001				
	780 ms - 885 ms	0.1	0.08	1.0	1				
	780 ms - 990 ms	0.2	0.08	3.0	0.080				
	780 ms - 1200 ms	0.6	0.08	7.6	<.0001				
	780 ms - 1485 ms	0.6	0.08	6.9	<.0001				
	885 ms - 990 ms	0.2	0.08	2.0	1				
	885 ms - 1200 ms	0.5	0.08	6.7	<.0001				
	885 ms - 1485 ms	0.5	0.08	6.0	<.0001				
	990 ms - 1200 ms	0.4	0.08	4.7	<.0001	-0.3	0.1	-2.9	0.022
	990 ms - 1485 ms	0.3	0.08	4.1	0.001	-0.9	0.1	-8.3	<.0001
	1200 ms - 1485 ms	0.0	0.08	-0.5	1.00	-0.6	0.1	-5.4	<.0001

 Note. All *p*-values have been Bonferroni corrected for 30 (Experiment 1) or 6 (Experiment 2)

 comparisons.

Data availability statement

All datafiles, and code used for data acquisition, data analysis, and figure creation, are available through https://osf.io/uwny8/.

Author contribution

Fleur L. Bouwer: Conceptualization, Methodology, Formal analysis, Investigation, Writing – Original Draft, Writing – Review & Editing, Visualization

Johannes J. Fahrenfort: Software, Formal analysis, Writing – Review & Editing

Samantha K. Millard: Methodology, Formal analysis, Investigation, Writing – Review & Editing

Niels A. Kloosterman: Software, Formal analysis, Writing – Review & Editing

Heleen A. Slagter: Conceptualization, Methodology, Resources, Writing – Review & Editing, Supervision, Funding acquisition

Acknowledgements

We would like to thank Babak Chawoush and Charlotte van Roijen for their assistance with the data collection, and fruitful discussions during the project.

Funding Information

FLB is supported by an ABC Talent Grant awarded by Amsterdam Brain and Cognition, a Veni grant awarded by the Dutch Research Council NWO (VI.Veni.201G.066), and the ERC starting grant awarded to HAS. HAS is supported by a European Research Council (ERC) starting grant (679399).

References

Arnal, L. H., Doelling, K. B., & Poeppel, D. (2014). Delta-beta coupled oscillations underlie temporal prediction accuracy. *Cerebral Cortex*, *25*(9), 3077–3085. https://doi.org/10.1093/cercor/bhu103

Assaneo, M. F., Ripollés, P., Orpella, J., Lin, W. M., de Diego-Balaguer, R., & Poeppel, D. (2019). Spontaneous synchronization to speech reveals neural mechanisms facilitating language learning. *Nature Neuroscience*, *22*(4), 627–632. https://doi.org/10.1038/s41593-019-0353-z

Auksztulewicz, R., Myers, N. E., Schnupp, J. W., & Nobre, A. C. (2019). Rhythmic temporal expectation boosts neural activity by increasing neural gain. *The Journal of Neuroscience*, *39*(49), 9806–9817. https://doi.org/10.1523/JNEUROSCI.0925-19.2019

Auksztulewicz, R., Schwiedrzik, C. M., Thesen, T., Doyle, W., Devinsky, O., Nobre, A. C., Schroeder, C. E., Friston, K. J., & Melloni, L. (2018). Not all predictions are equal: "What" and "when" predictions modulate activity in auditory cortex through different mechanisms. *The Journal of Neuroscience*, *38*(40), 8680–8693. https://doi.org/10.1523/JNEUROSCI.0369-18.2018

Bauer, A.-K. R., Jaeger, M., Thorne, J. D., Bendixen, A., & Debener, S. (2015). The auditory dynamic attending theory revisited: A closer look at the pitch comparison task. *Brain Research*, *1626*, 198–210. https://doi.org/10.1016/j.brainres.2015.04.032

Bouwer, F. L. (2022). Neural entrainment to auditory rhythms: Automatic or top-down driven? *Journal of Neuroscience*, *42*(11), 2146–2148. https://doi.org/10.1523/JNEUROSCI.2305-21.2022

Bouwer, F. L., Burgoyne, J. A., Odijk, D., Honing, H., & Grahn, J. A. (2018). What makes a rhythm complex? The influence of musical training and accent type on beat perception. *PLOS ONE*, *13*(1), e0190322. https://doi.org/10.1371/journal.pone.0190322

Bouwer, F. L., & Honing, H. (2015). Temporal attending and prediction influence the perception of metrical rhythm: Evidence from reaction times and ERPs. *Frontiers in Psychology*, *6*(July), 1094. https://doi.org/10.3389/fpsyg.2015.01094

Bouwer, F. L., Honing, H., & Slagter, H. A. (2020). Beat-based and memory-based temporal expectations in rhythm: Similar perceptual effects, different underlying mechanisms. *Journal of Cognitive Neuroscience*, *32*(7), 1221–1241. https://doi.org/10.1162/jocn_a_01529

Bouwer, F. L., Nityananda, V., Rouse, A. A., & ten Cate, C. (2021). Rhythmic abilities in humans and non-human animals: A review and recommendations from a methodological perspective. *Philosophical Transactions of the Royal Society B*, *376*, 20200335. https://doi.org/10.1098/rstb.2020.0335

Bouwer, F. L., Van Zuijen, T. L., & Honing, H. (2014). Beat processing is pre-attentive for metrically simple rhythms with clear accents: An ERP study. *PLoS ONE*, *9*(5), e97467. https://doi.org/10.1371/journal.pone.0097467

Bouwer, F. L., Werner, C. M., Knetemann, M., & Honing, H. (2016). Disentangling beat perception from sequential learning and examining the influence of attention and musical abilities on ERP responses to rhythm. *Neuropsychologia*, *85*, 80–90. https://doi.org/10.1016/j.neuropsychologia.2016.02.018

Breska, A., & Deouell, L. Y. (2012). Automatic bias of temporal expectations following temporally regular input independently of high-level temporal expectation. *Journal of Cognitive Neuroscience*, *26*(7), 1555–1571. https://doi.org/10.1162/jocn

Breska, A., & Deouell, L. Y. (2017). Neural mechanisms of rhythm-based temporal prediction: Delta phase-locking reflects temporal predictability but not rhythmic entrainment. *PLOS Biology*, *15*(2), e2001665. https://doi.org/10.1371/journal.pbio.2001665

Breska, A., & Ivry, R. B. (2016). Taxonomies of timing: Where does the cerebellum fit in? *Current Opinion in Behavioral Sciences*, *8*, 282–288. https://doi.org/10.1016/J.COBEHA.2016.02.034

Breska, A., & Ivry, R. B. (2018). Double dissociation of single-interval and rhythmic temporal prediction in cerebellar degeneration and Parkinson's disease. *Proceedings of the National Academy of Sciences*, *115*(48), 12283–12288. https://doi.org/10.1073/pnas.1810596115

Breska, A., & Ivry, R. B. (2020). Context-specific control over the neural dynamics of temporal attention by the human cerebellum. *Science Advances*, *6*(49), 1–16. https://doi.org/10.1126/sciadv.abb1141

Brysbaert, M., & Stevens, M. (2018). Power analysis and effect size in mixed effects models: A tutorial. *Journal of Cognition*, *1*(1), 1–20. https://doi.org/10.5334/joc.10

Cameron, D. J., & Grahn, J. A. (2014). Enhanced timing abilities in percussionists generalize to rhythms without a musical beat. *Frontiers in Human Neuroscience*, *8*(December), 1003. https://doi.org/10.3389/fnhum.2014.01003

Cannon, J. J. (2021). Expectancy-based rhythmic entrainment as continuous Bayesian inference. *PLoS Computational Biology*, *17*(6), e1009025. https://doi.org/10.1371/journal.pcbi.1009025

Cannon, J. J., & Patel, A. D. (2021). How Beat Perception Co-opts Motor Neurophysiology. *Trends in Cognitive Sciences*, 25(2), 137–150. https://doi.org/10.1016/j.tics.2020.11.002

Capilla, A., Pazo-Alvarez, P., Darriba, A., Campo, P., & Gross, J. (2011). Steady-State Visual Evoked Potentials Can Be Explained by Temporal Superposition of Transient Event-Related Responses. *PLoS ONE*, 6, e14543. https://doi.org/10.1371/journal.pone.0014543

Carifio, J., & Perla, R. (2008). Resolving the 50-year debate around using and misusing Likert scales. *Medical Education*, 42(12), 1150–1152. https://doi.org/10.1111/j.1365-2923.2008.03172.x

Chemin, B., Huang, G., Mulders, D., & Mouraux, A. (2018). EEG time-warping to study non-strictly-periodic EEG signals related to the production of rhythmic movements. *Journal of Neuroscience Methods*, 308, 106–115. https://doi.org/10.1016/J.JNEUMETH.2018.07.016

Christensen, R. H. B. (2019). *Ordinal—Regression models for Ordinal Data. R package version 2019.4-25.*

Cole, S., & Voytek, B. (2019). Cycle-by-cycle analysis of neural oscillations. *Journal of Neurophysiology*, 122, 849–861. https://doi.org/10.1152/jn.00273.2019

Cravo, A. M., Rohenkohl, G., Wyart, V., & Nobre, A. C. (2013). Temporal expectation enhances contrast sensitivity by phase entrainment of low-frequency oscillations in visual cortex. *The Journal of Neuroscience*, 33(9), 4002–4010. https://doi.org/10.1523/JNEUROSCI.4675-12.2013

Dalla Bella, S., Dotov, D., Bardy, B., & Cochen De Cock, V. (2018). Individualization of music-based rhythmic auditory cueing in Parkinson's disease. *Annals of the New York Academy of Sciences*, 1423, 308–317. https://doi.org/10.1111/nyas.13859

Damm, L., Varoqui, D., De Cock, V. C., Bella, S. D., & Bardy, B. (2019). Why do we move to the beat? A multi-scale approach, from physical principles to brain dynamics. *Neuroscience & Biobehavioral Reviews*. https://doi.org/10.1016/J.NEUBIOREV.2019.12.024

Damsma, A., Schlichting, N., & van Rijn, H. (2021). Temporal context actively shapes EEG signatures of time perception. *The Journal of Neuroscience*. https://doi.org/10.1523/JNEUROSCI.0628-20.2021

de Graaf, T. A., Gross, J., Paterson, G., Rusch, T., Sack, A. T., & Thut, G. (2013). Alpha-Band Rhythms in Visual Task Performance: Phase-Locking by Rhythmic Sensory Stimulation. *PLoS ONE*, 8(3), e60035. https://doi.org/10.1371/journal.pone.0060035

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, 134, 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Ding, N., Patel, A. D., Chen, L., Butler, H., Luo, C., & Poeppel, D. (2017). Temporal modulations in speech and music. *Neuroscience and Biobehavioral Reviews*, 81, 181–187. https://doi.org/10.1016/j.neubiorev.2017.02.011

Doelling, K. B., & Assaneo, M. F. (2021). Neural oscillations are a start toward understanding brain activity rather than the end. *PLoS Biology*, 19(5), 1–12. https://doi.org/10.1371/journal.pbio.3001234

Doelling, K. B., Assaneo, M. F., Bevilacqua, D., Pesaran, B., & Poeppel, D. (2019). An oscillator model better predicts cortical entrainment to music. *Proceedings of the National Academy of Sciences*, 116(20), 10113–10121. https://doi.org/10.1073/pnas.1816414116

Drake, C., Jones, M. R., & Baruch, C. (2000). The development of rhythmic attending in auditory sequences: Attunement, referent period, focal attending. *Cognition*, 77(3), 251–288. https://doi.org/10.1016/S0010-0277(00)00106-2

Elliott, M. T., Wing, A. M., & Welchman, A. E. (2014). Moving in time: Bayesian causal inference explains movement coordination to auditory beats. *Proceedings of the Royal Society B*, 281(May), 20140751.

Fahrenfort, J. J., van Driel, J., van Gaal, S., & Olivers, C. N. L. (2018). From ERPs to MVPA Using the Amsterdam Decoding and Modeling Toolbox (ADAM). *Frontiers in Neuroscience*, 12, 368. https://doi.org/10.3389/fnins.2018.00368

Fox, J., & Weisberg, S. (2019). *An R Companion to Applied Regression* (3rd ed.). Sage.

Geiser, E., Ziegler, E., Jancke, L., Meyer, M., Jäncke, L., Meyer, M., Jancke, L., Meyer, M., Jäncke, L., & Meyer, M. (2009). Early electrophysiological correlates of meter and rhythm processing in music perception. *Cortex*, 45(1), 93–102. https://doi.org/10.1016/j.cortex.2007.09.010

Grahn, J. A., & Brett, M. (2007). Rhythm and beat perception in motor areas of the brain. *Journal of Cognitive Neuroscience*, 19(5), 893–906. https://doi.org/10.1162/jocn.2007.19.5.893

Grahn, J. A., & Rowe, J. B. (2009). Feeling the beat: Premotor and striatal interactions in musicians and nonmusicians during beat perception. *The Journal of Neuroscience*, 29(23), 7540–7548. https://doi.org/10.1523/JNEUROSCI.2018-08.2009

Haegens, S., & Zion Golumbic, E. (2018). Rhythmic facilitation of sensory processing: A critical review. *Neuroscience and Biobehavioral Reviews*, 86(March), 150–165. https://doi.org/10.1016/j.neubiorev.2017.12.002

Hassall, C. D., Harley, J., Kolling, N., & Hunt, L. T. (2022). Temporal scaling of human scalp-recorded potentials. *Proceedings of the National Academy of Sciences*, *119*(43), e2214638119. https://doi.org/10.1073/pnas.2214638119

Heideman, S. G., van Ede, F., & Nobre, A. C. (2018). Early Behavioural Facilitation by Temporal Expectations in Complex Visual-motor Sequences. *Neuroscience*, *389*, 74–84. https://doi.org/10.1016/J.NEUROSCIENCE.2018.05.014

Henry, M. J., & Herrmann, B. (2014). Low-frequency neural oscillations support dynamic attending in temporal context. *Timing & Time Perception*, *2*(1), 62–86. https://doi.org/10.1163/22134468-00002011

Henry, M. J., Herrmann, B., & Obleser, J. (2014). Entrained neural oscillations in multiple frequency bands comodulate behavior. *Proceedings of the National Academy of Sciences*, *111*(41), 14935–14940. https://doi.org/10.1073/pnas.1408741111

Henry, M. J., & Obleser, J. (2012). Frequency modulation entrains slow neural oscillations and optimizes human listening behavior. *Proceedings of the National Academy of Sciences*, *109*(49), 20095–20100. https://doi.org/10.1073/pnas.1213390109

Herbst, S. K., Stefanics, G., & Obleser, J. (2022). Endogenous modulation of delta phase by expectation–A replication of Stefanics et al., 2010. *Cortex*, *149*, 226–245. https://doi.org/10.1016/j.cortex.2022.02.001

Hickok, G., Farahbod, H., & Saberi, K. (2015). The Rhythm of Perception: Entrainment to Acoustic Rhythms Induces Subsequent Perceptual Oscillation. *Psychological Science*, *26*(7), 1006–1013. https://doi.org/10.1177/0956797615576533

Honing, H. (2012). Without it no music: Beat induction as a fundamental musical trait. *Annals of the New York Academy of Sciences*, *1252*(1), 85–91. https://doi.org/10.1111/j.1749-6632.2011.06402.x

Honing, H., & Bouwer, F. L. (2019). Rhythm. In P. J. Rentfrow & D. Levitin (Eds.), *Foundations in Music Psychology: Theory and Research* (pp. 33–69). MIT Press.

Jamieson, S. (2004). Likert scales: How to (ab)use them. *Medical Education*, *38*(12), 1217–1218. https://doi.org/10.1111/j.1365-2929.2004.02012.x

JASP, T. (2019). *JASP* (0.11.1).

Jeffreys, H. (1961). *Theory of probability* (3rd ed.). Oxford University Press.

Jones, M. R., Moynihan, H., MacKenzie, N., & Puente, J. (2002). Temporal aspects of stimulus-driven attending in dynamic arrays. *Psychological Science*, *13*(4), 313–319. https://doi.org/10.1111/1467-9280.00458

Keele, S. W., Nicoletti, R., Ivry, R. B., & Pokorny, R. A. (1989). Mechanisms of perceptual timing: Beat-based or interval-based judgements? *Psychological Research*, *50*, 251–256. https://doi.org/10.1007/BF00309261

Keitel, C., Obleser, J., Jessen, S., & Henry, M. J. (2021). Frequency-Specific Effects in Infant Electroencephalograms Do Not Require Entrained Neural Oscillations: A Commentary on Köster et al. (2019). *Psychological Science*, *32*(6), 966–971. https://doi.org/10.1177/09567976211001317

King, J.-R., & Dehaene, S. (2014). Characterizing the dynamics of mental representations: The temporal generalization method. *Trends in Cognitive Sciences*, *18*(4), 203–210. https://doi.org/10.1016/J.TICS.2014.01.002

Kloosterman, N. A., Kosciessa, J. Q., Lindenberger, U., Fahrenfort, J. J., & Garrett, D. D. (2020). Boosts in brain signal variability track liberal shifts in decision bias. *ELife*, *9*, 1–22. https://doi.org/10.7554/ELIFE.54201

Kononowicz, T. W., & Penney, T. B. (2016). The contingent negative variation (CNV): Timing isn't everything. *Current Opinion in Behavioral Sciences*, *8*, 231–237. https://doi.org/10.1016/j.cobeha.2016.02.022

Kosciessa, J. Q., Kloosterman, N. A., & Garrett, D. D. (2020). Standard multiscale entropy reflects neural dynamics at mismatched temporal scales: What's signal irregularity got to do with it? *PLoS Computational Biology*, *16*(5), e1007885. https://doi.org/10.1371/journal.pcbi.1007885

Kösem, A., Bosker, H. R., Takashima, A., Meyer, A., Jensen, O., & Hagoort, P. (2018). Neural Entrainment Determines the Words We Hear. *Current Biology*, *28*(18), 2867-2875.e3. https://doi.org/10.1016/j.cub.2018.07.023

Large, E. W. (2008). Resonating to musical rhythm: Theory and experiment. In S. Grondin (Ed.), *Psychology of time* (pp. 189–231). Emerald Group Publishing. https://doi.org/10.1016/B978-0-08046-977-5.00006-5

Large, E. W., Herrera, J. A., & Velasco, M. J. (2015). Neural networks for beat perception in musical rhythm. *Frontiers in Systems Neuroscience*, *9*(November), 159. https://doi.org/10.3389/fnsys.2015.00159

Large, E. W., & Jones, M. R. (1999). The dynamics of attending: How people track time-varying events. *Psychological Review*, *106*(1), 119–159. https://doi.org/10.1037/0033-295X.106.1.119

Large, E. W., & Palmer, C. (2002). Perceiving temporal regularity in music. *Cognitive Science*, *26*(1), 1–37. https://doi.org/10.1207/s15516709cog2601_1

Law, L. N. C., & Zentner, M. (2012). Assessing Musical Abilities Objectively: Construction and Validation of the Profile of Music Perception Skills. *PLoS ONE*, *7*(12), e52508. https://doi.org/10.1371/journal.pone.0052508

Lenc, T., Keller, P. E., Varlet, M., & Nozaradan, S. (2018). Neural tracking of the musical beat is enhanced by low-frequency sounds. *Proceedings of the National Academy of Sciences*, *115*(32), 8221–8226. https://doi.org/10.1073/pnas.1801421115

Lenc, T., Merchant, H., Keller, P. E., Honing, H., Varlet, M., & Nozaradan, S. (2021). Mapping between sound, brain and behaviour: Four-level framework for understanding rhythm processing in humans and non-human primates. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *376*, 20200325. https://doi.org/10.1098/rstb.2020.0325

Lenth, R. (2019). *Emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.4.*

Leow, L., & Grahn, J. A. (2014). Neural mechanisms of rhythm perception: Present findings and future directions. In H. Merchant & V. de Lafuente (Eds.), *Neurobiology of Interval Timing* (pp. 325–338). Springer. https://doi.org/10.1007/978-1-4939-1782-2

Lin, W. M., Oetringer, D. A., Bakker-Marshall, I., Emmerzaal, J., Wilsch, A., ElShafei, H. A., Rassi, E., & Haegens, S. (2021). No behavioural evidence for rhythmic facilitation of perceptual discrimination. *European Journal of Neuroscience*. https://doi.org/10.1111/ejn.15208

London, J. (2012). *Hearing in time: Psychological aspects of musical meter.* (2nd ed.). Oxford University Press.

London, J., Polak, R., & Jacoby, N. (2017). Rhythm histograms and musical meter: A corpus study of Malian percussion music. *Psychonomic Bulletin & Review*, *24*(2), 474–480. https://doi.org/10.3758/s13423-016-1093-7

Manning, F. C., Harris, J., & Schutz, M. (2017). Temporal prediction abilities are mediated by motor effector and rhythmic expertise. *Experimental Brain Research*, *235*(3), 861–871. https://doi.org/10.1007/s00221-016-4845-8

Manning, F. C., & Schutz, M. (2013). "Moving to the beat" improves timing perception. *Psychonomic Bulletin & Review*, *20*, 1133–1139. https://doi.org/10.3758/s13423-013-0439-7

Mathewson, K. E., Prudhomme, C., Fabiani, M., Beck, D. M., Lleras, A., & Gratton, G. (2012). Making Waves in the Stream of Consciousness: Entraining Oscillations in EEG Alpha and Fluctuations in Visual Awareness with Rhythmic Visual Stimulation. *Journal of Cognitive Neuroscience*, *24*(12), 2321–2333. https://doi.org/10.1162/jocn_a_00288

Matthews, T. E., Thibodeau, J. N. L., Gunther, B. P., & Penhune, V. B. (2016). The Impact of Instrument-Specific Musical Training on Rhythm Perception and Production. *Frontiers in Psychology*, *7*(February), 69. https://doi.org/10.3389/fpsyg.2016.00069

Mento, G. (2013). The passive CNV: Carving out the contribution of task-related processes to expectancy. *Frontiers in Human Neuroscience*, *7*(December), 827. https://doi.org/10.3389/fnhum.2013.00827

Mento, G. (2017). The role of the P3 and CNV components in voluntary and automatic temporal orienting: A high spatial-resolution ERP study. *Neuropsychologia*, *107*, 31–40. https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2017.10.037

Merchant, H., Grahn, J. A., Trainor, L. J., Rohrmeier, M. A., & Fitch, W. T. (2015). Finding the beat: A neural perspective across humans and non-human primates. *Philosophical Transactions of the Royal Society B*, *370*, 20140093. https://doi.org/10.1098/rstb.2014.0093

Meyer, L., Sun, Y., & Martin, A. E. (2019). Synchronous, but not entrained: Exogenous and endogenous cortical rhythms of speech and language processing. *Language, Cognition and Neuroscience*, 1–11. https://doi.org/10.1080/23273798.2019.1693050

Morillon, B., Schroeder, C. E., Wyart, V., & Arnal, L. H. (2016). Temporal prediction in lieu of periodic stimulation. *The Journal of Neuroscience*, *36*(8), 2342–2347. https://doi.org/10.1523/JNEUROSCI.0836-15.2016

Müllensiefen, D., Gingras, B., Musil, J., & Stewart, L. (2014). The musicality of non-musicians: An index for assessing musical sophistication in the general population. *PLoS ONE*, *9*(2), e89642. https://doi.org/10.1371/journal.pone.0089642

Nobre, A. C., & van Ede, F. (2018). Anticipated moments: Temporal structure in attention. *Nature Reviews Neuroscience*, *19*, 34–48. https://doi.org/10.1038/nrn.2017.141

Novembre, G., & Iannetti, G. D. (2018). Tagging the musical beat: Neural entrainment or event-related

1481   potentials? *Proceedings of the National Academy of Sciences*, *115*(47), E11002–E11003.
1482   https://doi.org/10.1073/pnas.1815311115

1483 Nozaradan, S., Peretz, I., Missal, M., & Mouraux, A. (2011). Tagging the neuronal entrainment to beat and
1484   meter. *The Journal of Neuroscience*, *31*(28), 10234–10240.
1485   https://doi.org/10.1523/JNEUROSCI.0411-11.2011

1486 Nozaradan, S., Peretz, I., & Mouraux, A. (2012). Selective neuronal entrainment to the beat and meter
1487   embedded in a musical rhythm. *The Journal of Neuroscience*, *32*(49), 17572–17581.
1488   https://doi.org/10.1523/JNEUROSCI.3203-12.2012

1489 Obleser, J., & Kayser, C. (2019). Neural Entrainment and Attentional Selection in the Listening Brain. *Trends*
1490   *in Cognitive Sciences*, *23*(11), 913–926. https://doi.org/10.1016/J.TICS.2019.08.004

1491 Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open source software for
1492   advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational*
1493   *Intelligence and Neuroscience*, *2011*, 156869. https://doi.org/10.1155/2011/156869

1494 O'Reilly, J. X., McCarthy, K. J., Capizzi, M., & Nobre, A. C. (2008). Acquisition of the temporal and ordinal
1495   structure of movement sequences in incidental learning. *Journal of Neurophysiology*, *99*, 2731–
1496   2735. https://doi.org/10.1152/jn.01141.2007

1497 Palmer, C., & Krumhansl, C. L. (1990). Mental representations for musical meter. *Journal of Experimental*
1498   *Psychology. Human Perception and Performance*, *16*(4), 728–741.

1499 Pesnot Lerousseau, J., Trébuchon, A., Morillon, B., & Schön, D. (2021). Frequency selectivity of persistent
1500   cortical oscillatory responses to auditory rhythmic stimulation. *The Journal of Neuroscience*, *41*(38),
1501   7991–8006. https://doi.org/10.1523/JNEUROSCI.0213-21.2021

1502 Polak, R., London, J., & Jacoby, N. (2016). Both Isochronous and Non-Isochronous Metrical Subdivision
1503   Afford Precise and Stable Ensemble Entrainment: A Corpus Study of Malian Jembe Drumming.
1504   *Frontiers in Neuroscience*, *10*. https://www.frontiersin.org/articles/10.3389/fnins.2016.00285

1505 Povel, D.-J., & Essens, P. (1985). Perception of Temporal Patterns. *Music Perception*, *2*(4), 411–440.
1506   https://doi.org/10.2307/40285311

1507 Povel, D.-J., & Okkerman, H. (1981). Accents in equitone sequences. *Perception & Psychophysics*, *30*(6),
1508   565–572. https://doi.org/10.3758/BF03202011

1509 Praamstra, P., Kourtis, D., Kwok, H. F., & Oostenveld, R. (2006). Neurophysiology of implicit timing in
1510   serial choice reaction-time performance. *The Journal of Neuroscience*, *26*(20), 5448–5455.
1511   https://doi.org/10.1523/JNEUROSCI.0440-06.2006

1512 R Development Core Team. (2008). *R: A language and environment for statistical computing*. R Foundation
1513   for Statistical Computing.

1514 Repp, B. H. (2005). Sensorimotor synchronization: A review of the tapping literature. *Psychonomic Bulletin*
1515   *& Review*, *12*(6), 969–992. https://doi.org/10.3758/BF03206433

1516 Rimmele, J. M., Morillon, B., Poeppel, D., & Arnal, L. H. (2018). Proactive sensing of periodic and aperiodic
1517   auditory patterns. *Trends in Cognitive Sciences*, *22*(10), 870–882.
1518   https://doi.org/10.1016/J.TICS.2018.08.003

1519 Saberi, K., & Hickok, G. (2022a). A critical analysis of Lin et al.'s (2021) failure to observe forward
1520   entrainment in pitch discrimination. *European Journal of Neuroscience*, ejn.15778.
1521   https://doi.org/10.1111/ejn.15778

1522 Saberi, K., & Hickok, G. (2022b). Forward entrainment: Psychophysics, neural correlates, and function.
1523   *Psychonomic Bulletin & Review*. https://doi.org/10.3758/s13423-022-02220-y

1524 Schroeder, C. E., & Lakatos, P. (2009a). The Gamma Oscillation: Master or Slave? *Brain Topography*, *22*,
1525   24–26. https://doi.org/10.1007/s10548-009-0080-y

1526 Schroeder, C. E., & Lakatos, P. (2009b). Low-frequency neuronal oscillations as instruments of sensory
1527   selection. *Trends in Neurosciences*, *32*(1), 9–18. https://doi.org/10.1016/j.tins.2008.09.012

1528 Schultz, B. G., Stevens, C. J., Keller, P. E., & Tillmann, B. (2013). The implicit learning of metrical and
1529   nonmetrical temporal patterns. *The Quarterly Journal of Experimental Psychology*, *66*(2), 360–380.
1530   https://doi.org/10.1080/17470218.2012.712146

1531 Shalev, N., Nobre, A. C., & van Ede, F. (2019). Time for What? Breaking Down Temporal Anticipation.
1532   *Trends in Neurosciences*, *42*(6), 373–374. https://doi.org/10.1016/J.TINS.2019.03.002

1533 Stefanics, G., Hangya, B., Hernádi, I., Winkler, I., Lakatos, P., & Ulbert, I. (2010). Phase Entrainment of
1534   Human Delta Oscillations Can Mediate the Effects of Expectation on Reaction Speed. *The Journal*
1535   *of Neuroscience*, *30*(41), 13578–13585. https://doi.org/10.1523/JNEUROSCI.0703-10.2010

1536 Su, Y.-H., & Pöppel, E. (2012). Body movement enhances the extraction of temporal structures in auditory
1537   sequences. *Psychological Research*, *76*(3), 373–382. https://doi.org/10.1007/s00426-011-0346-3

1538 Sun, Y., Michalareas, G., & Poeppel, D. (2021). The impact of phase entrainment on auditory detection is

highly variable: Revisiting a key finding. *European Journal of Neuroscience*, *June*, 1–18. https://doi.org/10.1111/ejn.15367

Tal, I., Large, E. W., Rabinovitch, E., Wei, Y., Schroeder, C. E., Poeppel, D., & Golumbic, E. Z. (2017). Neural entrainment to the beat: The "missing-pulse" phenomenon. *The Journal of Neuroscience*, *37*(26), 6331–6341. https://doi.org/10.1523/JNEUROSCI.2500-16.2017

Teki, S., Grube, M., Kumar, S., & Griffiths, T. D. (2011). Distinct neural substrates of duration-based and beat-based auditory timing. *The Journal of Neuroscience*, *31*(10), 3805–3812. https://doi.org/10.1523/JNEUROSCI.5561-10.2011

Tichko, P., & Large, E. W. (2019). Modeling infants' perceptual narrowing to musical rhythms: Neural oscillation and Hebbian plasticity. *Annals of the New York Academy of Sciences*, *1453*(April), 125–139. https://doi.org/10.1111/nyas.14050

van Atteveldt, N., Musacchia, G., Zion-Golumbic, E., Sehatpour, P., Javitt, D. C., & Schroeder, C. E. (2015). Complementary fMRI and EEG evidence for more efficient neural processing of rhythmic vs. Unpredictably timed sounds. *Frontiers in Psychology*, *6*(OCT), 1–11. https://doi.org/10.3389/fpsyg.2015.01663

van Bree, S., Melcón, M., Kolibius, L. D., Kerrén, C., Wimber, M., & Hanslmayr, S. (2022). The brain time toolbox, a software library to retune electrophysiology data to brain dynamics. *Nature Human Behaviour*, *6*(10), Article 10. https://doi.org/10.1038/s41562-022-01386-8

van Bree, S., Sohoglu, E., Davis, M. H., & Zoefel, B. (2021). Sustained neural rhythms reveal endogenous oscillations supporting speech perception. *PLOS Biology*, *19*(2), e3001142. https://doi.org/10.1371/journal.pbio.3001142

van der Weij, B., Pearce, M. T., & Honing, H. (2017). A probabilistic model of meter perception: Simulating enculturation. *Frontiers in Psychology*, *8*(May), 824. https://doi.org/10.3389/fpsyg.2017.00824

Vuust, P., Pallesen, K. J., Bailey, C., Van Zuijen, T. L., Gjedde, A., Roepstorff, A., & Østergaard, L. (2005). To musicians, the message is in the meter: Pre-attentive neuronal responses to incongruent rhythm are left-lateralized in musicians. *NeuroImage*, *24*(2), 560–564. https://doi.org/10.1016/j.neuroimage.2004.08.039

Wagenmakers, E.-J., Love, J., Marsman, M., Jamil, T., Ly, A., Verhagen, J., Selker, R., Gronau, Q. F., Dropmann, D., Boutin, B., Meerhoff, F., Knight, P., Raj, A., van Kesteren, E.-J., van Doorn, J., Šmíra, M., Epskamp, S., Etz, A., Matzke, D., … Morey, R. D. (2018). Bayesian inference for psychology. Part II: Example applications with JASP. *Psychonomic Bulletin & Review*, *25*(1), 58–76. https://doi.org/10.3758/s13423-017-1323-7

Waschke, L., Kloosterman, N. A., Obleser, J., & Garrett, D. D. (2021). Behavior needs neural variability. *Neuron*, *109*(5), 751–766. https://doi.org/10.1016/j.neuron.2021.01.023

Wilsch, A., Mercier, M. R., Obleser, J., Schroeder, C. E., & Haegens, S. (2020). Spatial Attention and Temporal Expectation Exert Differential Effects on Visual and Auditory Discrimination. *Journal of Cognitive Neuroscience*, *32*(8), 1562–1576. https://doi.org/10.1162/jocn_a_01567

Wollman, I., & Morillon, B. (2018). Organizational principles of multidimensional predictions in human auditory attention. *Scientific Reports*, *8*(1), 13466. https://doi.org/10.1038/s41598-018-31878-5

Zalta, A., Petkoski, S., & Morillon, B. (2020). Natural rhythms of periodic temporal attention. *Nature Communications*, *11*(1), 1051. https://doi.org/10.1038/s41467-020-14888-8

Zhou, H., Melloni, L., Poeppel, D., & Ding, N. (2016). Interpretations of Frequency Domain Analyses of Neural Entrainment: Periodicity, Fundamental Frequency, and Harmonics. *Frontiers in Human Neuroscience*, *10*, 274. https://doi.org/10.3389/fnhum.2016.00274

Zoefel, B., ten Oever, S., & Sack, A. T. (2018). The Involvement of Endogenous Neural Oscillations in the Processing of Rhythmic Input: More Than a Regular Repetition of Evoked Neural Responses. *Frontiers in Neuroscience*, *12*, 95. https://doi.org/10.3389/fnins.2018.00095

Zoefel, B., & Vanrullen, R. (2017). Oscillatory mechanisms of stimulus processing and selection in the visual and auditory systems: State-of-the-art, speculations and suggestions. *Frontiers in Neuroscience*, *11*(May), 296. https://doi.org/10.3389/fnins.2017.00296