1

Fundamental resource specialization of herbivorous butterflies decreases toward

lower latitudes

Running title: Drivers of resource specialization

Ryosuke Nakadai[1,2,5*], Tommi Nyman[3], Koya Hashimoto[4,5], Takaya Iwasaki[6], Anu

Valtonen[1]

[*] Correspondence: R. Nakadai

r.nakadai66@gmail.com

[1] Department of Environmental and Biological Sciences, Faculty of Science and

Forestry, University of Eastern Finland, Yliopistokatu 7, FI-80101 Joensuu, Finland

[2] Department of Ecosystem Studies, Graduate School of Agricultural and Life Sciences,

The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan

[3] Department of Ecosystems in the Barents Region, Norwegian Institute of Bioeconomy

Research, Svanhovd 35, NO-9925, Svanvik, Norway

[4] Department of Environmental Management, Faculty of Agriculture, Kindai University,

Nakamachi, 3327-204, Nara, Nara, 631-8505, Japan

[5] Biodiversity division, National Institute for Environmental Studies, Onogawa 16-2,

Tsukuba, Ibaraki, 305-8506, Japan

[6] Natural Science Division, Faculty of Core Research, Ochanomizu University, 2-1-1

Otsuka, Bunkyo-ku, Tokyo 112-8610, Japan

ORCID: Ryosuke Nakadai (0000-0002-9512-8511)

ORCID: Tommi Nyman (0000-0003-2061-0570)

ORCID: Koya Hashimoto (0000-0003-3510-1453)

ORCID: Takaya Iwasaki (0000-0002-2194-4153)

ORCID: Anu Valtonen (0000-0003-1532-1563)

Acknowledgements

We are particularly grateful to the local entomologists and naturalists who collected the

butterfly data that we compiled for use in this study. We also thank Dr. Matthew

Forister, who provided R code associated with the alpha shape parameter. The

Biodiversity Center of Japan and the Japanese Ministry of the Environment kindly

provided us with datasets on butterfly distributions. Dr. Yasuhiro Sato provided

insightful comments on an early version of the manuscript. Financial support was

provided by the Japanese Society for the Promotion of Science (grant numbers

15J00601 and 18J00093, to R.N.) and the Academy of Finland (grant number 324392,

to A.V.).

Abstract

Aim

Revealing the drivers of variation in resource specialization is a long-standing goal of

ecological and evolutionary research. It is generally assumed that the degree of resource

specialization increases toward lower latitudes. Although herbivorous insects are one of

the best-studied consumer groups, the factors determining their degree of specialization

at large spatial scales are poorly understood. In this work, we investigated the processes

determining spatial variation in "fundamental" resource specialization, which we

defined as the host breadth when not limited by interspecific interactions at the same

trophic level in the species pool. More specifically, we aimed to reveal the drivers of

geographical variation in fundamental resource specialization through macroecological

approaches.

Location

The Japanese archipelago (22°N–45°N) including hemiboreal, temperate, and

subtropical zones.

Taxon

Herbivorous butterflies.

Methods

Species-specific fundamental host breadth was calculated based on both pooled

geographical occurrence and host-use records. We focused on the latitudinal pattern and

investigated significant drivers of the degree of specialization in regional species pools.

As potential drivers, we focused on geography, current climate, diversity and body size

of butterflies, and their changes along geographical gradients. Through Bayesian

structural equation modeling, we investigated the complicated relationships between

these variables and resource specialization with a comprehensive model fitted using

indices of host breadth within each community.

Results

We observed that fundamental resource specialization of butterfly communities at 10

km grid level tend to become more specialized toward higher latitudes. This observed

pattern is contrary to the presumed general trend found in studies based on knowledge

of realized resource specialization within local communities. We found that this pattern

is driven mainly by factors related to climate, butterfly diversity, and body size in each

community.

Main conclusions

Our findings suggest that fundamental resource specialization may show different

latitudinal patterns from the conventional prediction based on knowledge of realized

resource specialization. Our results emphasize the importance of the current climate as a

major factor regulating butterfly morphology and host breadth, regardless of whether

the impact is direct or indirect.

Keywords: Bayesian structural equation modeling, community assembly, continental

islands, host breadth, Pareto distribution, latitudinal gradient, macroecology, trait-based

approach

Introduction

Understanding the processes structuring communities across space and time is a central

subject of ecological research (Cavender-Bares et al., 2009). Resource specialization is

a major factor shaping interspecific interactions and, therefore, community composition

(Nylin et al., 2017; Volf et al., 2017; Dyer & Forister, 2018). The drivers of variation in

the degree of specialization have been intensively investigated in the context of

community ecology (Novotny & Basset, 2005; Forister et al., 2015). Such studies have

generally reported that species tend to be more specialized in the tropics than in the

temperate zone (MacArthur's latitude–niche breadth hypothesis; MacArthur, 1972;

Vázquez & Stevens, 2004; Dyer et al., 2007; reviewed by Dyer & Forister 2018). Using

a global dataset, Forister et al. (2015) addressed variation in resource specialization of

herbivorous insects along a geographical gradient, and showed that the mean and

variance of diet breadth in local communities decrease toward the equator.

　　Resource specialization is driven not only by geographical and climatic gradients, but

also by many other factors. Specifically, the higher number of species in the tropics

(Loder *et al*., 1998; Dyer & Forister, 2018) may intensify interspecific competition,

thereby favoring "species packing" (Dobzhansky, 1950; MacArthur, 1972) via

specialization (Coley & Kursar, 2014). The importance of body size and physiological

constraints imposed by climatic conditions has long been recognized (Bergmann's rule;

Bergmann, 1847). For example, average consumer body size and its variation increase

toward lower latitudes (Loder et al., 1998), although body size is generally positively

correlated with host breadth (Lindström et al., 1994; Pöyry et al., 2017). Identification

of the drivers of specialization is further complicated by the fact that numerous

potentially important factors change in parallel along latitudinal and climatic gradients

(Loder *et al*., 1998).

On the other hand, some studies have found no differences in the degree of

specialization between temperate and tropical regions (e.g., Novotny et al., 2006), and

the hypothesis has been referred to as a "zombie idea" that "still wanders the world

feasting on scientists' brains" after being refuted by data or replaced by a new theory

(Fox, 2011; Moles & Ollerton, 2016). To date, most studies on variation in resource

specialization have relied on data on local communities. However, resource use within

local communities can be shaped by direct or indirect interspecific interactions (Connell,

1980), and local processes of community assembly can produce "realized" resource

specialization, which may mask underlying fundamental resource specialization.

Shifting the focus to fundamental resource specialization can therefore facilitate

understanding processes driving variation in the degree of resource specialization

without considering the complicated effects of local interspecific interactions. In this

work, we investigated the processes driving spatial variation in "fundamental" resource

specialization, which we define as host breadth when not limited by interspecific

interactions at the same trophic level. More specifically, we aimed to reveal the drivers

of geographical variation in fundamental resource specialization using macroecological

approaches.

We focus on angiosperm-feeding butterflies across the Japanese archipelago to reveal

the drivers of spatial variation in resource specialization. We calculated species-specific

fundamental host breadths based on pooled geographical occurrence data and host-use

records. The latitudinal range of the archipelago (22°N–45°N) includes hemiboreal,

temperate, and subtropical zones, which makes it a useful system for assessing spatially

continuous changes in resource specialization, rather than a comparison between

disconnected areas. In fact, most previous studies have simply compared resource

specialization between tropical and temperate regions, which Dyer and Forister (2018)

noted is a major methodological problem. We focused on the latitudinal pattern and

aimed to reveal significant drivers of the degree of specialization in regional species

pools. As potential drivers, we focused on geography, current climate, diversity and

body size of butterflies, and their changes along geographical gradients (Table 1). We

identified drivers of resource specialization using Bayesian structural equation modeling

(Bayesian SEM), which has received increasing attention in ecology and evolution (e.g.,

Arhonditsis et al., 2007; Read et al., 2018; reviewed in Grace et al., 2012). Specifically,

we investigated the complicated relationships between the explanatory variables and

resource specialization using a comprehensive model fitted with a single index

reflecting the distribution of host breadth, and the mean and divergence of phylogenetic

host breadth within each community.

Table 1 Hypothetical trends observed in previous studies and considered in this study

Variable	Predictor		Explanation	References
Climate	Geography	(a)	Temperature, precipitation, and snow vary with latitude and elevation gradients (+/–).	Romero et al., 2018[G]
Butterfly diversity	Geography	(b)	Butterfly diversity varies with latitude (+/–) and elevation (+/–).	Sutton & Collins, 1991[G]; Gutiérrez, 2009[R]; Owens et al., 2017[G]
Butterfly diversity	Climate	(c)	Butterfly diversity varies with climatic conditions (+/–).	Currie et al., 2004[G]; Owens et al., 2017[G]
Butterfly body size	Geography	(d)	Butterfly body size varies with latitude and elevation (+/–).	Park, 1949[R]; Fischer et al., 1960[G]
Butterfly body size	Climate	(e)	Butterfly body size and divergence vary with climatic conditions (+/–).	Park, 1949[R]; Zeuss et al., 2017[R]
Butterfly body size	Butterfly diversity	(f)	Body size is larger in regions with lower butterfly diversity (+).	Hutchinson & MacArthur, 1959[T]
Host breadth	Geography	(g)	Mean and divergence of host breadth decrease toward lower latitudes (–).	Schemske et al., 2009[S]; Forister et al., 2015[G]
Host breadth	Climate	(h)	Mean and divergence of host breadth are lower in higher-energy regions (+).	Schemske et al., 2009[S]; Read et al., 2018[R]
Host breadth	Butterfly diversity	(i)	Higher butterfly diversity intensifies competition, and host breadths therefore	Dobzhansky, 1950[S]; MacArthur,

				become narrower (–).	1972[S]; Coley & Kursar, 2014[S]
Host breadth	Butterfly body size	*	(j)	There is a positive correlation between host breadth and butterfly body size (possibly related to the effectiveness of resource exploitation) (+).	Brown & Maurer, 1989[R]; Lindström et al., 1994[R]; Davis et al., 2013[R]

*Correlative relationship

Letters in superscript represent the focal scale or study type: L: Local, R: Regional, G: Global, S: Synthesis, and T: Theoretical study.

We also hypothesized correlations within each factor (e.g., mean and divergence of body size).

Materials and Methods

Study area

The Japanese archipelago is a long chain of continental islands located off the eastern coast

of Asia, and is recognized as a hotspot of biodiversity (Mittermeier et al., 2011). The mean

temperature of the coldest month ranges from −18.4°C to 22.3°C, while that of the warmest

month ranges from 6.2°C to 29.2°C, and annual precipitation is 700.4–4477.2 mm (Fig. S1a,

b: Japan Meteorological Agency, 1981–2010; http://www.jma.go.jp/jma/indexe.html).

Northern Japan, particularly Hokkaido, receives heavy snowfall during the winter. By

contrast, the Ryukyu Islands, a subtropical region in the southern part of the archipelago, are

characterized by warm temperatures year-round, with little seasonal variation. The western

(Sea of Japan) side of the archipelago typically receives more snow than the eastern (Pacific

Ocean) side (Fig. S1c). The Japanese archipelago belongs to a transitional region between the

paleotropical or Indo-Pacific and Holarctic regions (Cox, 2001), and harbors relict plants that

were distributed throughout a large portion of the Northern Hemisphere during much of the

Tertiary (Milne & Abbott, 2002). More than 280 species of butterflies in five families are

found in the region (Shiro□zu, 2006). With the exception of three lycaenid species, larvae of

Japanese butterfly species feed exclusively on plants (Honda, 2005). Their host plants are

diverse and include both dicot and monocot angiosperms, as well as gymnosperms (Saito et

al., 2016).

Evaluation of butterfly host breadth and body size

Our analyses were based on butterfly census data, including records of 273 species or

subspecies of butterflies throughout the entire Japanese archipelago, available from the

website of the Biodiversity Center of Japan, Ministry of the Environment

(http://www.biodic.go.jp/index_e.html), in grid cells of 5 min latitude and 7.5 min longitude

(about 10 km × 10 km; the Japanese Standard Second Mesh). We used the results of the

fourth and fifth censuses (1988–1991 and 1997–1998, respectively) of the National Survey of

the Natural Environment in Japan (http://www.biodic.go.jp/kiso/15/do_kiso4.html). This

database includes records from 273 species or subspecies of butterflies in grid cells

throughout the Japanese archipelago. We aggregated data at the species level and converted

all records to presence versus absence (1/0) of each species in each grid cell. Following

aggregation, 93,449 records, across all targeted species, were used in the study. For each

species, records were used in combination with relevant climate data to predict potential

distributions throughout Japan using Maxent ver. 3.3.3e (Phillips et al., 2006; analyzed in

Nakadai et al., 2018). To compensate for incomplete distributional data, distribution

probabilities were converted to potential distributions (0/1) using maximum training

sensitivity plus specificity logistic thresholds. Due to difficulties in estimating the potential

distributions of rare species using Maxent (Vaughan & Ormerod, 2005; Breiner et al., 2015),

the presence records of 18 butterfly species with four or fewer records each were used

directly in subsequent analyses.

Information on the host plants of each studied butterfly species was compiled for

evaluation of diet breadth. The host plants of 278 butterfly species or subspecies (3,573

records in total, after excluding duplicates) were obtained from Saito et al. (2016). We used

adult forewing length as a proxy for the body size of each butterfly species. The forewing

lengths of 284 species were derived from Shiro☐zu (2006; published in Nakadai et al., 2020).

For a total of 247 butterfly species, we assembled data on predicted distribution, host plants

and body size. However, nine species were excluded from the analyses for various reasons

(see Table S1). After the non-target species (those with taxonomic changes or non-

angiosperm hosts) were removed, the remaining 3,377 host plant records were selected and

converted into plant-genus resolution for further analyses.

To describe the overall degree to which species are specialized in their host usage, we

estimated the alpha parameter of the discrete truncated Pareto distribution fitted to numbers

of host genera for all butterfly species occurring within each grid cell (Forister et al., 2015;

Kozubowski et al., 2015). For highly nonsymmetrical distributions, the shape parameter (α)

of the discrete truncated Pareto distribution is more informative than measures of central

tendency such as the mean (Forister et al., 2015). Higher values of the Pareto alpha parameter

indicate increased specialization in the community, i.e., a higher proportion of species with

relatively narrow host breadth.

For each cell, we also calculated the grid-level community mean and divergence (based on

Rao's quadratic entropy; RaoQ) of both phylogenetic host breadth and butterfly body size

using the ade4 package (Chessel et al., 2013) in R (R Development Core Team 2017). The

community mean indicates the dominant trait value within an assemblage (Violle et al., 2007),

while community divergence represents the degree of dissimilarity between all possible pairs

of species within the assemblage (Mouchet et al., 2010). To describe the genus-level

phylogenetic host breadth of butterfly species, we applied Faith's phylogenetic diversity

index (cf., Davis et al., 2013). We calculated this index with the R package picante (Kembel

et al., 2010) based on a genus-level angiosperm phylogeny derived from Zanne et al. (2014)

using Phylomatic (Webb & Donoghue, 2005). The methods for estimating phylogenetic host

breadth and the distribution of host breadth followed those of Forister et al. (2015), which

provides detailed descriptions and statistical background information.

To determine whether the observed mean and divergence of each trait in each grid were

greater or less than the values obtained by drawing butterfly species at random from the

regional species pool, we calculated the standardized effect size (SES) using two types of null

models (i.e., null model 1 and 2). "Null model 1" was generated by sampling the grid cell-

specific number of species, without replacement, from the pool of all 247 focal species. This

null model standardizes the values of the indices by removing bias resulting from differences

in the number of species within grid cells (Mason et al., 2007). "Null model 2" was generated

by randomization of the grid cell–species matrix using an independent swap algorithm

(Gotelli, 2000), thereby retaining the original number of species in each grid cell, as well as

the frequency of occurrence of each species across all grid cells. The randomization

procedures for both null models were repeated 1,000 times to generate simulated datasets,

which were then used to compute cell-specific expected values for the mean and divergence

of host breadth and body size. We calculated SES values for each grid cell as the observed

value minus the mean of the null distribution, and the result was divided by the standard

deviation of the null distribution. Therefore, a negative RaoQ value indicates that trait values

within the cell are more similar than expected based on the null model (i.e., convergence

within the community), whereas a positive value indicates that species in the assemblage are

more dissimilar than expected (i.e., divergence within the community). For our main analyses,

we used datasets based on the simpler null model 1, and then re-fitted the final model using

null model 2 to explore the robustness of our results.

Climatic variables

We used annual mean temperature and precipitation at the 1 km grid scale from Mesh

Climate Data 2010 (Japan Meteorological Agency,

http://nlftp.mlit.go.jp/ksj/gml/datalist/KsjTmplt-G02.html). We also included annual

maximum depth of snow cover (Mesh Climate Data 2010), and elevation data from the

National Land Numerical Information database (Geospatial Information Authority of Japan).

All of these variables were downloaded at the 1 km grid scale. Finally, for all variables, we

calculated average values at the 10 km grid scale using ArcGIS Pro ver. 2.3.3 software

(Environmental Systems Resource Institute, Redlands, CA, USA).

Driver analyses for host breadth and body size

To reveal the drivers of resource specialization, we compiled a Bayesian SEM in which the

alpha parameter of the discrete truncated Pareto distribution and the two indices (i.e., mean

and divergence) of phylogenetic host breadth were explained by butterfly species richness,

two indices of body size (i.e., mean and divergence) and five environmental factors. The

environmental factors included as explanatory variables were latitude and elevation as

geographical factors, and annual precipitation, mean temperature and annual maximum depth

of snow cover as climatic factors. Although climatic and geographical factors are correlated

(Table S2), including both types of correlated variables may be informative. For example, a

strong negative correlation exists between latitude and annual mean temperature, but other

effects of latitude may include ecological factors that were not considered in this study, such

as variation in day length and the length of the active season.

Values of *fit* were calculated by the average least-squares differences between the values of

observed and predicted host breadth as $fit = \frac{\sum_{i=1}^{N}(B_{i.obs} - B_{i.pred})^2}{N}$, where $B_{i.obs}$ and $B_{i.pred}$ are

the relative numbers of species with host breadth *i* and N is the maximum host breadth in a

community. Specifically, $B_{i.obs}$ and $B_{i.pred}$ are calculated as the number of species with host

breadth i divided by total species richness in a community. If the total values of observed and

predicted species richness are almost the same, a *fit* ≥ 0.01 indicates that the deviation

between the observed and predicted number of species of each host breadth is more than ten

percent on average. Prior to analyses, two types of grid cells were excluded from the dataset

following Ulrich et al. (2016): cells with fewer than 10 butterfly species and cells with *fit* ≥

0.01. These cells (N = 378 and N = 695, respectively: N = 759 in total) were excluded

because their values were considered less reliable or due to difficulty in correctly estimating

the alpha parameter of the discrete truncated Pareto distribution. For the first criterion, we

repeated the analyses with the minimum number of species set to 15 or 20 to validate the

robustness of the main results.

We fitted the full Bayesian structural equation model in Mplus version 8.4 (Muthén &

Muthén, 1998–2017). Starting parameters were selected using the maximum likelihood

method (Muthén & Muthén, 1998–2017). We ran three Markov chain Monte Carlo analyses

with 1,000,000 iterations from the posterior distribution, using the Metropolis–Hastings

algorithm. Every 100th iteration was sampled from each chain, and the first half of the

sample set was discarded as burn-in, which is the default setting of Mplus. We visually

assessed the status of the posterior distributions and assessed convergence based on the

potential scale reduction factor (PSRF) for each parameter using a threshold of 1.05 (Gelman

& Rubin, 1992), which is also the default setting of Mplus; PSRF should be close to 1 if the

Markov chains have converged to the target posterior distribution (Gelman & Rubin, 1992).

We evaluated model fit based on the Bayesian variant of the root mean square error of

approximation (BRMSEA), the Bayesian comparative fit index (BCFI), and the Bayesian

Tucker-Lewis index (BTLI) (Hoofs et al., 2018, Garnier-Villarreal & Jorgensen, 2019,

Asparouhov & Muthén, 2019). For BCFI and BTLI, Asparouhov and Muthén (2019)

suggested criteria that required values above 0.95 for adequate fit. For BRMSEA,

Asparouhov and Muthén (2019) proposed three possible outcomes: the fit index is

inconclusive (lower limit $\leq 0.06 \leq$ upper limit), the index suggests that the model fits

reasonably well (both lower and upper limits < 0.06), and the index suggests a poor fit

between model and data (both lower and upper limits > 0.06).

After fitting the full model, we trimmed nonsignificant paths from the model until only

significant paths remained (Kline et al., 2011). We explored the robustness of the main

analysis results by re-fitting the final model for each response variable using datasets of

observed distributions rather than the fundamental distributions estimated by Maxent, null

model 2, and the two minimum numbers of butterfly species noted above (i.e., 15 and 20)

within each grid cell.

Results

Throughout the Japanese archipelago, butterfly species richness increased toward lower

latitudes (south) and higher elevations (Fig. 1a), and both the mean (Fig. 1b) and divergence

(Fig. 1c) of butterfly body size increased toward lower latitudes. The Pareto alpha parameter,

which indicates the proportion of species with narrower host breadths, increased toward

higher latitudes (north) (Fig. 1d), while the mean (Fig. 1e) and divergence (Fig. 1f) of

phylogenetic host breadth decreased toward higher latitudes (Table S2).

Figure 1 Patterns of (a) butterfly species richness based on stacked predicted species

distribution data (Nakadai et al., 2018), (b) mean body size of the butterflies, (c) body size

divergence, (d) alpha parameter of the discrete truncated Pareto distribution, (e) mean

phylogenetic host breadth, and (f) phylogenetic host breadth divergence across the Japanese

archipelago. Only grids that satisfied the criteria (see text) are shown in (d), (e), and (f).

Values in all panels except (a) represent standardized effect size (SES) estimates based on

null model 1.

Of the 4,668 total grid cells covering the Japanese archipelago, 3909 passed our filtering

criteria of good fit, as measured using BCFI, BTLI and BRMSEA. Only two of the 52 paths

in the full model were nonsignificant, and therefore were removed during the process of

model simplification (Table 2).

Of the many factors affecting host breadth, annual mean temperature was most strongly

associated with host breadth in the SEM (Table 2, Fig. 2). Specifically, communities

containing low concentrations of species with narrow host ranges (i.e., those with low Pareto

alpha values) increased with annual mean temperature, as did the mean and divergence for

phylogenetic host breadth at the community level (Table 2, Fig. 2). Negative correlations

were found between Pareto alpha and the mean and divergence of body size (Table 2, Fig. 2).

In addition, positive correlations were identified between the mean and divergence of

phylogenetic host breadth and the mean and divergence of body size (Table 2, Fig. 2).

Conversely, no significant correlation was found between body size and host breadth at the

species level (Fig. S2).

Butterfly species richness was driven by annual mean temperature, annual precipitation,

annual maximum depth of snow cover, latitude, and elevation. Based on standardized

estimates, latitude was the strongest (negative) driver of butterfly species richness, while

annual mean temperature was the second most effective (negative) predictor (Table 2, Fig. 2).

The mean of butterfly body size was strongly driven by annual mean temperature

(positively) and latitude (negatively), while the effects of annual precipitation, annual

maximum depth of snow cover and butterfly species richness were relatively weak (Table 2,

Fig. 2). Butterfly body size divergence was driven by latitude, annual mean temperature,

annual precipitation, annual maximum depth of snow cover, elevation and butterfly species

richness. Of these factors, latitude strongly negatively influenced body size divergence (Table

2, Fig. 2). Furthermore, a positive correlation was found between the mean and divergence of

butterfly body size (Table 2, Fig. 2).

The significance and the strength of the major relationships of both host breadth and

butterfly body size with the environmental predictors were consistent when the final models

were re-run with datasets of the observed distributions (Table S4) or null model 2 (Table S5),

and with two alternative criteria for the minimum number of butterfly species per grid cell

(Table S6).

Table 2 Final Bayesian SEM focusing on the alpha parameter of the discrete truncated Pareto

distribution fitted to butterfly host breadth. Abbreviations: LAT: latitude, ELV: elevation,

AP: annual precipitation, AMT: annual mean temperature, SNW: annual maximum depth of

snow cover, BSR: butterfly species richness, BSM: body size mean, BSD: body size

divergence, PAL: alpha parameter of the discrete truncated Pareto distribution, PHBM:

phylogenetic host breadth mean, and PHBD: phylogenetic host breadth divergence.

				95% CI		
	Response	Explanation	Estimate	Lower 2.5%	Upper 2.5%	Significance
Causal effect	AMT	LAT	−0.866	−0.872	−0.859	*
		ELV	−0.477	−0.489	−0.465	*
	AP	LAT	−0.566	−0.586	−0.545	*
		ELV	0.274	0.249	0.297	*
	SNW	LAT	0.504	0.481	0.527	*
		ELV	0.313	0.288	0.337	*
	BSR	AMT	−0.489	−0.638	−0.341	
		AP	−0.135	−0.166	−0.103	*
		SNW	−0.197	−0.227	−0.166	*
		LAT	−0.771	−0.906	−0.637	*
		ELV	0.401	0.326	0.476	*
	BSM	AMT	0.381	0.335	0.428	*
		AP	0.204	0.180	0.226	*
		SNW	−0.133	−0.155	−0.110	*
		LAT	−0.322	−0.375	−0.268	*
		ELV	−	−	−	
		BSR	−0.129	−0.152	−0.106	*
	BSD	AMT	0.118	0.042	0.193	
		AP	0.251	0.230	0.272	*
		SNW	−0.200	−0.220	−0.179	*
		LAT	−0.486	−0.559	−0.413	*
		ELV	−0.183	−0.219	−0.148	*

		BSR	−0.290	−0.312	−0.269	*
	PAL	AMT	−0.866	−0.919	−0.814	*
		AP	-0.234	-0.261	-0.207	*
		SNW	0.091	0.065	0.118	*
		LAT	−0.407	−0.470	−0.345	*
		ELV	–	–	–	
		BSR	−0.408	−0.435	−0.381	*
	PHBM	AMT	0.971	0.869	1.075	*
		AP	0.235	0.207	0.262	*
		SNW	−0.226	−0.253	−0.200	*
		LAT	0.647	0.549	0.747	*
		ELV	0.069	0.020	0.119	*
		BSR	0.470	0.443	0.498	*
	PHBD	AMT	1.030	0.911	1.15	*
		AP	0.293	0.264	0.323	*
		SNW	−0.078	−0.107	−0.048	*
		LAT	0.824	0.711	0.936	*
		ELV	0.105	0.046	0.163	*
		BSR	0.641	0.613	0.669	*
Co-variate	PAL	BSM	−0.172	−0.202	−0.141	*
		BSD	−0.233	−0.262	−0.204	*
		PHBM	−0.656	−0.673	−0.638	*
		PHBD	−0.589	−0.609	−0.568	*
	PHBM	BSM	0.288	0.259	0.317	*
		BSD	0.363	0.335	0.390	*
		PHBD	0.862	0.853	0.869	*
	PHBD	BSM	0.176	0.145	0.206	*
		BSD	0.234	0.205	0.264	*
	BSM	BSD	0.712	0.697	0.727	*
	AMP	SNW	0.329	0.300	0.356	*

	Model fit (N=3909)		

	Lower 5.0%	Upper 5.0%	
BCFI	0.998	0.999	††
BTLI	0.982	0.990	††
BRMSEA	0.051	0.068	†
BIC		171233.876	

††Satisfied the criteria for good model fit for BCFI and BTLI (both lower 5.0% and upper 5.0% > 0.95;

Asparouhov & Muthén, 2019), and BRMSEA (both lower 5.0% and upper 5.0% < 0.06; Asparouhov & Muthén,

2019).

†Satisfied only the criteria for inconclusive model fit for BRMSEA (lower limit ≤ 0.06 ≤ upper limit;

Asparouhov & Muthén, 2019).

*Confidence interval (CI) did not overlap zero

–Paths trimmed during model modification (original full model is provided in Table S3)

Figure 2 Path diagram of Bayesian SEM explaining host-breadth parameters. Parameter estimates are shown in Tables 2 and S3. Red arrows indicate significant positive effects or correlations, and blue arrows indicate significant negative ones. Line weights indicate the strength of the standardized estimates (see legend). Black dashed lines indicate the paths removed during the modification process. Abbreviations: LAT: latitude, ELV: elevation, AP: annual precipitation, AMT: annual mean temperature, SNW: annual maximum depth of snow

PAL: alpha parameter of the discrete truncated Pareto distribution, PHBM: phylogenetic host

breadth mean, and PHBD: phylogenetic host breadth divergence.

Discussion

Latitudinal pattern of fundamental host breadth is opposite to classic predictions based on

realized host breadth

One of the most widespread assumptions in ecology is that the degree of resource

specialization increases toward lower latitudes (i.e., toward the tropics) (Moles & Ollerton,

2016; Dyer & Forister, 2018). By contrast, our results show that the host breadth of butterfly

communities becomes more specialized at higher latitudes (i.e., toward the north) across the

Japanese archipelago. Thus, our finding apparently contradicts the widespread assumption of

how species combine to form local communities. According to our results, the host breadth of

butterfly communities was most strongly associated with climatic factors (i.e., annual mean

temperature, annual precipitation, annual maximum depth of snow cover), a geographic

factor (latitude), and butterfly species richness. Specifically, both mean and divergence of

host breadth of butterfly communities were higher where temperatures were higher.

Furthermore, grid cells with high butterfly diversity showed broader and more divergent host

ranges.

The trend of broader and more divergent host ranges within communities at lower

latitudes may be produced due to relaxing of physiological constraints with increased energy

availability, thereby reducing the relative importance of abiotic environmental filters and

allowing for wider variation in niches (Park, 1949; Fischer, 1960). Read et al. (2018)

suggested that relaxing of physiological constraints combined with stronger biotic

interactions among species toward the tropics leads to increased resource specialization,

which is the opposite trend to our findings. As mentioned in the Introduction, this study

focused on fundamental host breadth, which was defined as the host breadth when not limited

by interspecific interactions within a trophic level. Evidently, without considering

interspecific interactions, the relaxation of abiotic regulation toward lower latitudes allows

for the occurrence of butterfly species with broader host ranges. Thus, our results appear

contrary to the conventional assumption of latitudinal gradients in resource specialization.

However, previous studies have failed to clarify the constraints of interspecific interactions

on resource specialization among herbivorous insects. In fact, the existence of strong

interspecific resource competition among herbivorous insects, which would cause niche

partitioning, has repeatedly been questioned (Lawton & Strong, 1981; Strong et al., 1984;

Condon et al., 2014; Nakadai & Kawakita, 2017). Nakadai et al. (2018) showed that butterfly

species pairs that use similar host-plant taxa tend to have overlapping distributions in the

Japanese archipelago. Thus, the relevance of interspecific competition to the latitudinal

gradient in the diet breadth of herbivorous insects is not clear.

Along the focal latitudinal and climatic gradients, many variables other than host breadth

may change together or independently, including both the body size and diversity of

butterflies. Teasing apart the directionality of causalities among these patterns is a long-

standing research subject in ecology (Wasserman & Mitter, 1978; Loder et al., 1998).

Consistent with previous studies on plant-feeding insects (Brown & Maurer 1989; Pöyry et

al., 2017), we identified a strong positive relationship between the mean body size and mean

host breadth at the grid community level. Lindström et al. (1994) argued that larger body size

in lepidopterans allows for a larger "home range," and thereby increases encounters with a

wide range of host plants, but also requires more energy, which is more easily acquired if the

species is a broad generalist. In our dataset, both the mean and divergence of body size

increased toward lower latitudes and higher temperatures (Tables 2). This pattern of

diminishing body sizes toward higher latitudes is referred to as a converse Bergmann cline

(although most interspecific studies on insects have found no clines; Shelomi, 2012). The

southward increases in both the mean and divergence of body size are consistent with the

hypothesis based on relaxed physiological constraints (Table 1). Furthermore, mean body

size is positively correlated with mean phylogenetic host breadth. Increasing mean body size

within a community may therefore be one of the factors enhancing diet generalization in

communities of herbivorous insects (Table 1). At higher latitudes or elevations, shorter

growing seasons also limit the duration of the larval stage, restricting the distributions of

large-bodied or slow-developing insects (Blanckenhorn & Demont, 2004; Shelomi, 2012).

For example, Zeuss et al. (2017) reported that among European lepidopteran species, average

body size and the average number of generations per year decrease with latitude. However,

we did not detect significant correlations between body size and host breadth at the species

level (Fig. S2). Therefore, the relationship between mean body size and mean phylogenetic

host breadth at the grid level may simply reflect correlations driven by the responses of both

variables to the same underlying environmental predictors (e.g., annual mean temperature).

Potential drivers of contrasting patterns between fundamental and realized resource

specialization

The largest difference between our analytical approach and those of previous studies

concerns the properties in the herbivore community dataset. The analyses of Forister et al.

(2015) and many others (e.g., Novotny et al., 2006) were based on direct observations of

insect–plant associations in local communities. By contrast, our 'species-pool' approach

based on modeled distributions may combine the distributions of some species pairs that

would typically not occur at the same site at the same time, for reasons such as phenological

niche division. Our analyses did not focus on phenological patterns along the latitudinal

gradient, as our main focus was on "fundamental" patterns at a large spatial scale. If the

difference between our results and those of other studies (e.g., Forister et al., 2015) are due to

differences in the approach between fundamental and realized resource specialization, biotic

filtering across trophic levels (i.e., tri-trophic interactions; reviewed in Schemske et al., 2009)

may be an important local-scale factor (Cavender-Bares et al., 2009). For example, narrow

and overlapping host breadths may be strongly promoted by host-specific natural enemies

(e.g., Condon et al., 2014), and enemy pressure is thought to be greater in the tropics (Roslin

et al., 2017; Romero et al., 2018). Local adaptation of herbivorous butterflies is a frequent

phenomenon (e.g., Singer & Parmesan, 1993; reviewed in Gandon & Van Zandt, 1998), and

locally specialized enemies related to host plants and variation in plant resistance to

herbivory (i.e., coevolutionary interactions) may result in large variations in host plants

among local sites, in turn leading to wider host breadths in focal species at the regional level.

It seems possible for a species to have high local specialization in a lower latitude community,

even if its members tend to have a relatively broad host breadths throughout their range. Over

time, this should translate into diversification among these specialized local populations,

creating the expected positive latitudinal gradient in host breadth. However, because the

Japanese archipelago does not include stable tropical regions, it is unclear whether there has

been enough time for local plant species to adapt to the full butterfly fauna, or a subset

thereof.

Methodological considerations

To elucidate the detailed drivers of host breadth in Japanese butterflies, we attempted to

include as many factors as possible in our analyses. However, our results also reveal several

important limitations of current methods that must be addressed in future studies.

Unfortunately, we could not directly evaluate the effects of regional plant diversity on the

geographic distribution of resource specialization. Plant diversity is highest near the center (at

circa N 35°) of the Japanese archipelago (e.g., Kubota et al., 2015). In fact, Kubota et al.

(2015) demonstrated that overall vascular plant richness and endemism are highest in central

Japan, and that elevation and annual mean temperature are major drivers of the pattern,

indicating that the trend is driven by more than just latitude. Central Japan is characterized by

mountainous regions, and elevation may be an indirect predictor of plant diversity. In fact,

the patterns of the simple correlations in our study indicated that elevation is strongly

correlated with butterfly species richness but is less correlated with host breadth (Table S2).

Therefore, we doubt that plant diversity is the strongest driver of fundamental host breadth.

In general, the proportion of rare plant species increases toward the tropics (e.g., Ulrich et

al., 2016), possibly due to negative density-dependent survival effects (Janzen–Connell

hypothesis; Janzen, 1970). Such processes were suggested as a fundamental explanation for

the contrasting latitudinal pattern (i.e., more generalists toward lower latitudes) of resource

specialization found in the simulation results of Forister and Jenkins (2017). Specifically, a

high proportion of rare plants would favor the evolution of greater host breadth in insects,

causing the mean and divergence of host breadth to increase toward lower latitudes (Ghazoul

& Sheil, 2010). The evenness of the biomass of plant species is a potential driver of resource

specialization, although the evenness of plant abundance (e.g., number of individuals) was

used as an explanatory variable in previous simulation studies (Forister & Jenkins 2017). In

any case, data on both plant biomass and abundance were not available for use in our study or

in previous studies. Therefore, investigations focusing on the available biomass of resources

are needed to further explore the effects of plant community composition on herbivore host

breadth. Furthermore, we were unable to include phylogenetic information on butterfly

species in our analysis. Both life history traits and distributions may be affected by

evolutionary history (i.e., phylogenetic conservatism). This is a limitation of both our study

and previous investigations and may have introduced bias. Phylogenetic information

hopefully can be incorporated into models in the future, when the lepidopteran tree of life

becomes better resolved.

In addition, we focused mainly on the transitional region between the hemiboreal and

subtropical zones, and the patterns observed may not be generalizable to the global scale.

Regional studies focusing on narrower latitudinal gradients should consider geohistorical

effects (e.g., when the region was colonized by particular species or where hotspots of

endemism are located). Our findings also may differ from the patterns observed in previous

studies (e.g., Forister et al., 2015) because of our more limited geographic scope, i.e., the

absence of large tropical regions which have significant in-situ diversification rates. The

effects of co-evolutionary interactions may be absent as there has been relatively little in-situ

diversification within our study area. In fact, most butterfly species on the Japanese

archipelago are not endemics but rather are widely distributed across the Eurasian continent.

This pattern is largely explained by the fact that the Japanese archipelago was repeatedly

connected to the continent with land bridges throughout glacial stages of the Pliocene and

Pleistocene epochs, allowing for butterfly dispersal (details reviewed in Tojo et al., 2017). On

the other hand, detailed regional studies have the advantage of providing extensive and

accurate information on host use. Previous studies have suggested that sampling effort

directly affects the inferred degree of host specialization (Dixon et al., 1987), but such effects

are difficult, if not impossible, to address in a large-scale meta-analysis.

Historical and evolutionary perspectives

We did not address evolutionary processes in this study, as phylogenetic information was

not available for all of the studied butterfly species. However, our findings should be

generalizable to other regions and broader spatial patterns, as they are based on evolutionary

processes. For example, we could expand on the evolutionary concept of species association

dynamics recently proposed by Nylin et al. (2017) and the oscillation hypothesis proposed by

Janz et al. (2001, 2006), which focus on speciation and species distributions, by integrating

spatial perspectives by adding a latitudinal direction of oscillation dynamics. The oscillation

hypothesis proposed by Janz et al. (2001, 2006) postulates that, over time, herbivorous insect

lineages oscillate between a generalist state, during which they may expand their

geographical ranges, and various specialist states, which present outcomes of local adaptation

of individual populations. In this model, speciation occurs after host specialization. To

explore this issue further, we plotted minimum and maximum values of phylogenetic host

breadth and body size within butterfly communities against latitude (Fig. S3), and confirmed

that the ranges of these trait values become increasingly narrow toward higher latitudes. If

physical constraints on host breadth or body size in insect herbivores exist (Fig. S3), very

large or very small butterfly species should have difficulty extending their distribution ranges

from the south to the north, and therefore such extreme species would be filtered out. Our

findings are consistent with the proposal of Fischer (1960), who argued that the stable climate

in the tropics allows for a wider range of morphological and physiological adaptations than

can occur in temperate regions. On the other hand, the spread of butterfly species with

broader host ranges from south to north might be hindered by biogeographic limitations

(Kimura, 2004; Hirao et al., 2015). Therefore, we propose a novel prediction: butterfly

species with large host breadths are more likely to have expanded their distributions to the

south or to have originated in the south and then specialized and diversified there via

oscillation processes, and only descendant species with narrower host ranges could migrate

northward. Although biogeographic limitations and abiotic factors (e.g., temperature,

precipitation, and snow) are important determinants of butterfly distributions (Quinn et al.,

1998; Luoto et al., 2006), our predictions provide further insights into the importance of

geographical and morphological constraints to species association dynamics and the

oscillation hypothesis. These predictions may also be combined with general ecological

principles such as Rapoport's rule (Rapoport, 1982), which is related to latitudinal

distribution range and global climate history and their dramatic effects on species

distributions (Nyman et al., 2012). Specifically, Rapoport's rule would alter the distributional

range of each species, which may affect the possibility of local resource specialization via

northward migration. In addition, climatically driven distributional changes along a north–

south axis, as well as associated changes in the abundance of different plant groups, have

been proposed as factors influencing the patterns and processes of herbivore diversification

(Nyman et al., 2012). Therefore, the inclusion of dynamic distributional changes along a

latitudinal gradient may generalize the oscillation hypothesis for understanding the

mechanisms that drive biodiversity.

Conclusions

Butterfly communities become more specialized in their host use toward higher latitudes

(i.e., toward the north) across the Japanese archipelago, a pattern that is opposite of the

predictions arising from classic hypotheses. Our analyses clearly revealed how the variation

in resource specialization is distributed, as well as the potential influence of physiological

constraints resulting from variation in body size along the same geographical gradient. Since

the work of Ehrlich and Raven (1964), coevolutionary interactions between plants and

herbivores have been considered the main determinant of host breadth. However, our results

highlight the importance of the current climate as a factor regulating butterfly host breadth

and morphology (Fischer, 1960), regardless of whether the impact is direct or indirect. Future

studies should explore how latitudinal gradients in the phylodiversity and evenness of

herbivores and plant resources influence community-level host breadth. We emphasize that

the approach used here is applicable to many other areas and taxa for which reliable

information on species occurrences and niches are available. The continuous improvement of

public databases on species distributions and host use (e.g., Global Biodiversity Information

Facility and HOSTs: Robinson et al. 2010) will hopefully eventually allow testing of patterns

of fundamental resource specialization on a global scale. Importantly, comparative analyses

of fundamental and realized resource specialization in communities can allow important

insights into factors that determine the assembly and local realizations of eco-evolutionary

interaction networks across the globe.

Data accessibility statement

Information on the forewing lengths of Japanese butterflies was published by Nakadai et al.

(2020). Environmental and geographic data were obtained from Mesh Climate Data 2010

(Japan Meteorological Agency, http://nlftp.mlit.go.jp/ksj/gml/datalist/KsjTmplt-G02.html)

and the National Land Numerical Information database (Geospatial Information Authority of

Japan), respectively. The host plant dataset was taken from Saito et al. (2016). Occurrence

data for butterfly species are available from the website of the Biodiversity Center of Japan,

Ministry of the Environment (http://www.biodic.go.jp/index_e.html).

References

Arhonditsis, G.B., Stow, C.A., Paerl, H.W., Valdes-Weaver, L.M., Steinberg, L.J. &
Reckhow, K.H. (2007). Delineation of the role of nutrient dynamics and hydrologic
forcing on phytoplankton patterns along a freshwater-marine continuum. *Ecol. Modell.*,
208, 230–246.

Asparouhov, T. & Muthén, B. (2019). Advances in Bayesian model fit evaluation for
structural equation models. http://www.statmodel.com/download/BayesFit.pdf

Bergmann, C. (1847). Über die Verhältnisse der Wärmeökonomie der Thiere zu ihrer Grösse.
Gött. Stud., 1595–708.

Blanckenhorn, W.U., & Demont, M. (2004). Bergmann and converse Bergmann latitudinal
clines in arthropods: Two ends of a continuum? *Integr. Comp. Biol.* 44, 413–424.

Breiner, F. T., Guisan, A., Bergamini, A., & Nobis, M. P. (2015). Overcoming limitations of
modelling rare species by using ensembles of small models. *Methods Ecol. Evol.*, 6,
1210–1218.

Brown, J. H., & Maurer, B. A. (1989). Macroecology: The division of food and space among
species on continents. *Science*, 243, 1145–1150.

Cavender-Bares, J., Kozak, K.H., Fine, P.V.A. & Kembel, S.W. (2009). The merging of
community ecology and phylogenetic biology. *Ecol. Lett.*, 12, 693–715.

Chessel, D., Dufour, A.-B. & Dray, S. (2018). CRAN R package ade4. Analysis of ecological
data, version 1.7–13. R Foundation for Statistical Computing, Vienna. https://cran.r-
project.org/web/packages/ade4/.

Condon, M.A., Scheffer, S.J., Lewis, M.L., Wharton, R., Adams, D.C., & Forbes, A.A.
(2014). Lethal interactions between parasites and prey increase niche diversity in a
tropical community. *Science*, 343, 1240–1244.

Connell, J.H. (1980). Diversity and the coevolution of competitors, or the ghost of
competition past. *Oikos*, 35, 131–138.

Coley, P. D., & Kursar, T. A. (2014). On tropical forests and their pests. *Science*, 343, 35–36.

Cox, B. (2001). The biogeographic regions reconsidered. *J. Biogeogr.*, 28, 511–523.

Currie, D. J., Mittelbach, G. G., Cornell, H. V., Field, R., Guégan, J. F., Hawkins, B. A. et al.
(2004). Predictions and tests of climate-based hypotheses of broad-scale variation in
taxonomic richness. *Ecol. Lett.*, 7, 1121–1134.

Davis, R.B., Õunap, E., Javoiš, J., Gerhold, P. & Tammaru, T. (2013). Degree of

specialization is related to body size in herbivorous insects: A phylogenetic confirmation.

Evolution, 67, 583–589.

Dixon, A. F. G., Kindlmann, P., Leps, J., & Holman, J. (1987). Why there are so few species

of aphids, especially in the tropics. *Am. Nat.*, 129, 580–592.

Dobzhansky, T. (1950). Evolution in the tropics. *Am. Sci.*, 38, 209–221.

Dyer, L.A. & Forister, M.L. (2018). Challenges and advances in the study of latitudinal

gradients in multitrophic interactions, with a focus on consumer specialization. *Curr.*

Opin. Insect Sci., 1–9.

Dyer, L. A., Singer, M. S., Lill, J. T., Stireman, J. O., Gentry, G. L., Marquis, R. J. et al.

(2007). Host specificity of Lepidoptera in tropical and temperate forests. *Nature*, 448,

696.

Ehrlich, P.R. & Raven, P.H. (1964). Butterflies and plants: A study in coevolution. *Evolution*,

18, 586.

Fischer, A. G. (1960). Latitudinal variations in organic diversity. *Evolution*, 14, 64–81.

Forister, M.L. & Jenkins, S.H. (2017). A neutral model for the evolution of diet breadth. *Am.*

Nat., 190, E40–E54.

Forister, M.L., Novotny, V., Panorska, A.K., Baje, L., Basset, Y. & Butterill, P.T. (2015).

The global distribution of diet breadth in insect herbivores. *Proc. Natl. Acad. Sci. USA*,

112, 442–447.

Fox, J. W. 2011. Zombie ideas in ecology.

https://dynamicecology.wordpress.com/2011/06/17/zombie-ideas-in-ecology/.

Gandon, S., & Van Zandt, P.A. (1998). Local adaptation and host–parasite interactions.

Trends Ecol. Evol., 13, 214–216.

Garnier-Villarreal, M., & Jorgensen, T.D. (2019). Adapting fit indices for Bayesian structural

equation modeling: Comparison to maximum likelihood. *Psychol. Methods*, 25, 46–70

Gelman, A. & Rubin, D.B. (1992). Inference from iterative simulation using multiple

sequences. *Stat. Sci.*, 7, 457–472.

Ghazoul, J., & Sheil, D. (2010). Tropical rain forest ecology, diversity, and conservation.

Oxford University Press, Oxford.

Gotelli, N.J. (2000). Null model analysis of species co-occurrence patterns. *Ecology*, 81,

2606–2621.

Grace, J.B., Schoolmaster, D.R., Guntenspergen, G.R., Little, A.M., Mitchell, B.R., Miller,

K.M. et al. (2012). Guidelines for a graph-theoretic implementation of structural

equation modeling. *Ecosphere*, 3, 1–44.

Gutiérrez, D. (2009). Butterfly richness patterns and gradients. Ecology of butterflies in

Europe (ed. by J. Settele, T. Shreeve, M. Konvicka and H. van Dyck), pp. 250–280.

Cambridge University Press, Cambridge.

Hawkins, B.A., Rueda, M., Rangel, T.F., Field, R., & Diniz-Filho, J.A.F. (2014). Community

phylogenetics at the biogeographical scale: Cold tolerance, niche conservatism and the

structure of North American forests. *J. Biogeogr.*, 41, 23–38.

Hirao, T., Kubota, Y. & Murakami, M. (2015). Geographical patterns of butterfly species

diversity in the subtropical Ryukyu Islands: The importance of a unidirectional filter

between two source islands. *J. Biogeogr.*, 42, 1418–1430.

Honda K (2005). Larval feeding habit and host selection. In: Honda K, Kato Y (eds) Biology

of butterflies. University of Tokyo Press, Tokyo, pp 255–301 (in Japanese)

Hoofs, H., van de Schoot, R., Jansen, N.W.H. & Kant, Ij. (2018). Evaluating model fit in

Bayesian confirmatory factor analysis with large samples: Simulation study introducing

the BRMSEA. *Educ. Psychol. Meas.*, 78, 537–568.

Hutchinson, G.E., & MacArthur, R.H. (1959). A theoretical ecological model of size

distributions among species of animals. *Am. Nat.*, 93, 117–125.

Inomata T (1990). Keys to the Japanese butterflies in natural color. Hokuryukan, Tokyo (In

Japanese)

Janz, N., Nyblom, K. & Nylin, S. (2001). Evolutionary dynamics of host-plant specialization:

A case study of the tribe Nymphalini. *Evolution*, 55, 783–796.

Janz, N., Nylin, S. & Wahlberg, N. (2006). Diversity begets diversity: Host expansions and

the diversification of plant-feeding insects. *BMC Evol. Biol.*, 6, 4.

Kimura, M.T. (2004). Cold and heat tolerance of drosophilid flies with reference to their

latitudinal distributions. *Oecologia*, 140, 442–449.

Kembel, S.W., Cowan, P.D., Helmus, M.R., Cornwell, W.K., Morlon, H., Ackerly, D. D. et

al. (2010). Picante: R tools for integrating phylogenies and ecology. *Bioinformatics*, 26,

1463–1464.

Kline, R.B., (2011). Principles and practice of structural equation modeling, The Guilford

Press, New York, NY, USA, 3rd edition.

Kozubowski, T.J., Panorska, A.K. & Forister, M.L. A discrete truncated Pareto distribution.

Stat. Methodol., 26, 135–150.

Lawton, J.H., & Strong, D.R. (1981). Community patterns and competition in folivorous

insects. *Am. Nat.*, 118, 317–338

Lindström, J., Kaila, L., & Niemelä, P. (1994). Polyphagy and adult body size in geometrid

moths. *Oecologia*, 98, 130–132.

Loder, N., Gaston, K.J. & Warren, P.H. (1998). Body size and feeding specificity:

Macrolepidoptera in Britain.

Luoto, M., Heikkinen, R.K., Pöyry, J. & Saarinen, K. (2006). Determinants of the

biogeographical distribution of butterflies in boreal regions. *J. Biogeogr.*, 33, 1764–1778.

MacArthur, R. H. (1972). Geographical ecology: Patterns in the distribution of species

Princeton University Press, Princeton.

Mason, N.W.H., Lanoiselée, C., Mouillot, D., Irz, P. & Argillier, C. (2007). Functional

characters combined with null models reveal inconsistency in mechanisms of species

turnover in lacustrine fish communities. *Oecologia*, 153, 441–452.

Milne, R.I. & Abbott, R.J. (2002). The origin and evolution of tertiary relict floras. *Adv. Bot.*

Res., 38, 281–314.

Mittermeier, R.A., Turner, W.R., Larsen, F.W., Brooks, T.M., & Gascon, C. (2011). Global

biodiversity conservation: The critical role of hotspots. In: Zachos FE, Habel JC (eds)

Biodiversity hotspots. Springer, Berlin, pp 3–22

Moles, A.T. & Ollerton, J. (2016). Is the notion that species interactions are stronger and

more specialized in the tropics a zombie idea? *Biotropica*, 48, 141–145.

Mouchet, M.A., Villéger, S., Mason, N.W.H. & Mouillot, D. (2010). Functional diversity

measures: An overview of their redundancy and their ability to discriminate community

assembly rules. *Funct. Ecol.*, 24, 867–876.

Muthén, L.K. and Muthén, B.O. (1998–2017). Mplus user's guide. eighth edition. Los

Angeles, CA

Nakadai, R., Hashimoto, K., Iwasaki, T. & Sato, Y. (2018). Geographical co-occurrence of

butterfly species: The importance of niche filtering by host plant species. *Oecologia*,

186, 995–1005.

Nakadai, R., Kawakita, A. (2017). Patterns of temporal and enemy niche use by a community

of leaf cone moths (*Caloptilia*) coexisting on maples (*Acer*) as revealed by

metabarcoding. *Mol. Ecol.*, 26, 3309–3319

Nakadai, R., Kobayashi, T., Hashimoto, K. (2020). Dataset of forewing length of Japanese

and Taiwanese butterfly species. *Ecol. Res.*, 35: 780-786

Novotny, V. & Basset, Y. (2005). Host specificity of insect herbivores in tropical forests.

Proc. Roy. Soc. B., 272, 1083–1090.

Novotny, V., Drozd P., Miller S.E., Kulfan, M., Janda, M., Basset, Y. et al. (2006). Why are

there so many species of herbivorous insects in tropical rainforests? *Science* 313, 1115–

1118.

Nylin, S., Agosta, S., Bensch, S., Boeger, W.A., Braga, M.P., et al. (2017). Embracing

colonizations: A new paradigm for species association dynamics. *Trends Ecol. Evol.*, 33,

4–14.

Nyman, T., Linder, H.P., Peña, C, Malm., T. & Wahlberg, N. (2012). Climate-driven

diversity dynamics in plants and plant-feeding insects. *Ecol. Lett.* 15: 889–898

Owens, H., Lewis, D., Dupuis, J., Clamens, A., Sperling, F., Kawahara, A., et al. (2017). The

latitudinal diversity gradient in New World swallowtail butterflies is caused by

contrasting patterns of out-of- and into-the-tropics dispersal. *Glob. Ecol. Biogeogr.*,

1447–1458.

Park, O. (1949). Application of the converse Bergmann principle to the carabid beetle,

Dicaelus purpuratus. *Physiol. Zool.*, 22, 359–372.

Phillips, S.J., Anderson, R.P. & Schapire, R.E. (2006). Maximum entropy modeling of

species geographic distributions. *Ecol. Modell.*, 190, 231–259.

Pöyry J., Carvalheiro L.G., Heikkinen R.K., Kuhn I., Kuussaari M., Schweiger O. et al.

(2017). The effects of soil eutrophication propagate to higher trophic levels Global *Ecol.*

Biogeogr., 26, 18–30

Quinn, R.M., Caston, K.J., & Roy, D.B. (1998). Coincidence in the distributions of butterflies

and their foodplants. *Ecography*, 21, 279–288.

Rapoport, E. H. (1982). Areography. Geographical strategies of species. Pergamon Press,

Oxford, UK.

R Development Core Team (2017). R: A language and environment for statistical computing.

R Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org/.

Read, Q.D., Grady, J.M., Zarnetske, P.L., Record, S., Baiser, B., Belmaker, J., et al. (2018).

Among-species overlap in rodent body size distributions predicts species richness along

a temperature gradient. *Ecography*., 41, 1718–1727.

Robinson, G.S., P. R. Ackery, I.J. Kitching, G.W. Beccaloni & L.M. Hernández, (2010).

HOSTS - A Database of the World's Lepidopteran Hostplants. Natural History Museum,

London. http://www.nhm.ac.uk/hosts.

Romero, G.Q., Gonçalves-Souza, T., Kratina, P., Marino, N.A., Petry, W.K., Sobral-Souza,

T., et al. (2018). Global predation pressure redistribution under future climate change.

Nature Climate Change, 8, 1087.

Roslin, T., Hardwick, B., Novotny, V., Petry, W.K., Andrew, N.R., Asmus, A., et al. (2017).

Higher predation risk for insect prey at low latitudes and elevations. *Science*, 356, 742–

744.

Saito, M.U., Jinbo, U., Yago, M., Kurashima, O. & Ito, M. (2016). Larval host records of

butterflies in Japan. *Ecol. Res*., 31, 491–491.

Shelomi, M. (2012). Where are we now? Bergmann's rule sensu lato in insects. *Am. Nat*., 180,

511–519.

Shiro☐zu, T. (2006). The butterflies of Japan in color. Gakken Holdings, Tokyo (In

Japanese)

Singer, M.C., & Parmesan, C. (1993). Sources of variations in patterns of plant–insect

association. *Nature*, 361, 251–253.

Strong, D.R., Lawton, J.H., & Southwood, S.R. (1984). Insects on plants. Community

patterns and mechanisms. Harvard University Press, Cambridge

Sutton, S.L., & Collins, N.M. (1991). Insects and tropical forest conservation. The

conservation of insects and their habitats, 405–424. Academic Press

Tojo, K., Sekiné, K., Takenaka, M., Isaka, Y., Komaki, S., Suzuki, T., et al. (2017). Species
diversity of insects in Japan: Their origins and diversification processes. *Entomol. Sci.*,
20, 357–381.

Ulrich, W., Kusumoto, B., Shiono, T., & Kubota, Y. (2016). Climatic and geographic
correlates of global forest tree species–abundance distributions and community evenness.
J. Vege Sci, 27: 295–305.

Vaughan, I.P., & Ormerod, S.J. (2005). The continuing challenges of testing species
distribution models. *J. Appl. Ecol.*, 42, 720–730.

Vázquez, D.P., & Stevens, R.D. (2004). The latitudinal gradient in niche breadth: Concepts
and evidence. *Am. Nat.*, 164(1), E1-E19.

Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., et al. (2007). Let
the concept of trait be functional! *Oikos*, 116, 882–892.

Volf, M., Pyszko, P., Abe, T., Libra, M., Kotásková, N., Šigut, M., et al. (2017). Phylogenetic
composition of host plant communities drives plant-herbivore food web structure. *J.*
Anim. Ecol., 86, 556–565.

Wasserman, S.S., & Mitter, C. (1978). The relationship of body size to breadth of diet in
some Lepidoptera. *Ecol. Entomol.*, 3, 155–160.

Webb, C.O. & Donoghue, M.J. (2005). Phylomatic: Tree assembly for applied phylogenetics.
Mol. Ecol. Notes, 5, 181–183.

Zanne, A.E., Tank, D.C., Cornwell, W.K., Eastman, J.M., Smith, S.A., FitzJohn, R.G., et al.
(2014). Corrigendum: Three keys to the radiation of angiosperms into freezing
environments. *Nature*, 514, 394–394.

Zeuss, D., Brunzel, S., & Brandl, R., (2017). Environmental drivers of voltinism and body
size in insect assemblages across Europe *Glob. Ecol. Biogeogr.*, 26, 154–165

Biosketch

Ryosuke Nakadai is a postdoctoral researcher with broad interests in biodiversity and
biological interactions across temporal and spatial scales. His works include development of
both statistical and molecular tools for a wide range of taxa.

Author contributions

RN conceived of and designed the study and analyzed the data; RN and TN constructed the

figures; KH, RN, and TI compiled the butterfly dataset; RN and AV wrote the first draft; and

all authors significantly contributed to the final text.

Appendices

Figure S1 Geographical patterns of climatic variables and elevation: (a) annual mean

temperature, (b) annual precipitation, (c) annual maximum depth of snow cover, and (d)

elevation.

Figure S2 Relationships of species-specific body size with (a) the number of host-plant

genera ($r = -0.062$, $p = 0.156$) and (b) phylogenetic host breadth ($r = -44.82$, $p = 0.131$)

across Japanese butterflies. Points indicate species. Significant correlations were not

confirmed by linear regressions.

Figure S3 Latitudinal patterns of (a) minimum phylogenetic host breadth ($r = -0.022$, $p <$

0.001), (b) maximum phylogenetic host breadth ($r = -0.072$, $p < 0.001$), (c) minimum body

size ($r = 0.020$, $p < 0.001$), and (d) maximum body size ($r = -0.167$, $p < 0.001$). Values

represent standardized effect sizes (SES) estimated using null model 1.

Figure S4 Latitudinal patterns of (a) the alpha parameter of the discrete truncated Pareto

distribution based on predicted occurrence data ($r = 0.018$, $p < 0.001$), and (b) the alpha

parameter of the discrete truncated Pareto distribution based on observed occurrence data ($r =$

0.005, $p < 0.001$). Values represent non-standardized trends comparable to those of Forister

et al. (2015).

Table S1 Species analyzed in the study. Nine species were excluded from the analyses for

various reasons. First, the taxonomic status of three species changed during the interval

between the fourth and fifth biodiversity censuses (Inomata, 1990; Shirôzu, 2006); we

excluded those species because the identifications were unreliable. Second, we excluded

three non-herbivorous species (including an omnivore, *Shirozua jonasi*) and three

gymnosperm-feeding species.

Table S2 Correlations between environmental variables and calculated indices.

Table S3 Comparison of the results between the full model and final model.

Table S4 Comparison of the results based on predicted and observed distribution datasets.

The model structures are described in Table 2.

Table S5 Comparison of the results based on null model 1 and null model 2. The model

structures are described in Table 2.

Table S6 Comparison of the results based on different criteria of minimum butterfly species

richness. The model structures are described in Table 2.

Table S7 Environmental variables and calculated indices that were used in this study for all

10 km grids.

(a) Butterfly species richness
1   124

(b) Body size mean (SES)
−2.78   4.93

(c) Body size divergence (SES)
−2.85   7.53

(d) Alpha parameter of
the truncated Pareto distribution (SES)
−3.00   4.84

(e) Phylogenetic host breadth mean (SES)
−1.89   7.74

(f) Phylogenetic host breadth divergence (SES)
−1.56   4.19