Extensive genome-wide association analyses identify genotype-by-environment interactions of growth traits in Simmental cattle

Camila U. Braz[a], Troy N. Rowan[a,b], Robert D. Schnabel[a,b,c], Jared E. Decker[a,b,c*]

[a]Division of Animal Sciences, University of Missouri, Columbia, MO 65211, USA

[b]Genetics Area Program, University of Missouri, Columbia, MO 65211, USA

[c]Informatics Institute, University of Missouri, Columbia, MO 65211, USA.

*Corresponding author: Jared E. Decker, email: DeckerJE@missouri.edu

CUB: camila_urbano@yahoo.com.br

TNR: tnr343@mail.missouri.edu

RDS: schnabelr@missouri.edu

JED: deckerje@missouri.edu

Abstract

Background: Understanding the genetic basis of genotype-by-environment interactions (GxE) is crucial to understand environmental adaptation in mammals and improve the sustainability of agricultural production. In addition, GxE information could also be useful to predict the vulnerability of populations to climate change.

Results: Here, we present an extensive study investigating the interaction of genome-wide SNP markers with a vast assortment of environmental variables and searching for SNPs controlling phenotypic variance (vQTL) using a large beef cattle dataset. We showed that GxE contribute 10%, 4%, and 3% of the phenotypic variance of birth weight, weaning weight, and yearling weight, respectively. GxE genome-wide association analysis (GWAA) detected a large number of GxE loci affecting growth traits, which the traditional GWAA did not detect, showing that functional loci may have non-additive genetic effects between genotype classes regardless of differences in genotypic means. We also showed that variance-heterogeneity GWAA can detect loci enriched with GxE effects without requiring prior knowledge of the interacting environmental factors. Functional annotation and pathway analysis of GxE genes revealed biological mechanisms by which cattle respond to changes in their environment, such as neural signaling, metabolic, hypoxia-induced, and immune system pathways. Knowledge of these pathways will be important as climate change becomes a burden on animal health and productivity. In addition, ecoregion-specific GxE SNPs detected in this study may play a crucial role in identifying resilient and adapted beef cattle across divergent environments.

Conclusions: We detected novel trait associations with large GxE effects for birth weight, weaning weight, and yearling weight. Functional annotation and pathway analysis uncovered genomic regions involved in response to environmental stimuli. We unraveled the relevance and complexity of the genetic basis of GxE underlying growth traits, providing new insights into how different environmental conditions interact with specific genes influencing adaptation and productivity in beef cattle and potentially across mammals

Keywords: adaptation, beef cattle, environment, ecoregion, variance heterogeneity

Background

Genotype-by-environment interactions (GxE) refer to a variable response of genotypes across environments, resulting in different trait values [1,2]. GxE is a central issue in genetics, evolution, and ecology [3], and cattle represent a unique opportunity to address these questions due to a large number of genotyped animals [4] reared over a wide range of climatic and topographic regions. GxE may contribute to poor adaptation, which negatively affects the profitability and sustainability of the beef cattle production systems [1,5]. Studies using the relationship between breeding values and the environment have reported the existence of GxE for body weight in beef cattle [6,7]; however, limited research has focused on understanding the genetic loci of GxE underlying such traits. This knowledge could be used to better drive selection and mating decisions [8] as well as in statistical models incorporating biological information to accomplish more accurate

genomic predictions [9]. In addition, GxE information could also be useful to predict the

vulnerability of populations to climate change [3] and to understand environmental

adaptation in other mammalian species.

Here, we present a comprehensive study of GxE for birth weight (BW), weaning

weight (WW), and yearling weight (YW) using SNP marker information and over 13,000

Simmental cattle. First, we estimated the contribution of GxE variance on the total

phenotypic variances using a restricted maximum likelihood (REML) approach. Then, we

searched for SNPs enriched for GxE using four linear mixed model approaches: i) direct

GxE genome-wide association analyses (GxE GWAA) using continuous environmental

variables; ii) direct GxE GWAA using discrete environmental variables (United States

ecoregions); iii) variance-heterogeneity genome-wide association analyses (vGWAA),

which enable identification of interactions, using phenotypes adjusted for additive

relationship matrix or additive, dominance and epistatic relationship matrices; and iv)

meta-analysis of ecoregion-specific GWAA. Lastly, we performed enrichment analyses

using GxE candidate genes to investigate the biological mechanisms by which

environmental factors modulate such phenotypes. To our knowledge, this is the most

comprehensive study investigating the interactions of genome-wide SNP markers with a

vast assortment of environmental variables and searching for SNPs controlling

phenotypic variance using a large beef cattle dataset. Our results revealed novel trait

associations and alternative biological mechanisms involved in shaping the total

phenotypic variance of growth traits providing new insights into how the environment

influences growth traits and adaptation in beef cattle and potentially across mammals.

Results

Environmental variance component estimates

Phenotypes were pre-adjusted for fixed effect of sex and random effect of contemporary

groups. The amount of BW, WW, and YW variation explained by contemporary group

effects was 19%, 53%, and 61%, respectively, showing that the effect of farm

management on the animal body weight increases with age. After adjusting the

phenotypes, we estimated the contribution of GxE variance on the total phenotypic

variances using additive relationship matrix calculated based on SNP marker information,

and United States ecoregions as environmental factor. Ecoregions were determined based

on the combination of 30-year normal values of mean temperature, precipitation, and

elevation [10]. Each 1 km square of the United States was then assigned to one of 9

resulting ecoregions. We found that GxE variances contribute in 10.1%, 3.8%, and 2.8%

to the BW, WW, and YW total phenotypic variances, respectively, suggesting that there

is a decrease in the effect of GxE on the animal body weight throughout life. These

results indicate an inverse relationship between farm management and environmental

stress and that management practices (e.g. season of birth, forage quality and quantity,

feed supplementation, health programs) are more relevant for later weights than climate

environmental factors. One possible explanation for these findings is that birth weight

reflects intrauterine growth which is highly affected by environmental conditions [11];

however, as the animal ages physiological, morphological, endocrine, cellular and

molecular mechanisms are altered in order to cope up with environmental challenges

[12]. These results also suggest that as management inputs are decreased, animals

become more sensitive to environmental stress, which has ramifications for animal

performance as we seek to decrease inputs to make beef production more sustainable.

Genotype-by-environment GWAA

The GxE GWAA for BW, WW, and YW were performed by fitting univariate and

multivariate linear mixed models considering continuous or discrete environmental

variables. The continuous environmental variables used were minimum, mean,

maximum, and mean dew point temperatures; elevation; precipitation; and minimum and

maximum vapor pressure deficit, separately (Additional file 1: Figures S1–S8). Discrete

environmental variables included United States ecoregions (Desert & Arid Prairie,

Southeast, High Plains, Forested Mountains, Fescue Belt, and Upper Midwest &

Northeast – Fig. 1F), which are combinations of mean temperature, precipitation, and

elevation 30-year normal values (Additional file 1: Figures S9–S14). Descriptive

information for growth traits and environmental variables are included in Additional file

2: Table S1. Univariate and multivariate models using both continuous and ecoregions

environmental variables detected a total of 2,319 GxE SNPs (P < 1e-5, Table 1,

Additional files 3 and 4). Only 25% of the GxE SNPs detected by the ecoregion GxE

GWAA were also identified by the continuous GxE GWAA. These results indicate that

the use of ecoregion information as environmental variable allowed us to capture GxE

SNPs interacting with region-specific stressors that were not directly measured (e.g.

forage differences, water availability, pathogens) and the combination of multiple

stressors.

There was no overlap between the GxE SNPs identified for all traits using univariate

models, however, multivariate models detected several GxE SNPs associated with BW,

WW, or YW (P < 1e-5, Table 1). Multivariate analyses were able to detect GxE SNPs

that were not identifiable by standard univariate analyses, indicating advantages of

incorporating a multivariate statistical approach to GxE genome-wide association studies

of correlated traits. Consistent with our GxE variance component results, a larger number

of GxE SNPs was detected for BW than for WW or YW. BW was highly affected by all

continuous environmental variables and ecoregions analyzed (Table 1). However, mean

dew point temperature was the environmental variable that most interacted with SNPs for

BW (P < 1e-5, Fig. 1B), and consequently, the ecoregion that identified more GxE SNPs

for BW was the Desert & Arid Prairie (P < 1e-5, Fig. 1G). The Desert & Arid Prairie

ecoregion has both high temperatures and many locations with low absolute humidity

measured by dew point temperature (Fig. 1A, Additional file 2: Table S1), which in

concert both affect BW. The effect of higher temperature on calf birth weight may be due

to the reduced feed intake or the decreased uterine blood flow and placental function

often found in heat-stressed cows, factors that contribute to impaired cow-to-fetal nutrient

exchange, affecting fetal growth [13–15]. For WW and YW, elevation (Fig. 1C) was

found to be the continuous environmental variable that most interacted with SNPs (P <

1e-5, Fig. 1 D and E respectively). In agreement, Forested Mountains, the ecoregion with

the highest elevation (Additional file 2: Table S1), showed more SNP-by-environment

interaction for WW and YW (P < 1e-5, Fig. 1 *H* and *I* respectively), compared with other

ecoregions. Higher altitude regions are associated with reduced atmospheric oxygen

(hypoxia), which causes pulmonary hypertension and vasoconstriction leading to right-

side heart failure in cattle [16]. This condition is known as brisket disease and accounts

for significant losses in growth and reproductive performance, being one of the top

causes of mortality in cattle residing at high altitudes [17].

Only one GxE SNP (*rs133638007*, chromosome 18, 61,721,281 bp) was detected

interacting with two different ecoregions. The *rs133638007* T allele increase BW in 1.06

kg in the Desert & Arid Prairie ecoregion, on the other hand, results from the multivariate

analysis showed that the *rs133638007* T allele decrease BW by 0.62 kg in Forested

Mountains ecoregion. Interestingly, this SNP is located close to the gene *VSTM1,* a

member of the leukocyte receptor complex, which is involved in immune responses [18].

In addition, the fact that only one GxE SNP was identified for two different ecoregions

indicates that the loci interacting with abiotic and biotic stressors are different across

divergent environments, and consequently, the ecoregions have specific genetic

architecture involved in GxE.

We explored 10 kb and 100 kb sequence windows that flanked all significant GxE

SNPs (P < 1e-5) to scan for genes in their vicinity and to identify possible regulatory

elements (Additional file 5). Using these GxE gene sets, we performed functional

enrichment analysis based on Gene Ontology (GO) and KEGG pathways to get insights

into biological mechanisms by which beef cattle respond to changes in their environment.

Response to stimulus was the most statistically significant GO term, and its direct

descendants GO terms, cellular response to stimulus and response to stress, were also

enriched (Additional file 2: Table S2). Additionally, many biological processes involved

in metabolism, development, localization, gene expression, transport, signaling, and cell

adhesion were enriched. Enriched pathways were related to metabolism; cell proliferation

and differentiation; immune and inflammatory responses; hypoxia-induced processes;

and mechanisms of neurotransmission in central nervous system (Additional file 2: Table

S3). In addition, circadian entrainment, a pathway involved in environmental adaptation

to natural periodic changes, including light/dark, temperature, humidity, social activity,

and several other cycles [19], was enriched. All GO terms and pathways enriched and

their respective GxE genes are available in Additional files 6 and 7, respectively.

As the genetic architecture of GxE appeared to be region-specific, we investigated

whether they affect the same or different biological functions across ecoregions. To do

so, we calculated the percentage of candidate GxE genes for each ecoregion that

participate in each biological processes (GO) or pathways enriched (Fig. 2 and 3,

Additional file 1: Figures S15 and S16). Even though the ecoregions exhibit mostly

unique GxE QTL, the candidate genes associated with these QTLs point toward the same

biological processes and pathways, however in different magnitudes. The GxE genes

involved in response to stimulus were identified in all ecoregions. In the Southeast,

55.8% of the GxE genes detected play roles in the response to stimulus process,

suggesting that the combined effects of high temperature and humidity pose a major

challenge by affecting how the animal adjusts to external changes and maintains

homeostasis. This will become a greater burden on animal health and productivity as the

climate continues to warm. Indeed, studies have shown that heat and humidity stress

disrupt homeostasis and result in compromised health, growth, fertility, and milk

production in several species, including cattle, pig, sheep, mice and humans [12,20–23].

Cardiomyopathy related pathways were enriched predominantly in GxE genes

interacting with ecoregions in the Western United States, as High Plains, Forested

Mountains and Desert & Arid Prairie, ecoregions with higher elevation. Such pathways

were identified likely due to the hypoxia condition exposure, which cause pulmonary

arterial hyperplasia resulting in right ventricular hypertrophy and congestive/dilatory

cardiac failure in cattle living at high altitude [17]. Pathways involved in neural

development and neurotransmission processes were also enriched in GxE genes.

Although these pathways were identified as being affected by GxE genes interacting with

all ecoregions, they seem to be more affected in animals raised in the Fescue Belt and

Southeast ecoregions. This may be due to the consumption of tall fescue, a grass

commonly used in grazing cattle systems in the Southeastern United States [24], which

usually contains an endophytic fungus that produces ergot alkaloids [25]. Ergot alkaloids

have vasoconstrictive effects and interact with GABAergic, glutamatergic and

serotonergic receptors affecting multiple aspects of animal's physiology regulated by

these neurotransmitters [26–28]. Animals consuming the endophyte-infected tall fescue

forage often develop fescue toxicosis, which is characterized by reduced feed intake,

weight gain, and blood flow, as well as elevated body temperature associated with heat

stress [28–31]. Interestingly, the majority of GxE genes for the Fescue Belt and Southeast

ecoregions were also found interacting with temperature. Numerous GxE genes involved

in neural signaling were also detected in the Forested Mountains ecoregion. Hypoxic

environments may affect cells of the nervous system as well. [32] reported that

neuroactive ligand-receptor interaction participates in the regulation of neural stem cells

behavior in mice under hypoxia. Pathways involved in hormone-releasing (oxytocin and

GnRH signaling) related to acute stress in cattle [33] were also identified for high

elevation ecoregions.

GxE genes of all ecoregions drove enrichments in biological pathways related to

immune and inflammatory responses (Cytokine-cytokine receptor interaction, Fc epsilon

RI signaling, Epstein-Barr virus infection, Tuberculosis, and Hepatitis), except Fescue

Belt. This result is unexpected since studies have observed that fescue toxicosis causes

hypoprolactemia leading to decreases in immune responses [34,35]. However, our results

indicate that this effect may be due to the environmental condition of Fescue Belt

ecoregion affecting the activity of GABA and serotonin. Such neurotransmitters are

involved in the regulation of prolactin secretion [36,37] and, consequently, may affect the

immune system. Many of these immune GxE genes have been associated with diseases

caused by pathogens leading to reduced productivity in cattle [38–48]. This result is in

agreement that heterogeneity in the immune response to infectious diseases across

populations living in different environmental conditions is under genetic control [49]. In

addition, pathogen resistance increases host fitness and has driven the evolution of plant

and animals through time [50].

GxE genes also participate in pathways involved in the regulation of cell functions

including proliferation, growth, gene expression, differentiation, cell survival, and

apoptosis, which are crucial processes to proper animal growth and development [51].

Such pathways (PI3K-Akt signaling, Ras signaling, MAPK signaling, Hippo signaling,

Focal adhesion, and ErbB signaling) were identified by GxE genes from all ecoregions.

However, we identified more GxE genes involved in such pathways in the Southeast,

Desert & Arid Prairie, Forested Mountains and High Plains ecoregions.

The majority of GxE SNP associations were identified for BW in this study. Studies

have shown that loci identified in GWAA of BW could either represent direct effects of

the fetal genotype, indirect effects of maternal genotype (acting via the intrauterine

environment), or some combination of the two [52,53]. According to [54] the uterine

environment may even have a greater influence on fetal growth and birth weight than the

parental genome. Thus, we investigated whether GxE SNPs affecting BW could be

reflecting maternal uterine environmental conditions. To test this, we adjusted BW by

including random maternal effects in the model (MAM) and compared the results with

those from the GxE GWAA without maternal effects (MA). Results showed that 876

SNPs (out of 1923) were not associated with GxE effects when accounting for the effect

of dam, suggesting that those SNPs are interacting with the uterine environment.

Candidate maternally-influenced GxE genes (Additional file 8) participate, in particular,

in metabolic, insulin signaling, starch and sucrose metabolism, PI3K-Akt signaling and

cGMP-PKG signaling pathways; and response to stimulus, blastocyst development, and

regulation of signaling processes. These findings support previous investigations

suggesting that detrimental environmental factors experienced by the cow during

gestation may influence cow-to-fetal nutrient exchange limiting fetal growth and

development [13–15,54–56]. Several maternally-influenced GxE genes identified in this

study have been previously reported as maternal genes influencing embryonic

development and/or response to environmental stress in cows and also in other mammals

[57–66]. These results demonstrate that the bovine is an important model to understand

the interaction between the physical environment, uterus, and fetus, and could be used in

future research.

Variance-heterogeneity GWAA

The vGWAA identify differences in variance, rather than differences in mean, between

genotype classes, called vQTLs, which can be explained by GxE or GxG (epistatic)

interactions [67]. Therefore, we performed a REML analysis to estimate how much of the

BW, WW, and YW variance is explained by additive, dominance and epistasis effects.

We fitted models (M) using only additive relationship matrix (MA) or using additive,

dominance, and epistasis relationship matrices (MADE), see Additional file 2: Table S4.

Overall, the heritabilities were greater than the ratio of the variance explained by the non-

additive genetic effects to the phenotypic variances. Proportions of dominance variances

in WW and YW were low (2% and 5%, respectively), whereas no dominance variance

was estimated for BW. Epistatic relationship matrix accounted for 25%, 12% and 11% of

the phenotypic variances for BW, WW, and YW, respectively, suggesting that epistasis

has a substantial contribution to phenotypic variation for body weight traits in Simmental

cattle. Thus, to search for vQTLs we used squared normalized residuals (z^2), which is a

measure of phenotypic variation [68], as dependent variables fitting univariate and

multivariate models. Using z^2 from the MA, we expected to detect loci with both GxE

and GxG effects, since this phenotype was adjusted only for the additive relationship

matrix. However, using z^2 from the MADE, we should detect exclusively GxE loci as the

phenotype was adjusted for additive, dominance and epistatic relationship matrices. In

total, 44 SNPs with effect on z^2 (P < 1e-5) were identified on chromosomes 3, 5, 8, 10,

11, 13, 14, 16, 17, 18, 21, 27, and 29 (Additional file 1: Figures S17 and S18, Additional

file 2: Table S5). These 44 SNPs tag 24 putative vQTLs (grouped by LD patterns), with

seven, four, and eleven vQTLs for univariate BW, WW, and YW, respectively. The

multivariate analysis identified five vQTLs, of which two vQTLs (chromosome 8 at 4

Mb; and chromosome 29 at 39 Mb) were not identified by the univariate analyses.

Nine vQTLs were only significant using z^2 from MA (three vQTLs for univariate

BW, three vQTLs for univariate WW, two vQTLs for univariate YW, and one vQTL

using multivariate analysis). Among these vQTLs likely involved in GxG, one vQTL for

BW is located on chromosome 14 at 23 Mb and resides near or within the genes *LYN,*

RPS20, MOS, PLAG1, and *CHCHD7*. This region has been reported to have epistatic

effects on several chromosomes affecting post-weaning weight in beef cattle [69]. These

effects could be due to the gene *PLAG1* which encodes a transcription factor that

regulates the expression of several genes involved in a variety of cellular processes [70],

and has been previously associated with stature and growth traits in many beef cattle

breeds [71–73]. In addition, we detected a GxG vQTL on chromosome 3 at 106 Mb

within *TRIT1*, a gene that was also reported to harbor a vQTL involved in epistatic

interactions in humans [74].

Using z^2 from MADE, 15 vQTLs were detected (four vQTLs for univariate BW; one

vQTL for univariate WW; nine vQTLs for univariate YW; and four vQTLs using

multivariate analyses, of which three vQTLs were also identified by the univariate

analyses), which likely identify GxE. Based on GO information, genes near or within

these vQTLs are involved in response to stimulus (*NOS1, FBLN1, YWHAH,* and

DEPDC5), inflammation and immunity mechanisms (*LYZL1, KSR2, FBXO21, IL18RAP,*

XCL2, XCL1, TTYH1, LENG8, LENG9, CDC42EP5, LAIR1, OSCAR, and *TARM1*),

neurogenesis (*ZHX2,* and *NOS1*), regulation of transcription (*ZHX2, YWHAH* and

CNOT3), developmental process (*FBXW8,* and *CNOT3*), transmembrane transport

(SLC9A2, *SLC9A4, SLC5A1, SLC5A4,* and *TTYH1*), neurotransmission regulation

(*SHANK2*), and cell proliferation (*DPT, RPS9,* and *TACC1*). Some of those genes were

previously reported to be involved in environmental adaptation in cattle [*XCL2* [75],

SLC9A4 [76], *ZMAT4* [77]], and heat tolerance in buffalo [*XCL1* [78]] and dairy cattle

[*SHANK2* [79]].

Using z^2 as dependent variables in the analyses allowed the estimation of the

proportion of residual variance that is explained by the SNPs in GEMMA (PVE). For

BW z^2, WW z^2 and YW z^2, SNPs explained 4.9% (SD = 0.8), 4.7% (SD = 0.8), and 5.9%

(SD = 1.1), respectively, showing that residual variance is heritable and could be changed

by selection, supporting previous studies [80–82]. Residual variability may reflect

differences between animals in uniformity for a certain trait, so the reduction of variance

could improve economic merit [80]. Rönnegard et al. [81] reported advantages in using

breeding values explaining differences in residual variance (vEBV) for milk yield

uniformity in Holstein cattle. According to [80] the reduction of phenotypic variance by

selecting for reduced residual variance could be more beneficial for low heritability traits.

Meta-analysis of ecoregion-specific GWAA

We performed a meta-analysis of ecoregion-specific GWAA (GWAA for each ecoregion

separately) to look for significant differences (Cochran's Q statistics) in effect size of

SNPs between the ecoregions, which are interpreted as GxE [83]. In addition, we

calculated the m-value statistic for the significant SNPs (Cochran's Q statistic's *P*-value <

1e-5), which is the posterior probability of an effect being present in an ecoregion given

the observations from all other ecoregions [83] allowing us to identify which ecoregions

a given variant has an effect. Southeast and Desert & Arid Prairie ecoregions were not included in the meta-analysis due to their small sample sizes. Meta-analysis of ecoregion-specific GWAA detected 12 GxE SNPs for BW located on chromosome 6 (Additional file 1: Figure S19A); two GxE SNPs for WW on chromosome 15 (Additional file 1: Figure S19B); and two GxE SNPs for YW, one located on chromosome 13 and one on chromosome 15 (Additional file 1: Figure S19C, Additional file 9). Analyzing the PM-plot (P-values against m-values) and PB-plot (P-values against beta-values) of the GxE SNPs detected by the meta-analysis, we observed that some GxE SNPs on chromosome 6 at 30 Mb have an effect on animal's BW from Forested Mountain and Upper Midwest & Northeast ecoregions, however with different certainty (Additional file 1: Figures S20A and S21A). For WW and YW, despite that the analysis had shown a significant heterogeneity between the ecoregions, they did not show different posterior probabilities of an effect in any ecoregion (m-value), (Additional file 1: Figure S20B and C, respectively). However, large differences in the SNP effect size (beta) were noticed between ecoregions (Additional file 1: Figure S21B and C, respectively).

Traditional genome-wide association analyses

We also performed traditional GWAA using univariate and multivariate linear mixed models in order to compare with the GxE GWAA results. We detected major QTLs on chromosomes 6 (26-48 Mb), 7 (88-90 Mb), 14 (15-30 Mb), and 20 (41-78 Mb) for univariate BW, WW and YW (Additional file 1: Figure S22A-C and Additional file 10). In addition, significant SNPs (P < 1e-5) were identified on chromosomes 3 (53.6 Mb) and 17 (73 Mb) for univariate BW; and on chromosomes 1 (81.1-81.3 Mb), 2 (116.1 Mb),

and 5 (96.7 Mb) for univariate WW. The GWAA using multivariate model detected QTL

also identified in the univariate GWAA on chromosomes 5, 6, 7, 14, 17, and 20 in

addition to on chromosomes 4 (24.5 Mb), 9 (57.9 Mb), 13 (63.3 Mb), 19 (50.5 Mb), and

26 (45.4 Mb) (Additional file 1: Figure S22D and Additional file 11). All genes detected

by GWAA are included in the Additional file 12.

Out of 948 total significant SNPs using multivariate GWAA analysis for the growth

traits, 531 SNPs were located on chromosome 6 (from 26.8 to 48.8 Mb). Likewise,

several SNPs interacting with Forested Mountains ecoregion, elevation, precipitation and,

mainly, with the temperatures were also detected in the same positions on chromosome 6.

Given the importance of chromosome 6 underlying the growth traits, we repeated the

multivariate GWAA including significant SNPs (e.g. *rs109849093, rs464458177,*

rs110305942, rs43459303, rs109278547, and *rs137209027* located at 37,211,057;

37,234,136; 37,418,164; 37,695,352; 38,139,495; and 41,084,765 bp, respectively) as

fixed effects in the model. In this way, if those SNPs are causal mutations accounting for

QTL effect the loci association signals would be missed. Results showed that even

including six significant SNPs (not in LD) simultaneously as fixed effects in the model, a

small number of SNPs were still associated (six out of 531 SNPs), indicating that there

are possibly multiple causal mutations in this region.

Comparison of the GWA approaches

Comparing traditional GWAA with all GxE GWAA results, we verified that 312 SNPs

with additive effects (QTL) also have GxE effects (mainly on chromosome 6, but also on

chromosomes 7 and 20). Some SNPs showed GxE effects even greater than their additive

effects on the traits. For example, the SNP *rs136523580* (chromosome 6 at 30,358,905)

has an additive effect of -0.43 kg for BW in the total dataset, but its GxE effects is even

more impactful (-0.67 kg) on the BW of animals from the Forested Mountains ecoregion.

The SNP *rs109868461* (chromosome 6, at 36,902,704 bp) has additive effects of -0.69 kg

for BW in the total dataset, however it has a GxE effect of 1.11 kg on the BW of animals

born in the Southeast ecoregion. These findings show the importance of GxE effects on

quantitative traits and the need for precision, ecoregion-specific genomic predictions to

optimize animal performance.

Comparing vGWAA with the traditional GWAA results, only one vQTL, on

chromosome 14 at 23.1-23.3 Mb harboring the genes *LYN*, *MOS*, *RPS20*, *PLAG1,* and

CHCHD7, was also found as a direct effect QTL for BW, meaning that this locus affects

both mean and variance of BW in this population. Five vQTLs were also identified by the

GxE GWAA, which were found to interact with multiple continuous environmental

variables as well as interacting with ecoregions. These results agree with [84] in

suggesting that vQTL effects are much more enriched with GxE effects than additive

effects. However, GxE GWAA detected a greater number of GxE loci when compared

with the vGWAA results.

Only few GxE SNPs were detected using the meta-analysis of ecoregion-specific

GWAA (Cochran's Q statistic's *P*-value < 1e-5), which were also identified by the GxE

GWAA. In addition, using the meta-analysis approach the Southeast and Desert & Arid

Prairie ecoregions could not be included in the analysis due to their small sample sizes,

suggesting that this was not an optimal approach to detect GxE associations for our

dataset. However, creating PM-plots (contrast between environment-specific *P*-value and

posterior probability of an effect, m-value)[85], and PB-plots (environment-specific *P*-value and SNP effect size) helped interpret results of our GxE GWAA and vGWAA. For significant GxE SNPs detected by the GxE GWAA, there were clear differences in the SNP effect sizes between the significant ecoregion and the remaining ecoregions (Additional file 1: Figures S23 and S24). For SNPs with significant variance-heterogeneity effects (detected by vGWAA), patterns were not as consistent. While some vQTLs had differences in mean effects, others did not (Additional file 1: Figures S25 and S26).

Discussion

This GxE study provides a more comprehensive understanding of how GxE modulate growth traits in beef cattle and possibly in other mammalian species. We observed that GxE explain substantial proportions of growth traits phenotypic variances, which is likely due to a great number of GxE SNPs with large effects affecting such traits. Detection of large GxE SNP effects indicates that there is considerable opportunity to improve environmental resistance and performance in cattle. Our study also revealed the first evidence of vQTLs controlling phenotypic variation in body weights due to GxG or GxE effects in beef cattle. In addition, we uncovered several biological pathways affected by GxE that likely drive environmental adaptation.

One of our major findings is that the ecoregions showed specific GxE with large effects on the traits. In addition, ecoregion GxE GWAA detected GxE loci that were not identified by the continuous GxE GWAA. One possible explanation for these results is

that when analyzing ecoregions, we take into account not only climate and topography,

but also unmeasured variables like forage quality and local pathogens, as well as the

combination of them. This information allowed us to identify environmentally sensitive

genotypes. It indicates that the same trait can be influenced by different genes in the

various ecoregions which may affect particular physiological and behavioral functions

leading to local adaptation. This assumption is supported by the GO and pathways

enrichment analysis of the GxE genes identified for each ecoregion showing that the

diverse ecoregion environment affects different biological mechanisms and in different

magnitude. We also found that major additive QTLs also carry GxE effects on growth

traits and that some of their GxE effects are greater or in opposite directions when

compared with their additive effects. Additionally, novel GxE associations were detected

in multiple genomic regions. These findings support the importance of creating

ecoregion-specific genomic predictions to identify local resilient or adapted animals,

using the GxE SNPs detected in this study to assist selection decisions which would lead

to an increase in animal performance within an ecoregion.

Some vQTLs were also identified by the GxE GWAA, indicating that vGWAA can

detect loci enriched with GxE effects without requiring prior knowledge of the interacting

environmental factors. Such GxE loci showed interaction with several continuous and

discrete (ecoregions) environmental variables in the GxE GWAA, suggesting that

vGWAA may capture GxE loci that interact with multiple environmental variables,

possibly even those not included in the GxE GWAA model (e.g. water availability,

pasture condition, pathogens). According to [86] when multiple interacting factors induce

variance heterogeneity, the power of identification of any single one of them or all

together may be lower than the power of the variance heterogeneity test, which may

explain why vGWAA identified GxE loci that were not detected by direct GxE GWAA.

We also observed that some vQTLs have differences in means across ecoregions

(Additional file 1: Figure S27), however, not all do. This suggests that some vQTLs

could be the difference between robust genotypes (little variation, flat reaction norms)

across environments while other genotypes at the same locus are more variable across

environments, indicating that selection for low variance vQTL genotypes may provide

sustainability advantages. Many genes harboring the vQTL identified here were

previously reported to be involved in environmental adaptation in cattle [75–79]

supporting that phenotypic variability may be an adaptive evolutionary solution to

environmental changes [87]. However, our results support [88] which reported that when

an environmental factor of interest has been measured on all the genotyped individuals,

the direct GxE GWAA will be able to detect a larger number of GxE loci.

We found that environmental conditions modulate growth traits through genes that

participate in a variety of biological mechanisms that are activated to help the body cope

with external stimuli and return to or maintain homeostasis [89,90]. Among them, several

pathways involved in neural signaling and development were detected, which play crucial

roles in the regulation of response to stress [91,92], body temperature, vasoconstriction

[93,94], and circadian rhythms [95]. Specifically, for animals living in higher altitude

ecoregions, we identified pathways related to hypoxia response and secretion of

hormones [33]. Other environmental pathways included immune functions reflecting the

involvement of innate and adaptive immune systems in the activation of host defense

reactions to deal with pathogens [94]. We also identified many metabolic processes that

may be involved in acute stress responses and thermoregulation [12,33,94]. These

biological mechanisms are affected differently by the diverse ecoregion environments

and have been reported to be involved in resilience [12,92,94], and adaptation processes

[10,16,96–98], revealing the role that GxE plays in adaptation and evolution at the level

of the genome in cattle and potentially across mammals. It is important to understand

such biological mechanisms since the inability of an animal to cope with its environment

results in failure to produce optimally leading to lost potential profitability of livestock

operations [12,17,33,99]. Further, these biological insights may help us understand how

adaptation occurs in wild populations and the potential effect of climate change on

animal physiological functions.

Conclusions

In summary, GxE GWAA and vGWAA detected several loci affecting growth traits

that the traditional GWAA did not, showing that functional loci may have non-additive

genetic effects between genotype classes regardless of differences in genotypic means.

We revealed biological mechanisms by which beef cattle respond to changes in their

environment. Neural signaling, metabolic, hypoxia-induced, and immune system

pathways were highlighted, indicating that climate change may become a burden on

animal health and productivity. In addition, ecoregion-specific GxE SNPs detected in this

study may play a crucial role in resilience and adaptation of beef cattle to divergent

environments. Although conventional genomic selection has shown great promise in

improving genetic gain, it only considers additive effects and, according to our findings,

ecoregion-specific genomic predictions should be created to identify animals best suited

for given environments. This could help producers in making optimal decisions given

their geographical location. Our results revealed novel trait associations and alternative

genetic mechanisms involved in shaping the total phenotypic variance of growth traits

providing new insights into how the environment influences such traits and adaptation in

beef cattle and other mammalian species.

503 Methods

Phenotype, genotype and environmental data

Data from the American Simmental Association males and females born between 1975

and 2016 and genotyped for genomic-enhanced EPDs were transferred to the University

of Missouri. The traits studied were birth weight (BW), 205-d adjusted weaning weight

(WW) and 365-d adjusted yearling weight (YW), with the traits also adjusted for age of

dam as per breed association practice. The animals were assigned to contemporary

groups (CG), defined by the combination of farm, season (spring vs. fall) and year of

birth.

Phenotypes were pre-adjusted for fixed effect of sex estimated using '*--reml-est-fix*'

option, and random effect of contemporary groups (CG) was predicted by the BLUP

method using '*--reml-pred-rand*' flag while controlling for population stratification using

single-trait animal model implemented in GCTA software [100]. CG were included in the

univariate model as random effect since 19% of individuals analyzed belong to CG with

only a single animal [101]. The random maternal permanent environmental effect was

also included in the WW adjustment.

The DNA samples were genotyped with various low-density assays and were imputed

to the combination of Illumina BovineHD (Illumina, San Diego, CA) and the GeneSeek

Genomic Profiler F250 (GeneSeek, Lincoln, NE) according to [102]. Genotype filtering

was performed to remove non-autosomal variants, as well as both variants and

individuals with call rates less than 0.90, using PLINK 1.9 [103]. SNP positions were

based on the ARS-UCD1.2 Bovine reference genome assembly [104]. The genotypes

were then phased using Eagle 2.4 [105] and imputed using Minimac3 [106]. The final

imputed genotype data contained a total of 835,947 bi-allelic variants with imputation

accuracy of 99.6% [102]. Variants with minor allele frequency less than 0.01 were

removed using PLINK 1.9 for further analysis, leaving 710,202, 709,054, and 706,771

SNP markers for 13,427, 11,847, and 8,546 animals for BW, WW and YW, respectively.

Eight continuous environmental variables were used for the GxE GWA analyses,

including 30-year normal of minimum, mean, maximum and mean dew point

temperatures; elevation; precipitation; and minimum and maximum vapor pressure

deficit, which were drawn from the PRISM climate dataset [107]. The United States was

partitioned into nine regions (Fig. 1F *Top*) based on similar topographic and

environmental conditions (ecoregions), including mean temperature, elevation and

precipitation information, using k-means clustering implemented in '*RStoolbox*' R

package [108,109]. The optimal ecoregions were identified using '*pamk*' function from

the R package '*fpc*' [110]. The ecoregions were named Desert (D), High Plains (HP),

Arid Prairie (AP), Forested Mountains (FM), Upper Midwest & Northeast (UN),

Southeast (SE), Rainforest (R), Foothills (FH), and Fescue Belt (FB) [10]. Animals were

assigned to the ecoregions based on the zip code of their breeder (Fig. 1F *Bottom*), and

only ecoregions with more than 200 records were analyzed. Thus, animals located in the

Rainforest ecoregion were removed from the dataset, and those that were reared in the

Desert and Arid Prairie ecoregions were analyzed together (called DA) due to their

similar environmental conditions. No animal record was assigned to the Foothills

ecoregion. A complete description of the dataset, including weight records and

environmental variable measurements is shown in Additional file 2: Table S2.

Estimation of variance components

Variance components for pre-adjusted BW, WW, and YW were estimated using multi-

component restricted maximum likelihood (REML) approach, implemented in GCTA

software, in univariate linear mixed models that included only additive effects (MA), or

additive, dominance and epistasis effects (MADE) as follows:

$$y^* = 1\mu + Z_a a + e \text{ (MA)}$$

$$y^* = 1\mu + Z_a a + Z_d d + Z_{ep} ep + e \text{ (MADE)}$$

where y^* is the vector of pre-adjusted phenotypes; μ is the overall mean; a is the vector

of random additive genetic effects; is d the vector of random dominance effects; ep is the

vector of random epistatic effects; and is e the vector of random residuals. $\boldsymbol{Z}$ matrices are

incidence matrices relating observations to animals. Assuming that $a \sim N(0, G\sigma_a^2)$,

$d \sim N(0, D\sigma_d^2)$, $ep \sim N(0, E\sigma_{ep}^2)$, $e \sim N(0, I\sigma_e^2)$, where $\sigma_a^2, \sigma_d^2, \sigma_{ep}^2$, and σ_e^2 are the additive,

dominance, epistatic, and residual variances, respectively. $\boldsymbol{I}$ is the identity matrix; and $\boldsymbol{G}$,

D, and E are the additive, dominance and epistatic genetic relationship matrices,

respectively. The genomic (G) and the dominance (D) relationship matrices were

calculated using SNP marker information according to [111] as: $G = W_G W_G'/N$, and

$D = W_D W_D'/N$, where $w_{G(i)} = (x_{G(i)} - 2p_i)/\sqrt{2p_i(1 - p_i)}$ and ,

$w_{D(i)} = (x'_{D(i)} - 2p_i^2)/\lfloor 2p_i(1 - p_i)\rfloor$ with x_G being 0, 1, or 2 and x'_D being 0, $2p$, or

$(4p - 2)$ for genotypes AA, AB or BB, respectively; p_i is the allele frequency at SNP i;

and N is the number of SNPs. According to Henderson [112] the epistatic relationship

matrix (E) can be derived from the additive genomic relationship matrix as: $E = G \# G$,

where # denotes the Hadamard product operation.

The variance of GxE interaction was also estimated considering the ecoregions as an

environmental factor and treated as random effect in the model in order to estimate the

amount of phenotypes that are explained by the GxE interaction among the ecoregions, as

implemented in GCTA.

Genome-wide association analyses (GWAA)

Univariate linear mixed model analyses were performed for the pre-adjusted BW, WW or

YW using GEMMA software [113] as follows:

$$y^* = 1\mu + X\beta + u + e$$

where y^* is an n-vector of pre-adjusted phenotypes; μ is the overall mean; X is the

incidence matrix of genotypes; β is the genotype effects; u is an n-vector of random

additive genetic effects; and e is an n-vector of random residual. Assuming that $u \sim$

$MVN_n(0, GV_g)$ and $e \sim MVN_n(0, IV_e)$, G is the $n \times n$ genomic relationship matrix

(GRM); V_g is the residual additive genetic variance; V_e is the residual variance

component; I is an n x n identity matrix. MVN_n denotes the n-dimensional multivariate

normal distribution. GRM were estimated using the standardized genotypes of 710,202,

709,054, and 706,771 SNP markers for 13,427, 11,847, and 8,546 animals for BW, WW

and YW, respectively.

We also fitted a multivariate linear mixed model for the pre-adjusted BW, WW and

YW using GEMMA [114] in the following form:

$$Y^* = 1\mu + X\beta + U + E$$

where $Y\textbf{*}$ is an n x d matrix of d pre-adjusted phenotypes for n individuals; μ is the

overall mean; X is the incidence matrix of genotypes; β is a d vector of the genotype

effects for the d phenotypes; U is an n x d matrix of random additive genetic effects; and

E is an n x d matrix of random residuals. $U \sim MN_{nxd}\left(0, G, V_g\right)$ and

$E \sim MN_{nxd}\left(0, I_{nxn}, V_e\right)$, G is the n x n genomic relationship matrix (GRM); V_g is the d x

d matrix of additive genetic variance component; V_e is a d x d matrix of the residual

variance component; I is an n x n identity matrix. MN_{nxd} denotes the n x d matrix normal

distribution with mean 0, row covariance matrix G or I (n x n), and column variance

matrix V_g or V_e (d x d). For all GWA analyses performed, single-marker P-values were

used to generate Manhattan plots using '*manhattan*' function implemented in '*qqman*' R

package [115].

Genotype-by-environment GWA analyses

The GxE GWAA were performed using GEMMA through two approaches: using eight

continuous environmental variables (i.e. minimum, mean, maximum and mean dew point

temperatures; elevation; precipitation; and minimum and maximum vapor pressure

deficit) separately; and using the ecoregions as discrete environmental variable, where

each ecoregion was compared against the total dataset using 0 and 1 dummy coding. In

the GxE GWAA, for each SNP in turn, GEMMA fits a linear mixed model that controls

both the SNP main effect and environmental main effect, while tested for the interaction

effect and controlling for population stratification, evaluating the alternative hypothesis

(H$_1$: $\neq 0$) against the null hypothesis (H$_0$:$x_j \beta_{ij} = 0$) for each interaction, therefore the

resulting *P*-values correspond to the significance of the GxE interaction [71]. This model

can be described as:

$$y^* = 1\mu + x_i\beta_i + x_j\beta_j + x_i * x_j\beta_{ij} + u + e$$

where y^* is an *n*-vector of pre-adjusted phenotype; x_i is an *n*-vector of the genotypes of

the SNP i; β_i is the effect size of the SNP i; x_j is an *n*-vector of environmental variable j;

β_j is the fixed effect of the environmental variable j; β_{ij} is the interaction effects between

the genotypes of the SNP i and the environmental variable j; u is an *n*-vector of random

additive genetic effects; and e is an *n*-vector of random residual. GxE GWA analyses

using multivariate linear mixed model were also performed.

Variance-heterogeneity GWAA

We performed vGWAA to detect locus affecting the difference in variance between

genotypes (vQTL). Residual effects (e) from the MA and MADE models were

standardized to z score by the rank-based inverse-normal transformation and squared (z^2),

which is a measure of phenotypic variance [84,116]. The vGWAA were performed for

BW z^2, WW z^2, and YW z^2 using univariate and multivariate linear mixed models

implemented in GEMMA software as denoted in "Genome-wide association analyses"

section, where y^* is an n-vector of z^2; and β_i is the effect size of the SNP i on z^2. Linkage

disequilibrium (LD) analyses were performed using PLINK 1.9. All vQTL were defined

by SNP markers with r^2 greater than 0.6 [117].

Meta-analysis of ecoregion-specific GWAA

In order to evaluate how significantly the effect size of SNPs (Cochran's Q test) are

between the ecoregions (indicative of GxE) [83] we combined the outputs from

univariate within-ecoregion GWAA into a single meta-analysis for each phenotype using

METASOFT software [118]. Only ecoregions with more than 1000 animals were

analyzed. In addition, the statistic m-value for the significant SNPs (Cochran's Q

statistic's P-value < 1e-5) were calculated, which is an estimative of the posterior

probability of a locus having an effect in a particular ecoregion [119]. The results were

visualized through PM-plots and PB-plots, in which P-values are simultaneously

visualized with the m-values and beta-values at each tested locus, respectively.

False discovery rate (FDR) estimation

FDR of the nominal significant threshold (P < 1e-5) was estimated for all analyses

performed (Additional file 2: Table S6), based on [120], as:

$$FDR = P\left(1 - \frac{A}{T}\right)\Big/\left(\frac{A}{T}\right)(1 - P),$$

where P is the nominal significant threshold (e.g., 1e-5); A is the number of SNP that

were significant at the nominal significant threshold; and T is the total number of SNPs

tested. Although the true FDR cannot exceed 1, the estimated FDR can exceed 1 if the

number of significant SNPs is smaller than expected by chance, meaning that all SNPs

that met the nominal significant threshold are false positives [69].

Functional enrichment analyses

We explored 10 kb and 100 kb sequence windows that flanked the significant SNPs ($P <$

1e-5) to scan for GxE genes located in their vicinity and to identify possible regulatory

elements, respectively, based on the ARS-UCD1.2 bovine reference genomic positions.

Enrichment analysis using the GxE genes based on Gene Ontology (GO) and KEGG

pathways were performed using ShinyGO v0.60 [121], in which the analyses were

adjusted for FDR of 5%. GO annotations were also retrieved from the AmiGO browser

[122].

Supplementary information

Additional file 1: Supplementary Figures. Supplementary Figures S1-S26. (DOCX 5.0

MB)

Additional file 2: Supplementary Tables. Supplementary Tables S1-S6. (DOCX 272.9

KB)

Additional file 3: Significant SNPs (P < 1e-5) using univariate GxE GWAA for Birth

Weight, Weaning Weight and Yearling Weight. (XLSX 248.7 KB)

Additional file 4: Significant SNPs (P < 1e-5) using multivariate GxE GWAA for Birth

Weight, Weaning Weight and Yearling Weight. (XLSX 40.7 KB)

Additional file 5: Candidate genes identified by GxE GWAA using 10 kb and 100 kb

region flanking the significant SNPs. (XLSX 253.8 KB)

Additional file 6: Enrichment analyses based on Gene Ontology terms using GxE genes

for 10 kb and 100 kb region flanking the significant SNPs. (XLSX 1.2 MB)

Additional file 7: Enrichment analyses based on KEGG pathways using GxE genes for

10 kb and 100 kb region flanking the significant SNPs. (XLSX 86.1 KB)

Additional file 8: Candidate maternally-influenced GxE genes using 10 kb and 100 kb

region flanking the significant SNPs. (XLSX 77.3 KB)

Additional file 9: Significant SNPs (Cochran's Q statistic's P-value < 1e-5) using meta-

analysis of ecoregion-specific GWAA for Birth Weight, Weaning Weight and Yearling

Weight. (XLSX 5.4 KB)

Additional file 10: Significant SNPs (P < 1e-5) using univariate GWAA for Birth

Weight, Weaning Weight and Yearling Weight. (XLSX 128.8 KB)

Additional file 11: Significant SNPs (P < 1e-5) using multivariate GWAA for Birth

Weight, Weaning Weight and Yearling Weight. (XLSX 93.6 KB)

Additional file 12: Candidate genes identified by GWAA for Birth Weight, Weaning

Weight and Yearling Weight. (XLSX 19.6 KB)

Declarations

Ethics approval and consent to participate

Animal Care and Use Committee approval was not required or obtained for data that were extracted from the existing American Simmental Association database.

Consent for publication

Not applicable.

Availability of data and materials

The raw data used in this research are available from the American Simmental Association under a Data Use Agreement, but are not publicly available. Derived data (analytical results) are however available as supplementary files associated with this publication.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was supported by Agriculture and Food Research Initiative Competitive, Grant No. 2016-68004-24827, from the USDA National Institute of Food and Agriculture.

Author's contributions

CUB and JED conceived the study. CUB analyzed the data and interpreted the results and wrote the manuscript. TNR and RDS managed data and imputed variants. JED assisted with analyses, preparation of the manuscript and interpretation of results. All authors read and approved the final manuscript.

Acknowledgments

The authors would like to acknowledge the American Simmental Association for providing phenotypic and genotypic data. We also appreciate the farmers and ranchers who collected this data.

References

1. Kolmodin R, Strandberg E, Madsen P, Jensen J, Jorjani H. Genotype by environment interaction in Nordic dairy cattle studied using reaction norms. Acta Agric Scand - Sect A Anim Sci. 2002;52:11–24.

2. Falconer DS, Mackay TFC. Introduction to Quantitative Genetics. 4th ed. Longmans Green, Harlow; 1996.

3. Saltz JB, Bell AM, Flint J, Gomulkiewicz R, Hughes KA, Keagy J. Why does the magnitude of genotype-by-environment interaction vary? Ecol. Evol. 2018. p. 6342–53.

4. Decker JE. Agricultural Genomics: Commercial Applications Bring Increased Basic Research Power. PLOS Genet. 2015;11:e1005621.

5. Howard JT, Kachman SD, Snelling WM, Pollak EJ, Ciobanu DC, Kuehn LA, et al. Beef cattle body temperature during climatic stress: A genome-wide association study. Int J Biometeorol. 2014;58:1665–72.

6. Williams JL, Bertrand JK, Misztal I, Łukaszewicz M. Genotype by environment
interaction for growth due to altitude in United States Angus cattle. J Anim Sci.
2012;90:2152–8.

7. Bradford HL, Fragomeni BO, Bertrand JK, Lourenco DAL, Misztal I. Genetic
evaluations for growth heat tolerance in angus cattle. J Anim Sci. 2016;94:4143–50.

8. Guidolin DGF, Buzanskas ME, Ramos SB, Venturini GC, Lôbo B RB, Paz CCP, et al.
Genotype-environment interaction for post-weaning traits in Nellore beef cattle. Anim
Prod Sci. 2012;52:975–80.

9. MacLeod IM, Bowman PJ, Vander Jagt CJ, Haile-Mariam M, Kemper KE,
Chamberlain AJ, et al. Exploiting biological priors and sequence variants enhances QTL
discovery and genomic prediction of complex traits. BMC Genomics. 2016;17:144.

10. Rowan TN, Durbin HJ, Nilson SM, Seabury CM, Schnabel RD, Decker JE. Powerful
detection of polygenic selection and environmental adaptation in US beef cattle
populations. bioRxiv. 2020;2020.03.11.988121.

11. Ferrell CL. Factors Influencing Fetal Growth and Birth Weight in Cattle. Rom. L.
Hruska U.S. Meat Anim. Res. Cent. 1993. p. 132.

12. Collier RJ, Baumgard LH, Zimbelman RB, Xiao Y. Heat stress: Physiology of
acclimation and adaptation. Anim Front. 2019;9:12–9.

13. Reynolds LP, Ferrell CL, Nienaber JA, Ford SP. Effects of chronic environmental
heat stress on blood flow and nutrient uptake of the gravid bovine uterus and foetus. J
Agric Sci. 2020;104:289–97.

14. Jones HN, Powell TL, Jansson T. Regulation of Placental Nutrient Transport - A
Review. Placenta. 2007;28:763–74.

15. Dunlap KA, Brown JD, Keith AB, Satterfield MC. Factors controlling nutrient
availability to the developing fetus in ruminants. J Anim Sci Biotechnol. 2015;6:1–10.

16. Newman JH, Holt TN, Hedges LK, Womack B, Memon SS, Willers ED, et al. High-
altitude pulmonary hypertension in cattle (Brisket disease): Candidate genes and gene
expression profiling of peripheral blood mononuclear cells. Pulm Circ. 2011;1:462–9.

17. Holt TN, Callan RJ. Pulmonary Arterial Pressure Testing for High Mountain Disease
in Cattle. Vet. Clin. North Am. - Food Anim. Pract. 2007. p. 575–96.

18. Schwartz JC, Sanderson ND, Bickhart DM, Smith TPL, Hammond JA. The Structure,
Evolution, and Gene Expression Within the Caprine Leukocyte Receptor Complex. Front
Immunol. 2019;10:2302.

19. Golombek DA, Rosenstein RE. Physiology of Circadian Entrainment. Physiol Rev.
2010;90:1063–102.

20. Nardone A, Ronchi B, Lacetera N, Bernabucci U. Climatic effects on productive
traits in livestock. Vet Res Commun. 2006;30:75–81.

21. Seltenrich N. Between extremes: Health effects of heat and cold. Environ. Health
Perspect. 2015. p. 276–80.

22. Banks S, King SA, Irvine DS, Saunders PTK. Impact of a mild scrotal heat stress on
DNA integrity in murine spermatozoa. Reproduction. 2005;129:505–14.

23. Hanna EG, Tait PW. Limitations to thermoregulation and acclimatization challenge
human adaptation to global warming. Int J Environ Res Public Health. 2015;12:8034–74.

24. Stuedemann JA, Seman DH. Integrating Genetics, Environment, and Management to
Minimize Animal Toxicoses. Neotyphodium Cool Grasses. 2008. p. 305–24.

25. McCarty K, Bence Á, Maresca S, Lopez Valiente S, Rodriguez AM, Canton GJ, et al.
Exposure to toxic fescue during late gestation on beef cows: effects on cow performance
and offspring performance from birth to weaning. EC Vet Sci. 2020;5:01–15.

26. Haddjeri N, Seletti B, Gilbert F, De Montigny C, Blier P. Effect of ergotamine on
serotonin-mediated responses in the rodent and human brain. Neuropsychopharmacology.
1998;19:365–80.

27. Xue Y, Liao SF, Strickland JR, Boling JA, Matthews JC. Bovine neuronal vesicular
glutamate transporter activity is inhibited by ergovaline and other ergopeptines. J Dairy
Sci. 2011;94:3331–41.

28. Foote AP, Harmon DL, Brown KR, Strickland JR, McLeod KR, Bush LP, et al.
Constriction of bovine vasculature caused by endophyte-infected tall fescue seed extract
is similar to pure ergovaline. J Anim Sci. 2012;90:1603–9.

29. Jackson JA, Hemken RW, Boling JA, Harmon RJ, Buckner RC, Bush LP. Summer
fescue toxicity in dairy steers fed tall fescue seed. J Anim Sci. 1984;58:1057–61.

30. Hoveland CS. Importance and economic significance of the Acremonium endophytes
to performance of animals and grass plant. Agric Ecosyst Environ. 1993;44:3–12.

31. Hemken RW, Boling JA, Bull LS, Hatton RH, Buckner RC, Bush LP. Interaction of
Environmental Temperature and Anti-Quality Factors on the Severity of Summer Fescue
Toxicosis. J Anim Sci. 1981;52:710–4.

32. Shi Z, Wei Z, Li J, Yuan S, Pan B, Cao F, et al. Identification and Verification of
Candidate Genes Regulating Neural Stem Cells Behavior Under Hypoxia. Cell Physiol
Biochem. 2018;47:212–22.

33. Etim NN, Oguike MA. Environmental and Management Stressors: Implications for
Reproductive and Productive Performances of Farm Animals in the Tropics. 2014;5:153–
70.

34. Dawe DL, Stuedemann JA, Hill NS, Thompson FN. Immunosuppression in Cattle
with Fescue Toxicosis. Neotyphodium/Grass Interact. 1997. p. 411–2.

35. Filipov NM, Thompson FN, Stuedemann JA, Elsasser TH, Kahl S, Sharma RP, et al.
Increased responsiveness to intravenous lipopolysaccharide challenge in steers grazing
endophyte-infected tall fescue compared with steers grazing endophyte-free tall fescue. J
Endocrinol. 1999;163:213–20.

36. Krulich L. Central Neurotransmitters and the Secretion of Prolactin, GH, LH and
TSH. Annu Rev Physiol. 1979;41:603–15.

37. Majumdar A, Mangal N. Hyperprolactinemia. J Hum Reprod Sci. 2013;6:168.

38. Li RW, Li C, Gasbarre LC. The vitamin D receptor and inducible nitric oxide
synthase associated pathways in acquired resistance to Cooperia oncophora infection in
cattle. Vet Res. 2011;42:48.

39. Neupane M, Kiser JN, Neibergs HL. Gene set enrichment analysis of SNP data in
dairy and beef cattle with bovine respiratory disease. Anim Genet. 2018;49:527–38.

40. Foley C, Chapwanya A, Callanan JJ, Whiston R, Miranda-CasoLuengo R, Lu J, et al.
Integrated analysis of the local and systemic changes preceding the development of post-
partum cytological endometritis. BMC Genomics. 2015;16:811.

41. Murakami H, Kuroiwa T, Suzuki K, Miura Y, Sentsui H. Analysis of syk expression
in bovine lymphoma and persistent lymphocytosis induced by bovine leukemia virus. J
Vet Med Sci. 2011;73:41–5.

42. Tizioto PC, Kim JW, Seabury CM, Schnabel RD, Gershwin LJ, Van Eenennaam AL,
et al. Immunological response to single pathogen challenge with agents of the bovine
respiratory disease complex: An RNA-sequence analysis of the bronchial lymph node
transcriptome. PLoS One. 2015;10:e0131459.

43. Coussens PM, Pudrith CB, Skovgaard K, Ren X, Suchyta SP, Stabel JR, et al. Johne's
disease in cattle is associated with enhanced expression of genes encoding IL-5, GATA-
3, tissue inhibitors of matrix metalloproteinases 1 and 2, and factors promoting apoptosis
in peripheral blood mononuclear cells. Vet Immunol Immunopathol. 2005;105:221–34.

44. Verschoor CP, Pant SD, You Q, Schenkel FS, Kelton DF, Karrow NA.
Polymorphisms in the gene encoding bovine interleukin-10 receptor alpha are associated
with Mycobacterium avium ssp. paratuberculosis infection status. BMC Genet.
2010;11:23.

45. Khalifeh MS, Stabel JR. Clinical disease upregulates expression of CD40 and CD40
ligand on peripheral blood mononuclear cells from cattle naturally infected with
Mycobacterium avium subsp. paratuberculosis. Clin Vaccine Immunol. 2013;20:1274–
82.

46. Pant SD, Verschoor CP, Schenkel FS, You Q, Kelton DF, Karrow NA. Bovine
CLEC7A genetic variants and their association with seropositivity in Johne's disease
ELISA. Gene. 2014;537:302–7.

47. Jin X, Li G, Zhang X, Gong P, Yu Y, Li J. Activation of a Neospora caninum EGFR-
like kinase facilitates intracellular parasite proliferation. Front Microbiol. 2017;8:1980.

48. Brym P, Kamiński S. Microarray analysis of differential gene expression profiles in
blood cells of naturally BLV-infected and uninfected Holstein–Friesian cows. Mol Biol
Rep. 2017;44:109–27.

49. Domínguez-André J, Netea MG. Impact of Historic Migrations and Evolutionary
Processes on Human Immunity Examples of Evolutionary Pressures Providing Protection
in African Populations. Trends Immunol. 2019;40:1105–19.

50. Mclaren MR, Callahan BJ. Opinion piece Pathogen resistance may be the principal
evolutionary advantage provided by the microbiome. Philos Trans R Soc B.
2020;375:20190592.

51. Conlon I, Raff M. Size control in animal development. Cell. 1999;96:235–44.

52. Beaumont RN, Warrington NM, Cavadino A, Tyrrell J, Nodzenski M, Horikoshi M,
et al. Genome-wide association study of offspring birth weight in 86 577 women
identifies five novel loci and highlights maternal genetic effects that are independent of
fetal genetics. Hum Mol Genet. 2018;27:742–56.

53. Warrington NM, Beaumont RN, Horikoshi M, Day FR, Helgeland Ø, Laurin C, et al.
Maternal and fetal genetic effects on birth weight and their relevance to cardio-metabolic
risk factors. Nat Genet. 2019;51:804–14.

54. Sharma RK, Blair HT, Jenkinson CMC, Kenyon PR, Cockrem JF, Parkinson TJ.
Uterine environment as a regulator of birth weight and body dimensions of newborn
lambs1. J Anim Sci. 2012;90:1338–48.

55. Stein AD, Zybert PA, Van De Bor M, Lumey LH. Intrauterine famine exposure and
body proportions at birth: the Dutch Hunger Winter. Int Epidemiol Assoc. 2004;33:831–
6.

56. Tao S, Monteiro APA, Thompson IM, Hayen MJ, Dahl GE. Effect of late-gestation
maternal heat stress on growth and immune function of dairy calves. J Dairy Sci.
2012;95:7128–36.

57. Cross AJ, Keel BN, Brown-Brandl TM, Cassady JP, Rohrer GA. Genome-wide
association of changes in swine feeding behaviour due to heat stress. Genet Sel Evol.
2018;50:11.

58. Halstead MM, Ma X, Schultz RM, Ross PJ. Chromatin remodeling in bovine
embryos indicates species-specific regulation of genome activation. bioRxiv.
2019;2019.12.12.874479.

59. Moore T, Haig D. Genomic imprinting in mammalian development: a parental tug-of-
war. Trends Genet. 1991;7:45–9.

60. Peretz H, Luboshitsky R, Baron E, Biton A, Gershoni R, Usher S, et al. Cys 618 Arg
mutation in the RET proto☐oncogene associated with familial medullary thyroid
carcinoma and maternally transmitted Hirschsprung's disease suggesting a role for
imprinting. Hum Mutat. 1997;10:155–9.

61. Takacova M, Bullova P, Simko V, Skvarkova L, Poturnajova M, Feketeova L, et al.
Expression pattern of carbonic anhydrase IX in medullary thyroid carcinoma supports a
role for RET-mediated activation of the HIF pathway. Am J Pathol. 2014;184:953–65.

62. Leiser R, Krebs C, Ebert B, Dantzer V. Placental vascular corrosion cast studies: a
comparison between ruminants and humans. Microsc Res Tech. 1997;38:76–87.

63. Pfarrer CD, Ruziwa SD, Winther H, Callesen H, Leiser R, Schams D, et al.
Localization of Vascular Endothelial Growth Factor (VEGF) and its Receptors VEGFR-1
and VEGFR-2 in Bovine Placentomes from Implantation Until Term. Placenta.
2006;27:889–98.

64. Hoffert-Goeres KA, Batchelder CA, Bertolini M, Moyer AL, Famula TR, Anderson
GB. Angiogenesis in day-30 bovine pregnancies derived from nuclear transfer. Cloning
Stem Cells. 2007;9:595–607.

65. Goyal R, Van Wickle J, Goyal D, Matei N, Longo LD. Antenatal Maternal Long-
Term Hypoxia: Acclimatization Responses with Altered Gene Expression in Ovine Fetal
Carotid Arteries. PLoS One. 2013;8:e82200.

66. Stallings JD, Ippolito DL, Rakesh V, Baer CE, Dennis WE, Helwig BG, et al.
Patterns of gene expression associated with recovery and injury in heat-stressed rats.
BMC Genomics. 2014;15:1058.

67. Rönnegård L, Valdar W. Recent developments in statistical methods for detecting
genetic loci affecting phenotypic variability. BMC Genet. 2012;13:63.

68. Visscher PM, Posthuma D. Statistical power to detect genetic loci affecting
environmental sensitivity. Behav Genet. 2010;40:728–33.

69. Bolormaa S, Pryce JE, Zhang Y, Reverter A, Barendse W, Hayes BJ, et al. Non-
additive genetic variation in growth, carcass and fertility traits of beef cattle. Genet Sel
Evol. 2015;47:26.

70. Voz ML, Mathys J, Hensen K, Pendeville H, Van Valckenborgh I, Van Huffel C, et
al. Microarray screening for target genes of the proto-oncogene PLAG1. Oncogene.
2004;23:179–91.

71. Smith JL, Wilson ML, Nilson SM, Rowan TN, Oldeschulte DL, Schnabel RD, et al.
Genome-wide association and genotype by environment interactions for growth traits in
U.S. Gelbvieh cattle. BMC Genomics. BMC Genomics; 2019;20:926.

72. Saatchi M, Schnabel RD, Taylor JF, Garrick DJ. Large-effect pleiotropic or closely
linked QTL segregate within and across ten US cattle breeds. BMC Genomics.
2014;15:442.

73. Seabury CM, Oldeschulte DL, Saatchi M, Beever JE, Decker JE, Halley YA, et al.
Genome-wide association study for feed efficiency and growth traits in U.S. beef cattle.
BMC Genomics. 2017;18:386.

74. Brown AA, Buil A, Viñuela A, Lappalainen T, Zheng HF, Richards JB, et al. Genetic
interactions affecting human gene expression identified by variance association mapping.
Elife. 2014;2014:e01381.

75. Porto-Neto LR, Sonstegard TS, Liu GE, Bickhart DM, Da Silva MVB, Machado MA,
et al. Genomic divergence of zebu and taurine cattle identified through high-density SNP
genotyping. BMC Genomics. 2013;14:876.

76. Taye M, Lee W, Jeon S, Yoon J, Dessie T, Hanotte O, et al. Exploring evidence of
positive selection signatures in cattle breeds selected for different traits. Mamm Genome.
2017;28:528–41.

77. Pitt D, Bruford MW, Barbato M, Orozco☐terWengel P, Martínez R, Sevane N.
Demography and rapid local adaptation shape Creole cattle genome diversity in the
tropics. Evol Appl. 2019;12:105–22.

78. Liu S, Ye T, Li Z, Li J, Jamil AM, Zhou Y, et al. Identifying Hub Genes for Heat
Tolerance in Water Buffalo (Bubalus bubalis) Using Transcriptome Data. Front Genet.
2019;10:209.

79. Dikmen S, Cole JB, Null DJ, Hansen PJ. Genome-Wide Association Mapping for
Identification of Quantitative Trait Loci for Rectal Temperature during Heat Stress in
Holstein Cattle. PLoS One. 2013;8:e69202.

80. Mulder HA, Bijma P, Hill WG. Selection for uniformity in livestock by exploiting
genetic heterogeneity of residual variance. Genet Sel Evol. 2008;40:37–59.

81. Rönnegard L, Felleki M, Fikse WF, Mulder HA, Strandberg E. Variance component
and breeding value estimation for genetic heterogeneity of residual variance in Swedish
Holstein dairy cattle. J Dairy Sci. 2013;96:2627–36.

82. Gonzalez PN, Pavlicev M, Mitteroecker P, Pardo-Manuel de Villena F, Spritz RA,
Marcucio RS, et al. Genetic structure of phenotypic robustness in the collaborative cross
mouse diallel panel. J Evol Biol. 2016;29:1737–51.

83. Kang EY, Han B, Furlotte N, Joo JW, Shih D. Meta-Analysis Identifies Gene-by-
Environment Interactions as Demonstrated in a Study of 4,965 Mice. PLoS Genet.
2014;10:e1004022.

84. Wang H, Zhang F, Zeng J, Wu Y, Kemper KE, Xue A, et al. Genotype-by-
environment interactions inferred from genetic effects on phenotypic variability in the
UK Biobank. Sci Adv. 2019;5:eaaw3538.

85. Kang EY, Park Y, Li X, Segrè A V., Han B, Eskin E. ForestPMPlot: A flexible tool
for visualizing heterogeneity between studies in meta-analysis. G3 Genes, Genomes,
Genet. 2016;6:1793–8.

86. Struchalin M V., Dehghan A, Witteman JCM, van Duijn C, Aulchenko YS. Variance
heterogeneity analysis for detection of potentially interacting genetic loci: Method and its
limitations. BMC Genet. 2010;11:92.

87. Dumitrascu B, Darnell G, Ayroles J, Engelhardt BE. Statistical tests for detecting
variance effects in quantitative trait studies. Bioinformatics. 2019;35:200–10.

88. Paré G, Cook NR, Ridker PM, Chasman DI. On the Use of Variance per Genotype as
a Tool to Identify Quantitative Trait Interaction Effects: A Report from the Women's
Genome Health Study. PLoS Genet. 2010;6:e1000981.

89. Black PH. Stress and the inflammatory response: A review of neurogenic
inflammation. Brain. Behav. Immun. 2002. p. 622–53.

90. Burdick NC, Randel RD, Carroll JA, Welsh TH. Interactions between Temperament,
Stress, and Immune Function in Cattle. Int J Zool. 2011;2011:373197.

91. Joëls M, Baram TZ. The neuro-symphony of stress. Nat. Rev. Neurosci. 2009. p.
459–66.

92. Mora F, Segovia G, Del Arco A, De Blas M, Garrido P. Stress, neurotransmitters,
corticosterone and body-brain integration. Brain Res. 2012;1476:71–85.

93. Morrison SF. Central control of body temperature. F1000Research. 2016;5:880.

94. Colditz IG, Hine BC. Resilience in farm animals: Biology, management, breeding
and implications for animal welfare. Anim. Prod. Sci. 2016. p. 1961–83.

95. Korshunov KS, Blakemore LJ, Trombley PQ. Dopamine: A modulator of circadian
rhythms in the central nervous system. Front. Cell. Neurosci. 2017. p. 91.

96. Verma P, Sharma A, Sodhi M, Thakur K, Kataria RS, Niranjan SK, et al.
Transcriptome Analysis of Circulating PBMCs to Understand Mechanism of High
Altitude Adaptation in Native Cattle of Ladakh Region. Sci Rep. 2018;8:7681.

97. Kong Z, Zhou C, Chen L, Ren A, Zhang D, Basang Z, et al. Multi-omics analysis
reveals up-regulation of APR signaling, LXR/RXR and FXR/RXR activation pathways in
holstein dairy cows exposed to high-altitude hypoxia. Animals. 2019;9:406.

98. Carvalheiro R, Costilla R, Neves HHR, Albuquerque LG, Moore S, Hayes BJ.
Unraveling genetic sensitivity of beef cattle to environmental variation under tropical
conditions. Genet Sel Evol. 2019;51:29.

99. Bova TL, Chiavaccini L, Cline GF, Hart CG, Matheny K, Muth AM, et al.
Environmental stressors influencing hormones and systems physiology in cattle. Reprod
Biol Endocrinol. 2014;12:58.

100. Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, et al.
Common SNPs explain a large proportion of the heritability for human height. Nat Genet.
2010;42:565–9.

101. Ugarte E, Alenda R, Carabaño MJ. Fixed or Random Contemporary Groups in
Genetic Evaluations. J Dairy Sci. 1992;75:269–78.

102. Rowan TN, Hoff JL, Crum TE, Taylor JF, Schnabel RD, Decker JE. A multi-breed
reference panel and additional rare variants maximize imputation accuracy in cattle.
Genet Sel Evol. 2019;51:77.

103. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al.
PLINK: A tool set for whole-genome association and population-based linkage analyses.
Am J Hum Genet. 2007;81:559–75.

104. Rosen BD, Bickhart DM, Schnabel RD, Koren S, Elsik CG, Tseng E, et al. De novo
assembly of the cattle reference genome with single-molecule sequencing. Gigascience.
2020;9:1–9.

105. Das S, Forer L, Schönherr S, Sidore C, Locke AE, Kwong A, et al. Next-generation
genotype imputation service and methods. Nat Genet. 2016;48:1284–7.

106. Loh PR, Danecek P, Palamara PF, Fuchsberger C, Reshef YA, Finucane HK, et al.
Reference-based phasing using the Haplotype Reference Consortium panel. Nat Genet.
2016;48:1443–8.

107. PRISM Climate Group, Oregon State University, http://prism.oregonstate.edu. 2004.

108. Leutner B, Horning N, Schwalb-Willmann J, Hijmans RJ. Package "RStoolbox"
Title Tools for Remote Sensing Data Analysis. 2019.

109. R Core Team. R: A language and environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org.
(2020).

110. Henning C. Package "fpc" Title Flexible Procedures for Clustering. 2019.

111. Zhu Z, Bakshi A, Vinkhuyzen AAE, Hemani G, Lee SH, Nolte IM, et al.
Dominance genetic variation contributes little to the missing heritability for human
complex traits. Am J Hum Genet. 2015;96:377–85.

112. Henderson CR. Best Linear Unbiased Prediction of Nonadditive Genetic Merits in
Noninbred Populations. J Anim Sci. 1985;60:111–7.

113. Zhou X, Stephens M. Genome-wide efficient mixed-model analysis for association
studies. Nat Genet Vol. 2012;44:821–6.

114. Zhou X, Stephens M. Efficient multivariate linear mixed model algorithms for
genome-wide association studies. Nat Methods. 2014;11:407–9.

115. Turner SD. qqman: an R package for visualizing GWAS results using Q-Q and
manhattan plots. J Open Source Softw. 2018;3:731.

116. Yang J, Loos RJF, Powell JE, Medland SE, Speliotes EK, Chasman DI, et al. FTO
genotype is associated with phenotypic variability of body mass index. Nature.
2012;490:267–73.

117. Braz CU, Taylor JF, Bresolin T, Espigolan R, Feitosa FLB, Carvalheiro R, et al.
Sliding window haplotype approaches overcome single SNP analysis limitations in
identifying genes for meat tenderness in Nelore cattle. BMC Genet. 2019;20:8.

118. Han B, Eskin E. Random-effects model aimed at discovering associations in meta-
analysis of genome-wide association studies. Am J Hum Genet. 2011;88:586–98.

119. Han B, Eskin E. Interpreting meta-analyses of genome-wide association studies.
PLoS Genet. 2012;8:e1002555.

120. Bolormaa S, Hayes BJ, Savin K, Hawken R, Barendse W, Arthur PF, et al. Genome-
wide association studies for feedlot and growth traits in cattle. J Anim Sci.
2011;89:1684–97.

121. Ge SX, Jung D, Jung D, Yao R. ShinyGO: A graphical gene-set enrichment tool for
animals and plants. Bioinformatics. 2020;36:2628–9.

122. Carbon S, Ireland A, Mungall CJ, Shu S, Marshall B, Lewis S, et al. AmiGO: online
access to ontology and annotation data. Bioinformatics. 2009;25:288–9.

1045

List of figures

Fig. 1. Genotype-by-environment interaction genome-wide association analyses (GxE

GWAA) for growth traits in Simmental cattle. (*A*) Mean dew point temperature average

annual over the most recent three full decades covering the conterminous United States.

(*B*) Manhattan plot of GxE GWAA of mean dew point temperature for birth weight. (*C*)

Elevation of the conterminous United States. (*D*) Manhattan plot of GxE GWAA of

elevation for weaning weight. (*E*) Manhattan plot of GxE GWAA of elevation for

yearling weight. (*F*) Boundaries for ecoregion assignments in the United States; (*Top*)

United States partitioned into nine ecoregions based on similar topographic and

environmental conditions; (*Bottom*) location of beef farms for which data was retrieved.

(G) Manhattan plot of GxE GWAA of Desert & Arid Prairie ecoregion for birth weight.

(*H*) Manhattan plot of GxE GWAA of Forested Mountains ecoregion for weaning

weight. (*I*) Manhattan plot of GxE GWAA of Forested Mountains ecoregion for yearling

weight. In Manhattan plots, horizontal red line indicates a significant threshold ($P <$ 1e-

5). Environmental continuous variables were drawn from the PRISM climate dataset

(http://prism.oregonstate.edu). The United States was partitioned into nine regions using

k-means clustering.

1064

1065

1066 **Fig. 2.** Percentage of candidate genes with genotype-by-environment effects for each

1067 ecoregion that contribute to the enriched gene ontology (GO) terms. Genes were located

1068 100 kb from significant SNPs (P < 1e-5).

1069

1070

Fig. 3. Percentage of candidate genes with genotype-by-environment effects for each

ecoregion that contribute to the enriched biological pathways. Genes were located 100 kb

from significant SNPs (P < 1e-5).

1074 **Table 1.** Number of significant SNPs ($P < 1e\text{-}05$) and magnitude of the allele substitution
1075 effects by environmental variables/factors for each trait.

Env	Birth Weight		Weaning Weight		Yearling Weight		Mv	Total
	N	ASE	N	ASE	N	ASE	N	N
Elev	142	-0.001–0.001	28	-0.008–0.002	15	-0.005–0.006	50	216
Precip	338	-0.001–0.002	4	-0.004– -0.004	1	-0.02	12	349
MT	555	-0.170–0.201	1	0.49	4	-1.240–1.704	69	611
MinT	756	-0.159–0.212	6	-0.868–0.439	-	-	88	823
MaxT	408	-0.175–0.207	1	0.43	4	-1.267–1.848	23	427
MDPT	941	-0.178–0.168	4	-0.516–0.409	1	-1.11	94	1018
MinVPD	152	-1.399–1.831	2	-3.254– -3.242	1	-6.62	10	165
MaxVPD	224	-0.213–0.262	10	-0.546–0.556	1	1.08	17	236
SE	54	-2.078–1.649	-	-	2	-11.120– -10.500	7	59
HP	40	-1.381–1.132	16	-5.353–3.930	4	-22.900–6.722	23	63
FM	90	-3.135–1.215	31	-6.785–4.737	49	-9.397–17.690	148	221
FB	47	-2.145–0.953	1	-2.94	1	18.53	11	56
UN	24	-0.709–0.872	-	-	1	8.18	10	33
DA	125	-8.243–6.043	8	-11.570–6.456	6	-16.450–23.620	*	125
Total	1923		101		78		353	2319

Results of the genotype-by-environment genome-wide Association Analysis using
environmental variables (Env), such as elevation (Elev), precipitation (Precip), mean
temperature (MT), minimum temperature (MinT), maximum temperature (MaxT), mean
dew point temperature (MDPT), minimum vapor pressure deficit (MinVPD), and
maximum vapor pressure deficit (MaxVPD), and using the United Stated ecoregions
(Southeast [SE], High Plains [HP], Forested Mountains [FM], Fescue Belt [FB], Upper
Midwest & Northeast [UN], and Desert & Arid Prairie [DA]) as environmental factors
(Env. Factor) in univariate (Birth Weight, Weaning Weight, and Yearling Weight) and
multivariate (Mv) linear mixed models. The number of significant SNPs (N) at the
significant threshold of $P < 1e\text{-}05$ and the magnitude of the allele substitution effects
(ASE, kg) were also provided.