Chroococcidiorella tianjinensis, **gen. et sp. nov. (Trebouxiophyceae,**

Chlorophyta), a green alga arises from the cyanobacterium TDX16

Qing-lin Dong and Xiang-ying Xing

Department of Bioengineering, Hebei University of Technology, Tianjin, 300130, China

Corresponding author: qldong@hebut.edu.cn

ABSTRACT

We provide a taxonomic description of the first origin-known alga TDX16-DE that arises from

the *Chroococcidiopsis*-like endosymbiotic cyanobacterium TDX16 by de novo organelle

biogenesis after acquiring its green algal host *Haematococcus pluvialis*'s DNA. TDX16-DE is

spherical or oval, with a diameter of 2.9-3.6 μm, containing typical chlorophyte pigments of

chlorophyll a, chlorophyll b and lutein and reproducing by autosporulation, whose 18S rRNA

gene sequence shows the highest similarity of 99.8% to that of *Chlorella vulgaris*. However,

TDX16-DE is only about half the size of *C. vulgaris* and structurally similar to *C. vulgaris* only

in having a chloroplast-localized pyrenoid, but differs from *C. vulgaris* in that (1) it possesses a

double-membraned cytoplasmic envelope but lacks endoplasmic reticulum and Golgi apparatus;

and (2) its nucleus is enclosed by two sets of envelopes (four unit membranes). Therefore, based

on these characters and the cyanobacterial origin, we describe TDX16-DE as a new genus and

new species, *Chroococcidiorella tianjinensis* gen. et sp. nov.

KEYWORDS: Green algae; Cyanobacterial origin; *Chroococcidiorella tianjinensis*; 18S rRNA

gene sequence; Ultrastructure

INTRODUCTION

Chlorella vulgaris is the first documented *Chlorella* green alga discovered by Beijerinck in 1890

(Beijerinck,1890). Since then, about 1000 orbicular-shaped *Chlorella* and *Chlorella*-like green

algae have been isolated (Luo et al., 2010). Classification of these small coccoid chlorophytes is

a difficult task, because their origins and evolutionary histories/relationships, the crucial

information necessary for their accurate assignments, are unknown. In this circumstance, the

Chlorella and *Chlorella*-like green algae are classified traditionally by comparing the degrees of

resemblance of their morphological, biochemical, physiological and ultrastructural features (Fott

and Nováková 1969; Andreyeva 1975; Maruyama 1977; KÜmmel and Kessler 1980; Takeda

1991; Kessler and Huss 1992; Ikeda and Takeda 1995; Hanagata et al., 1998) and currently by

inferring their origins and evolutionary histories/relationships via phylogenetic analyses of the

molecular data, e.g., sequences of the SSU and ITS regions of nuclear-encoded ribosomal DNA

(Friedl 1995; Huss et al., 1999; Krienitz et al., 2004; Henley et al., 2004; Karsten et al., 2005;

Luo et al., 2010; Bock et al., 2011; Pröschold et al., 2011; Neustupa et al., 2013; Škaloud et al.,

2014; Song et al., 2018). As such, the taxonomy of *Chlorella* and *Chlorella*-like green algae is

problematic and unstable, as indicated by the frequent taxonomic revisions.

In our previous studies, we found that the endosymbiotic cyanobacterium TDX16 resembling

Chroococcidiopsis thermalis escaped from the senescent/necrotic cells of green alga

Haematococcus pluvialis (host) (Dong et al., 2011), and turned into a *Chlorella*-like green alga

TDX16-DE (Dong et al., 2016) by de novo organelle biogenesis after hybridizing the acquired

host's DNA with its own one and expressing the hybrid genome (Dong et al., 2017). Therefore,

the *Chlorella*-like green alga TDX16-DE arises form the *Chroococcidiopsis*-like cyanobacterium

TDX16, which is the first alga with known origin and developmental process, providing a

reference for investigating the origins and evolutionary histories of other algae and improving

our understanding of prokaryote-eukaryote connection. In this study, based upon TDX16-DE's

cyanobacterial origin in combination with its cell morphology, ultrastructure, photosynthetic

pigments and 18S rRNA gene sequence, we describe TDX16-DE as a new genus and species,

Chroococcidiorella tianjinensis, gen. et sp. nov.

METHODS

Strain and cultivation

TDX16-DE was obtained and prepared in the same way as described by Dong et al., (2017).

Microscopy observations

Light microscopy: TDX16-DE cells were examined with a light microscope BK6000 (Optec,

China). Photomicrographs were taken under the oil immersion objective lens (100×) using a DV

E3 630 camera.

Transmission electron microscopy: TDX16-DE cells were harvested by centrifugation (3000

rpm, 10 min) and fixed overnight in 2.5% glutaraldehyde (50 mM phosphate buffer, pH7.2) and

1% osmium tetroxide (the same buffer) for 2 h at room temperature. After dehydration with

ascending concentrations of ethanol, the fixed cells were embedded in Spurr's resin at 60℃ for

24 h. Ultrathin sections (60 to 80 nm) were cut with a diamond knife, mounted on a copper grid,

double-stained with 3% uranyl acetate and 0.1% lead citrate, and examined using a JEM1010

electron microscope (JEOL, Japan).

Pigment analyses

In vivo absorption spectrum: TDX16-DE cells were scanned with Ultraviolet-Visible

Spectrophotometer Cary 300 (Agilent, USA).

Pigment separation and identification: Chlorophyll b (Chl b) and lutein were separated by

thin-layer chromatography according to the method described by Lichtenthaler (1987). Pigments

were analyzed with Spectrophotometer Cary 300 (Agilent, USA), and identified by spectroscopic

matching with the published data.

18S rRNA gene sequence

DNA sample was prepared according to the method (Dong et al., 2017). 18S rRNA gene was

amplified using the primers RP8 (5'-ACCTGGTTGATCCTGCCAGTAG-3') and RP9 (5'-

ACCTTGTTACGACTTCTCCTTCCTC-3') under the conditions: 5 min at 95℃, 35 cycles of 30

s at 95℃, 30 s at 55℃ and 1min at 72℃ and a final 10 min extension step at 72℃. The

amplified product was sequenced on ABI 3730 DNA analyzer (PerkinElmer Biosystems, USA).

RESULTS

Morphology and ultrastructure

TDX16-DE cells are green, spherical or oval, about 2.9-3.6 μm in diameter (Figs 1-2),

containing one nucleus, one chloroplast, one or two mitochondria and one or several vacuoles,

Figs 1-2. Morphology of *Chroococcidiorella tianjinensis*. Scale bar = 2 μm. C, chloroplast.
Fig.1. The three-celled autosporangium (arrowhead).
Fig.2. The two-celled autosporangium (arrowhead).

but no endoplasmic reticulum, Golgi apparatus and peroxysome (Figs 3-6). The nucleus has two

sets of envelopes, which are in close apposition in most cases (Figs 4-6), but separated from each

other when the electron-dense vesicles bud from the nuclear envelope into the interenvelope

space, and migrate outside to the chloroplast envelope and cytoplasmic envelope (Fig.3). The

chloroplast is parietal and "e"-shaped, in which there are starch granules, plastoglobuli and a

prominent pyrenoid that is surrounded by two starch plates and bisected by two pairs of

thylakoids (Figs 3-6). Mitochondria lie between the chloroplast and nucleus in the chloroplast

cavity (Figs 3-4, 6). Vacuoles are encircled by two unit membranes, sequestering electron-

Figs 3-6. Ultrastructure of *Chroococcidiorella tianjinensis*. Scale bar = 0.5 μm. C, chloroplast; CE, cytoplasmic envelope; CW, cell wall; EDV, electron-dense vesicle; ES, extracytoplasmic space; FB, microfibrils; IES, interenvelope space; LD, lipid droplet; M, mitochondrion; N, nucleus; NE, nuclear envelope; OE, outer nuclear envelope; P, pyrenoid; PG, plastoglobuli; S, starch granule; SH, sheath; SP; starch plate; TD, trilaminar domain; V, vacuole.

Fig. 3. The two sets of nuclear envelops separate from each other, resulting in a broad interenvelop space.

Fig. 4. The two sets of nuclear envelopes and cytoplasmic envelope are in close apposition in the opening of chloroplast cavity.

Fig. 5. The pyrenoid is bisected by two thylakoids (large arrowhead), the vacuole is delimited by double membranes (small arrowhead), the outer sheath of cell wall detaches from the trilaminar domain, and microfibrils emanate from the cytoplasmic envelope.

Fig. 6. Lipid droplets form at the inner leaflet of cytoplasmic envelope that consists of two membranes.

transparent or electron-dense materials (Figs 3, 5-6). The cells are enclosed with a cytoplasmic

envelope (two unit membranes), on and from which lipid droplets form and microfibrils emanate

into the extracytoplasmic space respectively (Figs 5-6), indicating that the double-membraned

cytoplasmic envelope serves the functions of endoplasmic reticulum and Golgi apparatus for

synthesizing lipids and building materials of cell wall. The cell wall comprises an inner stable

Figs 7-10. Reproduction of *Chroococcidiorella tianjinensis*. Scale bar = 0.5 μm. AU, autosporangium; C,
chloroplast; CW, cell wall; LD, lipid droplet; M, mitochondrion; N, nucleus; V, vacuole.
Fig. 7. A cell contains two small chloroplasts but no nucleus.
Fig. 8. A two-celled autosporangium ruptures, resulting in an opening (arrowhead).
Fig. 9. An autosporangium sequesters three mature daughter cells.
Fig. 10. An autosporangium encloses four immature daughter cells.

trilaminar domain with dark-light-dark configuration and an outer loosely-compacted stratified

sheath (Figs 3-4, 6), the latter of which occasionally scale off owing to osmotic-shock during cell

fixation (Fig.5). It is noteworthy that the two unit membranes of cytoplasmic envelope and

vacuoles, like those of chloroplast and mitochondrial envelopes, are difficult to distinguish when

they are tightly appressed and easily misrecognized as the two layers of one unit membrane as

they are separated, while the two sets of nuclear envelopes are easily misinterpreted as the two

unit membranes of one nuclear envelope without being aware of their formation processes (Dong

et al., 2017).

Reproduction

TDX16-DE reproduces exclusively by autosporulation with the production of two (Figs 2, 8),

three (Figs 1, 9) and four daughter cells (autospores) (Fig.10) per autosporangium. The

autospores are liberated after apical rupture of the autosporangium (Fig.8). At the beginning of

autosporulation, the chloroplast splits into two small ones, while the nucleus disassembles and

vanishes (Fig.7). Consistently, nucleus appears only in one cell in the immature four-celled

autosporangium (Fig.10), but in all cells in the mature three-celled autosporangium (Fig.9); by

contrast chloroplast presents in each cells in both the mature and immature autosporangia (Figs

9-10). These results suggest that chloroplast develops by splitting the existing one, while nucleus

forms de novo in the daughter cells during autosporulation.

Photosynthetic pigments

In vivo absorption spectrum of TDX16-DE (Fig.11) exhibits two absorption peaks of chlorophyll

a (Chl a) at 440 and 680 nm, a conspicuous shoulder peak at 653 nm of chlorophyll b (Chl b)

(Govindjee and Rabinowitch, 1960; Wilhelm et al., 1982), and a merged peak of carotenoids

around 485 nm. Chl b and lutein separated from TDX16-DE display absorption peaks at 456 and

645 nm (Fig. 12), 420, 446 and 475nm (Fig. 13) respectively, identical to those of plant pigments

(Lichtenthaler, 1987). Hence, TDX16-DE contains chl a, chl b and lutein.

Figs 11-13. Photosynthetic pigments of *Chroococcidiorella tianjinensis*.
Fig. 11. In vivo absorption spectrum.
Fig. 12. Absorption spectrum of the separated Chl b.
Fig. 13. Absorption spectrum of the separated lutein.

18S rRNA gene sequence

18S rRNA gene sequence analysis of TDX16-DE resulted in 1687 nucleotides, which shows the

highest similarity of 99.8% (query cover 75%) to those of *Chlorella vulgaris* deposited in

GenBank.

DISCUSSION

The presence of typical chlorophyte pigments of Chl a, Chl b and lutein in TDX16-DE (Figs 11-

13) indicates that TDX16-DE belongs to the Chlorophyta, and the highest similarity of 99.8%

between the 18S rRNA gene sequences of TDX16-DE and *Chlorella vulgaris* suggests that

TDX16-DE is most similar to *C. vulgaris*. The results of cell morphology and ultrastructure,

however, demonstrate that TDX16-DE is distinct from *C. vulgaris*, because they share only two

common traits but differ in six characters. Their common features include (1) chloroplasts are

embedded with starch granules and a pyrenoid that is surrounded by two starch plates and

bisected by two pairs of thylakoids in TDX16-DE (Figs 3-6) and *C. vulgaris* (Němcová and

Kalina, 2000; Přibyl et al., 2013; Gärtner et al., 2015) and (2) reproduction is achieved by

autosporulation, generating two, three and four autospores in TDX16-DE (Figs1-2 and 8-10) and

C. vulgaris (Gärtner et al., 2015; Němcová and Kalina, 2000; Yamamoto et al., 2004). By

contrast, their different characters encompass (1) TDX16-DE (2.9-3.6 μm) (Figs1-10) is only

about half the size of *C. vulgaris* (5.0-7.0 μm) (Yamamoto et al., 2004; Gärtner et al., 2015); (2)

TDX16-DE's chloroplast is "e"-shaped (Figs 3-6); while the chloroplast of *C. vulgaris* is cup-

shaped and sometimes has two lobes (bilobed) (Bisalputra et al., 1966; Gärtner et al., 2015); (3)

TDX16-DE lacks endoplasmic reticulum and Golgi apparatus (Figs 3-6), while *C. vulgaris*

possesses both of the two organelles (Bisalputra et al., 1966; Griffiths and Griffiths, 1969;

Gerken et al., 2013; Přibyl et al., 2013); (4) TDX16-DE's nucleus is enclosed by two sets of

envelopes (Figs 3-6), while the nucleus of *C. vulgaris* has one envelope (Griffiths and Griffiths,

1969); (5) TDX16-DE has a double-membraned cytoplasmic envelope (Figs 3-6), while *C.*

vulgaris has a single-membraned cytoplasmic membrane (Yamamoto et al., 2004); and (6)

daughter cell nuclei form de novo in TDX16-DE (Figs 7-10), but develop by dividing the

existing one in *C. vulgaris* (Bisalputra et al., 1966; Yamamoto et al., 2004).

Since TDX16-DE is tiny (2.9-3.6 μm) and nearly pico-sized (< 3 μm), it is much similar in

structure, particularly organelle number and arrangement, to the pico-sized *Nannochloris*-like

green algae, which have a minimal set of organelles and differ from TDX16-DE in their absence

of pyrenoid, such as *Choricystis minor* (Krienitz, et al., 1996), *Pseudodictyosphaerium jurisii*

(Krienitz et al., 1999), *Mychonastes homosphaera* (Hanagata, et al., 1999), *Picocystis salinarum*

(Lewin et al., 2000), *Picochlorum oklahomensis* (Henley et al., 2004), *Chloroparva pannonica*

(Somogyi et al., 2011) and *Pseudochloris wilhelmii* (Somogyi et al., 2013). The structural

similarity between TDX16-DE and these *Nannochloris*-like green algae lies primarily in three

aspects: (1) absence of endoplasmic reticulum and Golgi apparatus (except the Golgi apparatus-

like structure in *P. salinarum, P. jurisii* and *P. wilhelmii*); (2) the nucleus is peripherally

localized in the opening of chloroplast cavity and appressed to the cytoplasmic

membrane/envelope; and (3) mitochondria situate between the chloroplast and nucleus and

usually attach to the chloroplast.

Phylogenetic analysis of SSU and ITS rDNA is now applied predominately for classifying

the origin-unknown algae by inferring their origins and evolutionary histories/relationships,

which is, however, not required and also not feasible for taxonomic assignment of TDX16-DE.

Because (1) the origin and developmental relationship of TDX16-DE is clearly known; and (2)

current molecular phylogenetic analyses are based on the fundamental assumption that all algae

descend directly from other algae, while TDX16-DE is not the case, which arise from the

cyanobacterium TDX16 by acquiring *H. pluvialis*'s DNA (Dong et al., 2017). So, TDX16-DE

descend, to some extent, only indirectly from the green alga *H. pluvialis* via the cyanobacterium

TDX16. Such a developmental relationship apparently can not be inferred by current molecular

phylogenetic analysis, or in other words, the inferred phylogenetic relationship can not reflect the

true origin and development relationship of TDX16-DE (it is also the case for the common

ancestor of algae that did not arise from algae).

In truth, the experimental results discussed above are sufficient for assigning TDX16-DE.

The *Chlorella* and *Chlorella*-like coccoid green algae are currently categorized into two different

clades within in Chlorellaceae: the *Chlorella*-clade and the *Parachlorella*-clade (krienitz et al.

2004). The highest similarity of TDX16-DE's 18S rRNA gene sequence to that of *C. vulgaris*,

and the typical *Chlorella* structure: a prominent pyrenoid surrounded by two starch plates and

bisected by two pairs of thylakoids in the single parietal chloroplast indicate that TDX16-DE

falls within the *Chlorella*-clade; while TDX16-DE's tiny cell size, compartmental characters:

two sets nuclear envelopes, double-membraned cytoplasmic envelope and vacuoles and absence

of endoplasmic reticulum, Golgi apparatus and peroxysome distinguish it from all the members

within the genera of *Chlorella*-clade (Luo et al., 2010; Bock et al., 2011; Pröschold et al. 2011;

krienitz et al. 2012). Therefore, we propose a new genus *Chroococcidiorella* within the

Chlorella-clade and TDX16-DE is a new species *Chroococcidiorella tianjinensis*.

Chroococcidiorella Q. Dong & X. Xing, gen. nov.

Cells solitary, spherical or oval, enclosed by a double-membraned cytoplasmic envelope.
Nucleus encircled by two sets of envelopes, endoplasmic reticulum and Golgi apparatus absent.
One parietal chloroplast with a pyrenoid surrounded by two starch plates and traversed by two
pairs of thylakoids. Asexual reproduction by autosporulation. Sexual reproduction unknown.

Etymology: The new genus *Chroococcidiorella* is named for its origin from the cyanobacterium

Chroococcidiopsis; plus ella, small or diminutive.

Type species: *Chroococcidiorella tianjinensis* Q. Dong & X. Xing

***Chroococcidiorella tianjinensis* Q. Dong & X. Xing, sp. nov. (Figs 1-10)**

Cells solitary, spherical or oval, with a diameter of 2.9-3.6 μm, One nucleus with two sets of
envelopes, one parietal "e"-shaped chloroplast with a pyrenoid surrounded by two starch plates
and bisected by double thylakoids, double-membraned cytoplasmic envelope and vacuoles. No
endoplasmic reticulum, Golgi apparatus and peroxysome. Chloroplast pigments comprise
chlorophyll a, b and lutein. Asexual reproduction via 2-4 autospores. Sexual reproduction not
observed.

Holotype: TDX16-DE has been preserved in the microbial collection of Hebei university of
technology, Tianjin, China.

Etymology: The name tianjinensis refers to the Chinese city, Tianjin, where formation of this
alga was found.

REFERENCES

Andreyeva V. M (1975) Rod *Chlorella*. Izdat. Nauka, Leningrad: 82p.

Beijerinck M. W (1890) Culturversuche mit Zoochlorellen, Lichen-engonidien und anderen

niederen Algen I–III. *Botanische Zeitung* 48: 726–740.

Bisalputra T., Ashton F and Weier, T (1966) Role of dictyosomes in wall formation during cell

division of *Chlorella vulgaris*. *American Journal of Botany* 53: 213-216

Bock C., Krienitz L and Pröschold T (2011) Taxonomic reassessment of the genus *Chlorella*

(Trebouxiophyceae) using molecular signatures (barcodes), including description of seven

new species. *Fottea* 11: 293-312.

Dong Q. L., Li Z.W., Xing X.Y and Chen B (2011) Discovery of an endophytic cyanobacterium

in *Haematococcus pluvialis*. *Journal of Hebei University of Technology* 40: 1-5

Dong Q. L., Xing X.Y., Wu H.X., Han Y., Wei X.L and Zhang S (2016) Transition of a

prokaryotic endosymbiotic cyanobacterium into a eukaryotic green alga. *Chemical*

Engineering (China) 44: 1-6

Dong Q. L., Xing X.Y., Han Y., Wei X. L and Zhang S (2017) De novo organelle biogenesis in

the cyanobacterium TDX16 released from the green alga *Haematococcus pluvialis*. *BioRxiv*,

http://dx.doi.org/10.1101/161463

Fott B and Nováková M (1969) A monograph of the genus *Chlorella*. The freshwater species. In:

Fott B. (Ed.), Studies in Phycology. Academia, Praha: 10–59.

Friedl, T (1995) Inferring taxonomic positions and testing genus level assignments in coccoid

green algae: a phylogenetic analysis of 18S ribosomal RNA sequences from *Dictyochloropsis*

reticulata and from members of the genus *Myrmecia* (Chlorophyta, Trebouxiophyceae cl.

nov.). *Journal of Phycology* 31:632–639.

Gärtner G., Uzunov B., Ingolic E., Kofler W., Gacheva G., Pilarski P., Zagorchev L., Odjakova

M and Stoyneva M (2015) Microscopic investigations (LM, TEM and SEM) and

identification of *Chlorella* isolate R-06/2 from extreme habitat in Bulgaria with strong

biological activity and resistance to environmental stress factors. *Biotechnology &*

Biotechnological Equipment 29:536–540

Gerken H., Donohoe B and Knoshaug E (2013) Enzymatic cell wall degradation of *Chlorella*

vulgaris and other microalgae for biofuels production. *Planta* 237: 239–253.

Govindjee and Rabinowitch E (1960)Two forms of chlorophyll a in vivo with distinct

photochemical functions. *Science* 132: 355-356.

Griffiths D and Griffiths D (1969) The fine structure of autotrophic and heterotrophic cells of

Chlorella vulgaris (Emerson strain). *Plant Cell Physiology* 10:11–19.

Hanagata N., Karube I., Chihara M and Silva P. C (1998) Reconsideration of the taxonomy of

ellipsoidal species of *Chlorella* (Trebouxiophyceae, Chlorophyta), with establishment of

Watanabea sen. nov. *Phycological Research* 46: 221–229.

Hanagata N., Malinsky-Rushansky N and Dubinsky Z (1999) Eukaryotic picoplankton,

Mychonastes homosphaera (Chlorophyceae, Chlorophyta), in Lake Kinneret, Israel.

Phycological Research 47: 263–269

Henley W., Hironaka J., Guillou L., Buchheim M., Buchheim J., Fawley M and Fawley K (2004)

Phylogenetic analysis of the '*Nannochloris*-like' algae and diagnoses of *Picochlorum*

oklahomensis gen. et sp. nov. (Trebouxiophyceae, Chlorophyta). *Phycologia* 43: 641–652.

Huss V., Frank C., Hartmann E., Hirmer M., Kloboucek A., Seidel B., Wenzeler P and Kessler E

(1999) Biochemical taxonomy and molecular phylogeny of the genus *Chlorella* sensu lato

(Chlorophyta). *Journal of Phycology* 35: 587–598.

Ikeda, T and Takeda, H (1995) Species-specific differences of pyrenoids in *Chlorella*

(Chlorophyta). *Journal of Phycology* 31:813–818.

Karsten, U., Friedl, T., Schumann, R., Hoyer, K and Lembcke, S (2005) Mycosporing-like

amino acids and phylogenies in green algae: *Prasiola* and ITS relatives from the

Trebouxiophyceae (Chlorophyta). *Journal of Phycology* 41: 557–566.

Kessler, E and Huss, V.A.R (1992) Comparative physiology and biochemistry and taxonomic

assignment of the *Chlorella* (Chlorophyceae) strains of the Culture Collection of the

University of Texas at Austin. *Journal of Phycology* 28: 550-553.

Krienitz L., Huss V and Hummer C (1996) Picoplanktonic *Choricystis* species (Chlorococcales,

Chlorophyta) and problems surrounding the morphologically similar '*Nannochloris*-like'

algae. *Phycologia* 35: 332–341

Krienitz L., Takeda H and Hepperle D (1999) Ultrastructure, cell wall composition, and

phylogenetic position of *Pseudodictyosphaerium jurisii* (Chlorococcales, Chlorophyta)

including a comparison with other picoplanktonic green algae. *Phycologia* 38: 100–107.

Krienitz L., Hegewald E., Hepperle D., Huss V., Rohr T and Wolf M (2004) Phylogenetic

relationship of *Chlorella* and *Parachlorella* gen. nov. (Chlorophyta, Trebouxiophyceae).

Phycologia 43: 529–542.

Krienitz L., Bock C., Kotut, K and Pröschold T (2012) Genotypic diversity of *Dictyosphaerium*-

morphospecies (Chlorellaceae, Trebouxiophyceae) in African inland waters, including the

description of four new genera. *Fottea* 12: 231–253.

KÜmmel, H and Kessler, E (1980) Physiological and biochemical contributions to the taxonomy

of the genus *Chlorella*. XIII. Serological studies. *Archives of Microbiology* 126:15–19.

Lewin R., Krienitz L., Goericke R., Takeda H and Hepperle D (2000) *Picocystis salinarum* gen.

et sp. nov. (Chlorophyta) – a new picoplanktonic green alga. *Phycologia* 39: 560–565.

Lichtenthaler, H (1987) Chlorophylls and carotenoids: pigments of photosynthetic

biomembranes. *Methods in Enzymology* 148: 350–382.

Luo W., Pröschold T., Bock C and krienitz L (2010) Generic concept in *Chlorella*–related

coccoid green algae (Chlorophyta, Trebouxiophyceae). *Plant Biology* 12: 545–553.

Maruyama, K (1977) Classification of *Chlorella* strains by cell appearance and group sera. *The*

Botanical Magazine Tokyo 90:57–66.

Němcová Y and Kalina T (2000) Cell wall development, microfibril and pyrenoid structure in

type strains of *Chlorella vulgaris, C kessleri, C sorokiniana* compared with *C luteoviridis*

(Trebouxiophyceae, Chlorophyta). *Algological Studies* 100: 95–105

Neustupa J., Němcová Y., Veselá J., Steinová J and Škaloud P (2013) *Leptochlorella corticola*

gen. et sp. nov. and *Kalinella apyrenoidosa* sp. nov.: two novel *Chlorella*-like green

microalgae (Trebouxiophyceae, Chlorophyta) from subaerial habitats. International Journal

of Systematic and Evolutionary Microbiology 63: 377–387

Přibyl P., Cepák V and Zachleder V (2013) Production of lipids and formation and mobilization

of lipid bodies in *Chlorella vulgaris*. *Journal of Applied Phycology* 25: 545–553.

Pröschold T., Darienko T., Silva P., Reisser W and Krienitz L (2011) The systematic of

Zoochlorella revisited employing an integrative approach. *Environmental Microbiology* 13:

350-364.

Škaloud P., Němcová Y., Pytela J., Bogdanov N.I., Bock, C and Pickinpaugh S.H (2014)

Planktochlorella nurekis gen. et sp. nov.(Trebouxiophyceae, Chlorophyta), a novel coccoid

green alga carrying significant biotechnological potential. *Fottea* 14: 53–62.

Somogyi B., Felföldi T., Solymosi K., Böddi B., Marialigeti K and Vörös L (2011) *Chloroparva*

pannonica gen. et sp. nov. (Trebouxiophyceae, Chlorophyta) – a new picoplanktonic green

alga from a turbid, shallow soda pan. *Phycologia* 50: 1–10.

Somogyi B., Felföldi T., Solymosi K., Flieger K., Márialigeti K., Böddi B and Vörös L (2013)

One step closer to eliminating the nomenclatural problems of minute coccoid green algae:

Pseudochloris wilhelmii, gen. et sp. nov. (Trebouxiophyceae, Chlorophyta). *European*

Journal of Phycology 48: 427–436

Song H., Liu X., Hu Y., Wang Q., Long J., Liu G and Hu Z (2018) *Coronacoccus hengyangensis*

gen. et sp. nov., a new member of Chlorellaceae (Trebouxiophyceae, Chlorophyta) with

radiococcacean morphology. *Phycologia* 57: 363–373

Takeda, R (1991) Sugar composition of the cell wall and the taxonomy of *Chlorella*

(Chlorophyceae). *Journal of Phycology* 27: 224-232

Wilhelm C., Eisenbeis G., Wild A and Zahn R (1982) *Nanochlorum eucayotum*: a very reduced

coccoid species of marine Chlorophyceae. *ZeitschriJt fur Natuiforschung* 37c: 107-114.

Yamamoto M., Fujishita M., Hirata A and Kawano S (2004) Regeneration and maturation of

daughter cell walls in the autospore-forming green alga *Chlorella vulgaris* (Chlorophyta,

Trebouxiophyceae). *Journal of Plant Research* 117: 257–264.