Phylogenetic diversity and activity screening of cultivable actinobacteria isolated

from marine sponges and associated environments from the western coast of

India

Ulfat Baig[1], Neelesh Dahanukar[1], Neha Shintre[2], Ketki Holkar[1], Anagha Pund[1], Uttara Lele[1],

Tejal Gujarathi[1], Kajal Patel[1], Avantika Jakati[1], Ruby Singh[1], Harshada Vidwans[1], Vaijayanti

Tamhane[3], Neelima Deshpande[2] and Milind Watve[4]

[1]Indian Institute of Science Education and Research, Pune (IISER-P), Dr. Homi Bhabha

Road, Pashan, Pune 411008, Maharashtra, India.

[2]Department of Microbiology, M.E.S. Abasaheb Garware College, Karve Road, Pune

411004, Maharashtra, India.

[3]Institute of Bioinformatics and Biotechnology, Savitribai Phule Pune University, Pune –

411007, Maharashtra, India

[4]Behavioural Intervention for Lifestyle Disorders (BILD) clinic, Deenanath Mangeshkar

Hospital and Research Centre, Erandwane, Pune 411004, Maharashtra, India.

Correspondence and requests for materials should be addressed to N. Dahanukar (email:

n.dahanukar@iiserpune.ac.in)

Keywords: Bacterial predation; Antibiotic production; Secondary metabolites; Enzyme

inhibition; Molecular phylogeny.

Abstract

Phylogenetic diversity of cultivable actinobacteria isolated from sponges (*Haliclona* spp.) and associated environments of intertidal zones, along the northern parts of west coast of India, were studied using 16S rRNA gene sequences. A subset of actinobacteria were screened for three activities, namely predatory behavior, antibacterial activity and enzyme inhibition. We recovered 237 isolates of actinobacteria belonging to 19 families and 28 genera, which could be attributed to 95 putative species using maximum likelihood partition and 100 putative species using Bayesian partition in Poisson Tree Processes. Although the trends in the discovery of actinobacterial genera isolated from sponges was consistent with previous studies from different study areas, we provide first report of nine actinobacterial species from sponges. We observed widespread non-obligate epibiotic predatory behavior in eight actinobacterial genera and we provide first report of predatory activity in *Brevibacterium*, *Glutamicibacter*, *Micromonospora*, *Nocardiopsis*, *Rhodococcus* and *Rothia*. Sponge associated actinobacteria showed significantly more predatory behavior than environmental isolates. While antibacterial activity by actinobacterial isolates mainly affected Gram-positive target bacteria with little to no effect on Gram-negative bacteria, predation targeted both Gram-positive and Gram-negative prey with equal propensity. Actinobacterial isolates from both sponge and associated environment produced inhibitors of serine proteases and angiotensin converting enzyme. Predatory behavior was strongly associated with inhibition of trypsin and chymotrypsin. Our study suggests that sponge and associated environment of western coast of India are rich in actinobacterial diversity with widespread predatory activity, antibacterial activity and production of enzyme inhibitors. Understanding diversity and associations among various actinobacterial activities, with each other and the source of isolation, can provide new insights in marine microbial ecology and provide opportunities to isolate novel therapeutic agents.

INTRODUCTION

The marine ecosystem is not only diverse with respect to microorganisms found in it but also the natural products being synthesized by these microorganisms (Ward and Bora, 2006; Taylor et al., 2007; Lam, 2006). Actinobacteria are among the taxa rich in secondary metabolites (Barka et al., 2016) and are widely distributed in diverse habitats including soil, marine and freshwaters and sediments (Ward and Bora, 2006; Taylor et al., 2007; Tan et al., 2015; Brasel et al., 2019; Mincer et al., 2002; Kokare et al., 2004). They are also not uncommon in extreme environments (Jose and Jebakumar, 2014; Pathom-Aree et al., 2006; Mohammadipanah and Wink, 2016; Shivlata and Tulasi, 2015; Riquelme et al., 2015; Yang et al., 2015) and are also found as endobiotic symbionts of higher organisms (Taylor et al., 2007; Li et al., 2015; Mahmoud and Kalendar, 2016; Trujillo et al., 2015). They belong to the phylum Actinobacteria and represent one of the major phyla within the bacterial domain (Goodfellow, 2015). They are aerobic, spore forming, Gram-positive bacteria, which often produce diffusible pigments, and occur as cocci or rods, branched filaments, aerial or substrate mycelium (Goodfellow, 2015). The marine ecosystems are believed to have a wide range of unexplored diversity of actinobacteria (Montalvo et al., 2005) and their metabolites (Taylor et al., 2007; Lam, 2006; Manivasagan et al., 2005) with diverse biological activities like anticancer (Olano et al., 2009), anti-inflammatory (Trischman et al., 1994), antibiotic (Pimentel-Elardo et al., 2010; Cheng et al., 2015; Gandhimathi et al., 2008), cytotoxic (Abdelfattah et al., 2016) and enzyme inhibitory (Manivasagan et al., 2015; Imada, 2005) activity. Watve et al. (2001) estimated that the genus *Streptomyces* alone is capable of producing up to 10^5 different metabolites, majority of which remain unexplored. Of 23,000 medicinally important metabolites produced by marine microorganisms 70% are contributed by actinobacteria (Mahapatra et al., in press). Till date, eight genera of actinobacteria have

been reported to produce secondary metabolites and 267 products have been reported from 96

marine actinobacteria (Subramani and Sipkema, 2019)[31].

Ecologically it is difficult to understand the production of extracellular metabolites or

enzymes by aquatic bacteria, since any molecule secreted outside the cell can be quickly

washed off. Extracellular products could be useful to the producer only in viscous or partially

enclosed environments. In the marine environment, sponges are likely to provide such closed

environment for bacteria. Sponges are filter feeders and collect small nutrient particles

including bacteria. This makes the environment locally nutrient rich in an otherwise

oligotrophic surroundings. Bacteria, especially actinobacteria, isolated from these sponges

may live in a symbiotic relationship that helps the host in defense against predation, sponge

skeleton stabilization, translocation of metabolites and help in nutritional process (Taylor et

al., 2007; Li et al., 2015; Montalvo et al., 2005; Pimentel-Elardo et al., 2010; Cheng et al.,

2015; Gandhimathi et al., 2008; Lee et al., 2009; Thomas et al., 2010). In addition, since

sponges are sessile and lack other anti-predator defenses, secondary metabolites of bacteria

can provide them with chemical defense (Lee et al., 2001). Therefore, we expect more

secondary metabolite related activities from sponge-associated actinobacteria.

Sponge-associated actinobacteria are likely to have another ecological role. Among

actinobacteria at least three genera, namely *Agromyces*, *Streptomyces* and *Streptoverticillium*,

are shown to be predators that kill and feed on other live bacterial cells (Casida, 1980; 1983;

Kumbhar et al., 2014). Kumbhar and Watve (2013) argued that antibiotic activity might have

evolved primarily as a weapon in predation. However, the expression of secondary

metabolites during predation may be independent of antibiotic expression in pure culture; the

latter is likely to have evolved for mutualism with higher animal or plant hosts (Harir et al.,

2018; Van der Meij et al., 2017). Further, for a niche of predation in association with sponge,

the predatory species needs to protect itself from the digestive enzymes of the sponge as well

as its own enzymes used for predation. Therefore, predatory actinobacteria are also expected

to have efficient inhibitors of lytic enzymes.

In this study, we prepared an inventory of cultivable actinobacteria from sponges and

associated environments of intertidal zones along the northern parts of west coast of India and

studied their molecular diversity based on 16S rRNA gene sequences. We screened a subset

of randomly selected cultures for predatory activity, antibiotic production and enzyme

inhibition and tested their associations with each other and with the isolation source to test

the hypotheses mentioned earlier.

MATERIALS AND METHODS

Sample collection

Small tissue samples (less than one gram) of marine sponges (*Haliclona* spp.) were collected

at the time of low tide along Maharashtra and Goa coast (18–15ºN and 73–74ºE) of India

during April 2014 to October 2018 without damaging the sponge or its associated

environment. Specimens were rinsed and flushed with sterile media to remove debris and

loosely attached microbes. Each sponge sample was collected in labeled polystyrene tubes

with lids containing sterile Poor Ravan Saline (Watve et al., 2000) and ZoBell Marine broth

(ZoBell, 1941). Sediment, water and air samples were collected from the same environment

as that of the sponge and were collectively considered as environmental samples. The

samples were brought to laboratory maintaining cold chain and were immediately processed

for microbial culturing.

Isolation and maintenance of cultivable actinobacteria

Each sample was subjected to pre-heat treatment at 60°C for 15 minutes to eliminate non-

sporulating bacteria. Sponge tissue (0.1 cm^3) was homogenized in sterile medium and

vortexed for 5 minutes. Tubes were left undisturbed for two minutes. From the resulting supernatant serial 10 fold dilutions upto 10^{-5} were made and 0.1 ml sample was spread into triplicates on petri plates containing sterile medium. We used two media the Zobell Marine Agar (ZMA) and Poor Ravan Saline Agar (PRSA) with and without antibiotic chloramphenicol (25 µg/ml). Plates were incubated at 30°C for 7 days in the case of ZMA and 21 days for PRSA. Plates were observed regularly for the growth of actinobacteria. Bacterial colonies that showed resemblance to actinobacteria under light microscope were purified several times on the respective media. In all 237 actinobacterial isolates were selected and were re-streaked for making pure cultures. Colonies were labeled as per Maharashtra Gene Bank (MGB) project code and preserved on ZMA slants at 4°C for further use. Similarly, glycerol (18%) stocks were prepared and maintained at -20°C for long term storage. Actinobacterial cultures are deposited in the Microbial Culture Collection (MCC) of National Centre for Microbial Resource, National Center for Cell Sciences, Pune, India (accession numbers are provided in the Supplementary Table S1).

Genetic identification, phylogeny and species delimitation

Actinobacterial isolates were outsourced for near complete 16S rRNA gene sequencing. Gene sequences used for the study are deposited in the GenBank database under the accession numbers MN339687–MN339897 and MT598037–MT598065 (Supplementary Table S1)). Sequences were checked in BLAST (Altschul et al., 1990) to find the closest sequences available in the GenBank database (http://www.ncbi.nlm.nih.gov). Four species of Firmicutes, namely *Bacillus paralicheniformis* (MCC 6306), *B. thuringiensis* (MCC 7835), *B. subtilis* (MCC 6386) and *B. halotolerans* (MCC 8381), were used as outgroups (GenBank accession numbers MN339894–MN339897 respectively).

Gene sequences were aligned using MUSCLE (Edgar, 2004) implemented in MEGA

7 (Kumar et al., 2016). Final aligned matrix had 1595 sites. Best nucleotide substitution

model was determined using ModelFinder (Kalyaanamoorthy et al., 2017) based on Bayesian

information criterion (Schwarz, 1978; Nei and Kumar, 2000). Maximum likelihood analysis

was performed in IQ-TREE (Nguyen et al., 2015) with ultrafast bootstrap support (Hoang et

al., 2018) for 1000 iterations. Phylogenetic tree was edited in FigTree v1.4.2 (Rambaut,

2009).

To understand putative number of actinobacterial species we performed species

delimitation based on Poisson Tree Processes (Zhang et al., 2013) with maximum likelihood

partitioning (mPTP) and Bayesian partitioning (bPTP). Maximum likelihood tree was used to

delimit species by setting the parameter values as follows: MCMC generations = 100,000,

Thinning = 100, Burn-in = 0.1 and seed = 123.

We have identified all isolates up to genus level, while operational taxonomic units, in

terms of putative species, are provided based on mPTP and bPTP methods (see

Supplementary Table S1). Only in the text, some isolates are assigned to known species

based on BLAST search and sequence identity more than 99%.

Screening for activities

Out of 237 actinobacterial isolates, 50 isolates were randomly selected for screening of three

activities, namely predation, antibiotic production and production of enzyme inhibition.

Target bacteria used for predation and antibiotic screening

Test bacteria, used for checking actinobacterial predation and antibiotic production, were

obtained from MCC and National Collection of Industrial Microorganisms (NCIM), National

Chemistry Laboratory, Pune, India. Fourteen bacteria, namely *Acetobacter pasterianus*

(NCIM 2317), *Alcaligenes fecalis* (NCIM 2262), *Bacillus subtilis* (NCIM 2063),

Enterobacter fecalis, *Escherichia coli* (NCIM 2184), *Klebsiella pneumonae* (NCIM 2957),

Micrococcus luteus (NCIM 2673), *Mycobacterium smegmatis* (NCIM 5138), *Proteus*

vulgaris (NCIM 2172), *Pseudominas aeruginosa* (NCIM 5029), *Salinicoccus roseus* (MCC

7574), *Salmonella enterica* (NCIM 2501), *Serretia marcescens* (NCIM 2919) and

Staphylococcus aureus (NCIM 2121), were used as target species for screening.

Screening for actinobacterial predatory growth

Growth of predator with the zone of clearance on prey cells was considered as predation as

defined earlier (Kumbhar et al., 2014). The method for the preparation of prey cells was

modified from Kumbhar et al. (2014). Pure cultures of the prey species were inoculated on

nutrient agar plates to check the purity and were later re-inoculated in nutrient broth.

Inoculated flasks were incubated at 37°C for 24 h. Broth was centrifuged at 7000 rpm for 10

minutes to concentrate cells using Eppendorf centrifuge 5810R. Cells were washed thrice

with sterile distilled water to remove traces of nutrient broth. Pellet was suspended in saline

to obtain a thick suspension of optical density of 1.0 at 600 nm. Lawn of prey cells was

spread on water agarose plate and plates were incubated at 37°C for 40 minutes.

Actinobacterial culture was spot inoculated on pre incubated plates. These plates were

incubated at room temperature for 48–72 h at 30°C. Plates with plaque were examined

visually and by using 4x and 45x magnification under light microscope. Prey and predator

control plates were used for comparison. Each experiment consisted of triplicate sets of

plates, as well as one predator control for testing growth of actinobacterial predator without

prey. In addition, there was a prey control to demonstrate viable and independent growth of

prey without predator. In either controls there was no zone of clearance indicating there was

no predation in the presence of predator or prey alone.

Screening for antibacterial activity using conventional cross streak method

Selected actinobacterial cultures were screened for antibacterial activity by cross streak method (Velho-Pereira and Kamat, 2011; Valli et al., 2012). Test organism was streaked as a straight line along the diagonal of the petri dish with sterile ZMA medium. The isolated pure colony of actinobacteria was inoculated as a single streak perpendicular to the central streak. Streaking was done from the edge of the plate to the test organism growth line. Plates were incubated at 37°C for 18 h. The microbial inhibition was observed by determining zone of clearance around the sensitive organisms. Control plates of the same medium with the streak of test bacteria and without the streak of actinobacteria growth was used to observe the normal growth of the test bacteria.

Screening for enzyme inhibitors

Actinobacterial cultures were screened for their ability to inhibit the activity of serine proteases and angiotensin converting enzyme (ACE). Three different serine proteases i.e., Subtilisin, Trypsin and α-Chymotrypsin were used for screening of inhibitory activity. Protease inhibitor activity was studied using unprocessed X-ray films and spot-test method (Cheung, 1991) with modifications. As described by Tripathi et al. (Tripathi et al., 2011), dilutions of pure enzyme were first spotted on gelatine coated films. Lowest dilution showing complete clearance (indicating complete digestion of gelatine) was chosen for further studies. Pure enzyme (100 µg/ml) was incubated with equal quantity of cell free supernatant of actinobacterial isolates for 10 minutes and transferred to untreated X-ray-Fuji Medical X-ray, HRU grade-films. The mixtures were allowed to react for 15 minutes at room temperature and results were recorded after washing the x-ray films under running water. Unprocessed X-ray films contain a layer of gelatine on their surface, which acts as a substrate for various

proteolytic enzymes. Degradation of gelatine gives a clear zone at the site of activity. Thus,

upon action of the proteases, clear zones were seen on unprocessed x-ray films, at the site of

inoculation, whereas, if the gelatine layer remains intact, no clearance is observed. No

clearance on the films indicated presence of protease inhibitors.

ACE acts on a specific substrate N-Hyppuryl-His-Leu (HHL) to liberate hippuric acid

and His-Leu. Liberated hippuric acid was detected spectrophotometrically. Upon reaction of

the enzyme with ACE inhibitors, the enzyme becomes inactive and this is measured in terms

of lower levels of hippuric acid released. Protocol suggested by Cushman and Cheung (1971)

was used with certain modifications and hippuric acid liberated was checked using method

suggested by Ng et al. (2008). Equal amount of ACE and cell free supernatants (10 µl each)

were allowed to react at 37°C. After 10 minutes 20 µl of HHL was added to the reaction

mixture and reaction was continued for 30 mins at 37°C. The reaction was stopped by

addition of 40 µl of 1 N HCl. Blank was prepared by addition of HCl before addition of the

substrate. Positive enzyme control was prepared by incubating enzyme with un-inoculated

broth. Liberated hippuric acid was extracted in 90 µl ethyl acetate by vigorous shaking. Ethyl

acetate layer was collected in a fresh vial and allowed to dry in water bath of 50°C. The

liberated hippuric acid was diluted in 150 µl distilled water and absorbance was checked at

228 nm. Zero was adjusted using distilled water. Test vials with more than 15% inhibition of

ACE were considered as positive for ACE inhibitor.

RESULTS

Actinobacterial phylogenetic diversity in sponge and associated environment

Actinobacteria from sponges and associated environments showed a rich phylogenetic

diversity (Figure 1). We obtained 237 actinobacterial isolates, from sponge and associated

environments, belonging to 19 families and 28 genera (Supplementary Table S1). Species

delimitation based on mPTP suggested that these isolates belong to 95 putative species, while

bPTP suggested 100 putative species. The two species delimitation methods, mPTP and

bPTP, differed in the groups of species under genera *Micrococcus*, *Rhodococcus* and

Streptomyces (Supplementary Table S1). Air was generally devoid of actinobacteria and we

recovered only three isolates from air, belonging to genera *Brachybacterium*, *Brevibacterium*

and *Rhodococcus*, as compared to 39 isolates from water, 105 isolates from sediment and 90

isolates from sponge.

256   From sponges, 18 genera under 14 families belonging to 56 putative species (Table

257   1). From the sponge-associated environment, 22 genera under 15 families were recorded

258   belonging to 64 putative species as per mPTP and 65 putative species as per bPTP. A total of

259   12 genera under 9 families and 28 putative species based on mPTP and 25 putative species

260   based on bPTP were common to both sponge and associated environment.

261   Six genera, namely *Gordonia*, *Jonesia*, *Mycolicibacterium*, *Pseudonocardia*, *Rothia*

and *Serinicoccus* were isolated only from sponges (Table 1), which could be identified to

species *Gordonia terrae* (MCC 6452), *Jonesia denitrificans* (MCC 7852), *Mycolicibacterium*

poriferae (MCC 6242), *Pseudonocardia kongjuensis* (MCC 7930) and *Rothia terrae* (MCC

7823), *Serinicoccus marinus* (MCC 7935) respectively. Although 12 genera, namely

Agrococcus, *Arthrobacter*, *Brachybacterium*, *Brevibacterium*, *Klenkia*, *Kocuria*,

Microbacterium, *Micrococcus*, *Micromonospora*, *Nocardiopsis*, *Rhodococcus* and

Streptomyces, were isolated from both sponges and associated habitats, most of these genera

had some putative species that were either exclusive to sponges or associated environments.

In particular, 7 species, *Brachybacterium muris* (MCC 7614), *Brevibacterium casei* (MCC

6140, MCC 6152, MCC 6176), *Kocuria rhizophila* (MCC 8384), *Nocardiopsis salina* (MCC

7931), *Rhodococcus zopfii* (MCC 7934), *Streptomyces smyrnaeus* (MCC 7924) and

Streptomyces viridobrunneus (MCC 7990), were recorded only from sponges.

With respect to both, the number of isolates and number of putative species,

Streptomyces was the most dominant genus, which was found in both sponges and associated

environments. *Nocardiopsis* was the second most common genus with two dominant species

Nocardiopsis alba (MCC 8385) followed by *N. dassonvillei* (MCC 7845). Among the genera

and species that were recorded only from the environment, we provide first record of species

such as *Aeromicrobium massiliense* (MCC 6739) and *Glutamicibacter mysorens* (MCC 7825)

from marine waters.

Non-obligate epibiotic predatory activity

Out of the total 50 actinobacterial isolates screened for non-obligate epibiotic predatory

activity, 26 isolates showed predation on at least one of the 14 target organisms

(Supplementary Table S2). Of the 26 isolates with predatory behavior, 17 preyed on Gram-

negative prey, 21 preyed on Gram-positive prey, while 12 preyed on both Gram- negative and

Gram-positive prey. There was no significant difference (Mann-Whitney U = 15, P = 0.2601)

in the frequency of actinobacterial predators on Gram-negative and Gram-positive prey

(Table 2). Most actinobacterial predators (n = 14) preyed on a single prey species while only

a few predators preyed on multiple prey species. A single predator of the genus *Streptomyces*

tpreyed on 8 prey species. There was a significant association between the source of isolation

(sponge or associated environment) and predatory behavior (χ^2 = 5265, P = 0.0218), where

the isolates from sponge showed proportionately more predatory behavior (Figure 2).

All eight isolates of *Streptomyces* used for screening showed predatory behavior and

preyed on both Gram-negative and Gram-positive prey (Supplementary Table S2). Out of 25

isolates of *Nocardiopsis*, 12 showed predatory behavior, out of which 5 preyed on Gram-

negative bacteria while 11 preyed on Gram-positive bacteria. Both the isolates of

Micromonospora preyed on Gram-positive prey while only one preyed on Gram-negative

prey. Isolates belonging to genera *Brevibacterium*, *Glutamicibacter* and *Rhodococcus* preyed

only on Gram-negative prey while *Rothia* preyed only on Gram-positive prey.

Antibiosis, antibacterial activity and growth inhibition

Of the 50 actinobacterial isolates screened for antibacterial activity, 25 showed antibiosis

against at least one target organism (Supplementary Table S2). Of these 25 isolates, all

showed antibiosis against at least one of the Gram-positive target species, while only five

showed antibiosis against at least one of the Gram-negative organisms. The frequency of

antibacterial activity against Gram-positive organisms was significantly higher (Mann-

Whitney U = 1.5, P = 0.003) than those against Gram-negative organisms (Table 2). Most

antibacterial activities were broad spectrum with respect to the target organisms that they

affected. There were 10 actinobacterial isolates that showed antibiosis against two target

organisms, 6 isolates that affected 4 target species and 2 isolates that affected 6 target species.

There was no association between antibacterial activity and the source (sponge or associated

environment) of the isolation ($\chi^2 = 2.0129$, P = 0.1560).

Out of eight isolates of *Streptomyces* that were screened for antibacterial activity, five

showed antibiosis, of which two showed antibiosis against Gram-negative target species,

while all showed antibiosis against Gram-positive organisms. In the case of *Nocardiopsis*, of

the 25 isolates used for screening 17 showed antibiosis, of which all affected growth of

Gram-positive organisms, while only two affected growth of Gram-negative organisms.

Genus *Kytococcus* showed antibiosis that affected both Gram-positive as well as Gram-

negative organisms, while *Glutamicibacter* and *Rothia* showed antibiosis against Gram-

positive organisms only.

Enzyme inhibition

324 Out of 50 actinobacterial isolates screened for inhibition of four enzymes, 30 isolates

325 inhibited at least one of the enzyme (Supplementary Table S2). Of these 30 isolates, 28

326 inhibited trypsin, 24 inhibited chymotrypsin, three inhibited angiotensin converting enzyme

327 (ACE) and only two inhibited subtilisin. Venn diagram of frequency of isolates inhibiting

328 different enzymes (Figure 3) suggested that five isolates inhibited only trypsin and one isolate

329 each inhibited chymotrypsin and ACE, while subtilisin inhibition was accompanied by

330 inhibition of other enzymes. No isolate inhibited all four enzymes. Out of 30 actinobacteria

331 that produced enzyme inhibitors, 19 produced two inhibitors, four produced three inhibitors

332 while seven produced only one of the four inhibitors. There was no association between the

333 enzyme inhibition and source of the actinobacterial isolate (χ^2 = 2.3386, P = 0.1262).

334  Out of eight isolates of *Streptomyces* seven produced enzyme inhibitors against

335 proteases, while 12 out of 25 isolates of *Nocardiopsis* produced enzyme inhibitors of which

336 11 produced against proteases and two produced against ACE (Table 3). One isolate of

337 *Actinomycetospora* inhibited activity of ACE.

339 **Associations between different activities**

340 Out of 50 actinobacterial isolates that were screened for activities, 39 showed at least one of

341 the three activities. Of these 39 isolates, 15 showed all three activities, while nine showed

342 predation as well as enzyme inhibition (Figure 4). There were only seven isolates that showed

343 predation and antibiotic production against the same target organism (Table 2) and all these

344 isolates belonged to genera *Streptomyces* and *Nocardiopsis*.

345  Antibiotic production showed no significant association with predation (χ^2 = 2.8846,

346 P = 0.0894) or any of the four enzyme inhibition (χ^2 = 2.0525, P = 0.1520). However, there

347 were significant associations between predation and protease inhibitors (Figure 5). There

348 were 24 isolates that showed both predation as well as inhibition of at least one enzyme and

there was a significant association between the two activities ($\chi^2 = 26.172$, P < 0.0001),

where predators proportionately produced more enzyme inhibitors than non-predators (Figure

5a). There were 23 actinobacterial isolates that showed predation as well as trypsin inhibition

and there was a significant association between the two ($\chi^2 = 23.165$, P < 0.0001) with

predators more likely to produce trypsin inhibitors than non-predators (Figure 5b). Similarly,

24 actinobacteria were predators as well as inhibited chymotrypsin activity and there was a

significant association between the two ($\chi^2 = 42.604$, P < 0.0001) with predators more likely

to produce chymotrypsin inhibitors than non-predators (Figure 5c).

DISCUSSION

Sponges and associated environment in northern parts of western coast of India are rich in

actinobacterial diversity with about 95 putative species under 19 families and 28 genera. We

recorded 13 species of actinobacteria only from sponges. Out of these, *Mycobacterium*

poriferae was originally described from marine sponge (Padgitt and Moshier, 1987), while

three species, *Gordonia terrae* (Elfalah et al., 2013; Santos et al., 2019; Montalvo et al.,

2005), *Brevibacterium casei* (Kiran et al., 2010) and *Koc uria rhizophila* (Palomo et al.,

2013), have been previously reported from sponges. To our knowledge, we provide first

report of nine species, namely *Brachybacterium murisi, Jonesia denitrificans, Nocardiopsis*

salina, *Pseudonocardia kongjuensis*, *Rhodococcus zopfii*, *Rothia terrae*, *Serinicoccus*

marinus, *Streptomyces smyrnaeus* and *Streptomyces viridobrunneus*, from marine sponges,

although some of them are known from marine habitats (Stach et al., 2003; Satheeja and

Jebakumar, 2011; Yi et al., 2004; Shinde et al., 2018).

Streptomyces was the most dominant genus among the isolates, which agrees with the

findings of Zhang et al. (2008). Genus *Nocardiposis*, with its two species *N. alba* and *N.*

dassonvillei, has been suggested (Bennur et al., 2015) as the second common genus after

Streptomyces and that too agrees with our findings. Further, report of most genera, including

Agrococcus, *Arthrobacter*, *Brevibacterium*, *Kocuria*, *Microbacterium* and *Micrococcus*, from

sponges in our study are consistent with previous reports from other study areas including

South China Sea (Li et al., 2015), Yellow Sea (Zhang et al., 2008), Mediterranean Sea

(Cheng et al., 2015), coast of Florida in USA (Montalvo, 2005) and northern coast of Brazil

(Menezes et al., 2010) indicating that there are common trends in the discovery of

actinobacteria from sponges.

Among the first reports from marine environment from our study, *Aeromicrobium*

massiliense and *Glutamicibacter mysorens* are known from human fecal microbiota

(Ramasamy et al., 2012) and sewage (Nand and Rao, 1972) respectively. Presence of these

two species in the sediments along the collection site Harne (17.81ºN, 73.09ºE) likely

suggests fecal pollution in this area.

Although predation is a widespread behavior in bacterial kingdom, δ-proteobacteria

of the orders *Myxococcales* and *Bdellovibrionales* have received more attention (Jurkevitch,

2007) as compared to other taxa, especially the Gram-positive bacteria such as actinobacteria.

Among actinobacteria only three genera, namely *Agromyces*, *Streptomyces* and

Streptoverticillium, are known to have predatory behavior against other bacterial species

(Casida, 1980; 1983; 1988; Kumbhar et al., 2014; Zeph and Casida, 1986; Ibrahimi et al.,

2020). In the current study, for the first time, we show predation in six other genera of

actinobacteria, namely *Brevibacterium*, *Glutamicibacter*, *Micromonospora*, *Nocardiopsis*,

Rhodococcus and *Rothia*. Kumbhar et al. (2014) argued that predatory behavior is widespread

in genus *Streptomyces* and even in the current study we observed that all the isolates of

Streptomyces used for screening showed predation on Gram-positive as well as Gram-

negative prey.

Since sponges are sessile and lack other anti-predator defenses, it has been suggested

that secondary metabolites of bacteria can provide sponges with chemical defense (Lee et al.,

2001; Kumbhar and Watve, 2013). However, we did not observe any significant association

between the source of actinobacterial isolation and antibiotic production, suggesting that

isolates even from environment were equally likely to produce antimicrobials as that of the

isolates recovered from sponges. However, there was a significant association between the

source of isolation and predatory activity, with proportionately more predators among the

isolates recovered from sponge. Ecologically this makes sense. As the sponges are filter

feeders and have regular intake of environmental bacteria, sponge associated actinobacteria

will have better predation opportunities. It is also possible that the predatory activity of

sponge associated actinobacteria, could have evolved as a mutualistic activity as it can defend

sponges from pathogenic bacterial invasions.

Actinobacteria are known to produce several enzyme inhibitors (Manivasagan et al.,

2015; Imada, 2005). However, for the first time we show a strong association between

predation and enzyme inhibition, specifically inhibition of trypsin and chymotrypsin, where

predators produced proportionality more enzyme inhibitors as compared to non predators.

Predators themselves are known to produce a variety of hydrolytic enzymes for degrading the

prey (Pérez et al., 2016). Therefore, it is possible that the production of enzyme inhibitors

safeguards their own cells from being target of the enzyme. It is also possible that enzyme

inhibitors also protect the actinobacteria from hydrolytic enzymes produced from the sponge

host and other microbiota.

An interesting observation that we made, when comparing the predation and antibiotic

production by actinobacteria, was that, while predation was equally effective against Gram-

positive as well as Gram-negative target species, antibiotic production was mainly effective

against Gram-positive bacteria. Recently, Ibrahimi et al. (2020) suggested that there are some

bio-active secondary metabolites that co-cultured actinobacteria produce in the presence of

prey cells. It is therefore possible that studying the predatory behavior of actinobacteria and

predation specific metabolites could lead to discovery of novel therapeutic agents that are

more broad-spectrum.

Although actinobacteria are known to be rich in secondary metabolites, extracellular

enzymes and enzyme inhibitors, the ecological role of these extracellular bioactive molecules

is little known. We suggest that studying the ecological correlates of bioactivity and the inter-

correlation patterns of different types of bioactivity can be a useful tool in understanding the

ecological origins of bioactivity and testing alternative ecological hypotheses.

CONCLUSION

Sponges and associated environments of intertidal zones, along the northern parts of west

coast of India, are rich in actinobacterial diversity with 19 families and 28 genera, which

could be attributed to 95 putative species using mPTP and 100 putative species based on

bPTP methods. Although, at the genus level, the trends in the discovery of actinobacteria

isolated from sponges was consistent with previous studies from different study areas, we

provide first report of nine species, namely *Brachybacterium murisi, Jonesia denitrificans*,

Nocardiopsis salina, *Pseudonocardia kongjuensis*, *Rhodococcus zopfii*, *Rothia terrae*,

Serinicoccus marinus, *Streptomyces smyrnaeus* and *Streptomyces viridobrunneus*. Non-

obligate epibiotic predatory behavior was widespread among actinobacterial genera and we

provide first report of predatory activity in *Brevibacterium*, *Glutamicibacter*,

Micromonospora, *Nocardiopsis*, *Rhodococcus* and *Rothia*. Sponges associated actinobacteria

showed significantly more predatory behavior than environmental isolates, and we

hypothesize that predatory actinobacteria might provide sponges with defense against

pathogenic bacteria. While antibiotic produced from actinobacterial isolates affected Gram-

positive target bacteria with little to no effect on Gram-negative bacteria, predation targeted

both Gram-positive and Gram-negative prey with equal propensity, suggesting that study of

predation specific metabolites might provide novel therapeutic agents with broad-spectrum.

Actinobacterial isolates from both sponge and associated environment produced inhibitors of

serine proteases and angiotensin converting enzyme. Predatory behavior was strongly

associated with inhibition of trypsin and chymotrypsin, which might be helpful for the

actinobacteria for overcoming effects of proteolytic enzymes produced by sponge host and

other microbiota. Understanding diversity and associations among various actinobacterial

activities, with each other and the source of isolation, can provide new insights in marine

microbial ecology and provide opportunities to isolate novel therapeutic agents.

DATA AVAILABILITY

Sequences of 16S rRNA gene of studied isolates are submitted to GenBank NCBI under the accession numbers MN339687–MN339897 and MT598037–MT598065. Actinobacterial cultures are deposited in the Microbial Culture Collection (MCC) of National Centre for Microbial Resource, National Center for Cell Sciences, Pune, India (accession numbers are provided in the Supplementary Table S1). All the data used for analysis is provided in supplementary information (Supplementary Table S1 and Table S2).

ACKNOWLEDGEMENTS

This work was funded by Maharashtra Gene Bank Programme (RGSTC/File-2007/DPP-054/CR-28) of Rajiv Gandhi Science and Technology Commission, Government of Maharashtra, India. We thank the Director and Chair of Biology, Indian Institute of Science Education and Research (IISER), Pune, and Principle, M.E.S. Abasaheb Garware College, Pune for providing infrastructural facilities. We are thankful to V. S. Rao, IISER Pune, for support an encouragement.

AUTHOR CONTRIBUTIONS

M.W., U.B. and N. Deshpande conceived and designed the study. U.B., N.S., K.H., A.P., U.L., T.G., K.P., A.J., R.S., H.V and V.T. performed the study. N. Dahanukar and M.W. analyzed the data. N. Dahanukar, U.B. and M.W. wrote the manuscript with inputs from other authors. All authors contributed to the proofreading of the manuscript.

Supplementary information: Supplementary Table S1 and Table S2 accompanies the online version of the paper.

Competing Interests: The authors declare no competing interests.

References

Abdelfattah, M. S., Elmallah, M. I. Y., Hawas, U. W., El-Kassema, L. T. A., and Eid, M. A.

G. (2016). Isolation and characterization of marine-derived actinomycetes with cytotoxic

activity from the Red Sea coast. *Asian Pacific J. Tropical Biomed.* 6, 651–657.

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990). Basic local

alignment search tool. *J. Mol. Biol.* 215, 403–410.

Barka, E. A., Vatsa, P., Sanchez, L., Gaveau-Vaillant, N., Jacquard, C., Meier-Kolthoff, J. P.,

Klenk, H. P., Clément, C., Ouhdouch, Y., and van Wezel, G. P. (2016). Taxonomy,

physiology, and natural products of Actinobacteria. *Microbiol. Mol. Biol. Rev.* 80, 1–43.

Bennur, T., Kumar, A. R., Zinjarde, S., and Javdekar, V. (2015). *Nocardiopsis* species:

Incidence, ecological roles and adaptations. *Microbiol. Res.* 174, 33–47.

Braesel, J., Lee, J. H., Arnould, B., Murphy, B. T., and Eustáquio, A. S. (2019).

Diazaquinomycin biosynthetic gene clusters from marine and freshwater actinomycetes.

J. Nat. Prod. 82, 937–946.

Casida, L. E. Jr. (1980). Bacterial predators of *Micrococcus luteus* in soil. *Appl. Environ.*

Microbiol. 39, 1035–1041.

Casida, L. E. Jr. (1983). Interaction of *Agromyces ramosus* with other bacteria in soil. *Appl.*

Environ. Microbiol. 46, 881–888.

Casida, L. E. (1988). Minireview: Nonobligate bacterial predation of bacteria in soil. *Microb.*

Ecol. 15, 1–8.

Cheng, C., MacIntyre, L., Abdelmohsen, U. R., Horn,H., Polymenakou, P. N., Edrada-Ebel,

R., and Hentschel, U. (2015). Biodiversity, anti-trypanosomal activity screening, and

metabolomic profiling of actinomycetes isolated from Mediterranean sponges. *PLoS ONE*

10, e0138528. doi: 10.1371/journal.pone.0138528.

Cheung, A. L., Ying, P., and Fischetti, V. A. (1991). A method to detect proteinase activity

using unprocessed X-ray films. *Anal. Biochem.* 193, 20–23.

Cushman, D. W., and Cheung, H. S. (1971). Spectrophotometric assay and properties of the

angiotensin-converting enzyme of rabbit lung. *Biochem. Pharmacol.* 20, 1637–1648.

Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acids Res.* 32, 1792–1797.

Elfalah, H. W., Usup, G., and Ahmad, A. (2013). Anti-microbial properties of secondary

metabolites of marine *Gordonia tearrae* extract. *J. Agric. Sci.* 5, 94–101.

Gandhimathi, R. Arunkumar, M., Selvin, J., Thangavelu, T., Sivaramakrishnan, S., Kiran, G.

S., Shanmughapriya, S., and Natarajaseenivasana, K. (2008). Antimicrobial potential of

sponge associated marine actinomycetes. *J. Mycol. Med.* 18, 16–22.

Goodfellow, M. (2015) "Actinobacteria phyl. nov.", in *Bergey's Manual of Systematics of*

Archaea and Bacteria, eds Whitman, W. B. et al., (New York: Wiley), 1–2. doi:

10.1002/9781118960608.pbm00002.

Harir, M., Bendif, H., Bellahcene, M., Fortas, Z. & Pogni, R. (2018). "*Streptomyces*

secondary metabolites", in *Basic Biology and Applications of Actinobacteria*, ed. Enany,

S. (London, IntechOpen), 99–122.

Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q., and Vinh, L.S. (2018).

UFBoot2: Improving the ultrafast bootstrap approximation. *Mol. Biol. Evol.* 35, 518–522.

Ibrahimi, M., Korichi, W., Hafidi, M., Lemee, L., Ouhdouch, Y., and Loqman, S. (2020).

Marine Actinobacteria: Screening for Predation Leads to the Discovery of Potential New

Drugs against Multidrug-Resistant Bacteria. *Antibiotics* 9(2), 91.

Imada, C. (2005). Enzyme inhibitors and other bioactive compounds from marine

actinomycetes. *Antonie Van Leeuwenhoek* 87, 59–63.

Jose, P. A., and Jebakumar, S. R. D. (2014). Unexplored hypersaline habitats are sources of

novel actinomycetes. *Front. Microbiol.* 5, 242. doi: 10.3389/fmicb.2014.00242.

Jurkevitch, E. (2007). Predatory behaviors in bacteria-diversity and transitions. *Microbe* 2,

67–73.

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A., and Jermiin, L.S.

(2017). ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nat.*

Methods 14, 587–589.

Kiran, G. S., Sabarathnam, B., and Selvin, J. (2010). Biofilm disruption potential of a

glycolipid biosurfactant from marine *Brevibacterium casei. FEMS Immunol. Med.*

Microbiol. 59, 432–438.

Kokare, C. R., Mahadik, K. R., and Kadam, S. S. (2004). Isolation of bioactive marine

actinomycetes from sediments isolated from Goa and Maharashtra coastlines (west coast

of India). *Indian J. Mar. Sci.* 33, 248–256.

Kumar, S., Stecher, G., and Tamura, K. (2016). MEGA7: molecular evolutionary genetics

analysis version 7.0 for bigger datasets. *Mol. Biol. Evol.* 33, 1870–1874.

Kumbhar, C., and Watve, M. (2013). Why antibiotics: A comparative evaluation of different

hypotheses for the natural role of antibiotics and an evolutionary synthesis. *Nat. Sci.* 5,

26–40.

Kumbhar, C., Mudliar, P., Bhatia, L., Kshirsagar, A., and Watve, M. (2014). Widespread

predatory abilities in the genus *Streptomyces. Arch. Microbiol.* 196, 235–248.

Lam, K. S. (2006). Discovery of novel metabolites from marine actinomycetes. *Curr. Opin.*

Microbiol. 9, 245–251.

Lee, O. O., Wong, Y. H., and Qian, P. Y. (2009). Inter- and intraspecific variations of

bacterial communities associated with marine sponges from San Juan Island, Washington.

Appl. Environ. Microbiol. 75, 3513–3521.

Lee, Y. K., Lee, J. H., and Lee, H. K. (2001). Microbial symbiosis in marine sponges. *J.*

Microbiol. 39, 254–264.

Li, Z., Sun, W., Zhang, F., He, L., and Loganathan, K. (2015). Actinomycetes from the South

China Sea sponges: isolation, diversity and potential for aromatic polyketides discovery.

Front. Microbiol. 6, 1048. doi: 10.3389/fmicb.2015.01048.

Mahapatra, G. P., Raman, S., Nayak, S., Gouda, S., Das, G., and Patra, J. K. (in press).

Metagenomics approaches in discovery and development of new bioactive compounds

from marine actinomycetes. *Curr. Microbiol.* doi: 10.1007/s00284-019-01698-5.

Mahmoud, H. M., and Kalendar, A. A. (2016) Coral-associated Actinobacteria: diversity,

abundance, and biotechnological potentials. *Front. Microbiol.* 7, 204. doi:

10.3389/fmicb.2016.00204.

Manivasagan, P., Kang, K. H., Sivakumar, K., Li-Chan, E. C., Oh, H. M., and Kim, S. K.

(2014). Marine actinobacteria: an important source of bioactive natural products. *Environ.*

Toxicol. Pharmacol. 38, 172–188.

Manivasagan, P., Venkatesan, J., Sivakumar, K., and Kim, S. K. (2015). Actinobacterial

enzyme inhibitors–A review. *Crit. Rev. Microbiol.* 41, 261–272.

Menezes, C. B., Bonugli-Santos, R. C., Miqueletto, P. B., Passarini, M. R., Silva, C. H.,

Justo, M. R., Leal, R. R., Fantinatti-Garboggini, F., Oliveira, V. M., Berlinck, R. G., and

Sette, L. D. (2010). Microbial diversity associated with algae, ascidians and sponges from

the north coast of São Paulo state, Brazil. *Microbiol. Res.* 165, 466–482.

Mincer, T. J., Jensen, P. R., Kauffman, C. A., and Fenical, W. (2002). Widespread and

persistent populations of a major new marine actinomycete taxon in ocean sediments.

Appl. Environ. Microbiol. 68, 5005–5011.

Mohammadipanah, F., and Wink, J. (2016). Actinobacteria from arid and desert habitats:

diversity and biological activity. *Front. Microbiol.* 6, 1541. doi:

10.3389/fmicb.2015.01541.

Montalvo, N. F., Mohamed, N. M., Enticknap, J. J., and Hill, R. T. (2005). Novel

actinobacteria from marine sponges. *Antonie Van Leeuwenhoek* 87, 29–36.

Nand, K., and Rao, D. V. (1972). *Arthrobacter mysorens*–a new species excreting L-glutamic

acid. *Zentralbl Bakteriol Parasitenkd Infektionskr Hyg.* 127, 324–331.

Nei, M., and Kumar, S. (2000). *Molecular evolution and phylogenetics*. UK, Oxford

University Press.

Ng, K.H., Lye, H.S., Easa, A.M., and Liong, M.T. (2008). Growth characteristics and

bioactivity of probiotics in tofu-based medium during storage. *Ann. Microbiol.* 58, 477–

487.

Nguyen, L. T., Schmidt, H. A., von Haeseler, A., and Minh, B. Q. (2015). IQ-TREE: a fast

and e□ective stochastic algorithm for estimating maximum likelihood phylogenies. *Mol.*

Biol. Evol. 32, 268–274.

Olano, C., Méndez, C., and Salas, J. (2009). Antitumor compounds from marine

actinomycetes. *Marine Drugs* 7, 210–248.

Padgitt, P. J., and Moshier, S. E. (1987). *Mycobacterium poriferae* sp. nov., a

scotochromogenic, rapidly growing species isolated from a marine sponge. *Int. J. Syst.*

Evol. Microbiol. 37, 186–191.

Palomo, S., González, I, de la Cruz, M., Martín, J., Tormo, J. R., Anderson, M., Hill, R. T.,

Vicente, F., Reyes, F., and Genilloud1, O. (2013). Sponge-derived *Kocuria* and

Micrococcus spp. as sources of the new thiazolyl peptide antibiotic kocurin. *Marine*

Drugs 11, 1071–1086.

Pathom-Aree, W., Stach, J. E., Ward, A. C., Horikoshi, K., Bull, A. T., and Goodfellow, M.

(2006). Diversity of actinomycetes isolated from Challenger Deep sediment (10,898 m)

from the Mariana Trench. *Extremophiles* 10, 181–189.

Pérez, J., Moraleda-Muñoz, A., Marcos-Torres, F. J., and Muñoz-Dorado, J. (2016). Bacterial

predation: 75 years and counting! *Environ. Microbiol.* 18, 766–779.

Pimentel-Elardo, S. M., Kozytska, S., Bugni, T. S., Ireland, C. M., Moll, H., and Hentschel,

U. (2010). Anti-parasitic compounds from *Streptomyces* sp. strains isolated from

Mediterranean sponges. *Marine Drugs* 8, 373–380.

Ramasamy, D., Kokcha, S., Lagier, J.-C., Nguyen, T.-T., Raoult, D., and Fournier, P.-E.

(2012). Genome sequence and description of *Aeromicrobium massiliense* sp. nov. *Stand.*

Genomic Sci. 7, 246–257.

Rambaut, A. (2009). FigTree. ver 1.4.3 http://tree.bio.ed.ac.uk/software/%ef%ac%81gtree/.

Riquelme, C., Hathaway, J. J. M., Enes Dapkevicius, M. de L. N., Miller, A. Z., Kooser, A.,

Northup, D. E., Jurado, V., Fernandez, O., Saiz-Jimenez, C., and Cheeptham, N., (2015).

Actinobacterial diversity in volcanic caves and associated geomicrobiological

interactions. *Front. Microbiol.* 6, 1342. doi: 10.3389/fmicb.2015.01342.

Santos, J. D., Vitorino, I., De la Cruz,M., Díaz,C., Cautain, B., Annang, F., Pérez-Moreno,

G., Martinez, I. G., Tormo, J. R., Martín, J. M., Urbatzka, R., Vicente, F. M., and Lage,

O. M. (2019). Bioactivities and extract dereplication of Actinomycetales isolated from

marine sponges. *Front. Microbiol.* 10, 727. doi: 10.3389/fmicb.2019.00727.

Satheeja, S. V., and Jebakumar, S. R. (2011). Phylogenetic analysis and antimicrobial

activities of *Streptomyces* isolates from mangrove sediment. *J. Basic Microbiol.* 51, 71–

79.

Schwarz, G. (1978). Estimating the dimension of a model. *Ann. Stat.* 6, 461–464.

Shinde, V.L., Meena, R.M., and Shenoy, B.D., (2018). Phylogenetic characterization of

culturable bacteria and fungi associated with tarballs from Betul beach, Goa, India. *Mar.*

Pollut. Bull. 128, 593–600.

Shivlata, L., and Tulasi, S. (2015). Thermophilic and alkaliphilic Actinobacteria: biology and

potential applications. *Front. Microbiol.* 6, 1014. doi: 10.3389/fmicb.2015.01014.

Stach, J. E. M., Maldonado, L. A., Masson, D. G., Ward, A. C., Goodfellow, M., Bull, A. T.

(2003). Statistical approaches for estimating actinobacterial diversity in marine

sediments. *Appl. Environ. Microbiol.* 69, 6189–6200.

Subramani, R., and Sipkema, D. (2019). Marine rare actinomycetes: A promising source of

structurally diverse and unique novel natural products. *Marine Drugs* 17, 249.

Tan, L. T. H., Ser, H. L., Yin, W. F., Chan, K. G., Lee, L. H., and Goh, B. H. (2015).

Investigation of antioxidative and anticancer potentials of *Streptomyces* sp. MUM256

isolated from Malaysia mangrove soil. *Front. Microbiol.* 6, 1316. doi:

10.3389/fmicb.2015.01316.

Taylor, M. W., Radax, R., Steger, D., and Wagner, M. (2007). Sponge-associated

microorganisms: evolution, ecology, and biotechnological potential. *Microbiol. Mol. Biol.*

Rev. 71, 295–347.

Thomas, T. R., Kavlekar, D. P., and LokaBharathi, P. A. (2010). Marine drugs from sponge-

microbe association - a review. *Marine Drugs* 8, 1417–1468.

Tripathi, V. R., Kumar, S., and Garg, S. K. (2011). A study on trypsin, *Aspergillus flavus* and

Bacillus sp. protease inhibitory activity in *Cassia tora* (L.) syn *Senna tora* (L.) Roxb. seed

extract. *BMC Complement. Altern. Med.* 11, 56. doi: 10.1186/1472-6882-11-56.

Trischman, J. A., Tapiolas, D. M., Jensen, P. R., Dwight, R., Fenical, W., McKee, T. C.,

Ireland, C. M., Stout, T. J., and Clardy, J. (1994). Salinamides A and B: anti-

inflammatory depsipeptides from a marine streptomycete. *J. American Chem. Soc.* 116,

757–758.

Trujillo, M. E., Riesco, R., Benito, P., and Carro, L. (2015). Endophytic actinobacteria and

the interaction of *Micromonospora* and nitrogen fixing plants. *Front. Microbiol.* 6, 1341.

doi: 10.3389/fmicb.2015.01341.

Valli, S., Suvathi, S. S., Aysha, O. S., Nirmala, P., Vinoth, K. P., and Reena, A. (2012).

Antimicrobial potential of Actinomycetes species isolated from marine environment.

Asian Pacific J. Tropical Biomed. 2, 469–473.

Van der Meij, A., Worsley, S. F., Hutchings, M. I., and van Wezel, G. P. (2017). Chemical

ecology of antibiotic production by actinomycetes. *FEMS Microbiol. Rev.* 41, 392–416.

Velho-Pereira, S., and Kamat, N.M. (2011). Antimicrobial screening of actinobacteria using a

modified cross-streak method. *Indian J. Pharm. Sci.* 73, 223–228.

Ward, A. C., and Bora, N. (2006). Diversity and biogeography of marine actinobacteria.

Curr. Opin. Microbiol. 9, 279–286.

Watve, M., Shejval, V., Sonawane, C., Rahalkar, M., Matapurkar, A., Shouche, Y., Patole,

M., Phadnis, N., Champhenkar, A., Damle, K., Karandikar, S., Kshirsagar, V., and Jog,

M. (2000). The 'K' selected oligophilic bacteria: A key to uncultured diversity? *Curr. Sci.*

78, 1535–1542.

Watve, M. G., Tickoo, R., Jog, M. M., and Bhole, B. D. (2001) How many antibiotics are

produced by the genus Streptomyces? *Arch. Microbiol.* 176, 386–390.

Yang, J., Li, X., Huang, L., and Jiang, H. (2015). Actinobacterial diversity in the sediments

of five cold springs on the qinghai-tibet plateau. *Front. Microbiol.* 6, 1345. doi:

10.3389/fmicb.2015.01345.

Yi, H., Schumann, P., Sohn, K. and Chun, J. (2004). Serinicoccus marinus gen. nov., sp. nov.,

a novel actinomycete with L-ornithine and L-serine in the peptidoglycan. *Int. J. Syst.*

Evol. Microbiol. 54(5), 1585–1589.

Zeph, L. R., and Casida, L. E. (1986). Gram-negative versus gram-positive (actinomycete)

nonobligate bacterial predators of bacteria in soil. *Appl. Environ. Microbiol.* 52, 819–823.

Zhang, H., Zhang, W., Jin, Y., Jin, M., and Yu, X. (2008). A comparative study on the

phylogenetic diversity of culturable actinobacteria isolated from five marine sponge

species. *Antonie van leeuwenhoek* 93, 241–248.

Zhang, J., Kapli, P., Pavlidis, P., and Stamatakis, A. (2013). A general species delimitation

method with applications to phylogenetic placements. *Bioinformatics* 29, 2869–2876.

ZoBell, C. E. (1941). Studies on marine bacteria. I. The cultural requirements of

heterotrophic aerobes. *J. Mar. Res.* 4, 41–75.

Table 1. Putative number of species of actinobacterial genera based on PTP and bPTP

methods isolated from sponge, associate environment and both sources.

Family	Genus	Sponge		Environment		Both	
		mPTP	bPTP	mPTP	bPTP	mPTP	bPTP
Actinomycetaceae	*Streptomyces*	23	23	24	25	12	11
Brevibacteriaceae	*Brevibacterium*	5	5	2	2	1	1
Cellulomonadaceae	*Cellulomonas*	0	0	1	1	0	0
Dermabacteraceae	*Brachybacterium*	1	1	2	2	0	0
Dietziaceae	*Dietzia*	0	0	1	1	0	0
Geodermatophilaceae	*Klenkia*	1	1	1	1	1	1
Gordoniaceae	*Gordonia*	1	1	0	0	0	0
Intrasporangiaceae	*Janibacter*	0	0	2	2	0	0
	Knoellia	0	0	1	1	0	0
	Terrabacter	0	0	1	1	0	0
Jonesiaceae	*Jonesia*	1	1	0	0	0	0
Kytococcaceae	*Kytococcus*	0	0	1	1	0	0
Microbacteriaceae	*Agrococcus*	1	1	1	1	1	0
	Curtobacterium	0	0	1	1	0	0
	Microbacterium	1	1	1	1	1	1
Micrococcaceae	*Arthrobacter*	1	1	1	1	1	1
	Glutamicibacter	0	0	2	2	0	0
	Kocuria	4	4	6	6	2	2
	Micrococcus	6	6	8	8	4	4
	Rothia	1	1	0	0	0	0
Micromonosporaceae	*Micromonospora*	1	1	1	1	1	1
Mycobacteriaceae	*Mycolicibacterium*	1	1	0	0	0	0
Nocardiaceae	*Rhodococcus*	2	2	2	2	2	1
Nocardioidaceae	*Aeromicrobium*	0	0	1	1	0	0
Nocardiopsaceae	*Nocardiopsis*	4	4	3	3	2	2
Ornithinimicrobiaceae	*Serinicoccus*	1	1	0	0	0	0
Pseudonocardiaceae	*Actinomycetospora*	0	0	1	1	0	0
	Pseudonocardia	1	1	0	0	0	0
	Total	56	56	64	65	28	25

694 **Table 2.** Predation and antibiotic production by actinobacteria against the Gram positive and

695 Gram negative target species.

Target species	Predation	Antibiotic	Predation and Antibiotic by same actinobacterial isolate
Gram positive			
Mycobacterium smegmatis	3	12	0
Micrococcus luteus	8	5	0
Bacillus subtilis	1	24	1
Staphylococcus aureus	17	9	4
Salinicoccus roseus	9	3	0
Enterococcus faecalis	3	20	1
Gram negative			
Acetobacter pasterianus	7	0	0
Alcaligenes faecalis	3	1	1
Escherichia coli	2	5	0
Klebsiella pneumoniae	3	0	0
Proteus vulgaris	8	0	0
Salmonella enterica	2	0	0
Serratia marcescens	3	0	0
Pseudomonas aeruginosa	1	0	0

699   **Table 3.** Frequency of actinobacterial isolates producing four different enzyme inhibitors.

Genus	Number of isolates	Frequency of isolates inhibiting				Isolates with at least one inhibition activity
		Subtilisin	Trypsin	Chymotrypsin	ACE	
Actinomycetospora	2	0	1	0	1	2
Agrococcus	1	0	0	0	0	0
Brevibacterium	1	0	1	1	0	1
Glutamicibacter	1	0	1	1	0	1
Jonesia	1	0	0	0	0	0
Kocuria	1	0	0	0	0	0
Kytococcus	1	0	1	0	0	1
Micrococcus	1	0	1	0	0	1
Micromonospora	2	0	2	2	0	2
Nocardiopsis	25	0	11	11	2	12
Pseudonocardia	1	0	0	0	0	0
Rhodococcus	4	0	2	1	0	2
Rothia	1	0	1	1	0	1
Streptomyces	8	2	7	7	0	7

FIGURE CAPTIONS

Figure 1. Maximum likelihood phylogenetic tree of actinobacterial isolates based on

TIM3+F+I+G4 nucleotide substitution model (lnL of consensus tree: -18684.58). Firmicutes

belonging to genus *Bacillus* were used as outgroups.

Figure 2. Association between source of actinobacterial isolation on their predatory behavior.

There was a significant association between the source (sponge or associated environment) of

actinobacterial isolation and predation ($\chi^2 = 5.265$, P = 0.0218).

Figure 3. Venn diagrams combination of enzyme inhibitors produced by actinobacterial

isolates. Venn diagrams is not to scale.

Figure 4. Venn diagrams of predation, antibiotic production and enzyme inhibition by

actinobacterial isolates. Venn diagrams is not to scale.

Figure 5. Association between enzyme inhibition and predation in actinobacterial isolates.

Predation was significantly associated with (a) inhibition of any one of the four enzymes

tested ($\chi^2 = 26.172$, P < 0.0001), (b) inhibition of trypsin ($\chi^2 = 23.165$, P < 0.0001) and (c)

inhibition of chymotrypsin ($\chi^2 = 42.604$, P < 0.0001).