Maternally derived sex steroid hormones impact sex ratios of loggerhead sea turtles

Authors: Emma C. Lockley[1*], Thomas Reischig[2], Christophe Eizaguirre[1]

[1]Queen Mary University of London, School of Biological and Chemical Sciences, Mile End

Road, London E14NS, United Kingdom.

[2]Turtle Foundation, An der Eiche 7a, 50678 Cologne, Germany.

Corresponding Author: Emma Lockley, Queen Mary University of London, School of

Biological and Chemical Sciences, Mile End Road, London E14NS, United Kingdom.

emma.lockley@qmul.ac.uk

Classification: Biological Sciences

Sub-Classification: Evolution

Keywords:

Temperature-dependent sex determination, Maternal effects, Loggerhead sea turtle, Hormone

transfer, Conservation, Climate Change

<u>Abstract</u>

Global warming could drive species with temperature-dependent sex determination to

extinction by persistently skewing offspring sex ratios. Evolved mechanisms that buffer these

biases are therefore paramount for their persistence. Here, we tested whether maternally-

derived sex steroid hormones affect the sex-determination cascade and provide a

physiological mechanism to buffer sex ratio bias in the endangered loggerhead sea turtle

(*Caretta caretta*). We quantified estradiol and testosterone in nesting females and their egg

yolks at oviposition, before incubating nests in situ at standardised temperatures. Upon

hatchling emergence, we developed a new, non-lethal method to establish the sex of

individuals. Despite standardised incubation temperatures, sex ratios varied widely among

nests, correlating non-linearly with the estradiol:testosterone ratio in egg yolks. Males were

produced at an equal ratio, with females produced either side of this optimum. This result

provides evidence that maternal hormone transfer forms a physiological mechanism that

impacts sex determination in this endangered species.

Introduction

Fifty years after the discovery of environmental sex determination, our understanding of its evolutionary significance, underlying mechanisms and ecological consequences in the light of environmental change remains incomplete[1–5]. Most reptile and some fish species undergo temperature-dependent sex determination (TSD), in which gonad differentiation is regulated by temperature at a critical period of embryogenesis[6,7]. Some species produce males at moderate temperatures and females at hot and cold extremes (e.g. the American alligator *Alligator mississippiensis*[8], Type II TSD), but, more commonly, TSD species produce an increasing proportion of a specific sex across a range of incubation temperatures (Type Ia: Males at low temperatures, e.g. the painted turtle *Chrysemys picta*[9]; Type Ib: Females at low temperatures, e.g. the tuatara *Sphenodon punctatus*[10]). In all cases, both sexes are produced across a transitional range of temperatures, centered on a pivotal temperature at which both sexes develop in equal proportions. While the pivotal temperature varies among clutches within a population[11], studies have generally focused on quantifying population-level means as important proxies to estimate sex ratios and predict population dynamics[4,12,13]. These population-level proxies suggest that rising global temperatures present the potential for extreme sex ratio biases in TSD species, with implications for population dynamics and extinction risk[4].

The adaptive value of TSD is still debated, but fitness advantages under sex-specific thermal environments are predicted by the more favoured Charnov-Bull theory of differential fitness[14,15]. The postulates of this theory have been demonstrated in eggs of the Jacky dragon (*Amphibolurus muricatus*) that were experimentally treated with an aromatase inhibitor, constraining embryos to develop as males at female producing temperatures. These males showed lower lifetime reproductive success than controls[3]. While demonstrating the adaptive value of TSD, the use of an aromatase inhibitor to manipulate sex in this study also highlights the role of sex steroid hormones on the TSD mechanism[16–18].

Exogenous application of estradiol (E_2) to the incubating eggs of some species can feminise TSD embryos incubated at male-producing temperatures[19–22]. In addition, the application of

testosterone (T), the precursor androgen of E_2, can also feminise embryos via the synthesis of

E_2 from T by the aromatase enzyme [20,23], and indeed the use of aromatase inhibitors can force

male development[24,25]. Both temperature and exogenous treatment with E_2 activate the same

molecular pathways, altering the transcription of the chromatin modifier gene *Kdm6b*, and

conferring sensitivity to a sex-determining gene, *Dmrt1*[2].

In model TSD species that exhibit Type 1a TSD, such as the slider (*Trachemys scripta*) and

the painted turtle, maternal transfer of sex steroid hormones into eggs varies seasonally[16,26].

Elevated concentrations of maternal investment in yolk E_2 and greater E_2:T ratios increase the

likelihood of feminisation at a given temperature, effectively reducing the pivotal temperature

of a clutch by providing substrate that will prime the activation of female-producing

molecular pathways[16,26]. Should these patterns be found in non-model species, variation in

maternal hormone transfer to eggs could be a universal mechanism to (i) change the threshold

at which temperature affects an individual's sex development, (ii) modify the sex ratio of the

clutch, and (iii) possibly buffer against the negative effects of rapid global temperature

increase.

There is a particular need to understand the impacts of climate change on the demographics of

threatened species. As a consequence of rising temperatures, extreme feminisation of sea

turtle populations has been forecast by the end of the century[4,27–29]. Some studies suggest

effects are already visible in adult populations[28]. There has been much study into how

behavioural responses, such as modified phenology or nest site selection, may mitigate the

effects of a warming environment[30,31], but no overall trends preventing extirpation are visible.

There has been, however, little consideration for physiological mechanisms that may increase

variation in the pivotal temperature, and therefore on sex ratio. Understanding possible

physiological mechanisms has been constrained in sea turtles in particular by the lack of non-

lethal methods to sex neonates (but see[32]). This issue is especially important for endangered

populations, where sacrificing individuals is not possible.

Here, we tested whether maternally-derived sex steroid hormones affect the sex-

determination cascade and the resulting offspring sex ratios in an endangered sea turtle

population. Focusing on loggerhead turtles (*Caretta caretta*) nesting in the Cabo Verde

archipelago, we standardised the thermal environment of clutches in an experimental field

hatchery, exposed to natural conditions. Should temperature be the sole driver of sex

determination, similar sex ratios among clutches would be expected under these standardised

thermal conditions. Alternatively, any inter-clutch variation would arise from intrinsic

characteristics of the eggs, such as maternally-derived hormones. To test these hypotheses,

we quantified E_2 and T concentrations in the plasma of nesting females, their egg yolks and

neonates. We developed a non-lethal sexing method using circulatory hormone profiles of

neonates, and determined the clutches' sex ratios. Inter-clutch variation in sex ratio was then

linked to yolk hormone concentrations. Finally, we illustrate how maternal hormone transfer

impacts sex ratio in the face of IPCC climate change predictions, by re-parameterising a

previously used mathematical model[4] to forecast the future population dynamics of this

endangered nesting aggregation.

Results

This study focused on loggerhead turtles nesting on the island of Boavista in the Cape Verde

archipelago. First, using enzyme-linked immunosorbent assays (ELISA – Enzo LifeSciences),

we quantified concentrations of the sex steroid hormones E_2 and T in both the blood plasma

of 26 nesting females and up to two of their eggs directly after oviposition. Clutch sizes were

recorded at this time. High levels of individual variation were observed in adult plasma

hormone levels (SI appendix, Table S1), with a mean T concentration of 1148.48 ± 148.63

(SE) pg/ml, a mean E_2 concentration of 235.79 ± 22.71 (SE) pg/ml, and a mean E_2:T ratio of

0.32 ± 0.05 (SE). Linear models (LM) showed positive correlations between E_2 and T in both

the female plasma (SI Appendix, Fig. S1A, $F_{1,16} = 4.608$, $p = 0.048$) and their egg yolks (SI

Appendix, Fig. S1B, $F_{1,23} = 7.338$, $p = 0.013$). In reptiles, maternally derived hormones are

constant across all eggs of a given clutch[33], which we confirmed with a subset of clutches

where two egg yolks were analysed (Paired t-tests: T: df = 11, t = 0.224, p = 0.827; E_2: df =

10, t = -0.885, p = 0.397; E_2:T: df = 9, t = -1.173, p = 0.271).

There was a significant non-linear correlation between T concentrations in adult plasma and

egg yolks (SI appendix, Fig. S2A: LM: $F_{1,14}$ = 5.263, p = 0.038), where concentrations of yolk

T were lowest in eggs originating from females with intermediate levels of plasma T, but did

not correlate with clutch size (SI appendix, Fig. S2B: $F_{1,14}$ = 0.032, p = 0.862). In contrast,

adult female plasma E_2 concentrations were not correlated with E_2 in the egg yolk (SI

appendix, Fig. S2C: LM: $F_{1,21}$ = 0.908, p = 0.351), but as clutch size increased, yolk E_2

concentrations significantly decreased (SI Appendix, Fig. S2D: LM: $F_{1,21}$ = 4.945, p = 0.037).

The maternal E_2:T ratio showed a non-linear relationship with the E_2:T ratio in the egg yolk

(SI Appendix, Fig. S2E: $F_{1,14}$ = 6.493, p = 0.023), and was not correlated with clutch size (SI

Appendix, Fig. S2F: $F_{1,14}$ = 1.682, p = 0.215).

Immediately after oviposition, the clutches of these 26 females and two others (n = 28) were

relocated into an *in-situ* experimental hatchery that was protected from terrestrial predation,

yet exposed to natural sand and weather conditions. We buried clutches at a depth of 55 cm to

standardise the thermal incubation environment. We confirmed the standardised thermal

environment using data loggers placed at the centre of the clutch (mean thermosensitive

period temperature = 30.02 ± 0.05 (SE) °C, SI Appendix, Fig. S3). The small amount of

temperature variation observed was explained by differences in clutch size ($F_{1,26}$ = 4.418, p =

0.045), resulting from increased metabolic heat produced from more developing embryos in

larger clutches[34]. Assuming the pivotal temperature of this population to be 29 °C, as has

previously been used for this population[4], this incubation temperature would produce 12.89 ±

0.01 (SE) % male offspring if temperature was the sole determinant of sex ratio (Fig. 1A).

Incubation duration, the time between oviposition and neonate emergence, is also often used

as a proxy to predict offspring sex ratios (e.g. r^2 = 0.73 in nests in Brazil[35]) and was recorded

for each clutch[35]. Using the established logistic relationship between incubation duration and

sex ratio observed in loggerhead turtles from Kyparissia, Greece (Fig. 1B, the closest location

where the relationship between incubation duration and offspring sex ratio has been

quantified for loggerhead turtles), the predicted sex ratio of our study clutches would be 47.5

± 6 (SE) % males[36]. This suggests that levels of sex ratio variation are far greater than those

we would expect from temperature alone.

While the incubation duration represents a reasonable proxy for estimating the sex ratio of sea

turtle offspring, currently the only accurate method to resolve individual sex requires

sacrificing neonates and histological examination - a limiting factor for endangered

populations[35,37]. However, we developed a new method to ascertain individual sex without the

need to sacrifice animals. After taking $100 - 150$ μl of blood from 365 offspring from 28

clutches after emergence (mean offspring per clutch = 13 ± 4 (SE)), we measured plasma

hormone concentrations using ELISA. Hatchling hormone levels varied among individuals

(SI Appendix, Table S1) and among clutches, with the average E_2:T ratio of clutches ranging

from 1.06 ± 0.13 (SE) to 3.56 ± 0.68 (SE). We used affinity propagation clustering (APC) on

hatchling E_2:T ratios guided by incubation duration to identify clusters of individuals with a

similar hormonal phenotype. APC iteratively considers the similarity of a data point to its

neighbours. Importantly, it does not require the number of possible clusters to be defined *a*

priori, as is necessary for other clustering approaches such as k-means[38]. We identified three

APC clusters (Fig. 2A). Two of these originate from clutches with short incubation durations,

the classic trait of female neonates, and were distinguished by differences in their mean E_2:T

ratio (SI Appendix, Fig. S4, Cluster 1: mean = 4.45 ± 0.26 (SE), Cluster 2: mean = 1.72 ±

0.05 (SE), t-test: df = 44.08, t = 10.273, p < 0.001). The third group is formed by individuals

from clutches with longer incubation durations (t-test: df = 299.3, t = -32.933, p < 0.001) and

a low E_2:T ratio (SI Appendix, Fig. S4, mean = 1.52 ± 0.06 (SE)), the characteristics of male

sea turtle neonates.

Several positive theoretical controls were used to confirm this method since neonate

sacrificing is not possible. First, linear mixed effect models (LMM) using clutch ID as a

random factor revealed significant differences in hormone levels between the two sexes, that

were directly comparable to previous studies in which individuals' sex was confirmed

through histology[39,40]. As expected, T levels were higher in males (Fig. 2Bi, LMM: $F_{1,60}$ =

10.673, p = 0.002, mean = 63.63 ± 2.89 (SE) pg/ml) than in females (mean = 52.54 ± 2.34

(SE) pg/ml), and conversely E_2 levels were higher in females (Fig. 2Bii, LMM: $F_{1,57}$ = 7.521,

p = 0.008, mean = 92.94 ± 3.06 (SE) pg/ml) than in males (mean = 81.66 ± 3.16 (SE) pg/ml),

as was the overall E_2:T ratio (Fig. 2Biii, LMM: $F_{1,48}$ = 28.652, p < 0.001, females: mean =

2.22 ± 0.09 (SE); males: mean = 1.52 ± 0.06 (SE)). LMMs did not detect any difference in

weight ($F_{1, 348}$ = 0.024, p = 0.878) or size ($F_{1, 218}$ = 0.766, p = 0.382) between the sexes, as

would be expected under these conditions by the Charnov-Bull theory[14]. Second, by

combining individual offspring sex into an estimate of clutch sex ratio, and comparing this to

the incubation duration, we found the specific logistic regression curve that characterises

incubation durations in Type Ia TSD species (Fig. 2C). The pivotal duration was fitted to a

value of 57.25 days (95% CIs: 57.09, 57.43), with a transitional range of incubation durations

of 2.15 days (95% CIs: 1.52, 2.77). Importantly, if individual sex were incorrectly assigned,

this distinctive logistic regression curve of TSD species would not be seen. With this method,

we determined that clutch sex ratios were on average 40.49 ± 8.98 (SE) % male. This

suggests 26.1% more males and far more variation in clutch sex ratio than would be expected

based on incubation temperatures alone. Our sex ratio estimate is slightly below (7.1%) that

estimated from parameters based on incubation durations in Kyparissia, suggesting

population differences in development rate exist, likely as a result of different average pivotal

temperatures among rookeries.

After establishing that the inter-clutch variation in sex ratio (and also in incubation duration,

see SI Appendix Supplementary Analysis) was too great to be produced by temperature alone,

we tested whether metabolic heat and/or maternal hormone transfer in the yolk predicted

incubation duration and the estimated sex ratio. Yolk T correlated negatively with both

incubation duration (LM, $F_{1,22}$ = 10.624, p = 0.003) and the proportion of males produced

within a clutch (Fig. 3A, Binomial generalised linear mixed effect models (GLMM), x^2 =

4.371, df = 1, p = 0.037), but metabolic heat had no detectable effect (incubation duration

model: $F_{1,22}$ = 2.436, p = 0.133, sex ratio model: x^2 = 2.111, df = 1, p = 0.146). There was no

relationship between yolk E_2 and incubation duration or clutch sex ratio (Fig. 3B, incubation

duration: $F_{1,23}$ = 3.169, p = 0.088, sex ratio: x^2 = 0.183, df = 1, p = 0.669), yet the yolk E_2:T

ratio showed a non-linear relationship with both incubation duration (Fig. 3C, $F_{1,21} = 12.882$,

p = 0.002) and sex ratio independently of temperature ($x^2 = 39.319$, df = 2, p < 0.001). A

maximum incubation duration of 57.2 days was observed at an equal hormone ratio (E_2:T of

1.05, $y = -7.8x^2 + 16.3x + 48.7$) with the highest levels of male offspring developing at this

point. Because of the presence of a possible outlier with a high yolk E_2:T ratio, we re-

analysed the data without this point. The same patterns remained with significant non-linear

relationships between yolk E_2:T ratios and incubation duration ($F_{1,20} = 6.292$, p = 0.021) as

well as between yolk E_2:T ratios and hatchling sex ratios ($x^2 = 7.584$, df = 2, p = 0.022).

Male offspring production was highest when maternal investment of E_2 and T to the yolk was

equal. Asking whether the production of either sex is more costly in terms of total hormone

investment, we compared the total hormone concentration (E_2 + T) with the overall E_2:T ratio.

This relationship was again non-linear, with total hormone investment being highest when the

E_2:T ratio was unequal (SI Appendix, Fig. S5A, LM: $F_{2,22} = 4.951$, p = 0.017), suggesting that

producing females requires more maternal investment than males. The total hormone

investment also showed a non-linear relationship with clutch size (SI Appendix, Fig S5B,

$\log(E_2 + T)$: $F_{2,22} = 4.306$, p = 0.026), with an initial increase in investment across clutch sizes

between 65 and 75 eggs, after which investment declined with increasing clutch size.

Finally, to illustrate how maternal hormone transfer could impact population dynamics, we

re-parameterised a previously published mathematical projection of neonate sex ratios for the

Cape Verde population[4], which assumed a fixed pivotal temperature of 29 °C. We made the

simple assumption that the effect of maternally derived hormones on sex ratio is constant

across a thermal gradient and applied the 26.1% observed difference in male offspring

production for the coming century (Fig. 4). With a mechanism of this possible strength, the

population is unlikely to reach the levels of extreme feminisation previously forecasted –

instead of female production reaching over 97% in 2100, it is likely to instead reach

approximately 71%. As it remains to be determined how maternal hormone transfer interacts

with different incubation temperatures, this model only illustrates the potential importance of

trans-generational hormone transfer for population dynamics.

Discussion

It is widely speculated that global warming can drive species with temperature-dependent sex

determination to extinction because of the over-production of one sex. But, given the many

considerable historical shifts in climate experienced by TSD species, they are likely to have

evolved behavioural and/or physiological mechanisms to avoid unviable biases in offspring

sex ratio[5]. By experimentally standardising the thermal environment of loggerhead sea turtle

nests *in-situ*, we investigated whether maternally derived hormones correlate with offspring

sex independently of temperature. First, we developed a non-lethal method to determine the

sex of neonates upon their nest emergence, using affinity propagation clustering based on

individual circulatory sex steroid hormones and their incubation duration. With this method,

we found a non-linear relationship between the clutch sex ratios and the ratios of maternally

derived E_2:T within the egg yolks under standardised thermal conditions. Low concentrations

of equal investment in both hormones within the yolk maximise the production of male

offspring, while increasing the concentration of either E_2 or T, along with overall hormone

investment, feminises the clutches. Re-parameterising an existing model that predicts sex

ratio biases in response to climate change demonstrated that this trans-generational

mechanism could prevent the predicted extreme feminisation of loggerhead turtles in Cabo

Verde.

To date, an inability to determine neonate sex non-lethally has constrained the study of TSD

mechanisms in endangered sea turtles (but see[32]). A clustering approach that identifies

individuals with similar phenotypes (here hormone profiles) that match control traits of male

and female offspring (incubation duration) overcame this problem. Using E_2:T thresholds to

define neonate sex has been verified with histological analysis in loggerhead[40] and green[39]

turtles, but as E_2:T levels vary considerably among clutches, it is difficult to delineate a

population level threshold *a priori*. Using an APC method guided by incubation duration to

group hormone profiles, a common proxy for sex ratio, we avoid the need to define thresholds

and, importantly, the need to sacrifice individuals[35]. Because ethically we cannot validate this

method further, we rely on strong indirect evidence, such as (i) the identified significant

difference in circulating E_2:T ratios of male and female offspring (ii) the pivotal duration both

match those reported in other studies[36,39–42], and (iii) the relationship between sex ratio and

incubation duration fits the known logistic regression curve observed in Type Ia TSD species.

We anticipate that this non-lethal approach will prove an invaluable tool for both research on

TSD in sea turtle species and also for wider conservation.

Despite the standardised thermal environment of clutches within this experiment, high levels

of variation in incubation duration and sex ratio were observed among nests, both of which

correlated with maternally derived hormones within the egg yolk. The relationship between

the yolk E_2:T ratio and clutch sex ratio was best described by a quadratic curve, centred on an

equal concentration of both hormones and ranging from 0.37 to 1.73. When maternal

investment of E_2 and T was equal, incubation durations were long, and males were produced.

If hormone investment was biased in either direction, sex ratios became increasingly

feminised. The effects of elevated levels of both E_2 and T on sex ratios in this study are

consistent with experimental manipulation of these hormones in other species[2,20,22]. In

addition, the non-linear relationship explains studies where exogenous application of E_2 has

unexpectedly produced male hatchlings (e.g.[43,44]). In such cases, E_2 application, combined

with existing maternal contributions, may have resulted in shifting the E_2:T ratio within the

eggs closer to one, forcing male development. Interestingly, in all other reptiles for which

data are available, the E_2:T ratios that are transferred to the yolk consistently remain below or

above the ratio of one[45]. Thus, our study shows how maternal transfer of both hormones can

influence the feminisation process of reptiles under natural conditions.

Total hormone concentrations within the yolk were lowest at an equal, male producing, E_2:T

ratio. If this ratio departed from 1:1 in either direction, total concentrations of yolk hormones

increased. As E_2 and T positively correlate within the egg, if investment in either hormone is

elevated, there is an associated increase in the other. The outcome is that greater investment is

required to skew E_2:T ratios in a manner that favours the production of female offspring. This

maternal investment provides initial hormonal substrate with which to prime the reactions

required to activate molecular pathways that result in female gonad development. When E_2:T

ratios are skewed, and total hormone concentrations are high, feminisation is easily achieved

through either the presence of E_2 directly, or by the synthesis of E_2 from its precursor, T, by

the aromatase enzyme. When E_2 and T are in equilibrium, low concentrations of E_2 are not

sufficient to feminise the clutch. However, product-feedback inhibition of aromatase

receptors likely prevents further E_2 being synthesised from T, and consequently male

offspring are produced.

There is no doubt that temperature is the primary determinant of sex in TSD species, but

studies have repeatedly recorded variation in sex ratios among clutches exposed to similar

temperature regimes[32,46]. Maternally-derived E_2 and T can affect the same developmental

pathways as temperature, and explain some of this variation by priming reactions required to

initiate female producing pathways[2,22]. Interestingly, the effects of these hormones on sex

determination vary by experiment and species, likely as the results of adaptation to local

nesting conditions (e.g.[43,44]). However, here we show that in sea turtles, a clutch specific

threshold exist for feminisation that is the product of an interaction between temperature and

maternal hormone transfer. A shift towards an equal E_2:T ratio and lower maternal investment

will increase the pivotal temperature away from the feminisation threshold, and consequently

warmer temperatures would be required to feminise a clutch. This aligns with the sex ratios

observed within this study, which contained 26.1% more males than expected from a pivotal

temperature of 29 °C. However, this mechanism will be constrained by physiological limits of

maternal hormone investment. Our findings provide mechanistic explanations for the high

inter-clutch variation that has been observed in TSD systems (eg[11,46]), and also clarify

occasions where female offspring have been produced at assumed male producing

temperatures, or vice versa[32].

Two maternal traits show a relationship with levels of hormone transfer to the clutch. Firstly,

T concentrations within the yolk correlated non-linearly with those in maternal plasma.

Disentangling the cause of such a relationship is complex as it is likely to result from multiple

physiological cascades[47]. As vitellogenesis and follicular development in sea turtles occurs

prior to migration, it is also likely that T concentrations in maternal plasma varies when yolks

is formed[48]. However, this relationship does allow us to link T investment to maternal state.

Should maternal T vary in response to environmental cues, as in the spined toad (*Bufo*

spinosus), it may allow nesting females to plastically match the individual development

threshold of feminisation to the ambient temperature, and maintain more constant sex ratios

across a nesting season[49]. Similar differences in the maternally-transferred E_2:T ratio in the

egg yolk of a population of painted turtles resulted in a seasonal shift of the pivotal

temperature, albeit in a direction that accelerated female production as temperatures

increased[16]. Secondly, total E_2 investment within eggs decreased as clutch size increased, and

total hormone concentrations were low in large clutches. Thus, in large clutches with more

metabolic heat production, the developmental threshold of feminisation is increased –

minimising sex bias. We infer from these results that there are two distinct mechanisms that

can affect the ratio of E_2:T within the yolk, which explains how elevated investment in either

hormone can lead to feminisation. There is considerable variation in circulating T and E_2

levels between sea turtle populations and species (SI Appendix, Table S1), which may

suggest an element of local adaptation in response to environmental conditions, and a

heritable component of baseline physiological levels[50].

TSD species will require behavioural and/or physiological responses to maintain viable sex

ratios in the face of future climate change. Here, we highlight a previously under-considered

physiological mechanism for individual variation in the TSD process within sea turtle species.

There is a need for management plans that use temperature-based models to predict future sex

ratios to account for maternal hormonal influence, as this will have considerable implications

for population dynamics.

Methods

Sample Collection

We studied nesting loggerhead sea turtles on the island of Boavista, part of the Cabo Verde

archipelago in the eastern Atlantic. The sampling site (15°58'18.6"N, 22°48'06.2"W) is a 400

m stretch of coastline on the southern tip of this island. Twenty-eight nesting females were

sampled between 17 July and 1 August 2017. Immediately after oviposition, females were

individually marked with PIT (AVID) and metal (Inconel) tags[51]. Blood samples of 1-4 ml in

volume were collected from the dorsal cervical sinus of 26 females using a 40 mm, 21-gauge

needle and 5 ml syringe, and stored within lithium heparin containers. Finally, curved

carapace length (CCL) and width (CCW) were measured (± 0.1 cm).

The clutches of these turtles (containing 83 ± 3 (SE) eggs) were relocated to an experimental

hatchery protected from predation, situated on the nesting beach. At this point, up to two eggs

from the 28 clutches were removed from each clutch for yolk hormone analysis, and the rest

of the clutch was buried at a depth of 55 cm. By using a standard depth, temperature was

controlled for, while maintaining an otherwise natural environment. A TinyTag™

temperature logger was placed at the centre of each clutch, programmed to take a reading

every 15 minutes throughout the incubation period (accuracy ± 0.2 °C). As anticipated, the

uniform depth standardised the incubation temperature of the nests to 30.05 ± 0.05 (SE) °C

during the middle third of incubation, the period where embryo sex is established. This

variation in temperature is extremely conserved, and is representative of the thermal variation

produced within treatments under controlled laboratory incubations[52,53].

Upon emergence, twenty hatchlings were randomly selected for blood sampling (100 – 150

µl) from the dorsal cervical sinus, using a 26-gauge needle and 1 ml syringe[54]. Samples were

stored within lithium heparin coated tubes. Notch-to-notch straight carapace length (SCL)

and, width (SCW) were measured using digital callipers (± 0.01 mm), and weight was

measured with a digital scale (± 0.1 g)).

The blood samples of both the adults and offspring were refrigerated for up to 48 h before

being centrifuged to extract plasma. Egg yolks were separated from the albumen, and all

samples were stored at -20 °C until extraction.

Hormone extraction

Commercially available Enzyme-Linked Immunosorbent Assay (ELISA) kits for both E_2

(Catalogue # ADI-900-174, ENZO Life Sciences) and T (Catalogue # ADI-900-065) were

used to measure steroid levels in all samples. Details for hormone extraction protocols are

given in SI methods. Not all blood samples had sufficient volume for hormone extraction.

Consequently, we extracted E_2 from 24 adults and 388 hatchling blood samples, and T from

19 adult and 367 hatchling blood samples. This provided us with E_2:T ratios for 18 adult

females, and 365 hatchlings. E_2, and T were successfully extracted from the yolks of 26 out of

the 28 sampled clutches. One yolk T measurement was removed as an outlier, being more

than three standard deviations from the mean (mean: 741.26 ± 502.29 (SD) pg/g, outlier:

2882.05 pg/g).

Statistical Analyses

All analyses were conducted with R 3.3.3, using the R packages *lme4* and *lmerTest* for fitting

linear mixed models (LMMs) and generalized linear mixed models (GLMMs). A paired t-test

was used to compare intra-clutch E_2 and T levels between two eggs in a subset of clutches (n

= 13), to test whether that there was variation in hormone investment between eggs in a

clutch. As there was no difference between eggs from the same clutch, for subsequent

analyses the average hormone was used where possible, while a single egg was used for the

remainder of the clutches. Correlations between E_2 and T in female plasma and yolks, the

effect of clutch size on temperature, and the effects of metabolic heat and maternally derived

hormones on incubation duration were tested using general linear models (LM). A non-linear

relationship between the E_2:T ratio on incubation duration was fitted using a quadratic curve.

Similarly, when considering the relationship between E_2:T and total hormone investment, we

also fit a quadratic model. LMs were also used to estimate the correlation of clutch size and

plasma hormone concentrations with yolk hormone concentrations.

We used Algorithm Propagation Clustering (APC) to identify individual sex, using the R

package *apcluster*[38]. Cluster assignment was made based on the plasma E_2:T ratio of

hatchlings, guided by their incubation duration. Determining neonate sea turtle sex using the

E_2:T ratio has previously been extremely accurate (96% and 96.7% respectively) for

artificially incubated eggs of loggerhead and green sea turtles[39,40] that were ultimately

sacrificed for verification. Since variation likely exists among rookeries, those thresholds

however cannot be blindly applied to new populations. T-tests were used to compare hormone

levels of putative male and female hatchlings. A response curve of these estimated sex ratios

to incubation duration was produced using the logistic equation function of the R package

embryogrowth to further verify the accuracy of our non-lethal sexing method.

After identifying the sex of individuals, LMMs were used to compare individual size and

weight between the sexes and the APC clusters. Finally, we used binomial GLMMs to

determine whether individual hatchling sex was predicted by maternal hormone investment or

temperature. For all LMM and GLMM analyses, clutch was included as a random factor to

account for individual variation. Model selection was based on AIC criteria, using a

likelihood ratio tests to select for the best models. P-values of the selected models were

obtained by with the *car* R package, and models were verified for over-dispersion.

Thermal estimates of sex ratio were calculated using the equation first presented by Girondot

in 1999 with the R package *embryogrowth*, under an assumed pivotal temperature of 29 °C[55].

Estimates of sex ratio based on incubation duration were made based on data from a study on

a neighbouring loggerhead sea turtle population that nests in Kyparissia, Greece, and was

confirmed with histology[36]. To generate an illustrative model that compared the results of our

study with future predictions based on temperature alone, we extracted data from a previously

published study predicting sex ratios until 2100 based on a fixed pivotal temperature of 29 °C

alone. We then compared our observed mean clutch sex ratio to that expected from a pivotal

temperature of 29 °C, and added the difference, along with 95% confidence intervals, to the

original prediction.

Acknowledgements

This work was carried out as part of The Turtle Project, a citizen science project dedicated to

sea turtle research and conservation. We thank volunteers and staff from Turtle Foundation in

particular for facilitating this research. Particular thanks go to S. Cameron for considerable

help during data collection. E.L was supported by a studentship from the London Natural

Environment Research Council Doctoral Training Partnership (grant NE/L002485/1). This

study was supported by a National Geographic grant (NGS-59158R-19), Natural

Environment Research Council (NE/V001469/1) as well as Queen Mary University of

London Funds allocated to CE. We also acknowledge the support from "The Future Ocean'

Excellence Initiative by the Deutsche Forschungsgemeinschaft (DFG) that granted a

'Capacity building grant' to C.E. The authors would like to thank the Eizaguirre lab as well as

Dr. José María Martín-Durán and Dr. Gail Schoefield for their feedback on previous versions

of the manuscript.

Author Contributions

E.L. and C.E. designed the experiment. T.R. facilitated the fieldwork. E.L. and C.E.

conducted the fieldwork. E.L. analysed the data and drafted the initial manuscript, with

feedback from C.E.. All authors approved the manuscript.

References

1.   Charnier, M. Action of temperature on the sex ratio in the Agama agama (Agamidae,

Lacertilia) embryo. *C R Seances Soc Biol Fil.* **160**, 620–622 (1966).

2.   Ge, C. *et al.* The histone demethylase KDM6B regulates temperature-dependent sex

determination in a turtle species. *Science* **360**, 645–648 (2018).

3.   Warner, D. A. & Shine, R. The adaptive significance of temperature-dependent sex

determination in a reptile. *Nature* **451**, 2006–2009 (2008).

4.   Laloë, J.-O. *et al.* Climate change induced extinction does not seem imminent at an

important sea turtle rookery. *Nat. Clim. Chang.* **4**, 1–7 (2014).

5.   Mitchell, N. J. & Janzen, F. J. Temperature-dependent sex determination and

contemporary climate change. *Sex. Dev.* **4**, 129–140 (2010).

6.   Bull, J. J. Sex determination in reptiles. *Q. Rev. Biol.* **55**, 3–21 (1980).

7.   Deeming, D. C. *et al.* Environmental Regulation of Sex Determination in Reptiles.

Philos. Trans. R. Soc. B Biol. Sci. **322**, 19–39 (1988).

8.   Ferguson, M. W. J. & Joanen, T. Temperature-dependent sex determination in

Alligator mississippiensis. *J. Zool.* **200**, 143–177 (1983).

9.   Bull, J. J. & Vogt, R. C. Temperature-Dependent Sex Determination in Turtles.

Science **206**, 1186–1188 (1979).

10.   Cree, A., Thompson, M. B. & Daugherty, C. H. Tuatara sex determination. *Nature*

375, 543 (1995).

11.   Dodd, K. L., Murdock, C. & Wibbels, T. Interclutch variation in sex ratios produced at

pivotal temperature in the red-eared slider, a turtle with temperature-dependent sex

determination. *J. Herpetol.* **40**, 544–549 (2006).

12.   Godley, B. J., Broderick, A. C., Glen, F. & Hays, G. C. Temperature-dependent sex

determination of Ascension Island green turtles. *Mar. Ecol. Prog. Ser.* **226**, 115–124

(2002).

13.   Lolavar, A. & Wyneken, J. Experimental assessment of the effects of moisture on

loggerhead sea turtle hatchling sex ratios. *Zoology* **123**, 64–70 (2017).

14.   Charnov, E. . & Bull, J. . When is sex environmentally determined? *Nature* **266**, 828–

830 (1977).

15.   Shine, R. Why is sex determined by nest temperature in many reptiles? *Trends Ecol.*

Evol. **14**, 186–189 (1999).

16.   Bowden, R. M., Ewert, M. A. & Nelson, C. E. Environmental sex determination in a
reptile varies seasonally and with yolk hormones. *Proc. R. Soc. B Biol. Sci.* **267**, 1745–
1749 (2000).

17.   Elf, P. K. Yolk steroid hormones and sex determination in reptiles with TSD. *Gen.*
Comp. Endocrinol. **132**, 349–355 (2003).

18.   Bowden, R. M. & Paitz, R. T. Temperature fluctuations and maternal estrogens as
critical factors for understanding temperature-dependent sex determination in nature *J.*
Exp. Biol. part A Ecol. Integr. Physiol. 1–8 (2018).

19.   Crews, D., Bull, J. J. & Wibbels, T. Estrogen and sex reversal in turtles: A dose-
dependent phenomenon. *Gen. Comp. Endocrinol.* **81**, 357–364 (1991).

20.   Crews, D., Wibbels, T. & Gutzke, W. H. N. Action of sex steroid hormones on
temperature-induced sex determination in the snapping turtle (*Chelydra serpentina*).
Gen. Comp. Endocrinol. **76**, 159–166 (1989).

21.   Merchant-Larios, H., Ruiz-Ramirez, S., Moreno-Mendoza, N. & Marmolejo-Valencia,
A. Correlation among thermosensitive period, estradiol response, and gonad
differentiation in the sea turtle *Lepidochelys olivacea. Gen. Comp. Endocrinol.* **107**,
373–385 (1997).

22.   Wibbels, T., Bull, J. J. & Crews, D. Synergism between temperature and estradiol: a
common pathway in turtle sex determination? *J. Exp. Zool.* **260**, 130–134 (1991).

23.   Wibbels, T. & Crews, D. Specificity of steroid hormone-induced sex determination in
a turtle. *J. Endocrinol.* **133**, 121–129 (1992).

24.   Rhen, T. & Lang, J. W. Phenotypic plasticity for growth in the common snapping
turtle: effects of incubation temperature, clutch , and their interaction. *Am. Nat.* **146**,
726–747 (1995).

25.   Wibbels, T. & Crews, D. Putative aromatase inhibitor induces male sex determination

in a female unisexual lizard and in a turtle with temperature-dependent sex

determination. *J. Endocrinol.* **141**, 295–299 (1994).

26.   Carter, A. W., Bowden, R. M. & Paitz, R. T. Seasonal shifts in sex ratios are mediated

by maternal effects and fluctuating incubation temperatures. *Funct. Ecol.* **31**, 876–884

(2017).

27.   Hawkes, L., Broderick, A., Godfrey, M. & Godley, B. Climate change and marine

turtles. *Endanger. Species Res.* **7**, 137–154 (2009).

28.   Jensen, M. P. *et al.* Environmental warming and feminization of one of the largest sea

turtle populations in the world. *Curr. Biol.* **28**, 154-159 (2017).

29.   Tanner, C., Marco, A., Martins, S., Abella-Perez, E. & Hawkes, L. Highly feminised

sex ratio estimations for the world's third-largest nesting aggregation of the

loggerhead sea turtle. *Mar. Ecol. Prog. Ser.* **621**, 209–219 (2019).

30.   Mazaris, A. D., Kallimanis, A. S., Pantis, J. D. & Hays, G. C. Phenological response

of sea turtles to environmental variation across a species' northern range. *Proc. Biol.*

Sci. **280**, 20122397 (2013).

31.   Reneker, J. L. & Kamel, S. J. The maternal legacy: female identity predicts offspring

sex ratio in the Loggerhead Sea Turtle. *Sci. Rep.* **6**, 29237 (2016).

32.   Tezak, B., Sifuentes-Romero, I., Milton, S. & Wyneken, J. Identifying sex of neonate

turtles with temperature-dependent sex determination via small blood samples. *Sci.*

Rep. **10**, 5012 (2020).

33.   Janzen, J., Wilson, M. E., Tucker, J. K. & Ford, S. P. Endogenous yolk steroid

hormones in turtles with different sex-determining mechanisms. *Gen. Comp.*

Endocrinol. **111**, 306–317 (1998).

34. DeGregorio, B. A. & Williard, A. S. Incubation temperatures and metabolic heating of

relocated and in situ loggerhead sea turtle (*Caretta caretta*) nests at a northern

rookery. *Chelonian Conserv. Biol.* **10**, 54–61 (2011).

35. Mrosovsky, N., Baptistotte, C. & Godfrey, M. H. Validation of incubation duration as

an index of the sex ratio of hatchling sea turtles. *Can. J. Zool.* **77**, 831–835 (1999).

36. Mrosovsky, N., Kamel, S., Rees, A. F. & Margaritoulis, D. Pivotal temperature for

loggerhead turtles (*Caretta caretta*) from Kyparissia Bay, Greece. *Can. J. Zool.* **80**,

2118–2124 (2002).

37. Ceriani, S. A. & Wyneken, J. Comparative morphology and sex identification of the

reproductive system in formalin-preserved sea turtle specimens. *Zoology* **111**, 179–

187 (2008).

38. Frey, B. J. & Dueck, D. Clustering by passing messages between data points. *Science.*

315, 972–976 (2007).

39. Xia, Z.-R., Li, P.-P., Gu, H.-X., Fong, J. J. & Zhao, E.-M. Evaluating noninvasive

methods of sex identification in green sea turtle (*Chelonia mydas*) hatchlings.

Chelonian Conserv. Biol. **10**, 117–123 (2011).

40. Gross, T. S., Crain, D. a, Bjorndal, K. a, Bolten, a B. & Carthy, R. R. Identification of

sex in hatchling loggerhead turtles (*Caretta caretta*) by analysis of steroid

concentrations in chorioallantoic/amniotic fluid. *General and comparative*

endocrinology **99**, 204–210 (1995).

41. Kaska, Y. *et al.* Sex ratio estimations of loggerhead sea turtle hatchlings by

histological examination and nest temperatures at Fethiye beach, Turkey.

Naturwissenschaften **93**, 338–343 (2006).

42. Marcovaldi, M., Godfrey, M. H. & Mrosovsky, N. Estimating sex ratios of loggerhead

turtles in Brazil from pivotal incubation durations. *Can. J. Zool.* **75**, 755–770 (1997).

43.   Warner, D. A., Addis, E., Du, W. guo, Wibbels, T. & Janzen, F. J. Exogenous

application of estradiol to eggs unexpectedly induces male development in two turtle

species with temperature-dependent sex determination. *Gen. Comp. Endocrinol.* **206**,

16–23 (2014).

44.   Warner, D. A., Mitchell, T. S., Bodensteiner, B. L. & Janzen, F. J. Sex and incubation

temperature independently affect embryonic development and offspring size in a turtle

with temperature-dependent sex determination. *Physiol. Biochem. Zool.* **93**, 62–74

(2020).

45.   Radder, R. S. Maternally derived egg yolk steroid hormones and sex determination:

review of a paradox in reptiles. *J. Biosci.* **32**, 1213–1220 (2007).

46.   LeBlanc, A. M. *et al.* Nest temperatures and hatchling sex ratios from loggerhead

turtle nests incubated under natural field conditions in Georgia, United States.

Chelonian Conserv. Biol. **11**, 108–116 (2012).

47.   Groothuis, T. G. G. & Schwabl, H. Hormone-mediated maternal effects in birds:

mechanisms matter but what do we know of them? *Philos. Trans. R. Soc. B Biol. Sci.*

363, 1647–1661 (2008).

48.   Hamann, M., Limpus, C. J. & Whittier, J. M. Patterns of lipid storage and mobilisation

in the female green sea turtle (*Chelonia mydas*). *J. Comp. Physiol. B Biochem. Syst.*

Environ. Physiol. **172**, 485–493 (2002).

49.   Brischoux, F., Lourdais, O., Boissinot, A. & Angelier, F. Influence of temperature,

size and confinement on testosterone and corticosterone levels in breeding male spined

toads (*Bufo spinosus*). *Gen. Comp. Endocrinol.* **269**, 75–80 (2018).

50.   Tschirren, B., Sendecka, J., Groothuis, T. G. G., Gustafsson, L. & Doligez, B.

Heritable variation in maternal yolk hormone transfer in a wild bird population. *Am.*

Nat. **174**, 557–564 (2009).

51.   Stiebens, V. A. *et al.* Living on the edge : how philopatry maintains adaptive potential.

Proc. R. Soc. **280**, 1–9 (2013).

52.   Mrosovsky, N., Bass, A., Corliss, L. A., Richardson, J. I. & Richardson, T. H. Pivotal

and beach temperatures for hawksbill turtles nesting in Antigua. *Can. J. Zool.* **70**,

1920–1925 (1992).

53.   Mrosovsky, N., Kamel, S. J., Díez, C. E. & van Dam, R. P. Methods of estimating

natural sex ratios of sea turtles from incubation temperatures and laboratory data.

Endanger. Species Res. **8**, 147–155 (2009).

54.   Wibbels, T., Hanson, J., Balazs, G., Hillis-Starr, Z.-M. & Phillips, B. Blood sampling

techniques for hatchling cheloniid sea turtles. *Herpetol. Rev.* **29**, 218–220 (1998).

55.   Girondot, M. Statistical description of temperature-dependent sex determination using

maximum likelihood. *Evol. Ecol. Res.* **1**, 479–486 (1999).

56.   Schwabl, H. Yolk is a source of maternal testosterone for developing birds. *Proc. Natl.*

Acad. Sci. U. S. A. **90**, 5 (1993).

57.   Monsinjon, J. *et al.* Embryonic growth rate thermal reaction norm of Mediterranean

Caretta caretta embryos from two different thermal habitats, Turkey and Libya.

Chelonian Conserv. Biol. **16**, 172–179 (2017).

58.   Houghton, J. D. R. *et al.* Protracted rainfall decreases temperature within leatherback

turtle (*Dermochelys coriacea*) clutches in Grenada, West Indies: Ecological

implications for a species displaying temperature dependent sex determination. *J. Exp.*

Mar. Bio. Ecol. **345**, 71–77 (2007).

59.   Lolavar, A. & Wyneken, J. Effect of rainfall on loggerhead turtle nest temperatures,

sand temperatures and hatchling sex. *Endanger. Species Res.* **28**, 235–247 (2015).

60.   Al-Habsi, A. A. *et al.* Plasma hormone levels in the green turtles *Chelonia mydas*

during peak period of nesting at Ras Al-Hadd-Oman. *J. Endocrinol.* **191**, 9–14 (2006).

61.   Rostal, D. C. *et al.* Changes in gonadal and adrenal steroid levels in the leatherback

sea turtle (*Dermochelys coriacea*) during the nesting cycle. *Gen. Comp. Endocrinol.*

122, 139–147 (2001).

62.   Rostal, D. C., Paladino, F. V., Patterson, R. M. & Spotila, J. R. Reproductive

physiology of nesting leatherback turtles (*Dermochelys coriacea*) at Las Baulas

National Park, Costa Rica. *Chelonian Conserv. Biol.* **2**, 230–236 (1996).

63.   Dobbs, K. A., Miller, J. D., Owens, D. W. & Landry Jr., A. M. Serum testosterone and

estradiol 17β levels during the peak period in a year-round hawksbill turtle,

Eretmochelys imbricata L. nesting population. *Pacific Conserv. Biol.* **13**, 151–157

(2007).

64.   Myre, B. L., Guertin, J., Selcer, K. & Valverde, R. A. Ovarian dynamics in free-

ranging loggerhead sea turtles (*Caretta caretta*). *Copeia* **104**, 921–929 (2016).

65.   Kobayashi, S. *et al.* The effects of nest incubation temperature on embryos and

hatchlings of the loggerhead sea turtle: Implications of sex difference for survival rates

during early life stages. *J. Exp. Mar. Bio. Ecol.* **486**, 274–281 (2017).

621 **Figures**

623 **Fig. 1: Sex ratios of study clutches A) as would be expected with a pivotal temperature**

624 **of 29 °C B) based on the relationship between incubation duration and clutch sex ratios**

625 **in Kyparassia**

Fig. 2: Individual sex as calculated by affinity propagation clustering (APC). A) APC identifies three different clusters based on individual E_2:T ratio and clutch incubation duration. These clusters equate to female (red and black) and male (blue) offspring; B) Significant differences in the concentrations of T ($F_{1,60} = 10.673$, $p = 0.002$), E_2 ($F_{1,57} = 7.521$, $p = 0.008$) and the E_2:T ratio ($F_{1,48} = 28.652$, $p < 0.001$) between male and female offspring (mean, 95% confidence intervals and raw data are shown); C) Frequency of male offspring estimated by APC in relation to incubation duration. The pivotal duration was estimated at 57.25 days.

Fig. 3: Relationship between maternally derived A) Testosterone T ($F_{1,22} = 10.624$, p = 0.003), B) Estradiol E_2 ($F_{1,23} = 3.169$, p = 0.088) and C) E_2:T ratio ($F_{1,21} = 12.882$, p = 0.002) and concentrations within egg yolks and incubation duration. Size of data points relates to the sex ratio as proportion of females as determined by APC.

Fig. 4: The population sex ratio of Cape Verde over the next century if it was determined by temperature alone [4] (black) and incorporating the effect of hormones observed here on the sex determining mechanism (red) along with the 95% confidence intervals.