

1 **Maternally derived sex steroid hormones impact sex ratios of loggerhead sea**
2 **turtles**

3
4 **Authors:** Emma C. Lockley¹, Thomas Reischig², Christophe Eizaguirre¹

5 ¹Queen Mary University of London, School of Biological and Chemical Sciences, Mile
6 End Road, London E14NS, United Kingdom.

7 ²Turtle Foundation, An der Eiche 7a, 50678 Cologne, Germany.

8 **Corresponding Author:** Emma Lockley, Queen Mary University of London, School
9 of Biological and Chemical Sciences, Mile End Road, London E14NS, United
10 Kingdom. e.lockley@qmul.ac.uk

11

12

13

14

15

16

17

18 **Keywords:**

19 Temperature-dependent sex determination, Maternal effects, Loggerhead sea turtle,
20 Hormone transfer, Conservation, Climate Change

21 **Abstract**

22 An optimal sex ratio is arguably one of the most important demographic traits of
 23 species. Rising global temperatures threaten temperature-dependent sex
 24 determination (TSD) species with extreme sex ratio bias and ultimately extinction.
 25 Because sex steroid hormones can impact sex determination in TSD reptiles,
 26 variation in their maternal transfer within the egg yolk may form a buffer mechanism
 27 against raising temperatures. We tested this hypothesis by quantifying the effect of
 28 maternal oestradiol (E_2) and testosterone (T) on offspring sex in a threatened TSD
 29 population of loggerhead sea turtles (*Caretta caretta*). Circulating levels of E_2 and T
 30 in nesting females, in egg yolks at oviposition and in neonates were measured.
 31 Immediately after oviposition, nests were relocated into an *in situ* experimental
 32 hatchery where temperatures were controlled by standardising the incubation depth.
 33 We used affinity propagation clustering on hormone profiles guided by incubation
 34 duration, to sex individuals from these nests in a non-lethal manner, offering a novel
 35 conservation tool for this endangered species. Despite standardised temperatures,
 36 we found a large level of variation in the sex ratio of clutches, which correlated in a
 37 non-linear manner with maternal investment of the E_2 :T ratio in egg yolks. Males
 38 were produced at equal levels of E_2 and T and females were produced on either side
 39 of this optimum. Modelling sex ratios for the coming century, we show that
 40 maternally-derived hormones form a trans-generational mechanism of TSD plasticity
 41 that adjusts offspring sex ratios in endangered sea turtles.

42

43 **Introduction**

44 Fifty years after the discovery of environmental sex determination, our understanding
 45 of its evolutionary significance, underlying mechanisms and ecological
 46 consequences in the light of environmental change remains incomplete¹⁻⁵. Most
 47 reptile and some fish species undergo temperature-dependent sex determination
 48 (TSD), in which gonad differentiation is determined by temperature at a critical period
 49 of embryogenesis^{6,7}. Some species produce males at moderate temperatures and
 50 females at hot and cold extremes (e.g. the American alligator *Alligator*
 51 *mississippiensis*⁸, Type II TSD), but, more commonly, TSD species produce an
 52 increasing proportion of a specific sex across a range of incubation temperatures
 53 (Type Ia: Males at low temperatures, e.g. the painted turtle *Chrysemys picta*⁹; Type
 54 Ib: Females at low temperatures, e.g. the tuatara *Sphenodon punctatus*¹⁰). In all
 55 cases, both sexes are produced across a transitional range of temperatures (TRT),
 56 centered on a pivotal temperature at which both sexes develop in equal proportions.
 57 Because of TSD, rising global temperatures present the potential for extreme sex
 58 ratio biases, with implications for population dynamics⁴ and increased risks of
 59 extinction.

60 The adaptive value of TSD is still debated, but fitness advantages under sex-specific
 61 thermal environments are predicted by the widely favoured Charnov-Bull theory of
 62 differential fitness^{11,12}. This has been demonstrated in eggs of the Jacky dragon
 63 (*Amphibolurus muricatus*) that were experimentally treated with an aromatase
 64 inhibitor, constraining embryos to develop as males at female producing
 65 temperatures. These males showed lower lifetime reproductive success than
 66 controls³. While demonstrating the adaptive value of TSD, the use of an aromatase
 67 inhibitor to manipulate sex in this study also highlights the role of sex steroid
 68 hormones on the TSD mechanism¹³⁻¹⁵.

69 Exogenous application of oestradiol (E₂) is known to feminise TSD embryos
 70 incubated at male-producing temperatures¹⁶⁻¹⁹. In addition, the application of
 71 testosterone (T), the precursor androgen of E₂, also feminises embryos via the
 72 synthesis of E₂ from T by the aromatase enzyme^{17,20}, and indeed the use of
 73 aromatase inhibitors can force male development^{21,22}. Both temperature and
 74 exogenous treatment with E₂ activate the same molecular pathways, altering the
 75 transcription of the chromatin modifier gene *Kdm6b*, and conferring sensitivity to a
 76 sex-determining gene, *Dmrt1*².

77 In model TSD species that exhibit Type 1a TSD, the slider (*Trachemys scripta*) and
 78 the painted turtle, maternal transfer of sex steroid hormones into eggs varies
 79 seasonally^{13,23}. Elevated concentrations of yolk E₂ and greater E₂:T ratios increase

the likelihood of feminisation at a given temperature, effectively reducing the pivotal temperature of a clutch^{13,23}. Should these patterns be found in non-model species, variation in maternal hormone transfer to eggs may be a universal mechanism to (i) change the threshold at which temperature affects an individual's sex, (ii) modify the sex ratio of the clutch, and (iii) buffer against the negative effects of rapid global temperature increase.

There is a particular need to understand the impacts of climate change on the demographics of threatened species. As a consequence of rising temperatures, extreme feminisation of sea turtle populations has been forecast by the end of the century^{4,24–26}. Some studies already suggest effects are already visible in adult populations²⁵. Yet, predictions mostly assume a fixed population-wide pivotal temperature with no account for physiological mechanisms that may increase variation in this trait and therefore on sex ratio. Understanding possible mechanisms has been constrained in sea turtles in particular by the lack of non-lethal methods to sex neonates. This issue is especially important for endangered populations, where sacrificing individuals is not possible.

Here, focusing on loggerhead turtles (*Caretta caretta*) nesting in the Cabo Verde archipelago, we standardised the thermal environment of clutches in an experimental field hatchery, exposed to natural conditions. Should temperature be the sole driver of sex determination, under standardised thermal conditions we would expect to observe similar sex ratios among clutches. Alternatively, any inter-clutch variation would likely arise from intrinsic characteristics of the eggs, such as maternally derived hormones. To test this alternative hypothesis, we quantified E₂ and T concentrations in the plasma of nesting females, their egg yolks and neonates. We developed a non-lethal method using circulatory hormone profiles of neonates, and determined the clutches' sex ratios. Inter-clutch variation in sex ratio was then linked to yolk hormone concentrations. Finally, we illustrate how maternal hormone transfer impacts sex ratio in the face of IPCC climate change predictions, by re-parameterising a previously used mathematical model⁴ to forecast the future population dynamics of this endangered nesting aggregation.

110

111 **Results**

This study focused on loggerhead turtles nesting on the island of Boavista in the Cape Verde archipelago. First, using enzyme-linked immunosorbent assays (ELISA – Enzo LifeSciences), we quantified concentrations of the sex steroid hormones E₂ and T in both the blood plasma of 26 nesting females and up to two of their eggs directly after oviposition. Clutch sizes were recorded at this time. High levels of

individual variation were observed in adult plasma hormone levels (SI appendix, Table S1), with a mean T concentration of 1148.48 ± 148.63 (SE) pg/ml, a mean E_2 concentration of 235.79 ± 22.71 (SE) pg/ml, and a mean $E_2:T$ ratio of 0.32 ± 0.05 (SE). Linear models (LM) showed positive correlations between E_2 and T in both the female plasma (SI Appendix, Fig. S1A, $F_{1,16} = 4.608$, $p = 0.048$) and their egg yolks (SI Appendix, Fig. S1B, $F_{1,23} = 7.338$, $p = 0.013$). In reptiles, maternally derived hormones are constant across all eggs of a given clutch²⁷, which we confirmed with a subset of clutches where two egg yolks were analysed (Paired t-tests: T: $df = 11$, $t = 0.224$, $p = 0.827$; E_2 : $df = 10$, $t = -0.885$, $p = 0.397$; $E_2:T$: $df = 9$, $t = -1.173$, $p = 0.271$).

There was a significant non-linear correlation between T concentrations in adult plasma and egg yolks (SI appendix, Fig. S2A: LM: $F_{1,14} = 5.263$, $p = 0.038$), where concentrations of yolk T were lowest in eggs originating from females with intermediate levels of plasma T, but did not correlate with clutch size (SI appendix, Fig. S2B: $F_{1,14} = 0.032$, $p = 0.862$). In contrast, adult female plasma E_2 concentrations were not correlated with E_2 in the egg yolk (SI appendix, Fig. S2C: LM: $F_{1,21} = 0.908$, $p = 0.351$), but as clutch size increased, yolk E_2 concentrations significantly decreased (SI Appendix, Fig. S2D: LM: $F_{1,21} = 4.945$, $p = 0.037$). The maternal $E_2:T$ ratio showed a non-linear relationship with the $E_2:T$ ratio in the egg yolk (SI Appendix, Fig. S2E: $F_{1,14} = 6.493$, $p = 0.023$), and was not correlated with clutch size (SI Appendix, Fig. S2F: $F_{1,14} = 1.682$, $p = 0.215$).

Immediately after oviposition, the clutches of these 26 females and two others ($n = 28$) were relocated into an *in-situ* experimental hatchery that was protected from predation, yet exposed to natural sand and weather conditions. We buried clutches at a depth of 55 cm to standardise the thermal incubation environment. We confirmed the standardised thermal environment using data loggers placed at the centre of the clutch (mean thermosensitive period temperature = 30.02 ± 0.05 (SE) °C, SI Appendix, Fig. S3). The small amount of temperature variation observed was explained by differences in clutch size ($F_{1,26} = 4.418$, $p = 0.045$), resulting from increased metabolic heat produced from more developing embryos in larger clutches²⁸. Assuming the pivotal temperature of this population to be 29 °C, as has previously been used for this population⁴, this incubation temperature would produce 12.89 ± 0.01 (SE) % male offspring if temperature was the sole determinant of sex ratio (Fig. 1A). Incubation duration, the time between oviposition and neonate emergence, is also often used as a proxy to predict offspring sex ratios (e.g. $r^2 = 0.73$ in nests in Brazil²⁹) and was recorded for each clutch²⁹. Using the established logistic relationship between incubation duration and sex ratio observed in loggerhead turtles

154 from Kyparissia, Greece (Fig. 1B, the closest location where the relationship between
155 incubation duration and offspring sex ratio has been quantified), the predicted sex
156 ratio of our study clutches would be 47.5 ± 6 (SE) % males³⁰. This suggests that
157 levels of sex ratio variation are far greater than those we would expect from
158 temperature alone.

159 While the incubation duration represents a reasonable proxy for estimating the sex
160 ratio of sea turtle offspring, currently the only accurate method to resolve individual
161 sex requires sacrificing neonates and histological examination - a limiting factor for
162 endangered populations^{29,31}. However, we developed a new method to ascertain
163 individual sex without the need to sacrifice animals. After taking 100 – 150 µl of blood
164 from 365 offspring from 28 clutches after emergence (mean offspring per clutch = 13 ± 4 (SE)), we measured plasma hormone concentrations using ELISA. Hatchling
165 hormone levels varied among individuals (SI Appendix, Table S1) and among
166 clutches, with the average $E_2:T$ ratio of clutches ranging from 1.06 ± 0.13 (SE) to
167 3.56 ± 0.68 (SE). We used affinity propagation clustering (APC) on hatchling $E_2:T$
168 ratios guided by incubation duration to identify clusters of individuals with a similar
169 hormonal phenotype. APC iteratively considers the similarity of a data point to its
170 neighbours. Importantly, it does not require the number of possible clusters to be
171 defined *a priori*, as is necessary for other clustering approaches such as k-means³².
172 We identified three APC clusters (Fig. 2A). Two of these originate from clutches with
173 short incubation durations, the classic trait of female neonates, and were
174 distinguished by differences in their mean $E_2:T$ ratio (SI Appendix, Fig. S4, Cluster 1:
175 mean = 4.45 ± 0.26 (SE), Cluster 2: mean = 1.72 ± 0.05 (SE), t-test: df = 44.08, t =
176 10.273, p < 0.001). The third group is formed by individuals from clutches with longer
177 incubation durations (t-test: df = 299.3, t = -32.933, p < 0.001) and a low $E_2:T$ ratio
178 (SI Appendix, Fig. S4, mean = 1.52 ± 0.06 (SE)), the characteristics of male sea
179 turtle neonates.
180

181 Several positive theoretical controls were used to confirm this method since neonate
182 sacrificing is not possible. First, linear mixed effect models (LMM) using clutch ID as
183 a random factor revealed significant differences in hormone levels between the two
184 sexes, that were directly comparable to previous studies in which individuals' sex
185 was confirmed through histology^{33,34}. As expected, T levels were higher in males
186 (Fig. 2Bi, LMM: $F_{1,60} = 10.673$, p = 0.002, mean = 63.63 ± 2.89 (SE) pg/ml) than in
187 females (mean = 52.54 ± 2.34 (SE) pg/ml), and conversely E_2 levels were higher in
188 females (Fig. 2Bii, LMM: $F_{1,57} = 7.521$, p = 0.008, mean = 92.94 ± 3.06 (SE) pg/ml)
189 than in males (mean = 81.66 ± 3.16 (SE) pg/ml), as was the overall $E_2:T$ ratio (Fig.
190 2Biii, LMM: $F_{1,48} = 28.652$, p < 0.001, females: mean = 2.22 ± 0.09 (SE); males:

mean = 1.52 ± 0.06 (SE)). LMMs did not detect any difference in weight ($F_{1, 348} = 0.024$, $p = 0.878$) or size ($F_{1, 218} = 0.766$, $p = 0.382$) between the sexes, as would be expected under these conditions by the Charnov-Bull theory¹¹. Second, by combining individual offspring sex into an estimate of clutch sex ratio, and comparing this to the incubation duration, we found the specific logistic regression curve that characterises incubation durations in Type Ia TSD species (Fig. 2C). The pivotal duration was fitted to a value of 57.25 days (95% CIs: 57.09, 57.43), with a transitional range of incubation durations of 2.15 days (95% CIs: 1.52, 2.77). Importantly, if individual sex were incorrectly assigned, this distinctive pattern of TSD species would not be seen. With this method we determined that clutch sex ratios were on average 40.49 ± 8.98 (SE) % male (Fig. 1C). This suggests 26.1% more males and far more variation in clutch sex ratio than would be expected based on incubation temperatures alone. Our sex ratio estimate is slightly below (7.1%) that estimated from parameters based on incubation durations in Kyparissia, suggesting population differences in development rate exist, likely as a result of different average pivotal temperatures among rookeries. After establishing that the inter-clutch variation in sex ratio (and also in incubation duration, see SI Appendix) was too great to be produced by temperature alone, we tested whether metabolic heat and/or maternal hormone transfer in the yolk predicted incubation duration and the estimated sex ratio. Yolk T correlated negatively with both incubation duration (LM, $F_{1,22} = 10.624$, $p = 0.003$) and the proportion of males produced within a clutch (Fig. 3A, Binomial generalised linear mixed effect models (GLMM), $\chi^2 = 4.371$, $df = 1$, $p = 0.037$), but metabolic heat had no detectable effect (incubation duration model: $F_{1,22} = 2.436$, $p = 0.133$, sex ratio model: $\chi^2 = 2.111$, $df = 1$, $p = 0.146$). There was no relationship between yolk E_2 and incubation duration or clutch sex ratio (Fig. 3B, incubation duration: $F_{1,23} = 3.169$, $p = 0.088$, sex ratio: $\chi^2 = 0.183$, $df = 1$, $p = 0.669$), yet the yolk E_2 :T ratio showed a non-linear relationship with both incubation duration (Fig. 3C, $F_{1,21} = 12.882$, $p = 0.002$) and sex ratio independently of temperature ($\chi^2 = 7.064$, $df = 2$, $p = 0.029$). A maximum incubation duration of 57.2 days was observed at an equal hormone ratio (E_2 :T of 1.05, $y = -7.8x^2 + 16.3x + 48.7$) with the highest levels of male offspring developing at this point. Male offspring production was highest when maternal investment of E_2 and T to the yolk was equal. Asking whether the production of either sex is more costly in terms of total hormone investment, we compared the total hormone concentration ($E_2 + T$) with the overall E_2 :T ratio. This relationship was again non-linear, with total hormone investment being highest when the E_2 :T ratio was unequal (LM: $F_{2,22} = 4.951$, $p =$

0.017), suggesting that producing females requires more maternal investment than males. The total hormone investment also showed a non-linear relationship with clutch size ($\log(E_2 + T)$: $F_{2,22} = 4.306$, $p = 0.026$), with an initial increase in investment across clutch sizes between 65 and 75 eggs, after which investment declined with increasing clutch size.

Finally, to illustrate how maternal hormone transfer could impact population dynamics, we re-parameterised a mathematical model of neonate sex ratio for the Cape Verde population⁴ based on IPCC climate emission prediction SRES2, from the Fourth Assessment Report released in 2007. We made the simple assumption that the effect of maternally derived hormones on sex ratio is constant across a thermal gradient and applied the 26.1% observed difference in male offspring production for the coming century (Fig. 4). With a mechanism of this possible strength, the population is unlikely to reach the levels of extreme feminisation previously forecasted – instead of female production reaching over 97% in 2100, it is likely to instead reach approximately 71%. As it remains to be determined how maternal hormone transfer interacts with different incubation temperatures, this model can only illustrate the potential importance of trans-generational hormone transfer for population dynamics.

Discussion

Given the many considerable historical climate shifts experienced by TSD species, they are likely to have evolved mechanisms to avoid unviable biases in offspring sex ratio⁵. By experimentally standardising the thermal environment of loggerhead sea turtle nests *in-situ*, we investigated whether maternally derived hormones correlate with offspring sex independently of temperature. First, we developed a non-lethal method to determine neonate sex upon nest emergence, using affinity propagation clustering, which considers individual circulatory sex steroid hormones in relation to their incubation duration. We found a non-linear relationship between the clutch sex ratios and the ratio of maternally derived $E_2:T$ within the egg yolk under standardised thermal conditions. Low concentrations of equal investment in both hormones within the yolk maximise the production of male offspring, while increasing the concentration of either E_2 or T , along with overall hormone investment, feminises the clutches. By re-parameterising an existing model that predicts sex ratio biases in response to climate change, we demonstrated that this trans-generational mechanism could prevent extreme feminisation loggerhead turtles in Cabo Verde.

To date, an inability to determine neonate sex non-lethally has constrained the study of TSD mechanisms in endangered sea turtles. A clustering approach that identifies

individuals with similar phenotypes (here hormone profiles) that match control traits of male and female offspring (incubation duration) overcame this problem. Using $E_2:T$ thresholds to define neonate sex has been verified with histological analysis in loggerhead³⁴ and green³³ turtles, but as $E_2:T$ levels vary considerably between clutches, it is difficult to delineate thresholds *a priori*. Using an APC method guided by incubation duration to group hormone profiles, a common proxy for sex ratio, we avoid the need to define thresholds and, importantly, the need to sacrifice individuals²⁹. There is strong evidence for the reliability of this method, as (i) circulating $E_2:T$ ratios of male and female offspring identified in this study and (ii) pivotal duration, both match those reported in other studies^{30,33–36}, and (iii) the relationship between sex ratio and incubation duration fits the known logistic regression curve observed in Type Ia TSD species. We anticipate that this non-lethal approach will prove an invaluable tool for both research on TSD in sea turtle species and also for wider conservation.

Despite the standardised thermal environment of clutches within this experiment, high levels of variation in incubation duration and sex ratio were observed, which correlated with maternally derived hormones within the egg yolk. The relationship between the yolk $E_2:T$ ratio and clutch sex ratio was best described by a quadratic curve, centred on an equal concentration of both hormones and ranging from 0.37 to 1.73. When maternal investment of E_2 and T was equal, incubation durations were long, and males were produced. If hormone investment was biased in either direction, sex ratios became increasingly feminised. The effects of elevated levels of both E_2 and T on sex ratio in this study are consistent with experimental manipulation of these hormones in other species^{2,17,19}. Interestingly, in all other reptiles for which data is available, the $E_2:T$ ratios that are transferred to the yolk consistently remain below or above the ratio of one³⁷. Thus, our study is the only example of both hormones influencing the feminisation process of reptiles under natural conditions.

Total hormone concentrations within the yolk were lowest at an equal $E_2:T$ ratio. If this ratio departed from one in either direction, total concentrations of yolk hormones increased. As E_2 and T positively correlate within the egg, if investment in either hormone is elevated, there is an associated increase in the other. The outcome is that greater investment is required to skew $E_2:T$ ratios in a manner that favours the production of female offspring. When $E_2:T$ ratios are skewed, and total hormone concentrations are high, feminisation is easily achieved through either the presence of E_2 directly, or by the synthesis of E_2 from its precursor, T, by the aromatase enzyme. When E_2 and T are in equilibrium, low concentrations of E_2 are not sufficient

301 to feminise the clutch. However, product-feedback inhibition likely prevents further E_2
302 being synthesised from T, and consequently male offspring are produced.

303 There is no doubt that temperature is the primary determinant of sex in TSD species,
304 yet there is growing evidence that E_2 and, indirectly, T affect the same developmental
305 pathways^{2,19}. Accordingly, a clutch specific threshold exists for feminisation that is the
306 product of both temperature and maternal hormone transfer. A shift towards an equal
307 E_2 :T ratio and lower maternal investment will increase the pivotal temperature away
308 from the feminisation threshold, and consequently warmer temperatures would be
309 required to feminise a clutch. This aligns with sex ratios observed here, which
310 contained 26.1% more males than expected from a pivotal temperature of 29 °C.
311 However, this mechanism will be constrained by physiological limits of maternal
312 hormone investment.

313 Two maternal traits show a relationship with levels of hormone transfer to the clutch.
314 Firstly, T concentrations within the yolk correlated non-linearly with those in maternal
315 plasma. Disentangling the cause of such a relationship is complex as it is likely to
316 result from multiple physiological cascades³⁸. However, this relationship does allow
317 us to link T investment to maternal state. Should maternal T vary in response to
318 environmental cues, as in the spined toad (*Bufo spinosus*), it may allow nesting
319 females to plastically match the individual development threshold of feminisation to
320 the ambient temperature, and maintain more constant sex ratios across a nesting
321 season³⁹. Similarly, differences in the maternally-transferred E_2 :T ratio in the egg yolk
322 of a population of painted turtles resulted in a seasonal shift of the pivotal
323 temperature¹³. Secondly, total E_2 investment within eggs decreased as clutch size
324 increased, and total hormone concentrations were low in large clutches. Thus, in
325 large clutches with more metabolic heat production, the developmental threshold of
326 feminisation is increased – minimizing sex bias. We infer from these results that
327 there are two distinct mechanisms that can affect the ratio of E_2 :T within the yolk,
328 which explains how elevated investment in either hormone can lead to feminisation.

329 There is considerable variation in circulating T and E_2 levels between sea turtle
330 populations and species (SI Appendix, Table S1), which may suggest an element of
331 local adaptation in response to environmental conditions, and a heritable component
332 of baseline physiological levels⁴⁰.

333 Overall, our work highlights a previously under-considered physiological mechanism
334 for individual variation in the TSD process within sea turtle species. There is a need
335 for management plans that use temperature-based models to predict future sex
336 ratios to account for maternal hormonal influence, as this will have considerable
337 implications for population dynamics.

338

339 **Methods**

340 *Sample Collection*

341 We studied nesting loggerhead sea turtles on the island of Boavista, part of the Cabo
342 Verde archipelago in the eastern Atlantic. The sampling site (15°58'18.6"N,
343 22°48'06.2"W) is a 400 m stretch of coastline on the southern tip of this island.
344 Twenty-eight nesting females were sampled between 17 July and 1 August 2017.
345 Immediately after oviposition, females were individually marked with PIT (AVID) and
346 metal (Inconel) tags⁴¹. Blood samples of 1-4 ml in volume were collected from the
347 dorsal cervical sinus of 26 females using a 40 mm, 21-gauge needle and 5 ml
348 syringe, and stored within lithium heparin containers. Finally, curved carapace length
349 (CCL) and width (CCW) were measured (± 0.1 cm).

350 The clutches of these turtles (containing 83 ± 3 (SE) eggs) were relocated to an
351 experimental hatchery protected from predation, situated on the nesting beach. At
352 this point, up to two eggs from the 28 clutches were removed from each clutch for
353 yolk hormone analysis, and the rest of the clutch was buried at a depth of 55 cm. By
354 using a standard depth, temperature was controlled for, while maintaining an
355 otherwise natural environment. A TinyTag™ temperature logger was placed at the
356 centre of each clutch, programmed to take a reading every 15 minutes throughout
357 the incubation period (accuracy ± 0.2 °C). As anticipated, the uniform depth
358 standardised the incubation temperature of the nests to 30.05 ± 0.05 (SE) °C during
359 the middle third of incubation, the period where embryo sex is established. This
360 variation in temperature is extremely conserved, and is representative of the thermal
361 variation produced within treatments under controlled laboratory incubations^{42,43}.

362 Upon emergence, twenty hatchlings were randomly selected for blood sampling (100
363 – 150 μ l) from the dorsal cervical sinus, using a 26-gauge needle and 1 ml syringe⁴⁴.
364 Samples were stored within lithium heparin coated tubes. Notch-to-notch straight
365 carapace length (SCL) and, width (SCW) were measured using digital callipers (\pm
366 0.01 mm), and weight was measured with a digital scale (± 0.1 g)).

367 The blood samples of both the adults and offspring were refrigerated for up to 48 h
368 before being centrifuged to extract plasma. Egg yolks were separated from the
369 albumen, and all samples were stored at -20 °C until extraction.

370

371 *Hormone extraction*

372 Commercially available Enzyme-Linked Immunosorbent Assay (ELISA) kits for both
373 E₂ (Catalogue # ADI-900-174, ENZO Life Sciences) and T (Catalogue # ADI-900-
374 065) were used to measure steroid levels in all samples. Details for hormone

375 extraction protocols are given in SI methods. Not all blood samples had sufficient
376 volume for hormone extraction. Consequently, we extracted E_2 from 24 adults and
377 388 hatchling blood samples, and T from 19 adult and 367 hatchling blood samples.
378 This provided us with E_2 :T ratios for 18 adult females, and 365 hatchlings. E_2 , and T
379 were successfully extracted from the yolks of 26 out of the 28 sampled clutches.

380

381 *Statistical Analyses*

382 All analyses were conducted with R 3.3.3, using the R packages *lme4* and *lmerTest*
383 for fitting linear mixed models (LMMs) and generalized linear mixed models
384 (GLMMs). A paired t-test was used to compare intra-clutch E_2 and T levels between
385 two eggs in a subset of clutches ($n = 13$), to test whether that there was variation in
386 hormone investment between eggs in a clutch. As there was no difference between
387 eggs from the same clutch, for subsequent analyses the average hormone was used
388 where possible, while a single egg was used for the remainder of the clutches.
389 Correlations between E_2 and T in female plasma and yolks, the effect of clutch size
390 on temperature, and the effects of metabolic heat and maternally derived hormones
391 on incubation duration were tested using general linear models (LM). A non-linear
392 relationship between the E_2 :T ratio on incubation duration was fitted using a
393 quadratic curve. Similarly, when considering the relationship between E_2 :T and total
394 hormone investment, we also fit a quadratic model. LMs were also used to estimate
395 the correlation of clutch size and plasma hormone concentrations with yolk hormone
396 concentrations.

397 We used Algorithm Propagation Clustering (APC) to identify individual sex, using the
398 R package *apcluster*³². Cluster assignment was made based on the plasma E_2 :T
399 ratio of hatchlings, guided by their incubation duration. Determining neonate sea
400 turtle sex using the E_2 :T ratio has previously been extremely accurate (96% and
401 96.7% respectively) for artificially incubated eggs of loggerhead and green sea
402 turtles^{33,34} that were ultimately sacrificed for verification. Since variation likely exists
403 among rookeries, those thresholds however cannot be blindly applied to new
404 populations. T-tests were used to compare hormone levels of putative male and
405 female hatchlings. A response curve of these estimated sex ratios to incubation
406 duration was produced using the logistic equation function of the R package
407 *embryogrowth* to further verify the accuracy of our non-lethal sexing method.

408 After identifying the sex of individuals, LMMs were used to compare individual size
409 and weight between the sexes and the APC clusters. Finally, we used binomial
410 GLMMs to determine whether individual hatchling sex was predicted by maternal
411 hormone investment or temperature. For all LMM and GLMM analyses, clutch was

412 included as a random factor to account for individual variation. Model selection was
413 based on AIC criteria, using a likelihood ratio tests to select for the best models. P-
414 values of the selected models were obtained by with the *car* R package, and models
415 were verified for over-dispersion.

416 Thermal estimates of sex ratio were calculated using the equation first presented by
417 Girondot in 1999, under an assumed pivotal temperature of 29 °C⁴⁵. Estimates of sex
418 ratio based on incubation duration were made based on data from a study on a
419 neighbouring loggerhead sea turtle population that nests in Kyparissia, Greece, and
420 was confirmed with histology³⁰. To generate an illustrative model that compared the
421 results of our study with future predictions based on temperature alone, we extracted
422 data from a previously published study predicting sex ratios until 2100 based on
423 temperature alone. We then compared our observed mean clutch sex ratio to that
424 expected from a pivotal temperature of 29 °C, and added the difference, along with
425 95% confidence intervals, to the original prediction.

426

427 **Acknowledgements**

428 We thank volunteers and staff from Turtle Foundation for facilitating this research.
429 Particular thanks go to S. Cameron for considerable help during data collection. E.L
430 was supported by a grant from the National Environment Research Council (grant
431 NE/L002485/1). This study was supported by a National Geographic grant (NGS-
432 59158R-19) as well as Queen Mary University of London Funds allocated to CE. We
433 also acknowledge the support from ‘The Future Ocean’ Excellence Initiative by the
434 Deutsche Forschungsgemeinschaft (DFG) that granted a ‘Capacity building grant’ to
435 C.E. The authors would like to thank the Eizaguirre lab as well as Dr. Jose M. Duran
436 and Dr. Gail Schoefield for their feedback on previous versions of the manuscript.

437

438 **References**

- 439 1. Charnier, M. Action of temperature on the sex ratio in the *Agama agama*
440 (*Agamidae*, *Lacertilia*) embryo. *C R Seances Soc Biol Fil.* **160**, 620–622
441 (1966).
- 442 2. Ge, C. *et al.* The histone demethylase KDM6B regulates temperature-
443 dependent sex determination in a turtle species. *Science (80-.).* **360**, 645–648
444 (2018).
- 445 3. Warner, D. A. & Shine, R. The adaptive significance of temperature-dependent
446 sex determination in a reptile. *Nature* **451**, 2006–2009 (2008).

- 447 4. Laloë, J.-O. *et al.* Climate change induced extinction does not seem imminent
448 at an important sea turtle rookery. *Nat. Clim. Chang.* **4**, 1–7 (2014).
- 449 5. Mitchell, N. J. & Janzen, F. J. Temperature-Dependent sex determination and
450 contemporary climate change. *Sex. Dev.* **4**, 129–140 (2010).
- 451 6. Bull, J. J. Sex Determination in Reptiles. *Q. Rev. Biol.* **55**, 3–21 (1980).
- 452 7. Deeming, D. C. *et al.* Environmental Regulation of Sex Determination in
453 Reptiles (and Discussion). *Philos. Trans. R. Soc. B Biol. Sci.* **322**, 19–39
454 (1988).
- 455 8. Ferguson, M. W. J. & Joanen, T. Temperature-dependent sex determination in
456 Alligator mississippiensis. *J. Zool.* **200**, 143–177 (1983).
- 457 9. Bull, J. J. & Vogt, R. C. Temperature-Dependent Sex Determination in Turtles.
458 *Science* (80-.). **206**, 1186–1188 (1979).
- 459 10. Cree, A., Thompson, M. B. & Daugherty, C. H. Tuatara sex determination.
460 *Nature* **375**, 543 (1995).
- 461 11. Charnov, E. . & Bull, J. . When is sex environmentally determined? *Nature*
462 **266**, 828–830 (1977).
- 463 12. Shine, R. Why is sex determined by nest temperature in many reptiles?
464 *Trends Ecol. Evol.* **14**, 186–189 (1999).
- 465 13. Bowden, R. M., Ewert, M. A. & Nelson, C. E. Environmental sex determination
466 in a reptile varies seasonally and with yolk hormones. *Proc. R. Soc. B Biol.*
467 *Sci.* **267**, 1745–1749 (2000).
- 468 14. Elf, P. K. Yolk steroid hormones and sex determination in reptiles with TSD.
469 *Gen. Comp. Endocrinol.* **132**, 349–355 (2003).
- 470 15. Bowden, R. M. & Paitz, R. T. Temperature fluctuations and maternal
471 estrogens as critical factors for understanding temperature-dependent sex
472 determination in nature STUDIES ARE THE FOUNDATION OF TSD
473 BETWEEN INCUBATION TEMPERATURES. *J. Exp. Biol. part A Ecol. Integr.*
474 *Physiol.* 1–8 (2018). doi:10.1002/jez.2183
- 475 16. Crews, D., Bull, J. J. & Wibbels, T. Estrogen and sex reversal in turtles: A

- 476 dose-dependent phenomenon. *Gen. Comp. Endocrinol.* **81**, 357–364 (1991).
- 477 17. Crews, D., Wibbels, T. & Gutzke, W. H. N. Action of sex steroid hormones on
478 temperature-induced sex determination in the snapping turtle (*Chelydra*
479 *serpentina*). *Gen. Comp. Endocrinol.* **76**, 159–166 (1989).
- 480 18. Merchant-Larios, H., Ruiz-Ramirez, S., Moreno-Mendoza, N. & Marmolejo-
481 Valencia, a. Correlation among thermosensitive period, estradiol response,
482 and gonad differentiation in the sea turtle *Lepidochelys olivacea*. *Gen. Comp.*
483 *Endocrinol.* **107**, 373–385 (1997).
- 484 19. Wibbels, T., Bull, J. J. & Crews, D. Synergism between temperature and
485 estradiol: a common pathway in turtle sex determination? *J. Exp. Zool.* **260**,
486 130–134 (1991).
- 487 20. Wibbels, T. & Crews, D. Specificity of steroid hormone-induced sex
488 determination in a turtle. *J. Endocrinol.* **133**, 121–129 (1992).
- 489 21. Rhen, T. & Lang, J. W. Phenotypic Plasticity for Growth in the Common
490 Snapping Turtle: Effects of Incubation Temperature, Clutch , and Their
491 Interaction. *Am. Nat.* **146**, 726–747 (1995).
- 492 22. Wibbels, T. & Crews, D. Putative aromatase inhibitor induces male sex
493 determination in a female unisexual lizard and in a turtle with temperature-
494 dependent sex determination. *J. Endocrinol.* **141**, 295–299 (1994).
- 495 23. Carter, A. W., Bowden, R. M. & Paitz, R. T. Seasonal shifts in sex ratios are
496 mediated by maternal effects and fluctuating incubation temperatures. *Funct.*
497 *Ecol.* **31**, 876–884 (2017).
- 498 24. Hawkes, L., Broderick, A., Godfrey, M. & Godley, B. Climate change and
499 marine turtles. *Endanger. Species Res.* **7**, 137–154 (2009).
- 500 25. Jensen, M. P. *et al.* Environmental warming and feminization of one of the
501 largest sea turtle populations in the world. *Curr. Biol.* **28**, 154–159.e4 (2017).
- 502 26. Tanner, C., Marco, A., Martins, S., Abella-Perez, E. & Hawkes, L. Highly
503 feminised sex ratio estimations for the world's third-largest nesting aggregation
504 of the loggerhead sea turtle. *Mar. Ecol. Prog. Ser.* **621**, 209–219 (2019).
- 505 27. Janzen, J., Wilson, M. E., Tucker, J. K. & Ford, S. P. Endogenous Yolk Steroid

- 506 Hormones in Turtles with Different Sex-Determining Mechanisms. *Gen. Comp.*
507 *Endocrinol.* **111**, 306–317 (1998).
- 508 28. DeGregorio, B. A. & Williard, A. S. Incubation Temperatures and Metabolic
509 Heating of Relocated and In Situ Loggerhead Sea Turtle (*Caretta caretta*)
510 Nests at a Northern Rookery. *Chelonian Conserv. Biol.* **10**, 54–61 (2011).
- 511 29. Mrosovsky, N., Baptistotte, C. & Godfrey, M. H. Validation of incubation
512 duration as an index of the sex ratio of hatchling sea turtles. *Can. J. Zool.* **77**,
513 831–835 (1999).
- 514 30. Mrosovsky, N., Kamel, S., Rees, A. F. & Margaritoulis, D. Pivotal temperature
515 for loggerhead turtles (*Caretta caretta*) from Kyparissia Bay, Greece. *Can. J.*
516 *Zool.* **80**, 2118–2124 (2002).
- 517 31. Ceriani, S. A. & Wyneken, J. Comparative morphology and sex identification of
518 the reproductive system in formalin-preserved sea turtle specimens. *Zoology*
519 **111**, 179–187 (2008).
- 520 32. Frey, B. J. & Dueck, D. Clustering by passing messages between data points.
521 *Science (80-.)*. **315**, 972–976 (2007).
- 522 33. Xia, Z.-R., Li, P.-P., Gu, H.-X., Fong, J. J. & Zhao, E.-M. Evaluating
523 Noninvasive Methods of Sex Identification in Green Sea Turtle (*Chelonia*
524 *mydas*) Hatchlings. *Chelonian Conserv. Biol.* **10**, 117–123 (2011).
- 525 34. Gross, T. S., Crain, D. a, Bjorndal, K. a, Bolten, a B. & Carthy, R. R.
526 Identification of sex in hatchling loggerhead turtles (*Caretta caretta*) by
527 analysis of steroid concentrations in chorioallantoic/amniotic fluid. *General and*
528 *comparative endocrinology* **99**, 204–210 (1995).
- 529 35. Kaska, Y. *et al.* Sex ratio estimations of loggerhead sea turtle hatchlings by
530 histological examination and nest temperatures at Fethiye beach, Turkey.
531 *Naturwissenschaften* **93**, 338–343 (2006).
- 532 36. Marcovaldi, M., Godfrey, M. H. & Mrosovsky, N. Estimating sex ratios of
533 loggerhead turtles in Brazil from pivotal incubation durations. *Can. J. Zool.* **75**,
534 755–770 (1997).
- 535 37. Radder, R. S. Maternally derived egg yolk steroid hormones and sex

- 536 determination: review of a paradox in reptiles. *J. Biosci.* **32**, 1213–1220
537 (2007).
- 538 38. Groothuis, T. G. G. & Schwabl, H. Hormone-mediated maternal effects in
539 birds: mechanisms matter but what do we know of them? *Philos. Trans. R.*
540 *Soc. B Biol. Sci.* **363**, 1647–1661 (2008).
- 541 39. Brischoux, F., Lourdais, O., Boissinot, A. & Angelier, F. Influence of
542 temperature, size and confinement on testosterone and corticosterone levels
543 in breeding male spined toads (*Bufo spinosus*). *Gen. Comp. Endocrinol.* **269**,
544 75–80 (2018).
- 545 40. Tschirren, B., Sendecka, J., Groothuis, T. G. G., Gustafsson, L. & Doligez, B.
546 Heritable Variation in Maternal Yolk Hormone Transfer in a Wild Bird
547 Population. *Am. Nat.* **174**, 557–564 (2009).
- 548 41. Stiebens, V. A. *et al.* Living on the edge □: how philopatry maintains adaptive
549 potential. *Proc. R. Soc.* **280**, 1–9 (2013).
- 550 42. Mrosovsky, N., Bass, A., Corliss, L. A., Richardson, J. I. & Richardson, T. H.
551 Pivotal and beach temperatures for hawksbill turtles nesting in Antigua. *Can.*
552 *J. Zool.* **70**, 1920–1925 (1992).
- 553 43. Mrosovsky, N., Kamel, S. J., Díez, C. E. & van Dam, R. P. Methods of
554 estimating natural sex ratios of sea turtles from incubation temperatures and
555 laboratory data. *Endanger. Species Res.* **8**, 147–155 (2009).
- 556 44. Wibbels, T., Hanson, J., Balazs, G., Hillis-Starr, Z.-M. & Phillips, B. Blood
557 Sampling Techniques for Hatchling Cheloniid Sea Turtles. *Herpetol. Rev.* **29**,
558 218–220 (1998).
- 559 45. Girondot, M. Statistical description of temperature-dependent sex
560 determination using maximum likelihood. *Evol. Ecol. Res.* **1**, 479–486 (1999).
- 561 46. Schwabl, H. Yolk is a source of maternal testosterone for developing birds.
562 *Proc. Natl. Acad. Sci. U. S. A.* **90**, 5 (1993).
- 563 47. Monsinjon, J. *et al.* Embryonic Growth Rate Thermal Reaction Norm of
564 Mediterranean *Caretta caretta* Embryos from Two Different Thermal Habitats,
565 Turkey and Libya. *Chelonian Conserv. Biol.* **16**, 172–179 (2017).

- 566 48. Al-Habsi, A. A. *et al.* Plasma hormone levels in the green turtles *Chelonia*
567 *mydas* during peak period of nesting at Ras Al-Hadd-Oman. *J. Endocrinol.*
568 **191**, 9–14 (2006).
- 569 49. Rostal, D. C. *et al.* Changes in gonadal and adrenal steroid levels in the
570 leatherback sea turtle (*Dermochelys coriacea*) during the nesting cycle. *Gen.*
571 *Comp. Endocrinol.* **122**, 139–147 (2001).
- 572 50. Rostal, D. C., Paladino, F. V., Patterson, R. M. & Spotila, J. R. Reproductive
573 Physiology of Nesting Leatherback Turtles (*Dermochelys coriacea*) at Las
574 Baulas National Park , Costa Rica. *Chelonian Conserv. Biol.* **2**, 230–236
575 (1996).
- 576 51. Dobbs, K. A., Miller, J. D., Owens, D. W. & Landry Jr., A. M. Serum
577 testosterone and estradiol 17B levels during the peak period in a year-round
578 hawksbill turtle, *eretmochelys imbricata* L. nesting population. *Pacific Conserv.*
579 *Biol.* **13**, 151–157 (2007).
- 580 52. Myre, B. L., Guertin, J., Selcer, K. & Valverde, R. A. Ovarian Dynamics in
581 Free-Ranging Loggerhead Sea Turtles (*Caretta caretta*). *Copeia* **104**, 921–
582 929 (2016).
- 583 53. Kobayashi, S. *et al.* The effects of nest incubation temperature on embryos
584 and hatchlings of the loggerhead sea turtle: Implications of sex difference for
585 survival rates during early life stages. *J. Exp. Mar. Bio. Ecol.* **486**, 274–281
586 (2017).

587

588

589

590

591

592

593

594

595


596

597

598

599

600 **Figures**



601

602 **Fig. 1: Sex ratios of study clutches A) as would be expected with a pivotal**
603 **temperature of 29 °C B) based on the relationship between incubation duration**
604 **and clutch sex ratios in Kyparassia (Greece)³⁰ and C) as experimentally**
605 **determined by hormone profiles and machine learning algorithm of individual**
606 **offspring.**

607




Fig. 2: Individual sex as calculated by affinity propagation clustering (APC). A) APC identifies three different clusters based on individual E₂:T ratio and clutch incubation duration. These clusters equate to female (red and black) and male (blue) offspring; B) Significant differences in the concentrations of T ($F_{1,60} = 10.673$, $p = 0.002$), E₂ ($F_{1,57} = 7.521$, $p = 0.008$) and the E₂:T ratio ($F_{1,48} = 28.652$, $p < 0.001$) between male and female offspring (mean, 95% confidence intervals and raw data are shown); C) Frequency of male offspring estimated by APC in relation to incubation duration. The pivotal duration was estimated at 57.25 days.




Fig. 3: Relationship between maternally derived A) T ($F_{1,22} = 10.624$, $p = 0.003$), B) E_2 ($F_{1,23} = 3.169$, $p = 0.088$) and C) $E_2:T$ ratio ($F_{1,21} = 12.882$, $p = 0.002$) and concentrations within egg yolks and incubation duration. Size of data points relates to the sex ratio as determined by APC.




Fig. 4: The population sex ratio of Cape Verde over the next century if it was determined by temperature alone ⁴ (black) and incorporating the effect of hormones observed here on the sex determining mechanism (red) along with the 95% confidence intervals.

