A Physiological Characterization in Controlled Bioreactors Reveals a Novel Survival
Strategy for *Debaryomyces hansenii* at High Salinity and Confirms its Halophilic Behavior.

Clara Navarrete[1], August T. Frost[1], Laura Ramos-Moreno[2], Mette R. Krum[1] and José L.
Martínez[1*]

[1]Technical University of Denmark, Department of Biotechnology and Biomedicine, Søltofts
Plads Building 223, 2800 Kgs. Lyngby, Denmark

[2]University of Córdoba, Department of Microbiology, Campus de Rabanales, Edificio Severo
Ochoa, 14071 Córdoba, Spain

*Corresponding author: Section of Synthetic Biology (DTU Bioengineering), Department of
Biotechnology and Biomedicine, Technical University of Denmark, Søltofts Plads Building 223,
2800 Kgs. Lyngby, Denmark. Tel: +45 45 25 26 30; E-mail: jlmr@dtu.dk

Keywords: *Debaryomyces hansenii*, non-conventional yeast, osmotic stress, salt tolerance, batch
fermentation, bioreactors

Abstract

Debaryomyces hansenii is traditionally described as a halotolerant non-conventional yeast, being
the model organism for the study of osmo- and salt tolerance mechanisms in eukaryotic systems
for the past 30 years.

However, unravelling of *D. hansenii´s* biotechnological potential has always been difficult due to
the persistent limitations in the availability of efficient molecular tools described for this yeast.
Additionally, there is a lack of consensus and contradictory information along the recent years
that limits a comprehensive understanding of its central carbon metabolism, mainly due to a lack
of physiological studies in controlled and monitored environments. Moreover, there is
controversy about the diversity in the culture conditions (media composition, temperature and
pH among others) used by different groups, which makes it complicated when trying to get
significant conclusions and behavioural patterns.

In this work, we present for the first time a physiological characterization of *D. hansenii* in batch
cultivations using highly instrumented and controlled lab-scale bioreactors. Our findings
contribute to a more complete picture of the central carbon metabolism and the external pH
influence on the yeast ability to tolerate high Na^+ and K^+ concentrations. Finally, the
controversial halophilic/halotolerant character of this yeast is further clarified.

Introduction

Debaryomyces hansenii is described as a halophilic/halotolerant non-conventional yeast. It is globally present in seawater and has been isolated from soil, air, materials of plant and animal origin as well as from polar waters and ice from Antarctic and Artic glaciers (Norkrans 1966; Breuer and Harms 2006; Gunde-Cimerman *et al*. 2009). *D. hansenii* has been a model for the study of osmo- and salt tolerance mechanisms in eukaryotic cells over the last 30 years (Adler *et al*. 1985; Prista *et al*. 1997, 2005). Potassium and sodium are crucial factors for yeast growth in high salt environments, and its halotolerant nature has been fully confirmed by the fact that the presence of sodium in the medium protects the yeast cells against oxidative stress and additional abiotic stresses like extreme pH or high temperature (Almagro *et al*. 2000; Papouskova and Sychrova 2007; Navarrete *et al*. 2009).

D. hansenii's genome was completely sequenced in 2004 (Dujon *et al*. 2004), and although more than 6500 genes have been annotated since then, the molecular characterization of this yeast is still far from being fully known. The majority of these annotated genes are mainly associated with salt and osmotic stress tolerance mechanisms (Prista *et al*. 2016). Besides, it has been described that *D. hansenii* has the highest coding capacity among yeasts (79.2% of the genome) with 6906 detected coding sequences (CDS) and a gene redundancy of almost 50% (Gènolevures online database, igenolevures.org/databases/).

As mentioned before, potassium and sodium fluxes (and their accumulation in cell organelles) play a key role in ion homeostasis and halotolerance in *D. hansenii*. Genes for K^+ influx (*DhTRK1, DhHAK1*) were identified and studied at a molecular, transcriptional and protein level by different authors (Prista *et al*. 2007; Martinez *et al*. 2011). Two different plasma membrane cation efflux systems (DhEna1/2, DhNha1) have also been described (Almagro *et al*. 2001; Velkova and Sychrova 2006) as well as two intracellular Na^+/H^+ antiporters (DhKha1 and DhNhx1) (Garcia-Salcedo *et al*. 2007; Montiel and Ramos 2007). Very importantly, the role of glycerol production and accumulation in the osmotic stress response to high osmolarity was also fully established by Gustafsson and Norkrans in 1976, and further confirmed by Adler *et al*. in 1985.

From a biotechnological point of view, *D. hansenii* is considered of great interest, due to its salt tolerant character (Gadanho *et al*. 2003; Butinar *et al*. 2005; Prista *et al*. 2005; Ramos *et al*. 2017 among others), especially in food industry where it is used in the ripening process of sausages by production of exopeptidases, development of flavor characteristics, and production of cheeses among others (Lopez Del Castillo-Lozano *et al*. 2007; Cano-Garcia *et al*. 2014). Moreover, *D. hansenii* is known to respire a broad range of carbon substrates and produce mycocins against other yeast species, like *Candida* (Banjara *et al*. 2016).

Nevertheless, only a few genes with high biotechnological relevance have been characterized so far in *D. hansenii*. It is the case of *DhJEN1*, coding for a monocarboxylic acid transporter (Casal *et al*. 2008), and genes coding for xylitol reductase, xylose dehydrogenase and xylose/H^+ transporter (Biswas *et al*. 2013; Ferreira *et al*. 2013), all of them used for the production of

second-generation bioethanol in the fermentation of pentoses by *Saccharomyces cerevisiae*
(Breuer and Harms 2006).

The difficulties when studying *D. hansenii*'s biotechnological potential have always been related
to the limitations in the availability of highly efficient molecular tools described for this yeast
(Prista *et al*. 2016). There is also a lack of information and full understanding of its carbon
metabolism and physiological characterization during biotechnological processes.

On top of that, the large variation in culture conditions used (e.g. media composition,
temperature, and pH) in previous studies impedes thorough understanding and solid conclusions
on this peculiar yeast's behaviour. To date, nobody has conducted an in-depth physiological
analysis of this yeast in a controlled environment (e.g. using bioreactors), and there is a lack of
consensus about its capacity to produce ethanol in high saline environments or its cell
performance in the presence of high salt concentrations. While some studies point to a beneficial
role of salts on *D. hansenii's* performance (Almagro *et al*. 2000; Papouskova *et al*. 2007;
Navarrete *et al*. 2009; Garcia-Neto *et al*. 2017), others claim that sodium is detrimental in terms
for the fitness of this yeast in general (Capusoni *et al*. 2019; Sánchez *et al*. 2018). In addition, the
osmo- and halotolerant terminology are often used incorrectly, as there are some studies in which
sodium is used for both purposes, despite there are reported evidences about sodium and
potassium triggering a different effect. Potassium does not cause a toxicity response, whereas
sodium has been described to have detrimental effects on cell growth at lower concentrations
than potassium, pointing to some authors to recently claim *D. hansenii* not being halophilic but
just halotolerant (Sánchez *et al*. 2018), whilst several previous studies report *D. hansenii* as an
halophilic yeast (Gonzalez-Hernandez *et al*. 2004; Chao *et al*. 2009; Martinez *et al*. 2011).

In relation to *D. hansenii*'s alcoholic fermentation capacity, the kinetics of cell inactivation in the
presence of ethanol at 20%, 22.5% and 25% (v/v), have been measured by progressive sampling
and viable counting, and used as an inference of the ethanol resistance status of different yeasts
(Pina *et al.* 2004). *D. hansenii* PYCC 2968T (a.k.a. CBS 767, which we use in our present study)
was found to be one of the most sensitive yeasts to ethanol in this study. Ethanol (or other
polyols) production has been only reported when a mix of sugars or complex substrates are used
as carbon source. For example, *D. hansenii* CCMI 941 was cultivated in xylose/galactose or
xylose/glucose (Tavares *et al*. 2008) for production of xylitol, although only ethanol and glycerol
were produced for a xylose/glucose ratio above 30%. Another strain background, *D. hansenii* B-
2, was grown in semi-synthetic banana peel-yeast extract-peptone broth for ethanol production,
where 40% (w/v) of glucose was fermented to 5.8% of ethanol (Brooks, 2008). In 2009,
Calahorra *et al.* stated the activation of glucose fermentation by salts in *D. hansenii* Y7426. In
this work, the authors incubated yeast cells in media containing 40mM of glucose, and
afterwards they made a protein extraction and measured the ethanol production *in vitro*, by the
enzyme extract. Only marginal ethanol production was reported though, within the range of
micromoles per gram of glucose. However a most recent work performed by Garcia-Neto *et al.*
in 2017, indicates otherwise: that the presence of high salts increases *D. hansenii's* respiratory
activity, in the same strain (Y7426). Overall, no ethanol production by *D. hansenii* has been
shown during the growth process using glucose as a sole carbon source.

In this work, we present for the first time a physiological characterization of *D. hansenii* during
batch cultivations in highly instrumented and controlled bioreactors. *D. hansenii*'s carbon
metabolism and the external pH influence on the yeast capacity to tolerate high Na^+ and K^+
concentrations are also shown. Finally, its capability of ethanol production and the controversial
halophilic/halotolerant character of this yeast are further discussed, and a novel survival strategy
at high saline environments suggested.

Materials and Methods

Strain and culture conditions

The *Debaryomyces hansenii* strain CBS767 (PYCC2968; Prista *et al*. 1997; Navarrete *et al*.
2009) was used in this study. For the micro-fermentation experiments and growth assay on plate,
IBT22 and IBT26 (CBS1101) strains were also used (from IBT culture collection of fungi from
DTU, Denmark). Glycerol stocks containing sterile 30% glycerol (Sigma-Aldrich, Germany)
were used to maintain the strain, and were preserved at -80⁰C.

Yeast extract Peptone Dextrose (YPD) medium plates with 2% agar, were used for growing the
cells from the cryostocks at 28°C. For the pre-cultures of yeast cells, synthetic complete medium
was used (6.7 g/L Yeast Nitrogen Base w/o amino acids, from Difco, plus 0.79 g/L complete
supplement mixture, from Formedium). Separately sterilized 2% D-(+)-glucose monohydrated
(VWR Chemicals, Germany) was added to the medium, and pH was adjusted to 6.0 with NaOH.
All the solutions were autoclaved at 121°C for 20 min. Cells were incubated in 500 mL baffled
Erlenmeyer shake flasks (culture volume 100 mL) at 28°C, 150 rpm for at least 24 hours.

For the growth curves in low glucose conditions, *D. hansenii* pre-cultures were prepared as
specified in the above paragraph. From those pre-cultures, cells were grown in the same medium
but with 0.2% of glucose, and with or without 1M/2M of NaCl or KCl, in order to get their
growth profile. Initial OD_{600} in the flask was 0.1 and samples were taken during eight days of
cultivation at 28°C, 150 rpm.

Growth assay on plate

Three *D. hansenii* strains (CBS767, IBT22 and IBT26) were pre-cultured for at least 24 h in
synthetic complete medium (pH 6) at 28°C as specified in "Strain and culture conditions"
section, and cell suspensions were prepared from those pre-cultures. Cell drops at OD_{600} 0.1 and
serial dilutions (1:10, 1:100) were spotted on YNB plates (pH 6) with or without KCl or NaCl at
different concentrations (0.5, 1 and 2M). Cell growth was monitored at 28°C for 144 h or 240 h
(when higher salt concentrations were used). Pictures were taken using an in-house build digital
camera set-up (Nikon D90 SLR camera).

Microscopy

D. hansenii cells were grown on synthetic complete medium (pH 6) with or without KCl or NaCl
(0.5, 1 and 2M) at 28°C as specified in "Strain and culture conditions" section. Undiluted cell

samples were taken and photographed at the microscope (BA310E & camera 580INT, In Vitro
DK) after 51 h of cultivation.

Bioreactor cultivations

Batch cultivations were performed in biological replicates (between 2-7 per condition) in 1.0 L
Biostat Qplus bioreactors (Sartorius Stedim Biotech, Germany). The temperature was controlled
at 28°C and pH was maintained at 6.0 (when desired) by the automatic addition of 2M NaOH /
2M H_2SO_4, and measured by pH sensors (Model EasyFerm Plus K8 160, Hamilton). The
volumetric flow rate (aeration) was set at 1 vvm and the stirring was constant at 600 rpm.
Dissolved oxygen concentration was also measured by DO sensors (Model OxyFerm FDA 160,
Hamilton). The working volume in the vessel was 0.5 L, using exactly the same medium
composition as in the pre-culture, containing either 2% or 0.2% of glucose when required. To
study the effect of salt in *D. hansenii* cells, NaCl or KCl (PanReac Applichem, ITW Reagents)
were added to the medium before autoclavation. The bioreactors were inoculated with 24 h
inoculum from the pre-culture to get an initial OD_{600} of 0.05-0.1. Samples for dry weight, optical
density and HPLC were taken after the CO_2 percentage values reached 0.1 and until stationary
phase.

Micro-fermentation experiments

All micro-fermentations were carried out in BioLector® I and II systems (m2p Labs, Germany)
using 48-well FlowerPlate® (MTP-48-B, m2p Labs, Germany) with a working volume of 1.5
mL. Synthetic complete medium (Yeast nitrogen base 6.7 g/L, Complete Supplement Mixture
0.79 g/L, D-(+)-Glucose monohydrated 2%) at different pH (4, 6 and 8) was used, and different
temperatures were tested (26, 28 and 30ºC). The pH was adjusted with NaOH, and potassium
phthalate monobasic ≥99.5% (20.4 g/L, Sigma-Aldrich, Germany) was added to the media and
used as a pH buffer for the experiments. Three *D. hansenii* strains (CBS767, IBT22 and IBT26)
were pre-cultured in YNB medium (pH 6) at 28°C for at least 24 h, washed with ddH_2O and
inoculated in the corresponding wells at an initial OD_{600} 0.1. The programmed settings were as
follows: filter; biomass (gain 10 and 3, respectively for the two BioLector® systems), humidity;
on (>85% using ddH_2O), temperatures; 26, 28, and 30 ºC, oxygen supply; 20.85% (atmospheric
air), agitation speed; 1000 rpm. The biomass formation was monitored for at least 70 h.

The light scattering data obtained from the BioLector® I and II systems were analyzed using R
version 3.6.1 (The R Foundation for Statistical Computing).

Metabolite analysis

The concentrations of glucose, glycerol and ethanol were measured by High Performance Liquid
Chromatography (Model 1100-1200 Series HPLC System, Agilent Technologies, Germany). The
injection volume was 20 µl, the eluent 5mM H_2SO_4 and the flow rate was set at 0.6 mL/h. The
temperature of a Bio-Rad Aminex HPX-87H column was kept at 60°C. A standard solution
containing glucose (20 g/L), glycerol (2 g/L) and ethanol (20 g/L) was used (Sigma-Aldrich,
Germany) for exo-metabolites concentration determination.

Off-gas and dissolved oxygen measurements

CO_2 and O_2 concentrations were continuously analyzed in real time by mass spectrometry coupled to the off-gas line (model Prima PRO Process MS, Thermo Scientific, UK). From the off-gas CO_2 emission data, given in percentage, the maximum specific growth rate was calculated. Off-gas CO_2 and O_2 emission data were also used to determine the Carbon Dioxide Evolution Rate (CER), the Oxygen Uptake Rate (OUR) and the Respiratory Quotient (RQ).

Dissolved oxygen values were measured by DO sensors (Model OxyFerm FDA 160, Hamilton) as previously described in "Bioreactor cultivations" section.

Analytical procedures

Specific growth rates in the different growth conditions were calculated based on the optical density (OD_{600}) and emitted CO_2 values.

Yield coefficients and carbon balances were used to describe the metabolite, by-products and biomass formation by the yeast cells, and were calculated based on the DW, accumulated CO_2 and HPLC data. The average minimal formula $CH_{1.79}O_{0.50}N_{0.20}$ for yeast dry cell biomass composition was used for the calculations as proposed by Roels in 1983. The specific glucose consumption rates were calculated based on the logarithmic method proposed in Görgens *et al.* in 2005.

Statistical analysis

Statistical analysis was performed with Microsoft Excel® 2016 (version 1903, 32-bit, USA). All values are represented as averages ± 95% confidence interval of independent biological replicate cultures. For regression analysis, the coefficient of determination (r^2) was used to determine the statistical significance of the fit, where a value above 95% was considered statistically significant.

Results

D. hansenii **growth rate under the effect of high salt concentrations**

Specific growth rates were calculated based on the optical density and the volumetric CO_2-production rates under the different growth conditions tested (Fig. 1). In general terms, a higher maximum specific growth rate was observed when a higher salt concentration is present in the media, except for concentrations of 2M of either salt, compared to control conditions. Sodium exhibit a significantly stronger positive impact than potassium, as the highest growth rate was reached in the presence of 0.5-1M NaCl (similar observations were made for KCl, where concentrations of 0.5M gave slightly higher values than 1M, both resulting in a higher growth rate compared to control, although lower than for NaCl). On the other hand, a decrease in specific growth rate is observed when the concentration of salts is above 1M, and up to 2M (Fig. 1, tables). However, even at concentrations of both sodium and potassium up to 1.5M, the growth rate values are still significantly higher than in control conditions (no salts added).

Surprisingly, at 2M NaCl the growth rate is lower, but still very close to control conditions, although a prolonged lag phase is observed in comparison (exponential growth phase starts around 30-35 hours in the control vs. 40-45 hours for 2M NaCl, as inferred from the CO_2 profiles in Fig.1). In contrast, the addition of 2M KCl results in a lower μ_{max} compared to the 2M NaCl and also lower than the control, reinforcing the fact that NaCl still exerts some beneficial effect in *D. hansenii*'s cell performance overall. Microscope samples were taken, randomly, from the liquid cultures at increasing salt concentrations and also from control cultures (no salts added), and neither morphological changes nor in cell size were appreciated among all the conditions (Suppl. Fig. S1).

The same conclusion can be inferred if we observe the carbon dioxide evolution rate (CER) of *D. hansenii* in the presence of salts over time (Fig. 1). The profiles show a higher CO_2 production (that can be translated into higher glucose consumption, hence a higher metabolic rate) when the cells grow in the presence of NaCl or KCl compared to control conditions with no salt. Once again, this effect begins to decrease when the salt concentrations are over 1.25M, nevertheless still higher than compared to the control conditions, except for 2M in which the production rate is lower than the control for both salts (Fig. 1).

Biomass yield on substrate and biomass titers show a differential effect among K^+ and Na^+

The observed decrease in the specific growth rate from above 1M of salt seems to be compensated by a slightly higher biomass yield upon glucose, as was observed for the dry weight measurements during the cultivation time, although only for the addition of potassium (Table 1). The final biomass titers were not significantly different though, with the exception of the 2M sodium which were slightly higher in comparison (Suppl. Fig. S2).

When the specific glucose consumption rate was calculated, *D. hansenii* showed higher rates of consumption in the presence of NaCl or KCl than in control conditions, being 1M the optimal concentration showing the highest values, and again this effect started to revert once the concentration of salts used was above 1M up to 1.5M, although still higher values than those observed in the control (Table 1), while at 2M NaCl the specific consumption rates were lower than the control. The specific glucose consumption values observed were higher for sodium than for potassium at lower concentrations (0.5M - 1M - 1.25M), as seen before for the specific growth rate, however the opposite is observed at higher concentrations (uptake rates are higher in the presence of potassium).

To further investigate the effect of salts on the specific glucose uptake rates, *D. hansenii* was grown in low glucose conditions (0.2%), preliminarily in shake flasks and later by using bioreactors, to further confirm the findings in flasks. In the shake flask experiments, it was observed that cells growing in the presence of 1M KCl adapted much faster to the nutrient limitation and were able to grow at a significantly higher rate compared to the other conditions tested (Suppl. Fig. S3). This effect was also observed in the presence of 1M NaCl, but at a lower level. Moreover, the presence of 2M of either salt in the medium had no detrimental effect for the cells, which still grew at a similar growth rate compared to the control. Strikingly, the presence of 2M of NaCl resulted in almost 3 times higher biomass concentration than the control

conditions. In this particular circumstance (high salts and low glucose), the effect of potassium
seems to be beneficial in terms of growth rate, but sodium seems to have a better effect in cell
performance in the long run, since the growth rate is slower but the final biomass concentration
is much higher in return (Suppl. Fig. S3). All together seem to confirm, without a doubt, the
halophilic character of *D. hansenii* in one hand, and the differential effect of Na^+ and K^+ salts, in
the other.

Based on these results in shake-flasks, and in order to obtain a more complete dataset that would
support such observations, we then conducted the same type of batch cultures with a reduced
initial glucose concentration (0.2%) in bioreactors without pH regulation (thus mimicking the
previous flask conditions). Our previous observations in flasks cultivations were confirmed,
however interesting differences were observed. The carbon dioxide evolution rates (CER) exhibit
a much faster adaptation (Fig. 2A), evidenced by a shorter lag phase, when cells grow in the
presence of 1M of either salt (NaCl or KCl), which confirms our observations in flasks. Further
analyses showed significantly higher specific glucose uptake rates and, surprisingly, much higher
growth rates at 1M of NaCl or KCl, than compared with the rest of the conditions (Fig. 2B). In
contrast, CO_2 yields and biomass yields are higher at concentrations of 2M, being 2M NaCl the
highest values of all conditions compared. Additional data showing the timeline of DW, OD_{600}
and glucose consumption, are shown in Suppl. Fig. S4.

As observed in the graph, the CER profiles are very similar between control conditions and cells
growing at 2M KCl, only for 2M NaCl the CO_2 production peak seems lower (Fig. 2A), however
when we look at the CO_2 yields on glucose, they are higher for both salts at 2M in comparison to
1M and also with the control. The biomass yields at 2M are also higher than the 1M
concentration, and much higher than the control, as shown in Figure 2B. Altogether, this
confirms our observations in the shake flasks, pointing that on reduced glucose conditions, the
positive effect of the presence of high salt concentration is even more accused. Moderate levels
of salt (1 M) result in higher glucose consumption rates, and higher specific growth rates, while
higher concentration of salts (2M) result in higher biomass yields and CO_2 yields, at the cost of a
lower growth rate. A metabolism switch that can be described as: from growing as fast as
possible to growing as much as possible. We also further confirmed that sodium exerts a more
positive impact than potassium, once again.

It is worth mentioning that the observed final OD values were higher for the control in bioreactor
experiments than in shake flasks. Still we observed a higher biomass when cells are growing in
2M NaCl, as we also saw in the shake flasks, however cell growth in control conditions was
arrested at a lower OD in flasks compared to bioreactors, so the difference in total biomass is less
significant for the latter (Suppl. Fig. S3). This is not surprising though, and simply illustrates the
importance of using well stirred reactors in physiology studies.

Dissolved oxygen concentration and RQ levels confirm a fully respiratory metabolism,
discarding a fermentative process in *D. hansenii* in our experimental conditions

Dissolved oxygen levels during bioreactor cultivations were observed to decrease faster when
Na^+ or K^+ is present in the medium, being those levels lower while the concentration of salt

increases, suggesting a higher oxygen demand (Fig. 3). This confirms a higher metabolic activity
at increasing salt concentrations, which again points to the halophilic character of *D. hansenii*,
and might already suggest a fully respiratory metabolism, regardless of the presence or absence
of (high) salt in the cultivation media. It is worth mentioning that in some of the tested conditions
the dissolved oxygen levels gets very closed to 0, however this does not mean that cells were
suffering from oxygen limitation: the airflow inlet throughout the whole bioreactor cultivation
was constant, so the cells were never deprived from oxygen, as can be seen (Fig. 3) in the very
stable RQ value represented by a constant flat line (RQ = 1) which coincides with the period of
higher oxygen utilization. Thus, the lower dissolved oxygen values simply illustrate a much
higher oxygen utilization by the cells, but never oxygen deprivation (which would show then an
RQ value > 1, and it is not our case).

The Respiratory Quotient (RQ) values calculated over the entire course of the culture, show to be
below or equal to 1 during the exponential growth phase, but never above this value (Fig. 3), this
further supports the absence of fermentation, and therefore confirms that *D. hansenii* is not
producing ethanol from glucose in our conditions, not even in the presence of high salts, as
reported by Calahorra *et al*. (2009). As a final confirmation of the absence of a fermentative
process, our off-gas data and the HPLC analysis show no trace of ethanol neither in the gas phase
nor in the liquid broth samples (Table 1).

Although it worths mentioning that this previous work (Calahorra *et al.* 2009) was performed
using a different *D. hansenii* strain in other culture conditions, and their measurements
correspond to ethanol production using enzyme extracts from previously cultured cells, a follow
up study performed by García-Neto *et al.* (2017) contradicted such observation using the same
strain. Our observations align well with the latter study, confirming that no ethanol production is
occurring in our strain either, thus further proving that *D. hansenii* is crabtree negative.

Additional limiting factors affect cell performance in non-controlled cultivation
environments

In order to test the influence of external pH regulation in the halotolerant / halophilic behavior of
D. hansenii, parallel bioreactor cultivations in normal glucose (2%) were run, in which no pH
control was set, and extracellular pH levels were measured on-line in real time. It was observed
that, when no pH control was exerted, the CO_2 profiles of *D. hansenii* evidenced a long
maintained plateau phase which cannot be seen in pH-controlled fermentations, both in control
conditions and under the effect of salts (Fig. 4). The changes in the external pH over time, for the
different conditions tested, are also shown in Supplementary Fig. S5.

This led us to determine, that no meaningful conclusion about metabolic patterns or behaviour
can be made, with a high degree of accuracy, in non-controlled cultivation environments:
whatever conclusion made is undoubtedly linked to other limiting factors. This means, that
previous studies reporting such conclusions, whose data was obtained from non-controlled
environments (such as shake flasks, for example) must be considered cautiously.

If we have a look at the maximum specific growth rates under no pH regulation, it can be
observed an increase in the μ_{max} when 1M NaCl and KCl are used. This points to a potential

summative effect of low pH and high salt concentrations and, once again, sodium seems to have
a higher positive impact on growth when compared to potassium. This had been already
suggested by Almagro *et al* (2000), so our results further confirm this observation. The specific
growth rates decrease once the concentration of both salts is close to 2 M, as also observed in
pH-controlled experiments, although this time there is a lower growth rate compared to the
control conditions, than when pH control is set in the bioreactors (Table S1 at Supp. material).

Interestingly, the previously described plateau in the CO_2 profiles is not observed for non-pH
controlled bioreactors run with glucose limiting conditions, where sharper and well defined
peaks can be seen in the CO_2 emission profiles (Figure 2A). Here, a faster pH decrease occurs in
optimal growth conditions, corresponding to 1M salt added (either NaCl or KCl) to the media
(Suppl. Fig. S6), compared to the control. This further supports the summative effect of low pH
and increasing salts, to be very beneficial for overall *D. hansenii'*s performance, as already
shown in Figure 2B.

About the type of limitation that is occurring, and that generates the long-plateau phase seen in
the CO_2 emission profiles at normal glucose (2%), and why we do not observe such limitation in
limited sugar (0.2%) we cannot elaborate further with the data that we have. This shall need to be
addressed by future, and more specific, experimental studies.

Spot-tests and micro-fermentation screening analyses confirm the advantages of high
salinity, especially sodium, for the optimal proliferation of *D. hansenii's* strains

In a follow up round of experiments, we aimed to confirm our observations in liquid media for
all the conditions tested (including controlled bioreactors and shake flask cultivations), that is: a
faster growth at medium-high salinity (1M salts) evidenced by higher growth rates, and slower
rate but longer proliferation capacity to higher biomass densities at higher salinity (up to 2M). To
that end, we performed spot tests in solid media plates with increasing salt concentrations
(Figure 5), this time including other *D. hansenii* strains than the model CBS767 to check if our
observations are strain dependent or otherwise if our conclusions can be extrapolated to other
strains of the same species, corresponding to other isolates from other international collections.

As shown in the figure, we confirmed in one hand that the presence of NaCl results more
beneficial than KCl as observed in the previous experiments in liquid media, and on the other
hand that moderate-to-high concentrations (0.5 – 1M) of sodium make yeast grow much faster
than the control, whilst high concentrations (2M) result in slower growth. However in the long
run (240 hours of cultivation), we can observe that the yeast proliferation capacity remains
unaltered, reaching similar cell densities than the control. This further confirms that sodium is
definitely not detrimental at all as it has no inhibitory effect on *Debaryomyces*' proliferation
capacity.

Finally, complementary tests were performed in controlled micro-fermentors (BioLector®)
including the other *D. hansenii* strains, to assess the effect of both salts in combination with other
abiotic parameters such as different pH conditions (4-6-8) and temperatures (26-28-30 °C). The
use of such micro-fermentation systems allows both to have controlled conditions and at the
same time screening several parameters in parallel reactions, thus studying the influence of such

combinations and offering the possibility of identification of behavioural trends among the yeast
strains tested, by using surface-response plots (Figure 6).

As a result, we can observe that although the three *Debaryomyces* strains tested show a
differential behaviour without the presence of salt in the culture media, once 1M of salt is present
we can clearly observe how the cell growth at the different conditions shows a strikingly similar
tendency for all the three strains (Figure 6). One important observation is that, as hypothesized
by the results obtained from the 1L bioreactors, there is a positive correlation between high Na^+
and low pH in terms of cell growth. This finally confirms what was already discussed in the
previous section of this study and previously suggested by Almagro *et al*. (2000), about the
summative effect of low pH and high salt. But the most important fact is that all these
observations are extensive for the different *Debaryomyces* strains tested, hence confirming that
the positive effect of high sodium is not strain dependent, but it can be inferred for *D. hansenii* in
general.

Discussion

The results obtained during this study, indicate that high salt concentrations in the culture media
are indeed needed for optimal levels of cell performance in *Debaryomyces hansenii*. Especially
in the case of NaCl, our results seem to finally confirm the halophilic behavior of this particular
yeast: sodium does not exert detrimental effects over *D. hansenii*, but on the contrary. Another
interesting finding is that NaCl exhibits a more significant positive impact than KCl, already
evidenced in the spot-tests in solid media plates, but especially clear as the optimal growth rates
obtained from the bioreactor experiments are reached in the presence of 1M of NaCl, closely
followed by the values reached at 1M of KCl. This effect is even higher when the glucose source
is scarce, as shown in the limited carbon (0.2%) bioreactor experiments.

There is, however, a longer lag phase when the yeast are grown in the presence of 2M NaCl,
both seen in normal (2%) and limited (0.2%) carbon, which may have led to the previous
conclusion by several other studies, performed in non-controlled environments, that the cell
growth is affected by high sodium concentrations (Capusoni *et al*., 2019). Although these
previous conclusions can of course be due to the fact that those studies have been performed
using different media and another yeast strain different than ours, and in non-controlled
conditions (we actually see in our study that 2M salts in non-controlled conditions might point to
a, mistakenly assessed, growth deficiency as observed in Fig. 4B and 4C), our data additionally
show that the decrease in growth rate is not that dramatic at high salts (2M), and moreover, we
also observed a slightly higher biomass yield in normal carbon, a much higher biomass yields in
nutrient-limiting conditions compared to control conditions, as shown in Figure 2B, that the
previous studies did not observe. This observation was again confirmed by spot-test studies with
increasing salinity in solid media plates, in which we clearly observe that the presence of up to
1M salt (and particularly sodium) results in a faster growth compared with control conditions (no
salt). Moreover, the presence of 2M sodium causes a slower growth compared to control,
however after 240 hours the yeast has grown to the exact same levels than the control, hence
there is no detrimental effect caused by high sodium concentrations: they simply grow slower
and there is no inhibition of growth. Considering all of the above, we propose this as a survival

strategy of *D. hansenii* to prevail in drying environments (e.g. during periods of drought). While
other microbial species chose, as surviving strategy, to enter a dormant state or sporulate while
environmental salinity levels are increasing (as a consequence of the lack of water),
Debaryomyces reacts by increasing μ_{max} at moderate-high salinity levels (over 1M is already
considered toxic for other yeast and bacterial species, so they have difficulties to proliferate (Yan
et al., 2015)), hence increasing the glucose uptake rate (as our data reveals) thus ensuring getting
the most of the available carbon for its survival in detriment to its competitors, and later when
the salt concentration in the media keeps on increasing, slowing down the growth rate while still
proliferating to overpopulate the area, changing the metabolic strategy from growing as fast as
possible to growing as much as possible. This suggested strategy would be in accordance with
the latest publications indicating that *Debaryomyces* are the most represented species in thawing
Arctic glacier samples and coastal environments (Butinar *et al*. 2011; Jaques *et al*. 2014).

The dissolved oxygen levels in the culture medium decrease faster in presence of higher salt
concentration, which suggests a higher metabolic activity. RQ values throughout the whole
cultivation period remain below or equal at a value of 1 (reached constantly during the
exponential phase), clearly supporting that there is no fermentative process, as stated previously.
Finally, no ethanol is observed neither in the exometabolite analysis by HPLC, nor in the
analysis in the off-gas by MS, therefore *D. hansenii* is herewith confirmed a crabtree negative
yeast.

Our data without pH control in the bioreactor vessels indicate additional limiting factors during
the cultivation, based on the CO_2 production profiles, compared with pH controlled cultivations,
evidenced by a long plateau phase, proving that shake flask experiments with non-controlled
environment are not the ideal setup to obtain accurate conclusions about physiological and/or
metabolic parameters, therefore we also suggest that previous studies providing conclusions
obtained by this means, must be taken cautiously. It is also worth mentioning that it is not
appropriate to choose randomly Na^+ salts or K^+ salts in order to study osmotolerance or
halotolerance in general, as our results show that there is a clear differential effect exerted by
either NaCl or KCl, as already suggested by Martinez *et al*. (2011 and 2012).

A very recent study, conducted by Ramos-Moreno *et al*. (2019) almost in parallel to ours,
interestingly hypothesizes about a link between salt stress and oxidative stress response in *D.*
hansenii. They cultured the yeast in the presence of high salt concentrations, reporting an
increase in the expression of ROS detoxification genes (sodium showing a higher effect than
potassium). Our findings align very well with their experimental results, and can complement
their observations adding additional value: our study shows that the presence of salts (specially
sodium) lead to a higher oxygen utilization. A higher respirative metabolism causes the
activation of oxidative stress responsive genes, which would confirm the link between salt stress
and the oxidative stress that Ramos-Moreno *et al*. suggest. In addition, both Ramos-Moreno *et*
al. and our results confirm earlier studies pointing to a protective effect of NaCl against oxidative
aggressions, such as the presence of H_2O_2 (Navarrete *et al.* 2009). In that case, the presence of
NaCl would activate oxidative stress defenses as described above, thus they would be already

active and therefore protective against the further aggression caused by addition of oxygen
peroxide.

All in all, our results shed light upon the behaviour of *D. hansenii* in controlled bioreactor
conditions, presenting this peculiar non-conventional yeast as a strain which is able to perform
very well in "standard" cultivation conditions (no stress added) but whose performance gets
significantly improved when environmental conditions get harsh: high salinity / osmotic
pressure, media acidification and nutrient scarcity, all in combination. This undoubtedly confers
to *D. hansenii* an incredibly strong potential for industrial production setups: those are the
conditions that, upon large scale bioproduction processes (meaning vessels of 1000 litre or
above) the microbial cells will encounter throughout the cultivation process, and that are limiting
the suitable performance of microbes in bioreactors (Takors, R. 2012). Therefore, having such a
strain with the above mentioned behaviour, and more importantly, obtaining sufficient
information for understanding it (and, consequently, being able to take advantage of that
knowledge) is of paramount interest for advancing in the field of cell factory design for industrial
bioprocesses, and therefore for the biotech industry overall.

One of the strongest outcomes of our current study is the possibility of using *D. hansenii* in
culture media containing relatively high salt concentrations, for industrial bioprocesses. On one
hand, there is no need of using pure water sources, which significantly decreases the production
costs as one could take advantage of desalination effluents, or even use directly sea water for the
media composition, while still increasing the production yields (salinity will not affect *D.*
hansenii, but will improve its cell performance, as shown by our research). On the other hand,
using saline environments has another strong advantage which is the reduced risk of
contaminations, hence sterilization costs would also be significantly reduced.

Conclusion

Altogether, our findings reveal the beneficial role of salts, and more particularly sodium, in the
cell performance of *Debaryomyces*, and open the need to further investigate how sodium and
potassium influence the cell metabolism at a molecular level. It is clear that salts are not just
tolerated in *D. hansenii*, but they play a crucial role in its survival strategy, to date
underestimated. The presence of salts is needed for optimal cell performance. Further research,
including a global expression analysis by RNAseq in steady-state continuous bioreactor
cultivations, will shed light upon what are the intracellular mechanisms that trigger such
metabolic changes, and how the discrimination between sodium and potassium occurs to trigger
the different behavioural patterns described within this study. It will also be interesting from a
biotechnological point of view, the identification of molecular elements that could potentially be
responsive to the presence of salts, as our observations suggest that there are undoubtedly
molecular switches which react to the presence or absence of sodium and/or potassium in the
environment, triggering a specific metabolic response.

Acknowledgements

We acknowledge the Novo Nordisk Fonden, within the framework of the Fermentation Based Biomanufacturing Initiative, for supporting this work. LRM received the fellowship "Ayuda de Movilidad Internacional para el Fomento de Tesis con Mención Internacional" from the University of Córdoba (Spain), to carry out her research work at DTU Bioengineering for three months. Prof. José Ramos (University of Córdoba, Spain) and Dr. Markus Bisschops (TU Delft, The Netherlands) are deeply acknowledged for critical reading of this manuscript, as well as Assoc. Prof. Christopher Workman for his assistance with the micro-fermentation data analysis. The authors would also like to thank the Fermentation Core at DTU Bioengineering, and Tina Johansen as well as Martin Nielsen for their technical support. This manuscript has been released as a pre-print at bioRxiv, (Navarrete *et al*.).

Author's contributions

JLM conceived the project. CN, ATF, LRM and MRK designed and performed the experiments. CN, ATF and MRK analyzed the data. CN and JLM wrote and revised the manuscript. All authors read, commented and approved the manuscript.

Competing interests

The authors declare that they have no competing interests.

References

Adler L, Blomberg A, Nilsson A. Glycerol metabolism and osmoregulation in the salt-tolerant yeast *Debaryomyces hansenii*. *J Bacteriol* 1985;**162**: 300-06.

Almagro A, Prista C, Castro S *et al*. Effects of salt on *Debaryomyces hansenii* and *Saccharomyces cerevisiae* under stress conditions. *Int J Food Microbiol* 2000;**56**: 191-97.

Almagro A, Prista C, Benito B *et al*. Cloning and expression of two genes coding for sodium pumps in the salt-tolerant yeast *Debaryomyces hansenii*. *J Bacteriol* 2001;**183**: 3251-55.

Banjara N, Nickerson KW, Suhr MJ *et al*. Killer toxin from several food-derived *Debaryomyces hansenii* strains effective against pathogenic *Candida* yeasts. *Int J Food Microbiol* 2016;**222**: 23-29.

Biswas D, Datt M, Aggarwal M *et al*. Molecular cloning, characterization, and engineering of xylitol dehydrogenase from *Debaryomyces hansenii*. *Appl Microbiol Biotechnol* 2013;**97**: 1613-23.

Breuer U, Harms H. *Debaryomyces hansenii* – an extremophilic yeast with biotechnological potential. *Yeast* 2006;**23**: 415-37.

Brooks AA. Ethanol production potential of local yeast strains isolated from ripe banana peels.
Afr J Biotechnol 2008;**7**: 3749-52.

Butinar L, Strmole T, Gunde-Cimerman N. Relative incidence of *Ascomycetous* yeasts in Arctic
coastal environments. *Microb Eco*l 2011;**61**: 832-43.

Butinar L, Zalar P, Frisvad JC *et al*. The genus Eurotium: members of indigenous fungal
community in hypersaline waters of salterns. *FEMS Microbiol Ecol* 2005;**51**: 155-66.

Calahorra M, Sánchez NS, Peña A. Activation of fermentation by salts in *Debaryomyces*
hansenii. *FEMS Yeast Res* 2009;**9**: 1293-301.

Cano-Garcia L, Belloch C, Flores M. Impact of *Debaryomyces hansenii* strains inoculation on
the quality of slow dry-cured fermented sausages. *Meat Sci* 2014;**96**: 1469-77.

Capusoni C, Arioli S, Donzella S *et al*. Hyper-osmotic stress elicits membrane depolarization
and decreased permeability in halotolerant marine *Debaryomyces hansenii* strains and in
Saccharomyces cerevisiae. *Front Microbiol* 2019;**10**: 64.

Casal M, Paiva S, Queiros O *et al*. Transport of carboxylic acids in yeasts. *FEMS Microbiol Rev*
2008;**32**: 974-94.

Chao HF, Yen YF, Ku MS. Characterization of a salt-induced DhAHP, a gene coding for an
alkyl hydroperoxide reductase, from the extremely halophilic yeast *Debaryomyces hansenii*.
BMC Microbiol 2009;**9**:182.

Dujon B, Sherman D, Fischer G *et al*. Genome evolution in yeasts. *Nature* 2004;**430**: 35-44.

Ferreira D, Nobre A, Silva ML *et al*. XYLH encodes a xylose/H$^+$ symporter from the highly
related yeast species *Debaryomyces fabryi* and *Debaryomyces hansenii*. *FEMS Yeast Res*
2013;**13**: 585-96.

Gadanho M, Almeida JM, Sampaio JP. Assessment of yeast diversity in a marine environment in
the south of Portugal by microsatellite-primed PCR. *Antonie Van Leeuwenhoek* 2003;**84**: 217-27.

García-Neto W, Cabrera-Orefice A, Uribe-Carvajal S *et al*. High osmolarity environments
activate the mitochondrial alternative oxidase in *Debaryomyces hansenii*. *PLOS One* 2017;
12(1):e0169621.

Garcia-Salcedo R, Montiel V, Calero F *et al*. Characterization of *DhKHA1*, a gene coding for a
putative Na(+) transporter from *Debaryomyces hansenii*. *FEMS Yeast Res* 2007;**7**: 905-11.

Gonzalez-Hernandez JC, Cardenas-Monroy CA, Peña A. Sodium and Potassium transport in the
halophilic yeast *Debaryomyces hansenii*. *Yeast* 2004;**21**:403-412.

Görgens JF, Van Zyl WH, Knoetze JH. Reliability of methods for the determination of specific
substrate consumption rates in batch culture. *Biochem Eng J* 2005;**25**: 109-12.

Gunde-Cimerman N, Ramos J, Plemenitas A. Halotolerant and halophilic fungi. *Mycol Res*
2009;**113**: 1231-41.

Gustafsson L, Norkrans B. On the mechanism of salt tolerance: production of glycerol and heat during growth of *Debaryomyces hansenii*. *Arch Microbiol* 1976;**110**: 177-83.

Jacques N, Zenouche A, Gunde-Cimerman N et al. Increased diversity in the genus *Debaryomyces* from Arctic glaciers samples. *Antonie Van Leeuwenhoek* 2015;**107**: 487-501.

Lopez Del Castillo-Lozano M, Delile A, Spinnler HE *et al*. Comparison of volatile sulphur compound production by cheese-ripening yeasts from methionine and methionine–cysteine mixtures. *Appl Microbiol Biotechnol* 2007;**75**: 1447-54.

Martinez JL, Sychrova H, Ramos J. Monovalent cations regulate expression and activity of the Hak1 potassium transporter in *Debaryomyces hansenii*. *Fungal Genet Biol* 2011;**48**: 177-84.

Martinez JL, Luna C, Ramos J. Proteomic changes in response to potassium starvation in the extremophilic yeast *Debaryomyces hansenii*. *FEMS Yeast Res* 2012;**12**: 651-61.

Montiel V, Ramos J. Intracellular Na and K distribution in *Debaryomyces hansenii*: cloning and expression in *Saccharomyces cerevisiae* of *DhNHX1*. *FEMS Yeast Res* 2007;**7**: 102-09.

Navarrete C, Frost AT, Ramos-Moreno L *et al*. A physiological characterization in controlled bioreactors reveals a novel survival strategy for *Debaryomyces hansenii* at high salinity and confirms its halophilic behavior. *bioRxiv* 2020. doi: https://doi.org/10.1101/2020.01.10.901843.

Navarrete C, Siles A, Martínez JL *et al*. Oxidative stress sensitivity in *Debaryomyces hansenii*. *FEMS Yeast Res* 2009;**9**: 582-90.

Norkrans B. Studies on marine occurring yeasts: growth related to pH, NaCl concentration and temperature. *Arch Mikrobiol* 1966;**54**: 374-92.

Papouskova K, Sychrova H. The co-action of osmotic and high temperature stresses results in a growth improvement of *Debaryomyces hansenii* cells. *Int J Food Microbiol* 2007;**118**: 1-7.

Pina C, Couto JA, Hogg T. Inferring ethanol tolerance of *Saccharomyces* and non-*Saccharomyces* yeasts by progressive inactivation. *Biotechnol Lett* 2004;**26**: 1521-27.

Prista C, Almagro A, Loureiro-Dias MC *et al*. Physiological basis for the high salt tolerance of *Debaryomyces hansenii*. *Appl Environ Microbiol* 1997;**63**: 4005-09.

Prista C, Loureiro-Dias MC, Montiel V *et al*. Mechanisms underlying the halotolerant way of *Debaryomyces hansenii*. *FEMS Yeast Res* 2005;**5**: 693-701.

Prista C, Gonzalez-Hernandez JC, Ramos J *et al*. Cloning and characterization of two K^+ transporters of *Debaryomyces hansenii*. *Microbiology* 2007;**153**: 3034-43.

Prista C, Michán C, Miranda IM *et al*. The halotolerant *Debaryomyces hansenii*, the Cinderella of non-conventional yeasts. *Yeast*. 2016;**33**: 523-33.

Ramos J, Melero Y, Ramos-Moreno L *et al*. *Debaryomyces hansenii* strains from Valle de Los Pedroches iberian dry meat products: isolation, identification, characterization, and selection for starter cultures. *J Microbiol Biotechnol* 2017;**27**:1576-85.

Ramos-Moreno L, Ramos J, Michán C. Overlapping responses between salt and oxidative stress
in *Debaryomyces hansenii*. *World J Microbiol Biotechnol*. 2019; **35**:170.

Roels JA. Energetics and kinetics in biotechnology. Amsterdam; New York: Elsevier Biomedical
Press, 1983, 1-330.

Sánchez NS, Calahorra M, Ramírez J *et al*. Salinity and high pH affect energy pathways and
growth in *Debaryomyces hansenii*. *Fungal Biol* 2018;**122**: 977-90.

Takors R. Scale-up of microbial processes: impacts, tools and open questions. *J Biotechnol*
2012;**160**:3-9.

Tavares JM, Duarte LC, Amaral-Collaco MT *et al.* The influence of hexoses addition on the
fermentation of D-xylose in *Debaryomyces hansenii* under continuous cultivation. *Enzyme*
Microb Technol 2000;**26**: 743-47.

Velkova K, Sychrova H. The *Debaryomyces hansenii NHA1* gene encodes a plasma membrane
alkali-metal-cation antiporter with broad substrate specificity. *Gene* 2006;**369**: 27-34.

Yan N, Marschner P, Cao W *et al*. Influence of salinity and water content on soil
microorganisms. *Int Soil and Water Conservation Research* 2015;**3**:316-323.

Table 1. Yield coefficients and specific glucose consumption rates from batch cultivations.

Conditions	Y_{SC}[1]	Y_{SX}[1]	Y_{SE}[1]	Y_{SG}[1]	C-balance[2]	SGCR (r_s)[3]
Control	0.63±0.01	0.35±0.00	ns[4]	ns	0.98±0.01	0.120±0.004
0.5M NaCl	0.57±0.03	0.32±0.02	ns	ns	0.90±0.05	0.157±0.001
1M NaCl	0.63±0.07	0.35±0.04	ns	ns	0.98±0.11	0.163±0.012
1.25M NaCl	0.60±0.02	0.34±0.01	ns	ns	0.94±0.04	0.159±0.002
1.5M NaCl	0.63±0.10	0.35±0.06	ns	ns	0.98±0.16	0.133±0.014
2M NaCl	0.63	0.36	ns	ns	0.99	0.098±0.014
0.5M KCl	0.61±0.00	0.34±0.00	ns	ns	0.95±0.00	0.153±0.009
1M KCl	0.68±0.05	0.38±0.03	ns	ns	1.05±0.08	0.159±0.012
1.25M KCl	0.63±0.05	0.35±0.03	ns	ns	0.98±0.08	0.148±0.004
1.5M KCl	0.68±0.04	0.38±0.02	ns	ns	1.06±0.06	0.144
2M KCl	0.66±0.14	0.37±0.08	ns	ns	1.02±0.21	0.138±0.001

Yield coefficients from batch cultivations of *Debaryomyces hansenii* at different salt concentrations are shown in the table. Additionally, the specific glucose consumption rates during the exponential phase calculated based on the logarithmic method are presented. The cells were grown in synthetic complete medium with or without salt at different concentrations, at 28°C and pH 6. Data shown are mean values ± 95% confidence interval of a number of replicates.

[1] Yield of CO_2 (C), Biomass (X), Ethanol (E), and Glycerol (G) from Glucose (S) in cmol/cmol.
[2] Carbon balance on a C-mol basis, where a value of 1 indicate a closed carbon balance.
[3] Specific Glucose consumption rate in cmol/cmol/h based on logarithmic method.
[4] ns: Not statistically significant. The amount of ethanol and glycerol was neglectable.

 Figure legends

Figure 1. Maximum specific growth rates and carbon dioxide evolution rate (CER) profiles from
batch cultivations with sodium (**A**) or potassium (**B**) chloride. Maximum specific growth rates
determined by optical density (OD) and off-gas CO_2 emission data from batch cultivations of
Debaryomyces hansenii in synthetic complete media at 28°C and pH 6 with varying
concentrations of sodium/potassium chloride are shown in the tables. The CER profiles are based
on off-gas CO_2 emission data over time from one replicate for each condition and are
representative of their associated replicate(s).

Figure 2. Carbon dioxide evolution rate (CER) profiles and yield coefficients, specific glucose
consumption and maximum specific growth rates from batch cultivations under limiting glucose
conditions and without pH regulation. Off-gas CO_2 emission data from batch cultivations of
Debaryomyces hansenii in synthetic complete media containing 0.2% of glucose at 28°C and
initial pH value of 6 with varying concentrations of sodium/potassium chloride are shown in the
figure (**A**). The CER profiles are based on off-gas CO_2 emission data over time from one
replicate for each condition and are representative of their associated duplicate. Yield
coefficients and maximum specific growth rates from batch cultivations of *Debaryomyces*
hansenii at different salt concentrations are shown in the table. Additionally, the specific glucose
consumption rates during the exponential phase calculated based on the logarithmic method are
presented (**B**). The cells were grown in synthetic complete medium containing 0.2% of glucose
and with or without salt at different concentrations, at 28°C and initial pH of 6. Data shown are
mean values ± 95% confidence interval of duplicates.

Figure 3. Dissolved oxygen (DO) and respiratory quotient (RQ) profiles from batch cultivations
with sodium (**A**) or potassium (**B**) chloride. Dissolved oxygen (%) levels measured over time (h)
in synthetic complete media at 28°C and pH 6 with varying concentrations of sodium/potassium
chloride, are represented in the figure. The DO profiles are based on one replicate for each
condition and are representative of their associated replicate(s). The RQ profiles are based on
OUR and CER values calculated from off-gas O_2 consumption and off-gas CO_2 emission data
from one replicate for each condition and are representative of their associated replicate(s).

Figure 4. Graphical comparison of CER profiles with and without pH regulation for batch
cultivations with varying concentrations of salts. The CER profiles are based on off-gas CO_2
emission data over time from one replicate for each condition and are representative of their
associated replicate(s). The cells were grown in synthetic complete media at 28°C with and
without pH regulation (pH 6 with regulation). The cells were either grown without salts (**A**),
NaCl (**B**) or KCl (**C**).

Figure 5. Growth assay of *D. hansenii* strains CBS767, IBT22 and IBT26 in YNB plates (pH 6)
control and supplemented with different concentrations of KCl or NaCl. Cells were grown in
flasks without salt for 24 h, and cell suspensions were prepared from those cultures. Cell drops at
OD_{600} 0.1 and serial dilutions (1:10, 1:100) were spotted on the plates. Growth was monitored
for 144 h, or 240 h when higher salt concentrations were used.

Figure 6. Maximum specific growth rates (μmax h^{-1}) from micro-fermentations of *D. hansenii*
strains CBS767, IBT22, and IBT26 in YNB control medium and supplemented with 1M NaCl
(both with 2% glucose) as a function of pH and temperature (ºC). The surface-response plots
were generated using R.

Figure 6

Figure 1

A

Maximum specific growth rates (μ_{max}).

Conditions	μ_{max} - OD[1]	μ_{max} - CO_2[1]
Control	0.103±0.006	0.105±0.004
0.5M NaCl	0.162±0.013	0.185±0.007
1M NaCl	0.170±0.005	0.176±0.000
1.25M NaCl	0.168±0.006	0.165±0.005
1.5M NaCl	0.150±0.008	0.143±0.017
2M NaCl	0.106±0.005	0.101±0.000

Note: Data shown are mean values ± 95% confidence interval of a number of replicates.
[1] The maximum specific growth rates (h^{-1}) on glucose determined by OD_{600} and off-gas CO_2.

B

Maximum specific growth rates (μ_{max}).

Conditions	μ_{max} - OD[1]	μ_{max} - CO_2[1]
Control	0.103±0.006	0.105±0.004
0.5M KCl	0.169±0.002	0.179±0.004
1M KCl	0.153±0.014	0.165±0.001
1.25M KCl	0.131±0.002	0.162±0.004
1.5M KCl	0.119±0.037	0.119±0.050
2M KCl	0.086±0.009	0.092±0.003

Note: Data shown are mean values ± 95% confidence interval of a number of replicates.
[1] The maximum specific growth rates (h^{-1}) on glucose determined by OD_{600} and off-gas CO_2.

Figure 2

A)

B)

Yield coefficients, specific glucose consumption and maximum specific growth rates from batch cultivations in glucose limiting conditions and without pH regulation.

Conditions	Y_{SC}[1]	Y_{SX}[1]	SGCR (r_s)[2]	μ_{max}[3]
Control	0.37±0.06	0.21±0.03	0.16±0.02	0.177±0.024
1M NaCl	0.53±0.03	0.30±0.02	0.38±0.13	0.214±0.008
1M KCl	0.51±0.07	0.29±0.04	0.46±0.03	0.214±0.005
2M NaCl	0.61±0.03	0.34±0.02	0.13±0.01	0.200±0.009
2M KCl	0.65±0.05	0.36±0.03	0.14±0.00	0.167±0.028

[1] Yield of CO_2 (C) and Biomass (X) from Glucose (S) in cmol/cmol.
[2] Specific Glucose consumption rate in cmol/cmol/h based on logarithmic method.
[3] The maximum specific growth rates (h^{-1}) on glucose determined by off-gas CO_2.

Figure 3

A

B

Figure 4

Figure 5