Contrasting population genetics of cattle- and buffalo- derived *Theileria annulata* causing tropical theileriosis

Umer Chaudhry[1‡]*, **Qasim Ali**[2‡], **Lynn Zheng**[1], **Imran Rashid**[3], **Muhammad Zubair Shabbir**[3], **Muhammad Nauman**[4], **Kamran Ashraf**[2], **Mike Evans**[1], **Shahzad Rafiq**[3], **Muhammad Oneeb**[3], **Ivan Morrison**[1], **Neil D. Sargison**[1]*

[1] University of Edinburgh, Royal (Dick) School of Veterinary Studies and Roslin Institute, UK
[2] Gomal University, Dera Ismail Khan, Khyber Pakhtunkhwa, Pakistan
[3] University of Veterinary and Animal Sciences Lahore, Pakistan
[4] Disease Diagnostic Laboratory, Livestock and Dairy Department, Okara, Pakistan

‡ Contributed equally

*Corresponding authors:

Umer Chaudhry
Email: uchaudhr@exseed.ed.ac.uk, Tel: 00441316519244
University of Edinburgh, The Roslin Institute, Easter Bush Veterinary Centre, UK, EH25 9RG

Neil Sargison
Neil.Sargison@ed.ac.uk, +44 (0)131 651 7300
University of Edinburgh, R(D)SVS, Easter Bush Veterinary Centre, UK, EH25 9RG

Abstract

The present study was designed to improve understanding of *Theileria annulata* in sympatric

water buffalo and cattle in the Punjab province of Pakistan. The prevalence of tropical theileriosis

is high, buparvaquone resistance is widespread, and vaccine protection is poor in the field. Better

understanding is, therefore, needed of the factors that influence the genetics of *T. annulata*

populations both within its hosts and in its overall populations. Here we utilise a panel of six

satellites and a mitochondrial cytochrome b marker to explore the multiplicity of *T. annulata*

infection and patterns of emergence and spread of different parasite genotypes. Parasite materials

were collected from infected animals in defined regions, where water buffalo and cattle are kept

together. Our results show that *T. annulata* is genetically more diverse in cattle- than in water

buffalo-derived populations (the mean numbers of unique satellite alleles were 13.3 and 1.8 and

numbers of unique cytochrome b locus alleles were 65 and 27 in cattle- and water buffalo- derived

populations, respectively). The data show a high level of genetic diversity among the individual

host-derived populations (the overall heterozygosity (H_e) indices were 0.912 and 0.931 in cattle,

and 0.874 and 0.861 in buffalo, based on satellite and cytochrome b loci, respectively). When

considered in the context of high parasite transmission rates and frequent animal movements

between different regions, the predominance of multiple *T. annulata* genotypes, with multiple

introductions of infection in the hosts from which the parasite populations were derived, may have

practical implications for the spread of parasite genetic adaptations; such as those conferring

vaccine cross-protection against different strains affecting cattle and buffalo, or resistance to

antiprotozoal drugs.

Key Words: Tropical theileriosis, *T. annulata*, buparvaquone resistance, antiprotozoal drugs.

1. Introduction

Vector-borne haemoprotozoa impact globally on the health, welfare, and production of livestock (Jabbar et al., 2015). The genus *Theileria* includes two particularly important species, *Theileria annulata* and *Theileria parva* (Lawrence, 1979). *T. parva* typically causes East Coast Fever, Corridor Disease, or January Disease in southern and eastern Africa (Uilenberg et al., 1982), whereas, *T. annulata* causes tropical theileriosis in North Africa and South Asia (Nourollahi-Fard et al., 2015). Tropical theileriosis is amongst the most important neglected tropical parasitic diseases of livestock (Sivakumar et al., 2014). Cattle and buffalo become infected with *T. annulata* following the transmission of sporozoites by ixodid ticks of the genus *Hyalomma*. These stages invade lymphocytes and develop into schizonts, which activate the infected cell to proliferate. Synchronous division of parasite and host cell results in rapid clonal expansion of the infected cell population. Some of the parasites undergo differentiation to produce merozoites, which are released from the lymphocytes enter erythrocytes where they develop to piroplasms. Piroplasms ingested during tick feeding, migrate to the arthropod gut where gametogeny and production of zygotes occurs. These enter the haemolymph and are carried to the salivary gland, where further intracellular development gives rise to sporozoites. The life cycle is completed when the next instar of the tick feeds (Gharbi and Darghouth, 2015). Clinically, infected cattle and buffalo show high fever and lymphadenopathy, sometimes accompanied by haemolytic anaemia, respiratory and ocular lesions (Al-Hosary et al., 2010; Mahmmod et al., 2011).

Measures used to control tropical theileriosis include avoidance, of infestation by *Hyalomma* ticks, including the application of acaricides, and prophylactic use of the theilericidal drug. However, these measures are expensive and difficult to apply. The unsustainability of these disease control measures highlights the potential for effective *T. annulata* vaccines. Attenuated live vaccines against *T. annulata* have been used in many countries where tropical theileriosis is endemic (Brown, 1990; Darghouth et al., 1999; Tait and Hall, 1990). However, there are practical constraints to the widespread use of live *T. annulata* vaccines in control programmes: (i) a requirement for distribution in liquid nitrogen, which accounts for approximately 30% of the cost (Bouslikhane et al., 1998); (ii) the need to use the vaccine immediately after thawing; (iii) the need for costly quality control measures to ensure consistent efficacy and the absence of other pathogens; (iv) in some cases, problems with post-vaccination reactions, which have been recorded in 3% of animals immunised with Chinese, Moroccan, Iranian or Tunisian stocks (Darghouth, 2008). To overcome these problems, research on the development of a killed subunit

vaccine against *T. annulata* has focused on antigens present on the surface of sporozoite and
merozoite stages of the parasite.

Genetic and antigenic diversity is a major constraint to the development of subunit vaccines for
some parasite species. For example, cross-protection between cattle- and buffalo-derived *T. parva*
strains is incomplete (Young et al., 1973). Studies of *T. parva* in African buffalo have shown a
high level of genetic diversity when compared to cattle-maintained *T. parva* (Oura et al., 2005).
While *T. annulata* can infect both cattle and Asian buffalo (Nourollahi-Fard et al., 2015), genetic
comparisons between strains infecting sympatric host populations have not been reported.
Understanding the genetic and antigenic diversity of *T. annulata* in buffalo and cattle and the
extent to which the parasites circulate between the two species is needed to inform the potential
applicability of vaccination.

Therapeutic control of *T. annulata* is heavily dependent on the use of a single theilericidal drug,
namely buparvaquone. However, the effective use of this compound in some regions is now
threatened by the emergence of drug resistance (Mhadhbi et al., 2015). Buparvaquone resistance
has been reported in several countries and now represents a serious challenge to efficient livestock
production (Mhadhbi et al., 2010). Although mutations in the candidate cytochrome b locus linked
to resistance phenotype have been demonstrated, a causal relationship has not been established
(Chatanga et al., 2019; Mhadhbi et al., 2015; Sharifiyazdi et al., 2012). Previous studies have
described multiple genotypes and high levels of genetic diversity in *T. annulata* (Al-Hamidhi et al.,
2015; Weir et al., 2011), although there is a lack of information on the genotypic diversity of drug-
resistant parasites and whether there are fitness costs associated with mutations of the cytochrome
b locus. It is, therefore, important to understand the genetic diversity and population sub-structure
of *T. annulata*, with reference to the emergence and spread of buparvaquone resistance mutations.

Population genetic studies of *T. annulata* have previously been performed using panels of
satellite DNA markers (Weir et al., 2007). More recently, deep amplicon sequencing of meta-
barcoded DNA has afforded a practical and high-throughput method for investigating the genetic
diversity between and within parasite populations. The Illumina MiSeq platform can provide
100,000 or more reads of up to 600 bp of loci of interest, depending on primer design. Recently,
we have used these methods to study the population genetics of *Calicophoron daubneyi* (Sargison
et al., 2019) and *Fasciola gigantica* infection in livestock keep in the United Kingdom and
Pakistan.

In this paper, we describe the results of an analysis of the genetic diversity of buffalo- and

cattle-derived *T. annulata,* by applying typing with six polymorphic satellite markers, combined

with the deep amplicon sequencing of the mitochondrial cytochrome b locus. The results provide a

comparison of the molecular profile of parasites in the two host species, which indicate greater

genetic diversity in cattle parasites. These data represent an essential basis for further analyses to

examine the emergence and spread of buparvaquone resistance in *T. annulata*.

2. Materials and Methods

2.1. Parasite resources, gDNA isolation and species identification

Four infected cell lines produced by the infection of blood mononuclear cells with laboratory-

maintained stocks of *T. annulata* held in the University of Edinburgh, originally isolated in Turkey,

India, Tunisia, and Morocco (Katzer et al., 1994), were used as reference controls. Blood samples

were collected from the piroplasm-positive cattle and buffalo in veterinary clinics throughout the

Punjab province of Pakistan between 2017 and 2019. Venous blood samples were collected into

EDTA tubes were stored at -20°C. Samples were collected by para-veterinary staff under the

supervision of local veterinarians following consent from the animal owners. The study was

approved by the Institutional Review Board of the University of Veterinary and Animal Sciences

(UVAS-24817). Peripheral blood smears were prepared and stained with 4% Giemsa, and

examined microscopically to detect piroplasms. Genomic DNA was isolated from positive

samples by lysis with GS buffer containing proteinase K as described in the TIANamp Blood

DNA Kit (TIANGEN Biotech Co. Ltd, Beijing) and stored at -20°C. 'Haemoprotobiome' high-

throughput sequencing (Chaudhry et al., 2019) was performed on piroplasm-positive blood

samples to confirm the presence of *T. annulata.*

2.2. Adapter and barcoded PCR amplification of T. annulata cytochrome b

A 517 bp region of the cytochrome b locus was selected for deep amplicon sequencing. One µl

gDNA of each of 31 buffalo, 54 cattle and the 4 positive control derived *T. annulata* samples was

used as templates for the 1st round adapter PCR amplification. A *de novo* primer set

(Supplementary Table S1) was used under the following conditions: 13.25 µl ddH$_2$O, 1 µl gDNA,

0.75 µl of 10 mM dNTPs mix, 5 µl of 5X HiFi Fidelity Buffer, 0.5 µl of 0.5U DNA polymerase

enzyme (KAPA Biosystems, USA), and 0.75 µl of 10 µM forward adaptor primers and reverse

adaptor primers. The thermocycling conditions were: 95°C for 2 min, 35 cycles at 98°C for 20 sec,

55°C for 15 sec, and 72°C for 2 min, followed by a final extension of 72°C for 2 min. The PCR

product was purified with AMPure XP Magnetic Beads (1X) (Beckman coulter Inc., USA). A

barcoded primer set was used in the 2nd round of PCR amplification to add a fragment of unique

sequence into each purified product (Supplementary Table S2) under the following conditions:

13.25 µl ddH$_2$O, 2 µl bead purified PCR products, 0.75 µl of 10 mM dNTPs mix, 5 µl of 5X HiFi

Fidelity Buffer, 0.75 µl of 0.5U DNA polymerase enzyme (KAPA Biosystems, USA), and 1.25 µl

of 10µM forward and reverse primers. The thermocycling conditions were: 98°C for 2 min, 7

cycles at 98°C for 20 sec, 63°C for 20 sec, and 72°C for 2 min. The PCR products were purified

with AMPure XP Magnetic Beads (1X) (Beckman coulter Inc., USA).

2.3. Illumina Mi-Seq run and data handling

A pooled library was prepared with 10 µl of barcoded bead purification PCR product from each

T. annulata sample and sent to Edinburgh Genomics, UK for deep amplicon sequencing. The size

of the amplicon was measured by qPCR library quantification (KAPA Biosystems, USA), before

running on an Illumina Mi-Seq sequencer using a 600-cycle pair-end reagent kit (Mi-Seq Reagent

Kits v2, MS-103-2003) at a concentration of 15 nM with addition of 15% Phix control v3

(Illumina, FC-11-2003). The Illumina Mi-Seq post-run processing uses the barcoded indices to

split all sequences by sample and generate FASTQ files. These were analysed using Mothur

v1.39.5 software (Schloss et al., 2009) with modifications in the standard operating procedures of

Illumina Mi-Seq (Kozich et al., 2013) in the Command Prompt pipeline. Briefly, the raw paired

read-ends were run into the 'make.contigs' command to combine the two sets of reads for each

sample. The command extracted sequence and quality score data from the FASTQ files, creating

the complement of the reverse and forward reads, and then joining the reads into contigs. After

removing the too long, or ambiguous sequence reads, the data were aligned with the *T. annulata*

cytochrome b reference sequence library prepared from the positive controls (for more details

Supplementary Data S1 and section 2.1) using the 'align.seqs' command. Any sequences that did

not match with the *T. annulata* cytochrome b reference library were removed and the

'summary.seqs' command was used to summarise the 517 bp sequence reads of the *T. annulata*

cytochrome b locus. The sequence reads were further run on the 'screen.seqs' command to

generate the *T. annulata* cytochrome b FASTQ file. Once the sequence reads were classified as *T.*

annulata, a count list of the consensus sequences of each population was created using the

'unique.seqs' command. The count list was further used to create FASTQ files of the consensus
sequences of each population using the 'split.groups' command (for more details Supplementary
Data S2).

2.4. Bioinformatics data analysis

The consensus sequences of *T. annulata* cytochrome b locus were aligned using the MUSCLE
alignment tool in Geneious v10.2.5 software (Biomatters Ltd, New Zealand) and then imported
into the FaBox 1.5 online tool (birc.au.dk) to collapse the sequences that showed 100% base pair
similarity after corrections into a single genotype. The genotype frequency of each sample was
calculated by dividing the number of sequence reads by the total number of reads. A split tree was
created in the SplitTrees4 software (bio-soft.net) by using the neighbour-joining method in the
JukesCantor model of substitution. The appropriate model of nucleotide substitutions for
neighbour-joining analysis was selected by using the jModeltest 12.2.0 program (Posada, 2008).
The tree was rooted with the corresponding cytochrome b sequence of *T. parva*. The branch
supports were obtained by 1000 bootstraps of the data. The genetic diversity of cytochrome b was
calculated within and between populations by using the DnaSP 5.10 software package (Librado
and Rozas, 2009), and the following values were obtained: Heterozygosity (H_e), the number of
segregating sites (S), nucleotide diversity (π), the mean number of pairwise differences (k), the
mutation parameter based on an infinite site equilibrium model, and the mutations parameter
based on segregating sites (S_θ).

2.5. Satellite genotyping and bioinformatics data analysis of T. annulata

Six satellite markers (TS6, TS8, TS20, TS31, TS12, TS16) were selected for use on each
population as previously described by Weir et al. (2007). A summary of primer sequences and
allele ranges is given in Supplementary Table S3. Eighteen buffalo and 35 cattle derived *T.*
annulata populations were successfully analysed from six markers. PCR amplification was
performed using a master mix containing 17.2 µl ddH$_2$O, 0.3 µl of 100 µM dNTPs, 2.5 µl of 1X
thermopol reaction buffer, 0.3 µl of 1.25U Taq DNA polymerase (New England Biolabs) and 0.3
µl of 0.1µM forward and reverse primers. The thermo-cycling parameters were 94^{o}C for 2 min
followed by 35 cycles of 94^{o}C for 1 min, staying at the annealing temperatures of 60^{o}C (TS6,
TS20), 55^{o}C (TS8, TS12, TS16), or 50^{o}C (TS31) for 1 min and 65^{o}C for 1 min, with a single final
extension cycle of 65^{o}C for 5 min. The forward primer of each microsatellite primer pair was 5'

end labelled with fluorescent dye (IDT, UK), and the ROX 400 internal size standard was used on

the ABI Prism 3100 genetic analyser (Applied Biosystems, USA).

Individual chromatograms were analysed using Peak Scanner software version 2.0 (Thermo

Fisher Scientific, USA) to determine the size of the alleles. These were combined across six

markers to generate a multilocus genotype (MLG). From the MLG data, allelic variation (A) and

heterozygosity (H_e) for individual populations were calculated using Arlequin 3.11 software (Guo

and Thompson, 1992). Significance levels were calculated using the sequential method of

Bonferroni correction for multiple comparisons in the same dataset (Rice, 1989). Analysis of

molecular variance (AMOVA) was estimated through the partition of satellite diversity between

and within populations (Excoffier et al., 1992) and fixation index (pairwise F_{ST}) values were

calculated using Arlequin 3.11 to provide a measurement of population genetic sub-structure.

Principal coordinate analysis (PCoA) was performed using GenALEx software to illustrate the

extent of genetically distinct features of individual populations with plot coordinates (Peakall and

Smouse, 2012).

The likelihood ratio test statistics (G-test) were calculated using Arlequin 3.11 software

(Excoffier et al., 2005) to estimate genetic linkage equilibrium and deviations from Hardy-

Weinberg equilibrium. There was no evidence to support linkage disequilibrium for any

combination of satellite loci in the individual *T. annulata* populations of buffalo and cattle,

indicating that alleles at these loci were randomly associating and not genetically linked (data on

file). There was some significant departure from Hardy-Weinberg equilibrium, even after

Bonferroni correction, in addition to relative P values for 108 loci combinations for buffalo and

210 loci combinations for cattle *T. annulata* populations. The bottleneck software (version 1.2.02)

was, therefore, used to search for the evidence of heterozygosity excess and mode-shift. The

Wilcoson signed-rank test was used to evaluate the statistical significance of any possible genetic

drift equilibrium (Lefterova et al., 2015). The data showed that there was no heterozygosity excess

according to the Sign Test and Wilcoxon Test (data on file). The mode shift analysis established

most populations had a normal L-shaped distribution.

3. Results

3.1. Allelic diversity between cattle- and buffalo- derived T. annulata populations

The allelic diversity data show that *T. annulata* is more diverse in cattle as compared to water

buffalo. The mean numbers (±SD) of satellite alleles in 35 cattle- and 18 buffalo- derived

populations were 34.7 (±9.1) and 23.7 (±5.2), respectively. The number of alleles (A_n) for each

satellite marker in the cattle- and buffalo-derived populations ranged from 22 to 46 and 17 to 32,

respectively. The mean numbers (±SD) of cattle- and buffalo-specific *T. annulata* satellite alleles

(A_u) per marker were 13.3 (±6.3) and 1.8 (±2.04), respectively (Table 1 and Fig. 1). Seventy-nine

and 41 cytochrome b locus alleles (A_n) were identified in the 54 cattle- and 31 buffalo-derived *T.*

annulata populations. The numbers of cattle- and buffalo-specific *T. annulata* alleles (A_u) were 65

and 27, respectively (Table 1 and Fig. 1). The high level of allelic variation described in *T.*

annulata might have practical implications for the development of a sporozoite and merozoite

antigen subunit vaccine and immunisation cross-protection.

3.2. Genetic diversity within T. annulata populations derived from individual cattle and buffalo

The heterozygosity (H_e) data reveal high levels of genetic diversity within each individual

populations (35 cattle and 18 buffalo) of *T. annulata*; with mean values (±SD) for the six satellite

markers ranging from 0.8 (±0.082) to 0.959 (±0.017) in cattle- and 0.738 (±0.196) to 0.956

(±0.034) in buffalo-derived populations (Table 2). Overall, the mean heterozygosities (H_e) (±SD)

of the cattle- and buffalo-derived *T. annulata* populations were 0.912 (±0.056) and 0.874 (±0.028)

respectively (Table 2). High levels of heterozygosity (H_e) were seen at the *T. annulata* cytochrome

b locus, ranging from 0.556 to 0.923 (overall mean value 0.931) in each of the 54 cattle-derived

populations and from 0.545 to 0.900 (overall mean value 0.861) in each of the 31 buffalo-derived

populations (Table 3). The high level of genetic diversity in *T. annulata* may have practical

implications for the emergence of buparvaquone drug resistance mutations.

3.3. Genetic sub-structure and phylogenetic relationship between T. annulata populations derived

from individual cattle and buffalo

Genetic differentiation was shown by comparing the fixation indices (F_{ST}) of two populations

with each other in a pairwise manner. The F_{ST} values indicated a low level of genetic

differentiation between *T. annulata* ranging from 0.001 to 0.114 in each of the 35 cattle-derived

populations and 0.003 to 0.132 in each of the 18 buffalo-derived populations, respectively,

respectively (Supplementary Table S4). An AMOVA was conducted using the panel of six

satellite markers to estimate the genetic variation within and between the populations. This

showed that genetic variation was distributed 97.91% and 97% and within; and 2.09% and 3%

between cattle- and buffalo-derived *T. annulata* populations, respectively (data on file). PCoA was

performed to illustrate as a two-dimensional plot the extent to which populations are genetically distinct. The two axes accounted for 72.5% (25.67 + 46.83) and 73.93% (27.6 + 46.33) of the variation in the cattle- and buffalo-derived populations, respectively; and showed that populations from different regions formed overlapping clusters, hence were not geographically sub-structured (Fig. 2).

Phylogenetic analysis of the seventy-nine cytochrome b genotypes of the 54 cattle-derived *T. annulata* populations, showed that two genotypes for 22.9% and 10.8% of the total number of the sequence reads, being present in 36 and 8, populations, respectively. Seven genotypes accounted for between 5.0% and 7.0% of the sequence reads and 45 genotypes accounted for less than 1.0% of the sequence reads (Fig. 3A). The split tree shows that eight genotypes are shared between populations derived from cattle holdings in multiple locations of Lahore, Chakwal, Gujranwala, Okara, and Sahiwal. Seven genotypes are shared between populations derived from cattle holdings in any two locations of Lahore, Gujranwala, Okara, and Sahiwal. Sixty-one genotypes are present in a single location (Gujranwala=24; Qadirabad=12; Okara =14; Lahore=11 genotypes) (Fig. 3A).

Phylogenetic analysis of the forty-one cytochrome b genotypes of the 31 buffalo-derived *T. annulata* populations, showed that three genotypes accounted for 39.1%, 10.8%, and 5.5% of sequence reads being present in 26, 15 and 3 populations, respectively. Eighteen genotypes accounted for between 1.0% and 3.0% of sequence reads and 20 genotypes accounted for less than 1.0% of sequence reads (Fig. 3B). The split tree shows that two genotypes are shared between *T. annulata* populations derived from buffalo holdings in multiple locations of Gujranwala, Hafizabad, Okara, and Sahiwal. Five genotypes are present in the populations derived from buffalo holdings in Gujranwala and Okara. One genotype is present in the populations derived from buffalo holdings in Hafizabad and Okara and one genotype from Okara and Sahiwal. Thirty-two genotypes are present in a single location (16 genotypes in Gujranwala; 13 genotypes in Okara; and 3 in Hafizabad) (Fig. 3B). The satellite and cytochrome b data provide evidence of a high level of gene flow predicted to occur due to livestock movements or translocations of ticks to a new region. This could potentially influence the spread of drug-resistant alleles.

3.4. Genotype distribution of T. annulata populations derived from individual cattle and buffalo

T. annulata satellite data were found to be highly polymorphic in each individual population (35 cattle and 18 buffalo), with overall numbers of genotypes per population ranging from 2 to 26

in cattle and 2 to 17 in buffalo. The mean numbers (±SD) of *T. annulata* genotypes in cattle- and
buffalo-derived populations were 8.60 (±2.52) and 6.25, respectively (Table 4).

Seventy-nine *T. annulata* cytochrome b genotypes from 54 individual cattle-derived
populations were analysed separately. Fifteen populations had a single genotype at high
frequencies ranging from 90 to 100%. These comprised of ten populations containing only one
genotype; two populations containing two genotypes; one population with three genotypes; one
population with four genotypes; and one population with seven genotypes (Fig. 4A). Thirty-nine
populations had multiple genotypes at high frequencies. These comprised of five populations
containing two genotypes; eight populations containing three genotypes; ten population containing
four genotypes; seven populations with five genotypes; four populations with six genotypes; one
population with seven genotypes; two populations with eight genotypes; one populations with 12
genotypes; and one with 14 genotypes (Fig. 4A).

Forty-one *T. annulata* cytochrome b genotypes from 31 individual buffalo-derived populations
were analysed separately. A single genotype predominated in 15 populations at a frequency of
between 90 and 100%. These comprised of eight populations containing only one genotype; three
populations containing two genotypes; one population with three genotypes; one population with
five genotypes; one population with six genotypes; and one population with seven genotypes (Fig.
4B). Sixteen populations had high frequencies of multiple genotypes. These comprised of seven
populations contained 2 genotypes; two populations contained 3 genotypes; one population
contained 4 genotypes; three populations had 5 genotypes; two populations had 7 genotypes and
one population contained a maximum of 8 genotypes (Fig. 4B). The satellite and cytochrome b
data provide evidence of the potential impact of the multiplicity of *T. annulata* infection and their
control strategies for tropical theileriosis.

4. Discussion

Theleria annulata is considered to be the most economically important protozoan parasite of
cattle and water buffalo in South Asia, the Middle East and North Africa, where livestock
production is critically important. *T. annulata* has a major impact on food production in low- and
middle-income countries (LMICs), where efficient agriculture is a priority and the disease causes
economic losses due to high mortality and morbidity of livestock, having a significant impact on
meat and milk production (Jabbar et al., 2015). Our results reveal a high level of genetic diversity
within *T. annulata* infecting individual hosts, and between different host-derived populations.

Although both cattle and buffalo parasite populations showed extensive diversity, the extent of

diversity was significantly greater in cattle compared to buffalo. The diversity and distribution of

Theileria annulata genotypes in the Punjab province of Pakistan are consistent with high parasite

transmission rates and frequent animal movements within the region.

The satellite and cytochrome b data reported in the present study show that *T. annulata* is

genetically more diverse, with more circulation of alleles, in cattle as compared to water buffalo.

The satellite data were informative due to the numbers of alleles present, showing more unique

alleles in cattle-derived than in Asian water buffalo-derived *T. annulata* populations. The

cytochrome b locus, although genotypically less polymorphic, provided further evidence of allelic

diversity and differences between cattle- and buffalo- populations. Similarly, high levels of

parasite heterogeneity have been reported in *T. parva*, although the diversity in cattle is much

more restricted compared to the buffalo host (Oura et al., 2011). In *T. parva*, there is evidence

that although most if not all buffalo-derived parasites are capable of infecting cattle, many are

incapable of undergoing tick transmission. Hence, differences in the capacity of different *T.*

annulata genotypes to undergo tick transmission in cattle and buffalo could potentially influence

the genotypic composition of the parasites that are maintained in the two host species.

Immunisation of cattle with different subunit versions of the sporozoite stage surface protein

(SPAG1) has provided a degree of protection against cattle-derived *T. annulata*, probably by

neutralising the ability of the parasite to infect host cells (Schnittger et al., 2002). Given the high

overall level of genetic diversity in *T. annulata*, it will be important to determine whether there is

polymorphism in candidate antigens that would influence the ability to generate broadly cross-

reactive immunity (Katzer et al., 1994).

Our satellite and cytochrome b locus data shows a high levels of genetic diversity in both

cattle- and buffalo-derived *T. annulata* populations, reflecting high effective population sizes (Al-

Hamidhi et al., 2015). *T. annulata* genome comprises 8.35 Mb spanning four chromosomes that

range from 1.9 to 2.6 Mb, with 3,792 putative protein-coding genes. In addition, a total of 49

transfer RNA and five ribosomal RNA genes were identified (Pain et al. (2005). High level of

genetic diversity in *T. annulata* rise the possibility of the emergence of potential drug resistance

mutations. Pakistani livestock is treated frequently with buparvaquone and anecdotal evidence

suggests that resistance is widespread, implying that there is likely to be strong selective pressure

for the emergence of drug resistance. The importance of this point is emphasised by the fact that

buparvaquone is the only drug available for the treatment of tropical theileriosis.

Fifteen cytochrome b genotypes were shared between the *T. annulata* populations of cattle and
nine were shared between cattle populations. The satellite data provide further evidence of low
levels of genetic differentiation among cattle-and buffalo-derived populations and show
overlapping clusters that are not geographically sub-structured, consistent with the high levels of
gene flow due to livestock movements or translocations of ticks to a new region. This could
influence the spread of drug-resistant alleles. Studies of the global genetic sub-structure of *T.*
annulata have shown a high level of genetic differentiation between Turkey and Tunisia (Weir et
al., 2007). Studies have also shown a low level of genetic differentiation in *T. annulata* within-
countries including Turkey, Tunisia, Oman, China and Portugal (Al-Hamidhi et al., 2015; Gomes
et al., 2016; Weir et al., 2011; Yin et al., 2018), reflecting high levels of animal movement, or
translocation of ticks.

The presence of a single cytochrome b genotype in 15 cattle and 9 buffalo- derived populations
suggests a single emergence of *T. annulata* infection. In contrast, the predominance and the high
proportions of multiple cytochrome b genotypes in 39 cattle- and 16 buffalo-derived populations
implies multiple emergences of infection. The numbers of satellite genotypes per population
provide further evidence for the pattern of the multiplicity of infection in genetically different
adapted strains. Random cross mating of gametes and genetic recombination of *T. annulata* in
ticks gives rise to the formation of new genotypes (Al-Hamidhi et al., 2015). The multiplicity of *T.*
annulata infection may be influenced by variations in the intensity of transmission due to levels of
different tick species infestation, or prevalence of tick infection (Yin et al., 2018). *Hyalomma*
scupense is the most common and economically important species and the major vector of *T.*
annulata, but *Hyalomma marginatum* and *Hyalomma anatolicum* are also found in Tunisia
(Bouattour et al., 1996). There are four *Hyalomma* species responsible for the transmission of *T.*
annulata in Turkey, of which *H. anatolicum* is the major vector for *T. annulata* (Aktas et al.,
2004). Two tick species, *Hyalomma lusitanicum*, and *H. marginatum* are the vectors of *T.*
annulata in Portugal (Estrada-Pena and Santos-Silva, 2005). There are a few reports that *H.*
anatolicum may be a vector of *T. annulata* in the Punjab province of Pakistan (Karim et al., 2017).
The multiplicity of *T. annulata* infection may also be influenced by population bottlenecking
effects, arising from seasonal effects of climatic conditions on tick transmission and completion of
the parasite's lifecycle. The *T. annulata* transmission season in Turkey is between May and
September, with the peak of clinical cases occurring in mid-summer (Sayin et al., 2003). In a
region of endemic stability of tropical theileriosis in Tunisia, there is a high level of multiplicity

of *T. annulata* infection, but clinical disease is rare; while in a region of endemic instability, a
proportion of the population becomes infected all year round, and the clinical disease occurs
particularly in adult cattle (Gharbi et al., 2011).

5. Conclusions

Control strategies for tropical theileriosis need to consider factors such as: the reproductive
isolation of parasite populations; management of host movements; control of tick vectors;
mitigation of the impacts of climate change; and consequences of parasite exposure to
antiprotozoal drugs. Understanding of the multiplicity of infection and of high levels of gene flow
is, therefore, important in the educational dissemination and implementation of advice on
sustainable parasite control.

442
443

Acknowledgement

The study was financially supported by the Carnegie Trust Scotland and Biotechnology and
Biological Sciences Research Council (BBSRC). Work at the University of Veterinary and
Animal Science Pakistan uses facilities funded by the Higher Education Commission of Pakistan.
The authors of this study would like to thank Vice-Chancellor of the University of Veterinary and
Animal Science Lahore Pakistan, for his great support in the arrangements for sample collections.

Conflict of interest

None
453

454 **References**
455
Aktas, M., Dumanli, N., Angin, M., 2004. Cattle infestation by Hyalomma ticks and prevalence of
Theileria in Hyalomma species in the east of Turkey. Vet Parasitol 119, 1-8.
Al-Hamidhi, S., M, H.T., Weir, W., Al-Fahdi, A., Johnson, E.H., Bobade, P., Alqamashoui, B., Beja-
Pereira, A., Thompson, J., Kinnaird, J., Shiels, B., Tait, A., Babiker, H., 2015. Genetic
Diversity and Population Structure of Theileria annulata in Oman. PLoS One 10, e0139581.
Al-Hosary, A., Abdel-Rady, A., Ahmed, L.S., Mohamed, A., 2010. Comparison between Using of
BUPAQUONE ® and Other Compounds in Treatment of Bovine Theileriosis. International
Journal for Agro Veterinary and Medical Sciences 4, 3-7.
Bouattour, A., Darghouth, M.A., Ben Miled, L., 1996. Cattle infestation by Hyalomma ticks and
prevalence of Theileria in H. detritum species in Tunisia. Vet Parasitol 65, 233-245.

Bouslikhane, M., M., K., H., O. 1998. La theilériose bovine au Maroc, investigations
épidémiologiques et étude de l'impact sur la productivité des élevages. In: Résumé de
communication orale, 15 ème congrès vétérinaire maghrébin.
Brown, C.G., 1990. Control of tropical theileriosis (Theileria annulata infection) of cattle.
Parassitologia 32, 23-31.
Chatanga, E., Mosssad, E., Abdo Abubaker, H., Amin Alnour, S., Katakura, K., Nakao, R., Salim, B.,
2019. Evidence of multiple point mutations in Theileria annulata cytochrome b gene
incriminated in buparvaquone treatment failure. Acta Trop 191, 128-132.
Chaudhry, U., Ali, Q., Rashid, I., Shabbir, M.Z., Abbas, M., Numan, M., Evans, M., Ashraf, K.,
Morrison, I., Morrison, L., Sargison, N.D., 2019. Development of a deep amplicon
sequencing method to determine the proportional species composition of piroplasm
haemoprotozoa as an aid in their control. bioRxiv.
Darghouth, M.A., 2008. Review on the experience with live attenuated vaccines against tropical
theileriosis in Tunisia: considerations for the present and implications for the future.
Vaccine 26 Suppl 6, G4-G10.
Darghouth, M.A., Bouattour, A., Kilan, M., 1999. Tropical theileriosis in Tunisia: epidemiology and
control. Parassitologia 41 Suppl 1, 33-36.
Estrada-Pena, A., Santos-Silva, M.M., 2005. The distribution of ticks (Acari: Ixodidae) of domestic
livestock in Portugal. Experimental & applied acarology 36, 233-246.
Excoffier, L., Laval, G., Schneider, S., 2005. Arlequin (version 3.0): an integrated software package
for population genetics data analysis. Evolutionary bioinformatics online 1, 47-50.
Excoffier, L., Smouse, P.E., Quattro, J.M., 1992. Analysis of molecular variance inferred from
metric distances among DNA haplotypes: application to human mitochondrial DNA
restriction data. Genetics 131, 479-491.
Gharbi, M., Darghouth, M.A., 2015. Control of tropical theileriosis (Theileria annulata infection in
cattle) in North Africa. Asian Pacific Journal of Tropical Disease 5, 505-510.
Gharbi, M., Touay, A., Khayeche, M., Laarif, J., Jedidi, M., Sassi, L., Darghouth, M.A., 2011.
Ranking control options for tropical theileriosis in at-risk dairy cattle in Tunisia, using
benefit-cost analysis. Revue scientifique et technique (International Office of Epizootics)
30, 763-778.
Gomes, J., Salgueiro, P., Inacio, J., Amaro, A., Pinto, J., Tait, A., Shiels, B., Pereira da Fonseca, I.,
Santos-Gomes, G., Weir, W., 2016. Population diversity of Theileria annulata in Portugal.
Infect Genet Evol 42, 14-19.
Guo, S.W., Thompson, E.A., 1992. Performing the exact test of Hardy-Weinberg proportion for
multiple alleles. Biometrics 48, 361-372.
Jabbar, A., Abbas, T., Sandhu, Z.U., Saddiqi, H.A., Qamar, M.F., Gasser, R.B., 2015. Tick-borne
diseases of bovines in Pakistan: major scope for future research and improved control.
Parasites & vectors 8, 283.
Karim, S., Budachetri, K., Mukherjee, N., Williams, J., Kausar, A., Hassan, M.J., Adamson, S., Dowd,
S.E., Apanskevich, D., Arijo, A., Sindhu, Z.U., Kakar, M.A., Khan, R.M.D., Ullah, S., Sajid,
M.S., Ali, A., Iqbal, Z., 2017. A study of ticks and tick-borne livestock pathogens in Pakistan.
PLoS neglected tropical diseases 11, e0005681.
Katzer, F., Carrington, M., Knight, P., Williamson, S., Tait, A., Morrison, I.W., Hall, R., 1994.
Polymorphism of SPAG-1, a candidate antigen for inclusion in a sub-unit vaccine against
Theileria annulata. Mol Biochem Parasitol 67, 1-10.

Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K., Schloss, P.D., 2013. Development of a
dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence
data on the MiSeq Illumina sequencing platform. Appl. Environ. Microbiol. 79, 5112-5120.
Lawrence, J.A., 1979. The differential diagnosis of the bovine theilerias of Southern Africa.
Journal of the South African Veterinary Association 50, 311-313.
Lefterova, M.I., Budvytiene, I., Sandlund, J., Farnert, A., Banaei, N., 2015. Simple Real-Time PCR
and Amplicon Sequencing Method for Identification of Plasmodium Species in Human
Whole Blood. J Clin Microbiol 53, 2251-2257.
Librado, P., Rozas, J., 2009. DnaSP v5: a software for comprehensive analysis of DNA
polymorphism data. Bioinformatics 25, 1451-1452.
Mahmmod, Y.S., Elbalkemy, F.A., Klaas, I.C., Elmekkawy, M.F., Monazie, A.M., 2011. Clinical and
haematological study on water buffaloes (Bubalus bubalis) and crossbred cattle naturally
infected with Theileria annulata in Sharkia province, Egypt. Ticks Tick Borne Dis 2, 168-
171.
Mhadhbi, M., Chaouch, M., Ajroud, K., Darghouth, M.A., BenAbderrazak, S., 2015. Sequence
Polymorphism of Cytochrome b Gene in Theileria annulata Tunisian Isolates and Its
Association with Buparvaquone Treatment Failure. PloS one 10, e0129678.
Mhadhbi, M., Naouach, A., Boumiza, A., Chaabani, M.F., BenAbderazzak, S., Darghouth, M.A.,
2010. In vivo evidence for the resistance of Theileria annulata to buparvaquone. Vet
Parasitol 169, 241-247.
Nourollahi-Fard, S.R., Khalili, M., Ghalekhani, N., 2015. Detection of Theileria annulata in blood
samples of native cattle by PCR and smear method in Southeast of Iran. Journal of
parasitic diseases : official organ of the Indian Society for Parasitology 39, 249-252.
Oura, C.A., Asiimwe, B.B., Weir, W., Lubega, G.W., Tait, A., 2005. Population genetic analysis and
sub-structuring of Theileria parva in Uganda. Mol Biochem Parasitol 140, 229-239.
Oura, C.A., Tait, A., Asiimwe, B., Lubega, G.W., Weir, W., 2011. Haemoparasite prevalence and
Theileria parva strain diversity in Cape buffalo (Syncerus caffer) in Uganda. Vet Parasitol
175, 212-219.
Pain, A., Renauld, H., Berriman, M., Murphy, L., Yeats, C.A., Weir, W., Kerhornou, A., Aslett, M.,
Bishop, R., Bouchier, C., Cochet, M., Coulson, R.M., Cronin, A., de Villiers, E.P., Fraser, A.,
Fosker, N., Gardner, M., Goble, A., Griffiths-Jones, S., Harris, D.E., Katzer, F., Larke, N.,
Lord, A., Maser, P., McKellar, S., Mooney, P., Morton, F., Nene, V., O'Neil, S., Price, C.,
Quail, M.A., Rabbinowitsch, E., Rawlings, N.D., Rutter, S., Saunders, D., Seeger, K., Shah, T.,
Squares, R., Squares, S., Tivey, A., Walker, A.R., Woodward, J., Dobbelaere, D.A., Langsley,
G., Rajandream, M.A., McKeever, D., Shiels, B., Tait, A., Barrell, B., Hall, N., 2005. Genome
of the host-cell transforming parasite Theileria annulata compared with T. parva. Science
309, 131-133.
Peakall, R., Smouse, P.E., 2012. GenAlEx 6.5: genetic analysis in Excel. Population genetic
software for teaching and research--an update. Bioinformatics 28, 2537-2539.
Posada, D., 2008. jModelTest: phylogenetic model averaging. Molecular biology and evolution 25,
1253-1256.
Rice, W., 1989. Analyzing tables of statistical tests. Evolution 43.
Sargison, N.D., Shahzad, K., Mazeri, S., Chaudhry, U., 2019. A high throughput deep amplicon
sequencing method to show the emergence and spread of Calicophoron daubneyi rumen
fluke infection in United Kingdom cattle herds. Veterinary parasitology.

Sayin, F., Dincer, S., Karaer, Z., Cakmak, A., Inci, A., Yukari, B.A., Eren, H., Vatansever, Z.,
Nalbantoglu, S., 2003. Studies on the epidemiology of tropical theileriosis (Theileria
annulata infection) in cattle in Central Anatolia, Turkey. Tropical animal health and
production 35, 521-539.
Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., Lesniewski, R.A.,
Oakley, B.B., Parks, D.H., Robinson, C.J., 2009. Introducing mothur: open-source,
platform-independent, community-supported software for describing and comparing
microbial communities. Appl. Environ. Microbiol. 75, 7537-7541.
Schnittger, L., Katzer, F., Biermann, R., Shayan, P., Boguslawski, K., McKellar, S., Beyer, D., Shiels,
B.R., Ahmed, J.S., 2002. Characterization of a polymorphic Theileria annulata surface
protein (TaSP) closely related to PIM of Theileria parva: implications for use in diagnostic
tests and subunit vaccines. Molecular and biochemical parasitology 120, 247-256.
Sharifiyazdi, H., Namazi, F., Oryan, A., Shahriari, R., Razavi, M., 2012. Point mutations in the
Theileria annulata cytochrome b gene is associated with buparvaquone treatment failure.
Vet Parasitol 187, 431-435.
Sivakumar, T., Hayashida, K., Sugimoto, C., Yokoyama, N., 2014. Evolution and genetic diversity of
Theileria. Infect Genet Evol 27, 250-263.
Tait, A., Hall, F.R., 1990. Theileria annulata: control measures, diagnosis and the potential use of
subunit vaccines. Rev Sci Tech 9, 387-403.
Uilenberg, G., Perie, N.M., Lawrence, J.A., de Vos, A.J., Paling, R.W., Spanjer, A.A., 1982. Causal
agents of bovine theileriosis in southern Africa. Tropical animal health and production 14,
127-140.
Weir, W., Ben-Miled, L., Karagenc, T., Katzer, F., Darghouth, M., Shiels, B., Tait, A., 2007. Genetic
exchange and sub-structuring in Theileria annulata populations. Mol Biochem Parasitol
154, 170-180.
Weir, W., Karagenc, T., Gharbi, M., Simuunza, M., Aypak, S., Aysul, N., Darghouth, M.A., Shiels, B.,
Tait, A., 2011. Population diversity and multiplicity of infection in Theileria annulata. Int J
Parasitol 41, 193-203.
Yin, F., Liu, Z., Liu, J., Liu, A., Salih, D.A., Li, Y., Liu, G., Luo, J., Guan, G., Yin, H., 2018. Population
Genetic Analysis of Theileria annulata from Six Geographical Regions in China,
Determined on the Basis of Micro- and Mini-satellite Markers. Frontiers in genetics 9, 50.
Young, A.S., Brown, C.G., Burridge, M.J., Cunningham, M.P., Kirimi, I.M., Irvin, A.D., 1973.
Observations on the cross-immunity between Theileria lawrencei (Serengeti) and
Theileria parva (Muguga) in cattle. Int J Parasitol 3, 723-728.
590

591 **Figure Legends**

592 **Fig. 1.** The mean number of alleles in *T. annulata* derived from buffalo and cattle based on a panel
593 of six satellite and a cytochrome b marker. The bar of each marker shows the proportion of alleles
594 in buffalo and cattle. The Y-axis shows the number of alleles of each marker. The unique and
595 shared alleles of buffalo and cattle are represented by a different colour.
596
597 **Fig. 2.** Principal coordinate analysis using a panel of six satellite markers to represent 35 cattle-
598 and 18 buffalo-derived *T. annulata* populations.
599

Fig. 3. Split tree analysis of 79 cattle-derived (3A) and 41 buffalo-derived (3B) *T. annulata* genotypes collected from the Lahore (A), Gujranwala (B), Chakwal (C), Qadirabad (D), Okara (E), Sahiwal (F), and Hafizabad (G) regions of the Punjab province of Pakistan. Each population (CY) is represented by a different colour in the individual genotype (CHA and BHA). The pie chart circles represent the distribution and percentage of sequence reads generated per genotype identified in 54 cattle- and 31 buffalo-derived populations, as indicated in the insert table.

Fig. 4. Relative genotype frequencies of the cytochrome b locus of 54 cattle-derived (4A) and 31 buffalo-derived (4B) *T. annulata* populations, collected from the Punjab province of Pakistan. Each genotype (CHA and BHA) is represented by a different colour in the individual population (CY). The population distribution and the frequency of sequence reads generated per population for each of the 79 cattle-derived and 41 buffalo-derived *T. annulata* genotypes is shown in the insert table.

Principal Coordinates Analysis

(A) Cattle

Coordinate 2 varience: 25.67

Coordinate 1 varience: 46.83

pop = CY2
pop = CY6
pop = CY79
pop = CY82
pop = CY89
pop = CY84
pop = CY20
pop = CY90
pop = CY26
pop = CY27
pop = CY28
pop = CY30
pop = CY34
pop = CY39
pop = CY38
pop = CY40
pop = CY41
pop = CY37
pop = CY160
pop = CY162
pop = CY154
pop = CY131
pop = CY130
pop = CY120
pop = CY119
pop = CY134
pop = CY150
pop = CY173
pop = CY124
pop = CY175
pop = CY177
pop = CY136
pop = CY151
pop = CY145
pop = CY33

Principal Coordinates Analysis

(B) Buffalo

Coordinate 2 varience: 27.60

Coordinate 1 varience: 46.33

pop = CY7
pop = CY16
pop = CY92
pop = CY109
pop = CY157
pop = CY103
pop = CY15
pop = CY98
pop = CY129
pop = CY142
pop = CY141
pop = CY138
pop = CY148
pop = CY126
pop = CY149
pop = CY149
pop = CY139
pop = CY133

(A) Cattle

Genotypes	Sequence reads	Number of populations
CHA1	306152 (22.9)	36 (CY1, CY2, CY3, CY6, CY20, CY25, CY26, CY27, CY28, CY30, CY31, CY32, CY34, CY35, CY79, CY80, CY82, CY83, CY84, CY86, CY87, CY88, CY89, CY120, CY130, CY130, CY131, CY134, CY135, CY136, CY140, CY145, CY153, CY154, CY158, CY162, CY163)
CHA7	144834 (10.8)	8 (CY6, CY36, CY37, CY38, CY39, CY40, CY41, CY42)
CHA2	93391 (6.9)	27 (CY1, CY2, CY3, CY20, CY25, CY27, CY28, CY30, CY31, CY32, CY33, CY35, CY79, CY82, CY83, CY87, CY88, CY89, CY119, CY120, CY131, CY134, CY135, CY140, CY145, CY177)
CHA23	80938 (6.0)	7 (CY36, CY37, CY38, CY39, CY40, CY41, CY42)
CHA35	70229 (5.2)	3 (CY42, CY151, CY160)
CHA65	84581 (6.3)	3 (CY119, CY124, CY175)
CHA47	73571 (5.5)	2 (CY90, CY150)
CHA8	92376 (6.9)	5 (CY20, CY26, CY38, CY42, CY120)
CHA66	80068 (5.9)	5 (CY119, CY131, CY134, CY135, CY173)
CHA43	12608 (0.9)	2 (CY84, CY86)
CHA74	11807 (0.8)	1 (CY150)
CHA48	10559 (0.7)	1 (CY90)
CHA40	10464 (0.7)	5 (CY79, CY84, CY88, CY90, CY162)
CHA75	13825 (1.0)	1 (CY150)
CHA5	7836 (0.5)	9 (CY6, CY33, CY35, CY79, CY84, CY89, CY120, CY136, CY177)
CHA4	6560 (0.4)	10 (CY2, CY20, CY27, CY29, CY79, CY82, CY130, CY131, CY140, CY145)
CHA41	5724 (0.4)	1 (CY83)
CHA49	6472 (0.4)	1 (CY90)
CHA27	7923 (0.5)	2 (CY39, CY40)
CHA44	6823 (0.4)	3 (CY84, CY89, CY145)
CHA11	4346 (0.3)	1 (CY26)
CHA37	5130 (0.3)	1 (CY79)
CHA13	4519 (0.3)	3 (CY26, CY41, CY86)
CHA55	4388 (0.3)	2 (CY90, CY162)
CHA34	5381 (0.4)	1 (CY42)
CHA19	5243 (0.3)	2 (CY32, CY89)
CHA28	5018 (0.3)	2 (CY39, CY40)
CHA68	4006 (0.2)	1 (CY136)
CHA50	3964 (0.2)	1 (CY90)
CHA12	3922 (0.2)	2 (CY26, CY86)
CHA20	4758 (0.3)	1 (CY33)
CHA51	4746 (0.3)	1 (CY90)
CHA67	3729 (0.2)	2 (CY130, CY145)
CHA76	4689 (0.3)	1 (CY150)
CHA77	5621 (0.4)	1 (CY150)
CHA52	3594 (0.2)	1 (CY90)
CHA24	4588 (0.3)	1 (CY37)
CHA14	4674 (0.3)	2 (CY27, CY28)
CHA9	3536 (0.2)	1 (CY20)
CHA10	4516 (0.3)	1 CY20)
CHA3	5489 (0.4)	1 (CY2)
CHA53	3476 (0.2)	1 (CY90)
CHA26	4474 (0.3)	2 (CY38, CY41)
CHA69	5449 (0.4)	1 (CY136)
CHA16	4441 (0.3)	1 (CY29)
CHA33	3397 (0.2)	6 (CY40, CY86, CY89, CY120, CY131, CY136)
CHA15	3373 (0.2)	2 (CY27, CY28)
CHA17	4354 (0.3)	1 (CY30)
CHA54	5347 (0.4)	1 (CY90)
CHA56	4315 (0.3)	1 (CY90)
CHA45	5308 (0.3)	2 (CY87, CY90)
CHA39	4273 (0.3)	4 (CY79, CY82, CY87, CY119)
CHA18	5225 (0.3)	1 (CY32)
CHA78	4223 (0.3)	1 (CY150)
CHA38	3210 (0.2)	1 (CY79)
CHA57	3510 (0.2)	1 (CY90)
CHA25	2206 (0.1)	1 (CY37)
CHA70	2200 (0.1)	1 (CY136)
CHA72	1994 (0.1)	1 (CY145)
CHA29	1888 (0.1)	1 (CY39)
CHA58	2186 (0.1)	1 (CY90)
CHA59	2985 (0.1)	1 (CY90)
CHA60	2185 (0.1)	1 (CY90)
CHA22	1674 (0.1)	1 (CY34)
CHA61	1873 (0.1)	1 (CY90)
CHA73	2162 (0.1)	1 (CY145)
CHA62	2459 (0.1)	1 (CY90)
CHA63	2647 (0.1)	1 (CY90)
CHA42	2143 (0.1)	2 CY83, CY84)
CHA36	2539 (0.1)	1 (CY42)
CHA30	1835 (0.1)	1 (CY39)
CHA31	1933 (0.1)	1 (CY39)
CHA6	2132 (0.1)	1 (CY6)
CHA71	2627 (0.1)	1 (CY136)
CHA79	2526 (0.1)	1 (CY177)
CHA32	2845 (0.1)	1 (CY39)
CHA46	1515 (0.1)	1 (CY87)
CHA64	1408 (0.1)	1 (CY90)
CHA21	1805 (0.1)	1 (CY33)

(B) Buffalo

Genotypes	Sequence reads	Number of populations
BHA1	278937 (39.1)	26 (CY91, CY92, CY96, CY98, CY103, CY105, CY107, CY110, CY111, CY112, CY113, CY117, CY123, CY126, CY128,CY129, CY133, CY138, CY139, CY149, CY157, CY7, CY8, CY15, CY16, CY19)
BHA2	77171 (10.8)	15 (CY19, CY98, CY113, CY117, CY128, CY129, CY133, CY138, CY148, CY149, CY7, CY8, CY15, CY16, CY19)
BHA16	39389 (5.5)	3 (CY109, CY128, CY142)
BHA31	14779 (2.0)	1 (CY129)
BHA23	19109 (2.6)	1 (CY123)
BHA17	18232 (2.5)	1 (CY109)
BHA34	19535 (2.7)	3 (CY138, CY8, CY19)
BHA35	19437 (2.7)	1 (CY138)
BHA24	20260 (2.8)	1 (CY123)
BHA25	10998 (1.5)	1 (CY123)
BHA4	12882 (1.8)	6 (CY91, CY133, CY138, CY141, CY148, CY19)
BHA36	14721 (2.0)	1 (CY138)
BHA11	13670 (1.9)	2 (CY98, CY133)
BHA18	15501 (2.1)	1 (CY109)
BHA38	12485 (1.7)	1 (CY19)
BHA33	14428 (2.0)	1 (CY133)
BHA37	12365 (1.7)	1 (CY148)
BHA15	8310 (1.1)	3 (CY107, CY117, CY126)
BHA41	7283 (1.0)	1 (CY94)
BHA3	8268 (1.1)	2 (CY91, CY138)
BHA28	6217 (0.8)	2 (CY128, CY129)
BHA39	7212 (1.0)	1 (CY19)
BHA40	5199 (0.7)	1 (CY19)
BHA12	4192 (0.5)	1 (CY98)
BHA13	5192 (0.7)	1 (CY98)
BHA20	4169 (0.5)	1 (CY110)
BHA9	5151 (0.7)	1 (CY92)
BHA27	3925 (0.5)	1 (CY128)
BHA19	3024 (0.4)	1 (CY109)
BHA32	3117 (0.4)	2 (CY129, CY157)
BHA29	4110 (0.5)	1 (CY128)
BHA5	3102 (0.4)	1 (CY91)
BHA30	2790 (0.3)	1 (CY128)
BHA10	2078 (0.2)	1 (CY96)
BHA6	2378 (0.3)	1 (CY91)
BHA7	2477 (0.3)	1 (CY91)
BHA21	2869 (0.4)	1 (CY110)
BHA14	1264 (0.1)	1 (CY98)
BHA26	1864 (0.2)	2 (CY123),
BHA22	1657 (0.2)	1 (CY128)
BHA8	1954 (0.2)	1 (CY91)

(A) Cattle

Populations	Sequence reads	Genotypes (%)
CY29	6442	CHA16(99.2),CHA4(0.2)
CY80	2997	CHA1(100)
CY124	3450	CHA65(100)
CY151	2032	CHA35(100)
CY153	4391	CHA1(100)
CY154	3483	CHA1(100)
CY158	1514	CHA1(100)
CY160	3621	CHA35(100)
CY163	2842	CHA1(100)
CY173	3393	CHA66(100)
CY175	3798	CHA65(1 00)
CY6	7361	CHA1(95.2), CHA5(3.0), CHA6(1.7), CHA7(0.1)
CY25	12422	CHA1(98.7), CHA2(1.2)
CY35	3780	CHA1(90.8), CHA2(3.4), CHA5(4.7)
CY136	109536	CHA1(97.6), CHA68(0.9), CHA69(0.4), CHA5(0.6), CHA33(0.1), CHA70(0.1), CHA71(0.1)
CY1	3055	CHA1(89), CHA2(11)
CY2	8032	CHA1(77.9), CHA2(10.6), CHA3(6.0), CHA4(5.3)
CY3	3573	CHA1(82.7), CHA2(17.2)
CY20	66615	CHA1 (60.8), CHA2(29.2), CHA4(6.9), CHA8(1.3), CHA9(0.8), CHA10(0.7)
CY26	8670	CHA1(69.6), CHA11(27.0), CHA12(1.6), CHA13(1.4), CHA8(0.1)
CY27	15741	CHA1(89.7),CHA2(6.4), CHA14(2.2), CHA15(1.5), CHA4(0.6)
CY28	9559	CHA1(90.3),CHA2(5.9), CHA14(2.3), CHA15(1.3)
CY30	8168	CHA1(81.1), CHA2(11.5), CHA17(6.3)
CY31	4219	CHA2(62.2), CHA1(37.7)
CY32	2304	CHA1(80.6), CHA2(8.4), CHA18(9.7), CHA19(1.1)
CY33	2162	CHA2(33.2), CHA5(43.6), CHA20(19.4), CHA21(4.5)
CY34	13252	CHA1(86.1), CHA22(13.8)
CY36	4812	CHA7(67.6), CHA23(32.3)
CY37	47941	CHA7(75.6), CHA23(22.6), CHA24(1.2), CHA25(0.4)
CY38	4812	CHA7(75.3), CHA23(24.0), CHA26(0.5), CHA8(0.2)
CY39	45065	CHA7(51.1), CHA23(42.0), CHA27(4.0), CHA28(1.4), CHA29(0.4), CHA30(0.2), CHA31(0.2), CHA32(0.2)
CY40	29245	CHA7(48.8), CHA23(44.0), CHA27(5.8), CHA28(1.2), CHA33(0.3)
CY41	77853	CHA7(83.7), CHA23(15.7), CHA13(0.1), CHA26(0.3)
CY42	57452	CHA7(62.4), CHA23(32.4), CHA34(4.3), CHA35(0.3), CHA36(0.2), CHA8(0.1)
CY79	48294	CHA1(85.4), CHA37(5.3), CHA2(4.6), CHA5(3.1), CHA4(0.7), CHA38(0.4), CHA39(0.1), CHA40(0.4)
CY82	23106	CHA1(79.2), CHA2(17.6), CHA4(2.8), CHA39(0.2)
CY83	64110	CHA1(90.8), CHA41(8.8), CHA2(0.1), CHA42(0.1)
CY84	77806	CHA1(86.3),CHA43(10.2), CHA5(2.0), CHA44(1.3), CHA40(0.3), CHA42(0.3)
CY86	60995	CHA1(85.0), CHA43(11.4), CHA13(2.1), CHA12(1.2), CHA33(0.1)
CY87	9526	CHA1(90.9), CHA45(4.1), CHA2(3.5), CHA39(0.1), CHA46(1.2)
CY88	5518	CHA1(89.1), CHA2(10.2), CHA40(0.5)
CY89	114872	CHA1(86.2), CHA2(11.5), CHA19(1.0), CHA44(0.7), CHA5(0.3), CHA33(0.1)
CY90	64051	CHA47(55.7), CHA48(16.4), CHA40(14.3), CHA49(5.4), CHA50(1.5), CHA51(1.1), CHA52(0.9), CHA53(0.7), CHA54(0.5), CHA55(0.5), CHA56(0.4), CHA57(0.3), CHA58(0.2), CHA59(0.2), CHA60(0.2), CHA61(0.2), CHA62(0.2), CHA63(0.2), CHA64(0.1), CHA45(0.1)
CY119	80952	CHA65(57.2), CHA66(23.2), CHA2(19.3), CHA39(0.1)
CY120	55003	CHA1(76.6), CHA2(21.4), CHA5(1.5), CHA33(0.7), CHA8(0.1)
CY130	26053	CHA1(78.8), CHA2(15.8), CHA67(0.7)
CY131	20896	CHA1(74.8), CHA2(24.9), CHA66(0.4), CHA4(0.1), CHA33(0.1)
CY134	3929	CHA66(68.3), CHA1(21.3),CHA2(10.3)
CY135	2083	CHA1(90.3), CHA66(9.5), CHA2(0.9)
CY140	4043	CHA1(50.3), CHA2(43.1), CHA4(6.5)
CY145	63270	CHA1(81.1), CHA2(15.7), CHA44(1.3), CHA67(0.8), CHA4(0.4), CHA72(0.3), CHA73(0.2)
CY150	25039	CHA74(48.1), CHA75(32.2), CHA47(7.4), CHA76(5.7), CHA77(5.4), CHA78(0.8)
CY162	3576	CHA1(34.8), CHA40(35.4),CHA55(29.6)
CY177	3917	CHA2(55.8), CHA5(40.1), CHA79(3.9)

(B) Buffalo

Populations	Sequence reads	Genotypes (%)
CY94	22831	BHA41(100)
CY103	14412	BHA1(100)
CY105	12823	BHA1(100)
CY111	51382	BHA1(100)
CY112	15822	BHA1(100)
CY139	18352	BHA1(100)
CY141	14892	BHA41(100)
CY142	12272	BHA16(100)
CY107	12712	BHA1(99), BHA15(0.3)
CY157	13062	BHA1(97.8), BHA32(2.1)
CY7	14892	BHA1(98.7), BHA2(1.2)
CY8	13682	BHA1(93.4), BHA2(2.9), BHA2(3.5)
CY91	29533	BHA1(95.1), BHA2(2.6), BHA3(0.5), BHA4(0.6), BHA6(0.2), BHA7(0.2), BHA8(0.1)
CY98	23295	BHA1(92.7), BHA2(4.0), BHA11(1.2), BHA12(0.8), BHA13(0.8), BHA14(0.2)
CY110	38681	BHA1(92.3), BHA20(4.3), BHA21(1.7), BHA22(1.4)
CY92	12361	BHA1(87.7), BHA9(12.2)
CY96	16971	BHA1(88.8), BHA10(11.9)
CY109	16723	BHA16(82.9), BHA17(13.3), BHA18(2.9), BHA19(0.7)
CY113	14081	BHA1(82.1), BHA2(17.8)
CY117	13502	BHA1(28), BHA15(69.8), BHA2(2.2)
CY123	43438	BHA1(87.5), BHA23(7.1),BHA24(2.9), BHA25(2.3), BHA26(0.1)
CY126	14262	BHA1(75.1), BHA15(15.2)
CY128	55010	BHA1(75.1), BHA2(21.7), BHA16(2.3), BHA26(0.3), BHA27(0.2), BHA28(0.2), BHA29(0.2), BHA30(0.1)
CY129	15470	BHA31(87.3), BHA1(9.7), BHA2(0.1), BHA28(1.9), BHA32(0.9)
CY133	49042	BHA1(79.7), BHA2(18.6), BHA4(0.2), BHA11(0.7), BHA33(0.8)
CY138	70250	BHA1(83.0), BHA2(13.2), BHA3(0.1), BHA4(0.2), BHA34(0.5), BHA35(2.0), BHA36(1.0)
CY148	12369	BHA2(77.2), BHA37(15.4), BHA4(7.3)
CY149	10771	BHA1(76.7), BAH2(23.2)
CY15	12621	BHA1(79.0), BHA2(20.9)
CY16	15922	BHA1(54.2), BHA2(45.7)
CY19	29321	BHA1(59.3), BHA2(33.6), BHA4(0.3), BHA34(3.9), BHA38(1.6), BHA39(0.7), BHA40(0.6)