A novel phylogenetic analysis and machine learning predict pathogenicity of human mtDNA variants

Bala Anı Akpınar[1][†], Paul O. Carlson[1], Ville O. Paavilainen[1], and Cory D. Dunn[1][†]

[1]Institute of Biotechnology, Helsinki Institute of Life Science, University of Helsinki, Helsinki, 00014, Finland

[†] Corresponding authors

Correspondence:

Bala Anı Akpınar, Ph.D.
P.O. Box 56
University of Helsinki
00014 Finland
Email: ani.akpinar@helsinki.fi
Phone: +358 50 311 9307

or

Cory Dunn, Ph.D.
P.O. Box 56
University of Helsinki
00014 Finland
Email: cory.dunn@helsinki.fi
Phone: +358 50 311 9307

ABSTRACT

Linking mitochondrial DNA (mtDNA) mutations to patient outcomes remains a formidable challenge. The multicopy nature and potential heteroplasmy of the mitochondrial genome, differential distribution of mutant mtDNAs among various tissues, genetic interactions among alleles, and environmental effects currently hamper clinical efforts to diagnose mitochondrial disease. Multiple sequence alignments are often used when estimating the potential significance of mitochondrial variants. However, factors including sample set bias, alignment errors, and sequencing errors currently limit the utility of multiple sequence alignments in pathogenicity prediction. Here, we describe a new approach to assessment of site-specific conservation and variant acceptability that minimizes previous limitations, and using the output of our methodology, we deploy machine learning in order to predict pathogenicity for thousands of human mtDNA variants. Our work demonstrates that a substantial fraction of mtDNA changes not yet characterized as harmful are, in fact, likely to be deleterious. Our findings will be of direct relevance to those at risk of mitochondria-associated illness, but the general applications of our methodology also extend beyond the context of mitochondrial disorders.

INTRODUCTION

Because of the critical role that mitochondria play in metabolism and bioenergetics, mutations of mitochondria-localized proteins and ribonucleic acids can adversely affect human health (1–4). Indeed, at least one in 5000 people (5) is estimated to be overtly affected by mitochondrial disease. While a very limited number of mitochondrial DNA (mtDNA) lesions can be directly linked to human illness, the clinical outcome for many other mtDNA changes remains ambiguous (6). Heteroplasmy among the hundreds of mitochondrial DNA (mtDNA) molecules found within a cell (7–9), differential distribution of disease-causing mtDNA among tissues (10), and modifier alleles within the mitochondrial genome (11, 12) magnify the problem of interpreting effects of different mtDNA alterations. Mito-nuclear interactions and environmental effects may also determine the outcome of mitochondrial DNA mutations (13–16). Beyond the obvious importance of determining the genetic etiology of symptoms in a clinical setting, the rapidly increasing prominence of direct-to-consumer genetic testing (17) calls for an improved understanding of which mtDNA polymorphisms might affect human health (18).

While our knowledge of mtDNA variants existing among humans continues to increase, tabulation of mtDNA variants found among healthy or sick individuals (19) can lack utility in predicting mitochondrial disease. Differing, strand-specific mutational propensities for mtDNA nucleotides at different locations within the molecule (20–22) should be taken into account when assessing population-wide data, yet allele frequencies are rarely, if ever, normalized in this way. Population sampling biases and recent population bottleneck effects can lead to misinterpretation of variant frequencies (23–27), and mildly deleterious variants arising in a population have not yet been removed by selection (28, 29). Finally, a lack of selection against variants that might act in a deleterious manner at the post-reproductive stage of life also makes likely the possibility that some mtDNA changes will contribute to age-related phenotypes while avoiding overt association with mitochondrial disease (30–34).

Examining evolutionary conservation by use of multiple sequence alignments offers important assistance when predicting a variant's potential pathogenicity (35, 36). However, caveats are also associated with predicting mutation outcome by the use of these alignments. First, while knowledge of amino acid physio-chemical properties is widely considered to be informative regarding whether an amino acid substitution may or may not have a damaging effect on protein function (37), the site-specific acceptability of a given substitution is ultimately decided within the context of its local protein environment (38). Second, sampling biases and improper clade selection may lead to inaccurate clinical interpretations regarding the relative acceptability of specific variants (23–26). Third, interactions among alleles and environmental effects can further complicate pathogenicity prediction (39–41). Finally, alignment (42, 43) and sequencing errors (44, 45) may falsely indicate the acceptability of a particular mtDNA substitution.

Here, we have deployed a novel methodology to calculate, based upon analyses of available mammalian genomes, the relative conservation of human mtDNA-encoded positions. Our conservation values are of potentially unlimited dynamic range, and are robust against sample bias, sequencing error, and alignment error. Moreover, we are able to identify ancestral direct substitutions that match deviations from the human reference sequence, providing further information about the potential pathogenicity of any human mtDNA substitution. By subsequent application of machine learning, we demonstrate that a surprising number of uncharacterized mtDNA mutations carried by humans are likely to promote disease.

RESULTS

Mapping apparent substitutions to a phylogenetic tree allows calculation of relative positional conservation

We previously introduced an improved, empirical approach for detection and quantification of substitutions within evolving mtDNA-encoded macromolecules (46). Here, we have extended our methods toward prediction of human mitochondrial variant pathogenicity. First, we retrieved full mammalian mtDNA sequences from the National Center for Biotechnology Information Reference Sequence (NCBI RefSeq) database and extracted specific sequences of each RNA or protein-coding gene within the *Homo sapiens* reference mtDNA. Next, we aligned the resulting protein, tRNA, and rRNA sequences (Figure 1a), concatenated the sequences of each species based upon molecule class, and generated phylogenetic trees using a maximum likelihood approach (Figure 1b). Following tree generation, we performed ancestral prediction (Figure 1c) to reconstruct the character values of each position at every bifurcating node. Using the sequences of extant species and the predicted ancestral node values, we subsequently analysed each edge of the tree for the presence or absence of substitutions at each aligned human position (Figure 1d). Importantly, we focused our attention upon mutations localized between internal nodes of maximum likelihood trees (Figure 1e), excluding edges leading directly to extant sequences and thereby minimizing the effects of alignment errors and sequencing errors that may lead to eventual misinterpretation of variant pathogenicity. Moreover, mutations mapped to internal edges are more likely to represent fixed changes informative for the purposes of disease prediction, while polymorphisms that have not yet been subject to selection of sufficient strength or duration might be expected to lack predictive power (28, 29). We subsequently sum the substitutions at internal positions to generate a new metric, the internal substitution score (ISS). The ISS should surpass metrics that consider positional character frequencies derived from multiple sequence alignments as a proxy of conservation, since character frequencies are highly sensitive to sample biases among extant sequences. Inferred ancestral and descendent characters at each edge were also extracted following generation of the substitution mapping matrix (Figure 1f), providing important information regarding which direct substitutions at each site are possible within a lineage.

When used during analysis of macromolecules encoded by mammalian mtDNAs, it is clear that the ISS provides an excellent readout of relative conservation at, and consequent functional importance of, each alignment position. When plotting increasing ISSs across all amino acid positions with gap values of 1% or less, a characteristic curve was generated (Figure 2a). At the beginning of the curve, we encountered many positions that were nearly or completely invariant, implying critical roles for the folding or function of each polypeptide. Next, we observed a slope of increasing ISSs associated with amino acids of decreasing evolutionary constraint. Finally, the tail of each curve dramatically increases in slope, suggesting aligned, yet degenerate positions under minimal or no selection. Conservation curves are also obtained when studying individual mtDNA-encoded proteins within each oxidative phosphorylation complex (Figure 2b-e). Similar conservation curves are generated when examining combined (Figure 2f) and individual (Figure 2g) rRNA molecules or when considering combined (Figure 2h) and individual (Figure 2i-j) tRNA molecules that are encoded by the human mitochondrial genome.

When comparing ISS data from different mtDNA-encoded proteins, our methodology supports the idea that the core subunits of cytochrome *c* oxidase (COX), and particularly the COI polypeptide, are the most highly conserved when considering mutability across amino acid positions (47, 48) (Figure 3a). To further demonstrate that low ISSs correspond with conservation, we plotted mammalian ISS sums for the

three mitochondria-encoded subunits of cytochrome c oxidase (COX) onto the high-resolution structure of the bovine enzyme (49). Indeed, according to ISSs extracted from the mammalian mtDNA tree, the most conserved positions were localized to the catalytic core of COX (Figure 3b-c and Movie S1). In contrast, more divergent amino acids clearly faced toward external regions of the complex, most prominent of which are the lipid-contacting regions of COX.

Internal substitution scores provide conservation measurements of unlimited dynamic range

Measures of conservation based upon character frequencies extracted from multiple sequence alignments, such as Shannon entropy, are inherently limited in dynamic range (50). Dynamic range limitations that can characterize frequency-based approaches to position analysis within alignments are especially prominent when considering nucleotides, where only four possibilities should occur at any given position. In contrast, summing positional substitutions across edges of a phylogenetic tree allows for unlimited dynamic range of calculated conservation values. To demonstrate the increased dynamic range provided by our method, we compared Shannon entropy values and ISSs for well-aligned protein (Figure S1a), rRNA (Figure S1b), and tRNA (Figure S1c) positions encoded by the full set of mammalian mtDNAs and by a smaller set of mtDNAs limited to primates. Since Shannon entropy values and ISSs can both report upon conservation at aligned positions, these two variables were, as expected, correlated. However, while the range of Shannon entropy values cannot increase when moving from primate-level analysis to mammal-level analysis, the range of ISSs did increase, supporting the idea that our approach to comparing selection at different sites benefits from the continuous accumulation of mtDNA sequence information.

We considered whether empirical measurements of substitution in mtDNA-encoded macromolecules, rather than a model-based approach to conservation, would poorly reflect conservation if the evolutionary divergence between two nodes is too large. To test the extent to which our method may undercount substitutions at aligned sites, we first selected alignment positions that were equal to or less than 1% gapped. Next, we selected protein, rRNA, or tRNA sites that were under the most minimal selective constraint, as reflected by their rank in the top 1% of ISSs. We then compared the number of apparent changes summed across all positions (regardless of gap %) along an internal tree edge to the number of substitutions occurring only at these degenerate sites. The binary nature of our edge-focused substitution analysis, like other approaches investigating substitution, obscures the development of a Poisson distribution that would reflect repeated changes to degenerate sites (38, 51). Therefore, we expected that the total number of degenerate sites mutated along an edge would plateau beyond a given level of evolutionary divergence. Indeed, when an edge contained ~150 - 200 mtDNA-encoded protein positions with substitutions, the rise in total substituted degenerate sites with increasing total substitutions along an edge comes to an end (Figure S2a). The relationship between rRNA (Figure S2b) and tRNA (Figure S2c) degenerate site substitutions and total edge substitutions is less clear-cut than for proteins, potentially as a result of a lower number of degenerate positions suitable for analysis. However, a similar limitation on degenerate site substitution detection beyond a given edge length remains apparent. Since most positions encoded by mtDNA appear to be evolutionarily constrained and cannot reach the ISSs characterizing the most degenerate sites (Figure 2), and because most edges analyzed are 'shorter' than the thresholds at which the total degenerate sites substituted reaches a maximum, we expect that any potential under-counting of substitutions along longer tree edges will minimally affect our analysis of site-specific conservation.

Positional substitution frequency and inferred direct substitutions can be linked to human mtDNA variant pathogenicity

Since summation of detected substitutions across a phylogenetic tree provides a robust measure of relative conservation at different macromolecular positions, we were confident that a phylogenetic analysis that includes ISSs would be informative about the pathogenicity of human mtDNA variants. To test this possibility, we focused our attention upon substitutions annotated within the MITOMAP database of pathogenic mtDNA alterations (52). Indeed, we detected a clear relationship between confirmed pathogenicity and conservation at mtDNA-encoded amino acid positions, as reflected by positional ISSs (Figure 4a). Similarly, there was a strong link between tRNA mutation pathogenicity and ISSs obtained from mammalian tRNA data (Figure 4b). The paucity of confirmed pathogenic mitochondrial rRNA mutations in the MITOMAP database made comparisons using this class of molecules impractical, yet we expect future confirmation of additional pathogenic mutations in mitochondria-encoded rRNAs to permit further analyses. Together, findings obtained by phylogenetic analysis of mitochondria-encoded proteins and tRNAs indicate that ISSs will be valuable when predicting which mtDNA mutations might lead to disease.

By ancestral prediction of nodes within our mammalian phylogenetic trees, we were able to infer ancestral and descendent characters along each tree edge. We reasoned that these data, while not totally uncoupled from positional ISSs, would provide additional information regarding what changes to mtDNA-encoded macromolecules might be harmful or harmless. Specifically, if direct substitution from the human reference character to the mutant character (or the inverse, assuming the time-reversibility of character substitutions) is predicted along the edge of a phylogenetic tree, then such a change at a given position might be expected to be less deleterious than an inferred direct substitution to or from the human character that was never encountered during evolution. In contrast, the simple presence or absence of a character at an alignment position, without the context of its ancestral character, will fail to reflect epistatic relationships that involve the position of interest (40, 53, 54). We again focus upon internal tree edges, rather than those leading to extant sequences, so that we might avoid the confounding effects of sequencing and alignment errors (44, 45) and of inadequate selection against newly arisen, mildly deleterious alleles (28, 29).

To test the possibility that inferred internal direct substitutions (IDSs), like ISSs, may predict pathogenicity, we compared human variants confirmed to cause disease to those variants currently labelled as polymorphisms within the MITOMAP database. Pathogenic human amino acid changes from the reference sequence are rarely encountered among the internal edges of a mammalian phylogenetic tree (Figure 5a). In contrast, reportedly polymorphic alleles were much more likely to have the amino acid substitution identifiable along internal edges of the mammalian tree. For nucleotide substitutions in tRNAs, a similar analysis comparing confirmed pathogenic variants and variants annotated as polymorphisms detected no significant relationship between pathogenicity and IDSs. However, subsequent evidence indicated that IDSs in tRNAs may indeed help to differentiate harmful from harmless variants (see below).

Widespread, uncharacterized pathogenicity among human mtDNA polymorphisms

A subset of substitutions currently annotated as 'polymorphic' in the MITOMAP database were distinguished by very low ISS scores at the relevant alignment position, suggesting that mutations at those positions could, in fact, be deleterious. Moreover, when considering polymorphisms, the absence

of IDSs within the mammalian phylogenetic tree that matched certain human variations raised questions regarding whether additional pathogenicity may exist among human mtDNA substitutions currently considered polymorphic. In order to explore whether additional pathogenic mutations might indeed be discovered among variants currently tabulated as 'polymorphic,' we tested whether protein polymorphisms tabulated in MITOMAP that cannot be directly linked to the human reference character along the edges of the mammalian phylogenetic tree would lead to purifying selection. Selection was assessed by examining the number of full-length mtDNA samples found in GenBank that are linked to a given variant. Measuring selection by assessment of population frequencies can be problematic due to potential sampling biases and population bottlenecks (55), timing of mutation arrival within an expanding population (56, 57), and the divergent nucleotide- and strand-specific mutational propensities of mtDNA (20–22). Even so, the distribution of variant frequencies among full-length sequences in GenBank was strikingly different for those mutations for which an IDS could be identified in our mammalian trees of proteins (Figure 5b), tRNAs (Figure 5c), and rRNAs (Figure 5d) when compared to those for which an IDS could not be identified. These results indicate that many human mtDNA variants currently considered to be polymorphic are, in fact, harmful.

A support vector machine predicts deleterious mtDNA polymorphisms

Given the clear presence of deleterious substitutions among thus far uncharacterized variants, we sought a high-throughput method that could, with high confidence, identify these potentially pathogenic changes. To achieve this, we turned to machine learning, and specifically a support vector machine (SVM) approach (58), to predict the risk of mtDNA variants in a manner dependent upon our phylogenetic analyses. Along with the mutations tabulated in MITOMAP, we also included mtDNA changes found in the recently established HelixMT database (HelixMTdb). This database was generated during a genetic survey, performed agnostic to phenotype, of nearly 200,000 independent human samples (59). Positive training sets for our SVM consisted of protein or tRNA mutations cataloged as confirmed pathogenic alterations within the MITOMAP database. Negative training sets included variants annotated as polymorphic in MITOMAP and characterized by the highest counts of homoplasmic alleles within HelixMTdb, which we interpreted as a sign of benignancy. Analyzed positions were gapped at 1.5% of sequences or less. As SVM features, we exploited ISSs and IDSs. We also expanded our feature set to include the Total Substitution Score (TSS), which includes edges leading to extant mtDNA sequences; the number of characters found at alignment positions (internal edges or all edges); direct substitution between the human reference character and the mutant human character along all edges; whether the mutant character is ever found at internal nodes or all nodes; and positional Shannon entropy.

For variants found within protein-coding genes, our SVM approach clearly permitted a robust and automated prediction of which polymorphisms not yet classified as pathogenic may cause or contribute to disease (Figure 6a and Table S1). Importantly, when examining the behavior of alleles within our negative training sets, no false-positives were predicted. Furthermore, few confirmed pathogenic mutations were found among the predicted negative set. No specific feature, all of which are related to site conservation in mammals, stood out as dominant in providing predictive value (Figure 6b). The robustness of our model in correctly identifying the true positives in the training set without corresponding loss in specificity is highlighted by the associated Receiver Operating Characteristic (ROC) curve [area under the receivership curve (AUC), 0.958] (Figure 6c).

For tRNAs, a SVM-based approach again provided notable separation between positive and negative training sets (Figure 6d and Table S2). During tRNA analysis, the TSS and the number of characters present at internal nodes appeared to play the strongest role in prediction (Figure 6e). The ROC curve for our tRNA analysis suggests high predictive proficiency of the SVM approach when focused upon tRNAs (AUC, 0.918), with a moderate reduction of predictive power when compared to that obtained during analysis of coding variants (Figure 6f). Due to the lack of confirmed pathogenic mutations in rRNA, we have not yet applied our predictive approach to this class of mtDNA-encoded macromolecules.

Beyond variants of confirmed pathogenicity used to train our SVM models, there are other changes provided in the MITOMAP database that are 'reported', yet not confirmed, to be pathogenic. We asked whether our SVM predictions might provide further evidence that some of these variants reported to be mitochondrial disease-associated are truly pathogenic. Toward this goal, we removed substitutions potentially related to carcinogenesis, ensuring a focus on metabolic disease caused by mitochondrial dysfunction, then assigned SVM prediction values to the remaining variants. We expected that pathogenic alterations are more likely to be detected as heteroplasmic within human samples, since overt mitochondrial disease may require a deleterious variant to make up a high proportion of the mtDNA molecules maintained by the cell (7, 8). Therefore, we asked whether variants in our two predicted classes could be differentiated by the percentage of samples reported as heteroplasmic within HelixMTdb. Indeed, there was a clear difference in propensity to heteroplasmy between the two classes when considering mutations affecting protein (Figure 7a) and tRNA (Figure 7b). With a continued focus on MITOMAP variants reported, but not confirmed, to be disease-associated, we next assessed purifying selection by assessing the number of total samples in HelixMTdb for each variant within our prediction classes. Indeed, there was a decrease in population prevalence among protein-coding (Figure 7c) and tRNA (Figure 7d) variants that we had predicted as pathogenic. Our findings demonstrate the general applicability of our pathogenicity predictions, while providing additional evidence supporting the deleterious nature of specific mtDNA changes suspected to be, but not yet confirmed as, harmful.

Following validation of our SVM prediction approach using a subset of variants enriched for pathogenicity, we turned to the thousands of mtDNA changes tabulated within MITOMAP or HelixMTdb that are not currently characterized as harmful. Coupling our evolutionary analysis of mtDNA-encoded macromolecules to machine learning again revealed the potential pathogenicity of hundreds of variants found within the human population. The heteroplasmy profile (Figure 7e) and population distribution (Figure 7f) were again clearly distinct when comparing those variants that we predict to be harmful and those variants that we predict to be benign. Similarly, although less prominently, tRNAs predicted to be harmful or harmless were also distinguished by heteroplasmy distribution (Figure 7g) and population frequency distribution (Figure 7h).

DISCUSSION

We describe here a methodology that allows improved quantification of the relative conservation of sites within and between genes, RNAs, and proteins. Mapping substitutions to phylogenetic tree edges can minimize errors in conservation calculation that results from sampling biases, from sequencing errors, and from alignment errors. Moreover, ISS scores provide a measure of conservation that is theoretically unlimited in dynamic range and, consequently, will continue to benefit from the ever-increasing mtDNA sequence information available in public and private databases. Even nearly identical

sequences can be utilized by our approach, allowing for an ever-increasing input dataset that can be deployed toward calculation of site-specific conservation.

Beyond the development of a novel quantitative measure of conservation, we also use ancestral predictions to generate site-specific substitution matrices. Researchers often consult generalized substitution matrices when predicting whether a change may be harmful or not (37, 60), yet amino acid exchangeability matrices change when moving across clades (61), and successful substitution of any given character occurs in the context of a very specific local environment (36, 38). By mapping substitutions to phylogenetic tree edges, ample sequence data can allow identification of IDSs within the context of a particular macromolecular position, thereby improving prediction of variant pathogenicity. We note that focusing upon IDSs, rather than the simple presence or absence of a character at a site, can indirectly integrate information about epistatic interactions that permit or block a substitution from being successfully established within a lineage.

Protein and tRNA variants of confirmed pathogenicity were clearly characterized by low ISSs and by the absence of IDSs. But how should one weigh these and other available factors when classifying, in an unbiased and high-throughput manner, other variants already encountered or waiting to be discovered among human mtDNAs? To address this question, we deployed a machine learning approach, trained on existing data present in the MITOMAP database, toward determination of mtDNA mutation pathogenicity. Other attempts have been made to predict the effect of mtDNA substitutions, with widely divergent prediction proficiency (62). Several of these approaches take advantage of machine learning (63, 64). However, our approach is a clear advance over previous methods, due to our avoidance of heavily biased character frequencies and our prioritization of node values and substitutions internal to the phylogenetic tree. Our apparent false-positive rate for protein coding variants appears to be extremely low, if not zero, which is essential when avoiding the harmful psychological effects of a false-positive test (65, 66). Specificity and sensitivity metrics resulting from our SVM predictions appear exemplary (AUC > 0.95 for protein variants, AUC > 0.91 for tRNA variants). Additional tests demonstrate that our approach is quite effective in uncovering harmful mtDNA variants. Even though population frequency is subject to bias (see below), variants that we now predict to be pathogenic were generally characterized by lower prominence in the population, suggesting purifying selection. Moreover, we detected a very strong bias against apparent homoplasmy when considering polypeptide mutations that we predicted to be harmful, and enrichment of detectable heteroplasmy is a common indicator of pathogenicity (67).

mtDNA variants predicted to be pathogenic may lead to cryptic and age-related mitochondrial disease

Our results are congruent with earlier analyses suggesting that harmful mtDNA substitutions may be common within the human population (12, 29, 67). So why have these deleterious changes not yet been classified as pathogenic within the clinic?

First, if a mutation is relatively common, clinicians may inappropriately determine that the mtDNA variant is unlinked to patient symptoms (19). Our findings do, in fact, support the idea that reduced population frequency of an alteration can correspond with pathogenicity. However, population frequency cannot currently be a highly reliable predictor of clinical outcome, since population counts require substantial correction for strand- and nucleotide-specific mtDNA mutational biases (20–22), and sampling biases are typically a hazard when carrying out population-wide studies (23–26).

Second, deleterious mtDNA variants often must rise beyond a certain threshold among the hundreds of mtDNA molecules potentially resident within a cell in order for overt symptoms to manifest (4, 7). Concordantly, our data suggest a strong propensity for heteroplasmy in the set of substitutions that we predict to be pathogenic, but are not yet clinically annotated as disease-associated. Differential distribution of mtDNA variants within a carrier, either established during development or resulting from bottleneck effects in renewable and non-mitotic tissues (7, 68), may generate clones with a high proportion of deleterious mutations and to complex, tissue-specific outcomes (69–72). Moreover, the phenomena described above may lead to age-related symptoms not easily classified as mitochondrial disease, since even relatively common mtDNA sequence variants have been suggested to contribute to diseases like diabetes, heart disease, and cancer (11, 73, 74). We are certainly tantalized by the prospect that pathogenic variants illuminated by our approach might impinge upon human lifespan.

METHODOLOGY

Mitochondrial DNA sequence acquisition and conservation analysis

Mammalian mtDNA sequences were retrieved from the National Center for Biotechnology Information database of organelle genomes (https://www.ncbi.nlm.nih.gov/genome/browse#!/organelles/ on September 26, 2019). These 1184 mammalian mtDNA genomes were aligned using MAFFT on the 'auto' setting (75). Four sequences that were egregiously misaligned were removed, and MAFFT alignment with the 'auto' setting was carried out again. Individual gene sequences were extracted from these alignments, based upon the annotated human mtDNA (NC_012920.1). After gap removal, translation of protein coding genes was performed using the vertebrate mitochondrial codon table in AliView (76). MAFFT alignments of each gene product were performed using the G-INS-i iterative refinement method, then individual concatenates for each species were generated from protein coding sequences, tRNAs, and rRNAs. Duplicates were removed from the protein, tRNA, and rRNA concatenates using seqkit v0.10.2 (77). Primate subsets containing concatenates from each molecule class were generated after analysis of species names using the taxize package (v0.9.9) in R (78).

Maximum likelihood trees for each molecule class concatenate were built using FastTreeMP (79) with four subtree-prune-regraft moves, two rounds of branch length optimization, slow nearest neighbor interchange, and by use of a generalized time-reversible model. Next, ancestral prediction was carried out using the PAGAN package (80), with concatenated alignments and phylogenetic trees used as input. The PAGAN output was then analysed using "binary-table-by-edges-v2.2" and "add-convention-to-binarytable-v1.1.py" (https://github.com/corydunnlab/hummingbird) (46) to allow for a sum of substitutions at alignment positions encoded by human mtDNA. All fluctuating edges were extracted using "report-on-F-values-v1.1" (https://github.com/corydunnlab/hummingbird). Correspondence files for the human mtDNA (NC_012920.1) convention and alignment positions were generated using "extract-correspondence-for-merged-alignment-v.1.1.py", followed by the construction of the look-up tables of amino acid direct substitution and presence using "direct-subst-lookup-table-proteins-v.1.1.py" and "AA-presence-lookup-table-v.1.1.py", respectively (https://github.com/corydunnlab/Edge_mapping_conservation). For IDS determination, only edges with standard characters at both connected nodes were considered. For ISS calculation, the PAGAN output files for tRNAs and rRNAs were, before inferring a substitution, processed so that each ambiguous IUPAC nucleotide code was replaced by one of the possible standard nucleotides represented by that IUPAC code within the entire tree at that given position. For positions that contained only N's and gaps,

N's were replaced by gaps. Replacement of ambiguous characters with standard characters was carried out using using script "replace-nonstandard.py" (https://github.com/corydunnlab/Edge_mapping_conservation), and the standard nucleotide was picked at random from available standard characters at a given alignment position using the random.choice() function of the Python random module. Gap percentage for each position, with mammal and primate concatenates handled separately, was calculated using trimAI v1.4.rev22 (81). Shannon entropy was calculated using the 'Shannon' script (https://gist.github.com/jrjhealey/). Custom scripts were used to merge various data tables. Software generated within the context of this study was written using Python 2.7.

Chimera v1.13.1 was used for plotting structural data (82), and statistical testing and graph production were performed using Prism 8.3.0 (https://www.graphpad.com). Heat maps were generated using Matrix2png (83). When plotting data on a logarithmic scale, variant population counts of zero were converted to one.

Database utilization

MITOMAP data used in this study (52) were downloaded on October 1, 2019. HelixMTdb data used in this study (59) were downloaded on October 15, 2019.

Support vector machine development

Our SVM classifier (58) was developed using the R-language (https://www.R-project.org/) package e1071 (84), whose implementation is based on libsvm (85). A supervised binary classification model was built using positive and negative training sets. Each negative training set included alleles with the highest number of homoplasmic samples in HelixMTdb and also annotated as a polymorphism within MITOMAP. For proteins, the negative training sets consisted of 100 variants, and for tRNAs, the negative training sets consisted of 50 variants. The positive training sets for proteins consisted of variants confirmed as pathogenic in MITOMAP. Analyzed positions for protein and tRNA included those gapped at 1.5% or less for mammalian or primate alignments. Optimal parameters for model fitting were identified using the tune.svm wrapper in e1071. The best parameters for the radial basis kernel "rbfdot" were found by performing a $\log_2$ grid search of gamma and cost values during 5-fold cross-validation. Cross-validation error rate averaged from 10 independent repeats was used as the metric in choosing the best parameters. Subsequently, the full training set was used to train the SVM using these best parameters. Features were scaled internally during training and when applying the model in order to predict the full data sets of mtDNA mutations. ROC curves were drawn with the mgraph function from rminer package (86), which uses ksvm from kernlab package (87). Feature importance was measured using the Importance function from rminer (88).

ACKNOWLEDGEMENTS

C.D.D. is funded by a ERC Starter Grant (RevMito 637649) and by the Sigrid Jusélius Foundation. V.O.P. is supported by the Academy of Finland (grants 289737, 314672, and 330255), U.S. National Institute of General Medical Sciences (Project 1R01GM132649-01), the Sigrid Jusélius Foundation, and by the Runar Bäckström Foundation. We thank Vivek Sharma, Tamara Somborac, and Gülayşe İnce Dunn for helpful comments on the manuscript.

DISCLOSURES

Figure 1: A workflow for generation of internal substitution scores and inferred internal direct substitutions. (**a**) Sequences for each macromolecule class (protein, rRNA, tRNA) are selected, aligned, and concatenated. (**b**) From these concatenates, phylogenetic trees are produced, followed by (**c**) ancestral prediction performed using alignments and phylogenetic trees as input. (**d**) Next, a binary matrix is generated, in which the presence or absence of mutations at each aligned position along each tree edge is tabulated with reference to a specific species convention. (**e**) From this matrix, substitutions at internal tree edges, which are less subject to sequence errors, alignment errors, and unselected polymorphisms, can be summed. (**f**) Using the predicted internal node values, inferred direct substitutions between ancestral and descendant characters are determined.

Figure 2: Most well-aligned positions encoded by the human mitochondrial genome are under selective constraint. Sequences of each macromolecule class were analyzed as described in Figure 1. ISS distributions are provided for all alignment positions equal to or less than 1% gapped in their respective alignments of mammalian sequences: (**a**) ISS analysis of all mtDNA-encoded proteins considered together as a concatenate (83% of total positions gapped 1% or less), (**b**) individual Complex I proteins, (**c**) individual Complex III proteins, (**d**) individual Complex IV proteins, (**e**) individual Complex V proteins, (**f**) concatenated, mitochondria-encoded rRNAs (76% of total positions gapped 1% or less), (**g**) individual rRNAs, (**h**) concatenated, mtDNA-encoded tRNAs (70% of total positions gapped 1% or less), (**i**) individual tRNAs encoded on the human H-strand, (**j**) individual tRNAs encoded on the human L-strand.

Figure 3: Internal substitution scores reflect conservation associated with important structural and catalytic residues. (**a**) Subunits of COX exhibit the most conservation, as reflected by ISSs across positions. The same data examined above are displayed using a box and whiskers plot, with whiskers indicating minimum and maximum values, box encompassing 25 through 75th percentile, and the line internal to the box representing the median. (**b-c**) Side chains of mtDNA-encoded COX subunits that are associated with high ISSs face away from core, catalytic residues. Mitochondria-encoded subunits COI, COII, and COIII from one half of the dimeric crystal structure of *Bos taurus* cytochrome *c* oxidase (PDB: 1OCO) (89) are represented in colored, ribbon format, with the other subunits depicted in grey. Bound heme groups are displayed in red and orange, and the conserved, catalytic E242 residue is colored green. ISSs from analysis of mammalian phylogenetic data, without respect to gap percentage, are reflected as follows: 0 substitutions, black; 1-6 substitutions, dim grey; 7-12 substitutions, blue; 13-24 substitutions, cornflower blue; 25 substitutions and above, purple. In (**b**), the mitochondrial matrix lies at the bottom of the model. In (**c**), the viewer looks down toward the mitochondrial matrix.

Figure 4: Confirmed pathogenic mutations are associated with low internal substitution scores. ISSs for positions of variants annotated in the MITOMAP database as either confirmed pathogenic or as polymorphisms are plotted for (**a**) protein positions analyzed using mammalian data and (**b**) tRNA positions analyzed using mammalian data. All analyzed positions are gapped at equal to or less than 1%. ****, Kolmogorov-Smirnov test approximate P-value of <0.0001. Dotted horizontal lines, median, with solid horizontal lines providing interquartile range.

Figure 5: Analysis of inferred internal direct substitutions indicates that variants currently annotated as polymorphic may be deleterious. (**a**) Inferred internal direct substitutions within the mammalian phylogenetic tree correspond to a lack of confirmed pathogenicity for mitochondrial protein

coding variants. Confirmed pathogenic protein substitutions and substitutions currently annotated as polymorphisms, both retrieved from MITOMAP, were tested for the presence of an IDS. ****, Fisher's two-sided exact test P-value of <0.0001. All analyzed positions are gapped at equal to or less than 1%. (**b-d**) Population frequency distributions differ when MITOMAP-annotated substitutions from the human reference character in protein (**b**), tRNAs (**c**), or rRNAs (**d**) are classified based upon the presence or absence of an IDS. Variants found within the MITOMAP database and encoding protein substitutions were classified based on whether an IDS between the human reference character and the mutant (**b**) protein, (**c**) tRNA, or (**d**) rRNA character could be detected. All analyzed positions are gapped at equal to or less than 1%. ****, Kolmogorov-Smirnov test approximate P-value of <0.0001.

Figure 6: A machine-learning approach predicts pathogenicity of mtDNA variants currently annotated as polymorphisms. (**a**) SVM-based prediction of pathogenicity for protein-coding variants found in MITOMAP and HelixMTdb using features taking advantage of mammalian alignment positions of gap value 1.5% or less. Positive and negative training sets are superimposed on the curve of variants and decision values. (**b**) Feature importance is calculated from the increase of the model's prediction error after shuffling the values for each feature. 10 test iterations are shown. (**c**) The receiver operating characteristic (ROC) curve demonstrates very high sensitivity is achieved with no loss in specificity for mammalian protein predictions. The true positive rate (sensitivity) of the predictive model is plotted against the false positive rate (1 - specificity) at various decision value thresholds. (**d**) As for (**a**), but for tRNAs. (**e**) As for (**b**), but for tRNAs. (**f**) As for (**c**), but for tRNAs.

Figure 7: SVM-based pathogenicity predictions are supported by separable distributions of heteroplasmy profiles and allelic population frequencies. (**a**) For mitochondria-encoded protein substitutions reported to be disease-associated within the MITOMAP database, but not confirmed as pathogenic, the percent of HelixMTdb samples for a given variant that were heteroplasmic is plotted according to pathogenicity prediction by SVM. (**b**) As for (**a**), except tRNA variants are plotted. (**c**) For protein substitutions reported to be disease-associated within the MITOMAP database, but not confirmed as pathogenic, the total number of HelixMTdb samples for a given variant is plotted according to pathogenicity prediction by SVM. (**d**) As for (**c**), except tRNA substitutions are plotted. (**e**) For protein substitutions in MITOMAP and HelixMTdb not associated with disease, the percent of heteroplasmic HelixMTdb samples for a given variant is plotted according to pathogenicity prediction. (**f**) For protein substitutions in MITOMAP and HelixMTdb not associated with disease, the number of samples with the given variant is displayed according to pathogenicity prediction. (**g**) As in (**e**), except that tRNA changes are analyzed. (**h**) As in (**f**), except that tRNA substitutions are considered. ****, Kolmogorov-Smirnov test approximate P-value of <0.0001.

SUPPLEMENTAL FIGURE LEGENDS

Figure S1: Internal substitution scores provide unlimited dynamic range and can scale with level of taxonomy. ISSs obtained from positional analysis of (**a**) protein, (**b**) rRNA, or (**c**) tRNA sequence concatenates and primate and mammal phylogenetic trees are compared to Shannon entropy values for the same alignment positions. All positions shown here are gapped at 1% or less.

Figure S2: Most edges from trees of mammalian mtDNA-encoded macromolecules are not saturated for substitutions. Degenerate positions characterized by the top 1% of ISSs and gapped at 1% or less were extracted during analysis of mammalian macromolecule concatenates. The number of those degenerate sites mutated along each edge of the mammalian protein tree is plotted against the total number of substitutions along the tree edge. Superimposed upon these data is a restricted cubic spline curve. Analysis was performed for (a) proteins (b) rRNAs and (c) tRNAs.

SUPPLEMENTAL TABLE LEGENDS

Table S1: SVM output analyzing protein-encoding variants found in the MITOMAP and HelixMTdb by use of mammalian sequence data. Details of SVM analyses and features are found in the methodology section.

Table S2: SVM output analyzing tRNA-encoding variants found in the MITOMAP and HelixMTdb by use of mammalian sequence data. Details of SVM analyses and features are found in the methodology section.

REFERENCES

1. C. L. Alston, M. C. Rocha, N. Z. Lax, D. M. Turnbull, R. W. Taylor, The genetics and pathology of mitochondrial disease. *J. Pathol.* **241**, 236–250 (2017).

2. A. Suomalainen, B. J. Battersby, Mitochondrial diseases: the contribution of organelle stress responses to pathology. *Nat. Rev. Mol. Cell Biol.* **19**, 77–92 (2018).

3. S. Khan, G. Ince-Dunn, A. Suomalainen, L. L. Elo, Integrative omics approaches provide biological and clinical insights: examples from mitochondrial diseases. *J. Clin. Invest.* **130**, 20–28 (2020).

4. O. M. Russell, G. S. Gorman, R. N. Lightowlers, D. M. Turnbull, Mitochondrial Diseases: Hope for the Future. *Cell* **181**, 168–188 (2020).

5. G. S. Gorman, *et al.*, Prevalence of nuclear and mitochondrial DNA mutations related to adult mitochondrial disease. *Ann. Neurol.* **77**, 753–759 (2015).

6. J. M. Vento, B. Pappa, Genetic counseling in mitochondrial disease. *Neurotherapeutics* **10**, 243–250 (2013).

7. J. B. Stewart, P. F. Chinnery, The dynamics of mitochondrial DNA heteroplasmy: implications for human health and disease. *Nat. Rev. Genet.* **16**, 530–542 (2015).

8. A. Hahn, S. Zuryn, The Cellular Mitochondrial Genome Landscape in Disease. *Trends Cell Biol.* **29**, 227–240 (2019).

9. W. Wei, P. F. Chinnery, Inheritance of mitochondrial DNA in humans: implications for rare and common diseases. *J. Intern. Med.* (2020) https:/doi.org/10.1111/joim.13047.

10. L. Boulet, G. Karpati, E. A. Shoubridge, Distribution and threshold expression of the tRNA(Lys) mutation in skeletal muscle of patients with myoclonic epilepsy and ragged-red fibers (MERRF). *Am. J. Hum. Genet.* **51**, 1187–1200 (1992).

11. W. Wei, A. Gomez-Duran, G. Hudson, P. F. Chinnery, Background sequence characteristics influence the occurrence and severity of disease-causing mtDNA mutations. *PLoS Genet.* **13**, e1007126 (2017).

12. H. R. Elliott, D. C. Samuels, J. A. Eden, C. L. Relton, P. F. Chinnery, Pathogenic mitochondrial DNA mutations are common in the general population. *Am. J. Hum. Genet.* **83**, 254–260 (2008).

13. J. N. Wolff, E. D. Ladoukakis, J. A. Enríquez, D. K. Dowling, Mitonuclear interactions: evolutionary consequences over multiple biological scales. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **369**, 20130443 (2014).

14. G. E. Hill, *et al.*, Assessing the fitness consequences of mitonuclear interactions in natural populations. *Biol. Rev. Camb. Philos. Soc.* **94**, 1089–1104 (2019).

15. O. Matilainen, P. M. Quirós, J. Auwerx, Mitochondria and Epigenetics - Crosstalk in Homeostasis and Stress. *Trends Cell Biol.* **27**, 453–463 (2017).

16. D. M. Turnbull, R. W. Taylor, R. McFarland, Pickett S, Grady JP, Ng YS, Gorman GS, Schaefer AM, Wilson IJ, Cordell HJ. *Annals of Clinical and Translational Neurology* **5**, 333–345 (2018).

17. K. A. Phillips, P. A. Deverka, G. W. Hooker, M. P. Douglas, Genetic Test Availability And Spending: Where Are We Now? Where Are We Going? *Health Aff.* **37**, 710–716 (2018).

18. M. Blell, M. A. Hunter, Direct-to-Consumer Genetic Testing's Red Herring: "Genetic Ancestry" and Personalized Medicine. *Front. Med.* **6**, 48 (2019).

19. N. Whiffin, *et al.*, Using high-resolution variant frequencies to empower clinical genome interpretation. *Genet. Med.* **19**, 1151–1158 (2017).

20. M. Tanaka, T. Ozawa, Strand asymmetry in human mitochondrial DNA mutations. *Genomics* **22**, 327–335 (1994).

21. J. J. Faith, D. D. Pollock, Likelihood analysis of asymmetrical mutation bias gradients in vertebrate mitochondrial genomes. *Genetics* **165**, 735–745 (2003).

22. A. Reyes, C. Gissi, G. Pesole, C. Saccone, Asymmetrical directional mutation pressure in the mitochondrial genome of mammals. *Mol. Biol. Evol.* **15**, 957–966 (1998).

23. O. Zuk, *et al.*, Searching for missing heritability: designing rare variant association studies. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E455–64 (2014).

24. H. Chheda, *et al.*, Whole-genome view of the consequences of a population bottleneck using 2926 genome sequences from Finland and United Kingdom. *Eur. J. Hum. Genet.* **25**, 477–484 (2017).

25. A. Keinan, A. G. Clark, Recent explosive human population growth has resulted in an excess of rare genetic variants. *Science* **336**, 740–743 (2012).

26. L. G. Landry, N. Ali, D. R. Williams, H. L. Rehm, V. L. Bonham, Lack Of Diversity In Genomic Databases Is A Barrier To Translating Precision Medicine Research Into Practice. *Health Aff.* **37**, 780–785 (2018).

27. N. Pirastu, *et al.*, Genetic analyses identify widespread sex-differential participation bias. *bioRxiv*, 2020.03.22.001453 (2020).

28. M. W. Nachman, Deleterious mutations in animal mitochondrial DNA. *Genetica* **102-103**, 61–69 (1998).

29. M. W. Nachman, W. M. Brown, M. Stoneking, C. F. Aquadro, Nonneutral mitochondrial DNA variation in humans and chimpanzees. *Genetics* **142**, 953–963 (1996).

30. A. A. Maklakov, L. Rowe, U. Friberg, Why organisms age: Evolution of senescence under positive pleiotropy? *Bioessays* **37**, 802–807 (2015).

31. P. B. Medawar, *An Unsolved Problem of Biology: An Inaugural Lecture Delivered at University College, London, 6 December, 1951* (H.K. Lewis and Company, 1952).

32. R. Cui, *et al.*, Relaxed Selection Limits Lifespan by Increasing Mutation Load. *Cell* **178**, 385–399.e20 (2019).

33. G. C. Williams, Pleiotropy, Natural Selection, and the Evolution of Senescence. *Evolution* **11**, 398–411 (1957).

34. D. C. Wallace, Mitochondrial DNA mutations in diseases of energy metabolism. *J. Bioenerg. Biomembr.* **26**, 241–250 (1994).

35. S. Raychaudhuri, Mapping rare and common causal alleles for complex human diseases. *Cell* **147**, 57–69 (2011).

36. H. Tang, P. D. Thomas, Tools for Predicting the Functional Impact of Nonsynonymous Genetic

Variation. *Genetics* **203**, 635–647 (2016).

37. M. Dayhoff, R. Schwartz, B. Orcutt, "22 a model of evolutionary change in proteins" in *Atlas of Protein Sequence and Structure*, (National Biomedical Research Foundation Silver Spring MD, 1978), pp. 345–352.

38. E. Zuckerkandl, L. Pauling, Evolutionary Divergence and Convergence in Proteins in *Evolving Genes and Proteins*, (1965), pp. 97–166.

39. T. N. Starr, J. W. Thornton, Epistasis in protein evolution. *Protein Sci.* **25**, 1204–1218 (2016).

40. A. S. Kondrashov, S. Sunyaev, F. A. Kondrashov, Dobzhansky–Muller incompatibilities in protein evolution. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 14878–14883 (2002).

41. S. Haut, *et al.*, The deleterious G15498A mutation in mitochondrial DNA-encoded cytochrome b may remain clinically silent in homoplasmic carriers. *Eur. J. Hum. Genet.* **12**, 220–224 (2004).

42. A. Kawrykow, *et al.*, Phylo: a citizen science approach for improving multiple sequence alignment. *PLoS One* **7**, e31362 (2012).

43. S. Iantorno, K. Gori, N. Goldman, M. Gil, C. Dessimoz, Who watches the watchmen? An appraisal of benchmarks for multiple sequence alignment. *Methods Mol. Biol.* **1079**, 59–73 (2014).

44. L. Chen, P. Liu, T. C. Evans Jr, L. M. Ettwiller, DNA damage is a pervasive cause of sequencing errors, directly confounding variant identification. *Science* **355**, 752–756 (2017).

45. D. R. Smith, Revisiting published genomes with fresh eyes and new data: Revising old sequencing data can yield unexpected insights and identify errors. *EMBO Rep.* **20**, e49482 (2019).

46. C. D. Dunn, B. A. Akpınar, V. Sharma, An unusual amino acid substitution within hummingbird cytochrome c oxidase alters a key proton-conducting channel. *bioRxiv*, 610915 (2019).

47. R. R. da Fonseca, W. E. Johnson, S. J. O'Brien, M. J. Ramos, A. Antunes, The adaptive evolution of the mammalian mitochondrial genome. *BMC Genomics* **9**, 119 (2008).

48. B. Nabholz, H. Ellegren, J. B. W. Wolf, High levels of gene expression explain the strong evolutionary constraint of mitochondrial protein-coding genes. *Mol. Biol. Evol.* **30**, 272–284 (2013).

49. S. Yoshikawa, *et al.*, Redox-coupled crystal structural changes in bovine heart cytochrome c oxidase. *Science* **280**, 1723–1729 (1998).

50. C. E. Shannon, A Mathematical Theory of Communication. *Bell System Technical Journal* **27**, 379–423 (1948).

51. N. Goldman, Maximum Likelihood Inference of Phylogenetic Trees, with Special Reference to a Poisson Process Model of DNA Substitution and to Parsimony Analyses. *Syst. Biol.* **39**, 345–361 (1990).

52. M. T. Lott, *et al.*, mtDNA Variation and Analysis Using Mitomap and Mitomaster. *Curr. Protoc. Bioinformatics* **44**, 1.23.1–26 (2013).

53. M. Kimura, The role of compensatory neutral mutations in molecular evolution. *J. Genet.* **64**, 7 (1985).

54. N. J. Marini, P. D. Thomas, J. Rine, The use of orthologous sequences to predict the impact of amino acid substitutions on protein function. *PLoS Genet.* **6**, e1000968 (2010).

55. P. L. Auer, G. Lettre, Rare variant association studies: considerations, challenges and opportunities. *Genome Med.* **7**, 16 (2015).

56. S. E. Luria, M. Delbrück, Mutations of Bacteria from Virus Sensitivity to Virus Resistance. *Genetics* **28**, 491–511 (1943).

57. W. A. Rosche, P. L. Foster, Determining Mutation Rates in Bacterial Populations. *Methods* **20**, 4–17 (2000).

58. C. Cortes, V. Vapnik, Support-vector networks. *Mach. Learn.* **20**, 273–297 (1995).

59. A. Bolze, *et al.*, Selective constraints and pathogenicity of mitochondrial DNA variants inferred from a novel database of 196,554 unrelated individuals. *bioRxiv*, 1151 (2019).

60. D. T. Jones, W. R. Taylor, J. M. Thornton, The rapid generation of mutation data matrices from protein sequences. *Comput. Appl. Biosci.* **8**, 275–282 (1992).

61. Z. Zou, J. Zhang, Amino acid exchangeabilities vary across the tree of life. *Sci Adv* **5**, eaax3124 (2019).

62. C. Bris, *et al.*, Bioinformatics Tools and Databases to Assess the Pathogenicity of Mitochondrial DNA Variants in the Field of Next Generation Sequencing. *Front. Genet.* **9**, 632 (2018).

63. A. Martín-Navarro, *et al.*, Machine learning classifier for identification of damaging missense mutations exclusive to human mitochondrial DNA-encoded polypeptides. *BMC Bioinformatics* **18**, 158 (2017).

64. S. Castellana, *et al.*, High-confidence assessment of functional impact of human mitochondrial non-synonymous genome variations by APOGEE. *PLoS Comput. Biol.* **13**, e1005628 (2017).

65. S. Stewart-Brown, A. Farmer, Screening could seriously damage your health. *BMJ* **314**, 533–534 (1997).

66. Committee on Bioethics, Ethical issues with genetic testing in pediatrics. *Pediatrics* **107**, 1451–1455 (2001).

67. K. Ye, J. Lu, F. Ma, A. Keinan, Z. Gu, Extensive pathogenicity of mitochondrial heteroplasmy in healthy human individuals. *Proceedings of the National Academy of Sciences* **111**, 10654–10659 (2014).

68. H. Zhang, S. P. Burr, P. F. Chinnery, The mitochondrial DNA genetic bottleneck: inheritance and beyond. *Essays Biochem.* **62**, 225–234 (2018).

69. L. C. Greaves, *et al.*, Mitochondrial DNA mutations are established in human colonic stem cells, and mutated clones expand by crypt fission. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 714–719 (2006).

70. E. Nekhaeva, *et al.*, Clonally expanded mtDNA point mutations are abundant in individual cells of human tissues. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 5521–5526 (2002).

71. A. Bratic, N.-G. Larsson, The role of mitochondria in aging. *J. Clin. Invest.* **123**, 951–957 (2013).

72. G. Fayet, *et al.*, Ageing muscle: clonal expansions of mitochondrial DNA point mutations and deletions cause focal impairment of mitochondrial function. *Neuromuscul. Disord.* **12**, 484–493 (2002).

73. P. F. Chinnery, A. Gomez-Duran, Oldies but Goldies mtDNA Population Variants and

Neurodegenerative Diseases. *Front. Neurosci.* **12**, 883 (2018).

74. S. Marom, M. Friger, D. Mishmar, MtDNA meta-analysis reveals both phenotype specificity and allele heterogeneity: a model for differential association. *Sci. Rep.* **7**, 43449 (2017).

75. K. Katoh, D. M. Standley, MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

76. A. Larsson, AliView: a fast and lightweight alignment viewer and editor for large datasets. *Bioinformatics* **30**, 3276–3278 (2014).

77. W. Shen, S. Le, Y. Li, F. Hu, SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File Manipulation. *PLoS One* **11**, e0163962 (2016).

78. S. A. Chamberlain, E. Szöcs, taxize: taxonomic search and retrieval in R. *F1000Res.* **2**, 191 (2013).

79. M. N. Price, P. S. Dehal, A. P. Arkin, FastTree 2--approximately maximum-likelihood trees for large alignments. *PLoS One* **5**, e9490 (2010).

80. A. Löytynoja, A. J. Vilella, N. Goldman, Accurate extension of multiple sequence alignments using a phylogeny-aware graph algorithm. *Bioinformatics* **28**, 1684–1691 (2012).

81. S. Capella-Gutiérrez, J. M. Silla-Martínez, T. Gabaldón, trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973 (2009).

82. E. F. Pettersen, *et al.*, UCSF Chimera—a visualization system for exploratory research and analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).

83. P. Pavlidis, W. S. Noble, Matrix2png: a utility for visualizing matrix data. *Bioinformatics* **19**, 295–296 (2003).

84. D. Meyer, *et al.*, e1071: Misc functions of the Department of Statistics (e1071), TU Wien. *R package version* **1** (2014).

85. C.-C. Chang, C.-J. Lin, LIBSVM: A library for support vector machines. *ACM transactions on intelligent systems and technology (TIST)* **2**, 27 (2011).

86. P. Cortez, "A tutorial on using the rminer R package for data mining tasks" (Universidade do Minho. Escola de Engenharia (EEng), 2015).

87. A. Karatzoglou, A. Smola, K. Hornik, A. Zeileis, kernlab-an S4 package for kernel methods in R. *J. Stat. Softw.* **11**, 1–20 (2004).

88. P. Cortez, M. J. Embrechts, Using sensitivity analysis and visualization techniques to open black box data mining models. *Inf. Sci.* **225**, 1–17 (2013).

89. T. Tsukihara, *et al.*, The whole structure of the 13-subunit oxidized cytochrome c oxidase at 2.8 A. *Science* **272**, 1136–1144 (1996).

Figure 1

Figure 2

a

b   **c**

side view   **top view**

Figure 3

Figure 4

a

MITOMAP confirmed
pathogenic protein
n = 34

****

MITOMAP protein
'polymorphism'
n= 2588

b

GenBank full-length samples

100000
10000
1000
100
10

****

1000   2000

MITOMAP annotated
protein 'polymorphism'

c

10000
1000
100
10

****

100   200   300

MITOMAP annotated
tRNA 'polymorphism'

d

GenBank full-length samples

10000
1000
100
10

****

200   400   600

MITOMAP annotated
rRNA 'polymorphism'

Figure 5

Figure 6

Figure 7