Classification

BIOLOGICAL SCIENCES: Microbiology

Title

Chaperone mediated coupling of subunit availability to activation of proton motive

force-driven flagellar Type III Secretion

Running title

Subunit-coupled activation of flagellar export

Authors

Owain J. Bryant[a], Betty Y-W. Chung[a] and Gillian M. Fraser[a,1]

Affiliations

[a]Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge,

CB2 1QP, United Kingdom.

Corresponding author

[1]Gillian M. Fraser

Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge,

CB2 1QP, United Kingdom.

Phone: +44 1223 330245

Email: gmf25@cam.ac.uk

Keywords

bacterial flagella biogenesis; Type III Secretion System; proton motive force;

protein export

Abstract

Bacterial flagella are assembled from thousands of protein subunits that are unfolded

and exported across the cell membrane by a specialized flagellar Type III Secretion

System (fT3SS). Export of subunit cargo is fuelled by the proton motive force (pmf),

facilitated by an export ATPase that is evolutionarily related to the F1 ATPase. The

FliJ stalk component of the ATPase binds the flagellar export gate, enabling it to

utilise the pmf electric potential ($\Delta\Psi$) to drive efficient export of subunit cargo. What

is unclear is how FliJ activation of the export gate is regulated to prevent constitutive

proton influx and wasteful dissipation of pmf when no subunit cargo is available. FliJ,

in addition to activating the export gate, recruits unladen export chaperones and

transfers them to cognate subunit cargo, creating a local cycle of chaperone-subunit

binding and release at the fT3SS. To investigate whether chaperones might regulate

FliJ activation of the export gate, we isolated chaperone variants that were defective

for FliJ binding yet still able to bind cognate subunits and other export components.

Disruption of the chaperone-FliJ interaction attenuated motility and cognate subunit

export. *In vitro* and *in vivo* competitive binding assays showed that chaperones and

the export gate component FlhA compete for binding FliJ. Our data indicate that

unladen chaperones, which would be present in the cell when subunit levels are low,

sequester FliJ to prevent activation of the export gate. This suggests a mechanism

whereby chaperones couple availability of export cargo to utilization of pmf by the

fT3SS.

Significance

Type III Secretion Systems (T3SS) are used by bacteria to build flagella, rotary

nanomachines for cell motility, and related injectosomes, specialised molecular

syringes that deliver bacterial effector proteins into host cells to subvert their function.

T3SS activity uses two power sources – the high-energy ATP molecule and the

proton motive force (pmf), chemical and electric potential stored in a proton gradient

across the bacterial membrane. Here, we show that use of pmf by the flagellar T3SS

is regulated by cytoplasmic chaperones that bind and deliver cargo proteins to the

T3SS for export. We demonstrate that when cargo availability is low, the unladen

chaperones bind the T3SS to reduce its activity, minimising energy use and

preventing wasteful dissipation of the pmf.

Introduction

Bacterial flagella are large macromolecular rotary motors that enable cell motility. In

Salmonella, the long helical flagellar filament propeller that extends from the cell

surface is connected to a flexible hook attached to a basal body spanning the cell

envelope. These flagellar substructures are assembled in a strict order, with the

basal body being constructed first, followed by the hook and then the filament (1, 2).

The filament comprises thousands of copies of the major filament subunit, FliC

(flagellin), together with three minor filament-class subunits: eleven copies each of

FlgK and FlgL form a junction connecting the semi-rigid filament to the hook, and five

FliD subunits assemble at the distal tip of the nascent flagellum to form a cap that

facilitates folding and incorporation of flagellin into the growing filament (3).

Export of the filament subunits is facilitated by the cytoplasmic chaperones FlgN, FliT

and FliS, which bind the C-terminal domains of their cognate structural subunits

(FlgN binds FlgK and FlgL, FliT binds FliD, and FliS binds FliC) (4-6). The

chaperones then pilot their cognate subunits to the specialized flagellar Type III

secretion system (fT3SS) machinery at the base of each flagellum (7-11).

Chaperone-subunit complexes initially dock at the FliI component of the flagellar

export ATPase, which is evolutionarily related to the F1-ATPase (12). Chaperoned

subunits are then thought to interact with the integral membrane export gate

component FlhA, where chaperones are released and the subunit cargo is unfolded

for translocation across the inner membrane, *via* the FlhAB-FliPQR export gate, into

a narrow central channel in the growing flagellum (9-13). Once released from their

cognate subunit cargo, the unladen chaperones FlgN and FliT - but not the flagellin

chaperone, FliS - are recruited by the FliJ stalk component of the ATPase, which

transfers them to newly synthesised cognate cargo to create a local cycle of

chaperone-subunit binding at the membrane fT3SS (8). This is thought to promote

export of the minor filament subunits that form the hook-filament junction and the

filament cap, which are required for initiation of filament assembly (14).

Export of flagellar subunit cargo across the membrane is powered by the pmf and the

flagellar ATPase complex(15, 16). Specifically, the ATPase is thought to facilitate

unfolding of subunit cargo (17). In addition, analogous to F- and V-type ATPases,

ATP hydrolysis is proposed to drive rotation of the ATPase stalk, FliJ, which interacts

with the nonameric export gate component, FlhA, converting the gate into a highly

efficient proton-protein antiporter that utilises the $\Delta\Psi$ electric component of the pmf

(18). What is unclear is how FliJ activation of the export gate is regulated in the

absence of subunit cargo to prevent constitutive proton influx and wasteful

dissipation of the pmf. One way in which export gate activation could be regulated is

by sequestration of FliJ by other binding partners, specifically the export chaperones

FlgN and FliT. A role for chaperones in the regulation of T3SS activity is supported

by the finding that the *Psuedomonas* virulence T3SS chaperone PcrG binds the FliJ

homologue, PscO, and that loss of this interaction results in upregulation of effector

secretion and renders the T3SS more resistant to pmf collapse (19). These

observations suggest that the chaperone-FliJ interaction is central to the regulation of

T3SS activity, however, the mechanistic basis of this regulation remains obscure.

Here, we identify point mutant variants of the chaperones FlgN and FliT that are

specifically defective for binding to the FliJ ATPase stalk yet retain the ability to bind

cognate subunits and the FlhA component of the fT3SS export gate. We show, *in*

vitro and *in vivo*, that the FlgN and FliT chaperones compete with FlhA for binding to

FliJ, and that chaperone sequestration of FliJ blocks the activating interaction

between FliJ and the export gate. This suggests a mechanism whereby binding of

FliJ by unladen chaperones acts as a signal to the export machinery that subunit

cargo is unavailable and, accordingly, export activity is limited to prevent wasteful

dissipation of the pmf. This model is supported by ribosome profiling data, which

show that the cytoplasmic ratios of cognate subunits to chaperones are low for FlgN

and FliT, relative to the ratio of flagellin to its chaperone, FliS, suggesting that levels

of unladen FlgN and FliT would be a sensitive proxy measure of intracellular flagellar

subunit levels. Our data reveal the mechanistic basis by which chaperones modulate

the FliJ-FlhA interaction to regulate energy use by the T3SS in response to subunit

availability.

Results and Discussion

Isolation of chaperone variants that are specifically defective for FliJ binding

To investigate the function of chaperone binding to the ATPase stalk, FliJ, we sought

to identify variants of the FlgN and FliT chaperones that did not bind FliJ yet could

interact with cognate subunits and the FlhA component of the fT3SS export gate. To

do this, we first constructed mutant alleles of *flgN* encoding variants with three-

residue deletions in the FlgN α3 helix (residues 70-102), which contains the

overlapping binding sites for cognate subunits and FliJ (Fig. 1; *SI Appendix*, Fig. S1)

(8). Screening of the FlgN deletion variants using affinity chromatography pull-down

assays with GST- FlgK subunit or GST-FliJ identified a single deletion variant,

FlgNΔ76-78, that could bind its cognate subunit but not the ATPase stalk (Fig. 1*B*).

Protein sequence comparisons revealed FlgN W_{78} to be highly conserved and

replacement of this residue with alanine (FlgN-W_{78}A) severely reduced FlgN binding

to FliJ but did not affect binding to FlgK subunit (Fig. 1*C*; FlgNΔ90-100, which binds

neither cognate subunit nor FliJ, and wildtype FlgN were included as negative and

positive controls, respectively) (8). Affinity chromatography pull-down assays

confirmed that FlgNΔ76-78 and FlgN-W$_{78}$A could still bind the cytoplasmic domain of

the export gate component FlhA (GST-FlhAc; Fig. 1*C*).

Having identified residues in the FlgN α3 helix that, when mutated, could decouple

chaperone binding of cognate subunits and FliJ, we targeted conserved residues in

the equivalent α3 helix of FliT for site directed mutagenesis to alanine (Fig. 1*D*; *SI*

Appendix, Fig. S1). To facilitate screening for FliJ binding, mutations were introduced

into a *fliT* allele encoding a truncated variant, FliT$_{94}$, which lacks the α4 helix and

shows enhanced binding to FliJ *in vitro* (20). Affinity chromatography pull-down

assays identified five FliT$_{94}$ variants (I$_{68}$A, L$_{72}$A, N$_{74}$A, E$_{75}$A and L$_{78}$A) that showed

severely reduced binding to GST-FliJ (Fig. 1*E*). Three of these FliJ non-binders -

FliT$_{94}$-I$_{68}$A, FliT$_{94}$-L$_{72}$A and FliT$_{94}$-N$_{74}$A - had previously been shown to bind cognate

subunit FliD (20). For this reason, and after assessing the expression and stability of

these three FliT$_{94}$ variants in *Salmonella* (not shown), we chose to further investigate

the function of truncated FliT$_{94}$-L$_{72}$A and full-length FliT-L$_{72}$A.

Binding of FliT-L$_{72}$A and truncated FliT$_{94}$-L$_{72}$A to cognate subunit FliD was confirmed

using pull-down assays (GST-FliD; *SI Appendix*, Fig. S2). To assess binding to the

export gate component FlhA, pull-down assays were carried out using GST-FlhAc

that had been incubated with cell lysates containing FliD cognate subunit, which is

required for high affinity binding of FliT to FlhA, and either wildtype FliT or full-length

FliT-L$_{72}$A (Fig. 1*F*) (9). The FliT-Y$_{106}$A variant, which cannot bind FlhA, was included

as a negative control (Fig. 1*F*) (11). Data from the pull-down assays indicated that,

when bound to cognate subunit FliD, the FliT-L$_{72}$A variant can bind the FlhA export

gate (Fig. 1*F*).

Attenuation of cell motility and cognate subunit export by specific disruption

of the chaperone-FliJ interaction

Having identified chaperone variants (FlgNΔ76-78, FlgN-W$_{78}$A, FliT-L$_{72}$A) that were

specifically defective for FliJ binding yet retained the ability to bind cognate subunits

and the fT3SS export gate, we went on to investigate the importance of the

chaperone-FliJ interaction for cell motility and subunit export. To enable expression

of chaperones at physiological levels, *Salmonella* strains were engineered to carry

variant chaperone genes (*flgNΔ76-78*, *flgN-W$_{78}$A* or *fliT-L$_{72}$A*) at the natural genetic

loci, replacing the wildtype chaperone gene. As negative controls, isogenic

Salmonella strains in which the chaperone genes were either deleted (Δ*flgN* or Δ*fliT*)

or replaced with variant genes encoding export defective chaperones unable to bind

either cognate subunit (*flgNΔ90-100*) or the FlhA export gate (*fliT-Y$_{106}$A*) were also

constructed. These strains were assessed for swimming and swarming motility, and

for subunit export efficiency relative to wildtype *Salmonella* (Fig. 2).

Cell populations producing the FlgNΔ76-78 or FlgN-W$_{78}$A variants, which cannot bind

FliJ, showed slightly decreased swimming (70-90%) and swarming (75-90%) motility

relative to wildtype *Salmonella*, though their motility was significantly better than that

of the *flgN* null strain or cells producing FlgNΔ90-100, which binds neither cognate

subunits nor FliJ (Fig. 2*A* and *SI Appendix*, Fig. S2). Secretion of cognate subunit

FlgK was also reduced to *c.*50% of wildtype in cells producing either FlgNΔ76-78 or

FlgN-W$_{78}$A, and this was accompanied by a concomitant increase in soluble

extracellular FliC (to 140-150% of wildtype), suggesting that assembly of the FlgKL

hook-filament junction was impaired, causing inefficient initiation of filament

assembly and increased release of unpolymerized FliC into culture supernatants

(Fig. 2*B*) (4). As expected, this secretion defect was more severe in the *flgN* null

strain and in cells producing FlgNΔ90-100 (Fig. 2*B*).

Analysis of motility and export phenotypes associated with the FliT chaperone is

complicated by the additional function of FliT as a negative regulator of the flagellar

master transcriptional activator $FlhD_4C_2$ (21-23). Cells lacking FliT show increased

cellular levels of $FlhD_4C_2$, and a concomitant increase in flagellar gene expression

that partially alleviates the motility and export defects typically associated with loss of

chaperone activity (4). The increased flagellar gene expression in the *fliT* null is

reflected by an increase in the cellular level of non-cognate subunit FlgK, which was

observed to be 1.4-fold higher than in wildtype (*SI Appendix*, Fig. S3). Similarly, the

level of FlgK was 1.4-fold higher in the strain producing the export defective FliT-

$Y_{106}A$ (*SI Appendix*, Fig. S3), indicating increased flagellar gene expression that may

be caused by sequestration of FliT-$Y_{106}A$ by cytosolic FliD, which would prevent FliT-

$Y_{106}A$ binding to FlhC and targeting it for degradation (21-23). However, cellular

levels of FlgK in cells producing FliT-$L_{72}A$ were similar to wildtype, suggesting that

while FliT-$L_{72}A$ cannot bind FliJ, it retains the ability to bind FlhC and thus regulate

flagellar gene expression (*SI Appendix*, Fig. S3).

Cells producing FliT-$L_{72}A$ displayed wildtype swimming motility but showed a

significant defect in population swarming motility relative to wildtype *Salmonella*

(*c.*50% of wildtype; Fig. 2A). Moreover, export of cognate subunit FliD was reduced

to *c.*50% of wildtype in the FliT-$L_{72}A$ strain. These data suggest that the FliT-FliJ

interaction may be required for export of sufficient FliD to promote the increased

flagellation required for swarming (24, 25). Cells producing FliT-$Y_{106}A$, which cannot

bind FlhA, showed only a slight defect in swimming and swarming motility, while the

fliT null strain displayed a marginal reduction in swimming to *c.*80% of wildtype, yet

an increase in swarming motility (to *c.*110% of wildtype; Fig. 2B). Both the *fliT* null

and FliT-$Y_{106}A$ strain showed a significant reduction in FliD export, to 50-60% of

wildtype, accompanied by concomitant release of unpolymerized FliC to culture

supernatants (to 110% of wildtype; Fig. 2B). The confounding influence of increased

flagellar gene expression in these strains, however, makes it difficult to interpret the

motility and export phenotypes of the *fliT* null and FliT-Y_{106}A controls, particularly as

overexpression of *flhDC* is known to overcome motility defects associated with

chaperone loss (26, 27).

Notwithstanding the difficulties in interpreting the motility and flagellar export

phenotypes associated with loss or mutation of *fliT*, the data for FlgNΔ76-78, FlgN-

W_{78}A and FliT-L_{72}A indicate that specific loss of chaperone interactions with FliJ,

which would break the local chaperone cycle at the fT3SS, result in attenuation of

cognate subunit export and cell motility.

Regulation of fT3SS export gate activation by chaperone sequestration of the

FliJ ATPase stalk

The identification and characterisation of FlgN and FliT chaperone variants that were

specifically unable to bind the FliJ ATPase stalk opened up the possibility of further

investigations into whether chaperones might regulate FliJ activation of the fT3SS.

To test whether unladen chaperones could dock at FliJ and block FliJ binding to the

FlhA export gate, we developed an *in vitro* competition assay in which FlhA$_C$ binding

to cobalt bead-immobilized (His)$_6$-FliJ was assessed in the presence of chaperone

variants that could interact with FliJ but not FlhA$_C$ (FliT$_{94}$ or FlgN-Y_{122}A) or were

defective in binding both FliJ and FlhA$_C$ (FliT$_{94}$-L_{72}A or FlgN-Y_{122}AΔ76-78; Fig. 3*A*).

As anticipated, we found that FlhA$_C$ could bind (His)$_6$-FliJ in the presence of

chaperone variants that could not bind FliJ (Fig. 3*B*). However, when assays were

performed with chaperones that could bind FliJ, the amount of FlhA$_C$ pulled down by

FliJ was severely reduced (Fig. 3*B*).

The *in vitro* competition assay established that the FlgN and FliT chaperones could

block FliJ binding to $FlhA_C$, but did not show if activation of the $FlhA_C$ export gate was

prevented. To investigate this, an *in vivo* competition assay was developed in which

the effect of FlgN-FliJ binding on $FlhA_C$ export activity was tested under conditions

where the availability of cognate subunits for FlgN was limited. Specifically, FlgN

variants that could bind FliJ but not $FlhA_C$ (FlgN-Y_{122}A) or were defective in binding

both FliJ and $FlhA_C$ (FlgN-Y_{122}AΔ76-78) were expressed *in trans* in Δ*flgE* strain, in

which flagellar assembly does not progress beyond completion of the rod and the

FlgN cognate subunits, FlgK and FlgL, are produced at very low levels (28-30).

Under these conditions, we hypothesised that unladen FlgN might disrupt the FliJ-

FlhA interaction, preventing activation of the flagellar export gate and inhibiting

subunit export. Assays of subunit export in a strain producing FlgN-Y_{122}A, which

binds FliJ, resulted in a *c.*75% reduction in FlgD export compared to the strain

producing FlgN-Y_{122}AΔ76-78, which cannot bind FliJ (Fig *3C*).

Taken together, the data from the *in vitro* and *in vivo* competitive binding assays

indicate that when cognate subunit availability is low, unladen chaperones block the

FliJ-FlhA interaction and prevent FliJ-dependent activation of the flagellar export

machinery.

Ribosome profiling reveals low cytoplasmic ratios of minor filament subunits

to cognate chaperones, FlgN and FliT

Our accumulating data suggested that, in addition to piloting subunit cargo to the

fT3SS, the FlgN and FliT chaperones might be conditional regulators of fT3SS

activity that sense and respond to the availability of subunit cargo. To function

effectively in the cell, this mechanism - in which unladen chaperones are a proxy

measure for subunit availability - would likely require low cytoplasmic ratios of FlgN

and FliT to their cognate subunits. To obtain an indication of the cytoplasmic ratios of

chaperones and their cognate subunits, we used RNA sequencing (RNA-Seq) and

ribosome profiling (Ribo-Seq) (31-34) to assess the *Salmonella* global transcript

abundance and translatome, respectively, at late-log growth phase when flagellar

gene expression is at its peak. Analysis of these data confirmed that flagellar mRNAs

are amongst the most abundant and highly translated in *Salmonella* (Fig. 4*A*). The

ribosome profiling data indicated that the ratio of FlgN to its cognate subunits (FlgK

and FlgL) was 1:3, and the ratio of FliT to its cognate subunit FliD was 1:4. These

chaperone-subunit ratios are low when compared to the 1:43 ratio of FliS chaperone

to its cognate subunit FliC (Fig. 4*B*), suggesting that the presence of unbound FlgN

and FliT in the cell would provide a sensitive proxy measure of cytoplasmic flagellar

subunit levels.

Taken together with our other findings, the low ratios of FlgN and FliT to their

cognate subunits strengthen the view that these unladen chaperones couple subunit

availability to pmf-driven export by modulating FliJ-dependent activation of the $FlhA_c$

export gate.

Conclusions

The synthesis and operation of bacterial flagella is energetically expensive, using an

estimated 2% of the cell's energy, and numerous regulatory mechanisms have

evolved to increase efficiency of flagella biogenesis and function. Here, we have

uncovered the mechanism by which fT3SS chaperones couple the availability of

subunit cargo to the energy efficiency of flagellar export. By identifying point mutant

variants of the FlgN and FliT chaperones that were specifically unable to bind the FliJ

stalk component of the fT3SS ATPase, we showed that the chaperone-FliJ

interaction prevents activation of the FlhA export gate component when subunit

availability is low, reducing wasteful dissipation of the pmf by the fT3SS machinery.

Chaperone modulation of fT3SS activity may be important when *Salmonella*

encounters environmental conditions in which flagellar gene expression is down-

regulated, including during the early stages of host infection (35-38). Our proposed

mechanism may also function in related virulence T3SS (vT3SS), as it is compatible

with previous observations showing that vT3SS FliJ homologues bind a subset of

secretion chaperones, and that loss of this interaction increases effector secretion

and renders the T3SS more resistant to pmf collapse (19, 39, 40). Our data suggest

a model for a local chaperone cycle at the T3SS machinery that (i) facilitates efficient

docking and export of cognate cargo when subunit/effector availability is high, and (ii)

prevents constitutive proton influx and wasteful dissipation of the pmf when cargo

availability is low (Fig. 5).

Materials and Methods

Bacterial strains, plasmids and growth conditions

Wildtype *Salmonella enterica* serovar Typhimurium (*S. typhimurium*) SJW1103 is

motile (41). The Δ*fliT*::K_m^R mutant in which the *fliT* gene was replaced by a

kanamycin resistance cassette was constructed using the λ Red recombinase

system (42). Strains containing chromosomally encoded FliT or FlgN variants were

constructed by *aph*-I-SceI Kanamycin resistance cassette replacement using

pWRG730 (42). Bacteria were cultured at 30–37°C in Luria-Bertani (LB) broth

containing, where appropriate, ampicillin (100 mg/ml) or chloramphenicol (20 mg/ml)

and collected by centrifugation (6,000g for 10 min). Recombinant proteins were

expressed in *Salmonella* from the isopropyl β-D-thiogalactoside-inducible (IPTG)-

inducible plasmid pTrc99a (43).

Recombinant proteins for purification were expressed in *E.coli* C41 (44) from (IPTG)-

inducible pACT7 (45) or pGEX-4T3 (46). To construct recombinant plasmids

encoding wildtype or derivative genes, *Salmonella* genes were amplified from

chromosomal DNA by PCR or overlap-extension PCR using Q5 High-Fidelity DNA

polymerase. PCR products were inserted BamHI/XhoI into pGEX-4T3 or NdeI/BamHI

into pACT7 and pTrc99a. Inserts were verified by DNA sequencing (Department of

Biochemistry, University of Cambridge). A full list and description of strains and

plasmids used in this study can be found in *SI Appendix*, Tables S1 and S2.

Affinity chromatography co-purification assays.

Co-purification of proteins expressed in *E.coli* C41 was performed with glutathione-

sepharose 4B. Cells were resuspended in buffer A (50 mM sodium phosphate pH7.4,

150 mM NaCl, 1mM β–mercaptoethanol) and mechanically lysed with a Constant

Systems cell disruptor at 30,000 kPa. Cleared lysates containing GST-tagged

proteins were incubated with affinity resin for 1 hour. After extensive washing with

buffer A, cell lysates containing overexpressed untagged prey proteins were then

incubated with the resin pre-bound GST-bait. Resins were washed extensively with

buffer A and proteins eluted by boiling in SDS-PAGE loading buffer. Proteins were

separated by SDS-PAGE and immunoblotted.

Flagellar subunit export assay

Salmonella strains were cultured at 37 °C in LB broth containing ampicillin and 100

µM IPTG to mid-log phase (OD_{600nm} 0.6-0.8). Cells were centrifuged (6000g, 5 min),

resuspended in fresh media and grown for a further 60 min at 37 °C. Cells were

pelleted by centrifugation (16,000g, 5 min) and the supernatant passed through a 0.2

µm nitrocellulose filter. Supernatant proteins were precipitated with 10%

trichloroacetic acid (TCA) and 1% Triton-X100 on ice for 1 hour, pelleted by

centrifugation (16,000g, 10 min), washed with ice-cold acetone and resuspended in

SDS-PAGE loading buffer (volumes calibrated according to cell densities). Fractions

were analysed by immunoblotting.

Motility assays

For swimming motility, *Salmonella* strains were grown in LB broth to A600nm 1. Two microliters of culture were inoculated into soft tryptone agar (0.25% agar, 10 g/L tryptone, 5g/L NaCl). Plates were incubated at 37 °C for between 4 and 6 hours. For swarming motility, one microliter of overnight cultures grown in LB broth was inoculated onto dried tryptone agar plates (0.6% agar, 10g/L tryptone, 5g/L NaCl) supplemented with 0.3% glucose and incubated at 30 °C for 16 hours.

Chaperone-FliJ *in vitro* competition assay

Recombinant proteins were expressed in *E.coli* C41 and cells were mechanically lysed as described for the co-purification assays. Cleared lysates containing His-tagged FliJ were incubated with cobalt affinity resin for 1 hour. After extensive washing, cell lysates containing overexpressed untagged FlhA$_C$ was added to each resin fraction and an equal volume of wildtype FliT94, FliT94 L72A, wildtype FlgN or FlgN W78A was added. Proteins were incubated for 1 hour, the resin was washed extensively with buffer A and proteins were eluted by boiling in SDS-PAGE loading buffer. Proteins were separated by SDS-PAGE and immunoblotted.

FlgN-FliJ *in vivo* competition assay

Salmonella flgE null strains carrying either pTrc99a encoding *flgN*-Y$_{122}$AΔ76-78 or *flgN*-Y$_{122}$A were grown in LB broth containing ampicillin and inducing agent (50 µM IPTG) to A600 1.0. Cells were centrifuged (6000g, 5 min) and resuspended in fresh media and grown for a further 60 min at 37°C. Cells were pelleted by centrifugation (16,000g, 5 min) and the supernatant passed through a 0.22 µm nitrocellulose filter (Sartorius). Proteins were precipitated with 10% trichloroacetic acid (TCA) and 1% Triton- X100 on ice for 1 hour, pelleted by centrifugation (16,000g, 10 min, 4°C), washed with ice-cold acetone and resuspended in SDS-PAGE loading buffer

(volumes calibrated according to cell densities). Fractions were analysed by

immunoblotting.

Ribosome profiling and RNA-Seq

Libraries were prepared essentially as described previously (47, 48) with the

following modifications: Cells were grown in LB at 37°C with 180 RPM shaking to

A600 1.0. Chloramphenicol was added to the culture to a final concentration of 1500

μg/ml, followed by rapid cooling of the culture and harvesting of cells by

centrifugation (6,000 g, 1 min, 4°C). The cell pellet was quickly resuspended in 1 ml

ice cold bacterial profiling buffer (20 mM Tris-Cl pH 7.5, 140 mM KCl, 5 mM $MgCl_2$,

1500 μg/ml chloramphenicol, 0.5 mM dithiothreitol (DTT), 0.5% NP40, 1% Triton X-

100, 2.5% Sucrose) and flash-frozen for cryo-grinding in liquid nitrogen. The frozen

powder was thawed and clarified by centrifugation (13,000 g, 2 mins, 4°C) followed

by adjustment of $A_{254\ nm}$10. Lysates were either snap frozen in liquid nitrogen for

storage at -80°C or nuclease treated with RNase 1 (700U, Ambion) followed by

ribosomal RNA depletion using a bacterial ribozero kit (Illumina) prior to library

preparation for ribosome profiling. For parallel RNA-Seq, total RNA was extracted

from corresponding lysates followed ribosomal RNA depletion using the bacterial

ribozero kit (Illumina) prior to library preparation. For Ribosome profiling, rRNA was

further depleted with duplex-specific nuclease as described previously (47).

Ribosome profiling and RNA-Seq libraries were pooled and sequenced using the

NextSeq® 500/550 platform and data were trimmed and mapped to the

transcriptome assembly of *Salmonella enterica* subsp. *enterica* serovar Typhimurium

strain ST4/74 (GenBank accession CP002487.1). Paired ribosome profiling and

RNA-Seq analysis was performed with riboSeqR as previously described (48, 49).

Acknowledgements

We thank Colin Hughes and Lewis Evans for useful discussions, and Sangita Ahmed for technical assistance. This work was funded by grants from the Biotechnology and Biological Sciences Research Council (BB/M007197/1) to G.M.F, Medical Research Council [MR/R021821/1] to B.Y-W.C., and a University of Cambridge John Lucas Walker Studentship to O.J.B.

Footnotes

The authors declare no conflict of interest.

Author contributions

O.J.B, B.Y-W.C. and G.M.F. designed the research; O.J.B performed the research; O.J.B, B.Y-W.C. and G.M.F. analyzed the data and wrote the paper.

References

1.   L. D. B. Evans, C. Hughes, G. M. Fraser, Building a flagellum outside the bacterial cell. *Trends. Microbiol*. **22**, 566-572 (2014)

2.   L. D. B. Evans, C. Hughes, G. M. Fraser. Building a flagellum in biological outer space. *Microb. Cell* **1**, 64-66 (2014)

3.   K. Yonekura, S. Maki, D. G. Morgan, D. J. DeRosier, F. Vonderviszt *et al.* The bacterial flagellar cap as the rotary promoter of flagellin self-assembly. *Science* **290** (2000)

4.   J. C. Bennett, J. Thomas, G. M. Fraser, C. Hughes, Substrate complexes and domain organization of the *Salmonella* flagellar export chaperones FlgN and FliT. *Mol. Microbiol.* **39**, 781–791 (2001).

5.   G. M. Fraser, J. C. Bennett, C. Hughes, Substrate-specific binding of hook-associated proteins by FlgN and FliT, putative chaperones for flagellum assembly. *Mol. Microbiol.* **32**, 569–580 (1999).

6.   F. Auvray, J. Thomas, G. M. Fraser, C. Hughes, Flagellin polymerisation

control by a cytosolic export chaperone. *J. Mol. Biol.* **308**, 221–229 (2001).

7.   J. Thomas, G. P. Stafford, C. Hughes, Docking of cytosolic chaperone-

substrate complexes at the membrane ATPase during flagellar type III protein

export. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 3945-3950 (2004).

8.   L. D. B. Evans, G. P. Stafford, S. Ahmed, G. M. Fraser, C. Hughes, An escort

mechanism for cycling of export chaperones during flagellum assembly. *Proc.*

Natl. Acad. Sci. U.S.A. **103**, 17474–9 (2006).

9.   G. Bange, N. Kümmerer, C. Engel, G. Bozkurt, K. Wild *et al*. FlhA provides the

adaptor for coordinated delivery of late flagella building blocks to the type III

secretion system. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 11295–11300 (2010).

10.   T. Minamino, M. Kinoshita, N. Hara, S. Takeuchi, A. Hida, *et al.* Interaction of

a bacterial flagellar chaperone FlgN with FlhA is required for efficient export of

its cognate substrates. *Mol. Microbiol.* **83**, 775–788 (2012).

11.   M. Kinoshita, N. Hara, K. Imada, K. Namba, T. Minamino, Interactions of

bacterial flagellar chaperone-substrate complexes with FlhA contribute to co-

ordinating assembly of the flagellar filament. *Mol. Microbiol.* **90**, 1249–1261

(2013).

12.   T. Ibuki, K. Imada, T. Minamino, T. Kato, T. Miyata *et al.* Common architecture

of the flagellar type III protein export apparatus and F- and V-type ATPases.

Nat. Struct. Mol. Biol. **18**, 277–282 (2011).

13.   L. Kuhlen, P. Abrusci, S. Johnson, J. Gault, J. Deme *et al.* Structure of the

core of the type iii secretion system export apparatus. *Nat. Struct. Mol. Biol.*

25, 583-590 (2018).

14.   M. Homma, K. Kutsukake, T. Iino, S. Yamaguchi, Hook-associated proteins

essential for flagellar filament formation in *Salmonella* typhimurium. *J.*

Bacteriol. **157**, 100-108 (1984).

15.   T. Minamino, K. Namba, Distinct roles of the FliI ATPase and proton motive

471  force in bacterial flagellar protein export. *Nature* **451**, 485–488 (2008).

472 16. K. Paul, M. Erhardt, T. Hirano, D. F. Blair, K. T. Hughes, Energy source of

473  flagellar type III secretion. *Nature* **451**, 489-492 (2008).

474 17. Y. Akeda, J. E. Galán, Chaperone release and unfolding of substrates in type

475  III secretion. *Nature* **437**, 911–915 (2005).

476 18. T. Minamino, Y. V. Morimoto, N. Hara, K. Namba, An energy transduction

477  mechanism used in bacterial flagellar type III protein export. *Nat. Commun.* **2**,

478  475 (2011).

479 19. P. C. Lee, S. E. Zmina, C. M. Stopford, J. Toska, A. Rietsch, Control of type III

480  secretion activity and substrate specificity by the cytoplasmic regulator PcrG.

481  *Proc. Natl. Acad. Sci. U.S.A.* **111**, E2027–E2036 (2014).

482 20. K. Imada, T. Minamino, M. Kinoshita, Y. Furukawa, K. Namba, Structural

483  insight into the regulatory mechanisms of interactions of the flagellar type III

484  chaperone FliT with its binding partners. *Proc. Natl. Acad. Sci. U.S.A.* **107**,

485  8812–8817 (2010).

486 21. S. Yamamoto, K. Kutsukake, FliT acts as an anti-$FlhD_2C_2$ factor in the

487  transcriptional control of the flagellar regulon in *Salmonella enterica* serovar

488  typhimurium. *J. Bacteriol.* **188**, 6703–6708 (2006).

489 22. K. Kutsukake, T. Ikebe, S. Yamamoto, Two novel regulatory genes, fliT and

490  fliZ, in the flagellar regulon of *Salmonella*. *Genes. Genet. Syst.* **74**, 287–292

491  (1999).

492 23. Y. Sato, A. Takaya, C. Mouslim, K. T. Hughes, T. Yamamoto, FliT selectively

493  enhances proteolysis of FlhC Subunit in $FlhD_4C_2$ complex by an ATP-

494  dependent protease, ClpXP. *J. Biol. Chem.* **289**, 33001–33011 (2014).

495 24. G. M. Fraser, C. Hughes, Swarming motility. *Curr. Opin. Microbiol.* **2**, 630-635

496  (1999).

497 25. D. B. Kearns, A field guide to bacterial swarming motility. *Nat. Rev. Microbiol.*

498  **8**, 634-644 (2010).

26.   A. Dufour, R. B. Furness, C. Hughes, Novel genes that upregulate the *Proteus*

mirabilis flhDC master operon controlling flagellar biogenesis and swarming.

Mol. Microbiol. **29**, 741-751 (1998).

27.   P. Aldridge, J. Karlinsey, K. T. Hughes, The type III secretion chaperone FlgN

regulates flagellar assembly via a negative feedback loop containing its

chaperone substrates FlgK and FlgL. *Mol. Microbiol.* **49**, 1333-1345 (2003).

28.   T. Hirano, T. Minamino, K. Namba, R. M. Macnab, Substrate specificity

classes and the recognition signal for *Salmonella* type III flagellar export. *J.*

Bacteriol. **185**, 2485-2492 (2003).

29.   K. T. Hughes, K. L. Gillen, M. J. Semon, J. E. Karlinsey, Sensing structural

intermediates in bacterial flagellar assembly by export of a negative regulator.

Science **19**, 1277-1280 (1993).

30.   K. Kutsukake, Excretion of the anti-sigma factor through a flagellar

substructure couples flagellar gene expression with flagellar assembly in

Salmonella typhimurium. *Mol. Gen. Genet.* **15**, 605-612 (1994).

31.   R. Lister, R. C. O'Malley, J. Tonti-Filippini, B. D. Gregory, C. C. Berry *et al*.

Highly Integrated Single-Base Resolution Maps of the Epigenome in

Arabidopsis. *Cell* **133**, 523-536 (2008).

32.   A. Mortazavi, B. A. Williams, K. McCue, L. Schaeffer, B. Wold, Mapping and

quantifying mammalian transcriptomes by RNA-Seq. *Nat. Methods* **5**, 621-628

(2008).

33.   U. Nagalakshmi, Z. Wang, K. Waern, C. Shou, D. Raha, *et al*. The

transcriptional landscape of the yeast genome defined by RNA sequencing.

Science 320, 1344-1349 (2008).

34.   N. T. Ingolia, S. Ghaemmaghami, J. R. Newman, J. S. Weissman, Genome-

wide analysis in vivo of translation with nucleotide resolution using ribosome

profiling. *Science* **324**, 218-223 (2009).

35.   I. Spöring, S. Felgner, M. Preuße, D. Eckweiler, M. Rohde *et al.* Regulation of

Flagellum Biosynthesis in Response to Cell Envelope Stress in *Salmonella*

enterica Serovar Typhimurium. *MBio* **9**, e00736-17 (2018).

36. P. Adams, R. Fowler, N. Kinsella, G. Howell, M. Farris *et al*. Proteomic

detection of PhoPQ- and acid-mediated repression of *Salmonella* motility.

Proteomics **1**, 597-607 (2005).

37. E. A. Miao, J. V. Rajan, *Salmonella* and Caspase-1: A complex interplay of

detection and evasion. *Front. Microbiol.* **2**, 85 (2011).

38. B. Ilyas, D. T. Mulder, D. J. Little, W. Elhenawy, M. M. Banda *et al.* Regulatory

Evolution Drives Evasion of Host Inflammasomes by *Salmonella* Typhimurium.

Cell Rep. **25**, 825-832 (2018).

39. L. D. B. Evans, C. Hughes, Selective binding of virulence type III export

chaperones by FliJ escort orthologues InvI and YscO. *FEMS Microbiol.*

Lett. **293**, 292-297 (2009)

40. Y. Cherradi, A. Hachani, A. Allaoui, Spa13 of *Shigella flexneri* has a dual role:

Chaperone escort and export gate-activator switch of the type III secretion

system. Microbiology. **160**, 130–141 (2014).

41. S. Yamaguchi, H. Fujita, T. Taira, K. Kutsukake, M. Homma, *et al*. Genetic

analysis of three additional fla genes in *Salmonella* typhimurium. *J. Gen.*

Microbiol. **130**, 3339-3342 (1984).

42. S. Hoffmann, C. Schmidt, S. Walter, J. K. Bender, R. G. Gerlach, Scarless

deletion of up to seven methylaccepting chemotaxis genes with an optimized

method highlights key function of CheM in *Salmonella* Typhimurium. *PLoS*

One **12**, e0172630. (2017).

43. E. Amann, B. Ochs, K. J. Abel, Tightly regulated tac promoter vectors useful

for the expression of unfused and fused proteins in *Escherichia coli*. *Gene* **69**,

301-315 (1988).

44. B. Miroux, J. E Walker, Over-production of proteins in *Escherichia coli*: Mutant

hosts that allow synthesis of some membrane proteins and globular proteins at

high levels. *J. Mol. Biol.* **260**, 289-298 (1996).

45.   T. Thanabalu, E. Koronakis, C. Hughes, V. Koronakis, Substrate-induced

assembly of a contiguous channel for protein export from *E.coli*: reversible

bridging of an inner-membrane translocase to an outer membrane exit pore.

EMBO J. **17**, 6487-6496 (1998).

46.   W. G. Kaelin, W. Krek, W. R. Sellers, J. A. DeCaprio, F. Ajchenbaum, *et al.*

Expression cloning of a cDNA encoding a retinoblastoma-binding protein with

E2F-like properties. *Cell* **70**, 351-364 (1992).

47.   B. Y. W. Chung, M. J. Deery, A. J. Groen, J. Howard, D. C. Baulcombe,

Endogenous miRNA in the green alga *Chlamydomonas* regulates gene

expression through CDS-targeting. *Nat. Plants* **3**, 787-794 (2017).

48.   B. Y. W. Chung, T. J. Hardcastle, J. D. Jones, N. Irigoyen, A. E. Firth *et al.*

The use of duplex-specific nuclease in ribosome profiling and a user-friendly

software package for Ribo-seq data analysis. *RNA* **21**, 1731-1745 (2015).

49. T. J. Hardcastle, Data from "Analysis of sequencing data from ribosome

profiling experiments." Bioconductor. Available at

doi:10.18129/B9.bioc.riboSeqR.

Figure Legends

Figure 1 Identification of chaperone variants defective for FliJ binding

A. Atomic resolution structure of the 140-residue *Salmonella* FlgN (PDB: 5B3D)

indicating the region (residues 70-102; red) in which three-residue scanning deletions

were introduced.

B. Affinity chromatography of glutathione sepharose-bound GST-FliJ or GST-FlgK

incubated with cell lysates containing wildtype FlgN or its three-amino acid deletion

variants. After washing, proteins were eluted in SDS-loading buffer, separated by

SDS(15%)-PAGE and stained with Coomassie Brilliant Blue. Apparent molecular

weights are in kilodaltons (kDa).

C. Affinity chromatography of glutathione sepharose-bound GST-FliJ, GST-FlgK or

GST-FlhAc incubated with cell lysates containing wildtype FlgN or its derivative

variants FlgNΔ76-78, FlgN-W78A or FlgNΔ90-100. After washing, proteins were

eluted in SDS-loading buffer, separated by SDS(15%)-PAGE and stained with

Coomassie Brilliant Blue. Apparent molecular weights are in kilodaltons (kDa).

D. Atomic resolution structure of the 122-residue *Salmonella* FliT (PDB: 3A7M)

indicating the position of conserved residues (red) within the region essential for FliJ

binding. *In vitro* binding to FliJ is enhanced by truncating FliT at residue 94 (dashed

red line).

E. Affinity chromatography of glutathione sepharose-bound GST-FliJ with cell lysates

containing FliT94 or its point mutant variants. After washing, proteins were eluted in

SDS-loading buffer, separated by SDS(15%)-PAGE and stained with Coomassie

Brilliant Blue. Apparent molecular weights are in kilodaltons (kDa).

F. Affinity chromatography of glutathione sepharose-bound GST-FlhAc incubated

with cell lysates containing FliD and wildtype FliT or its point mutant variants FliT-

L72A or FliT-Y106A. After washing, proteins were eluted in SDS-loading buffer,

separated by SDS(15%)-PAGE and stained with Coomassie Brilliant Blue (top panel)

or were analysed by immunoblotting with anti-FliT or anti-FliD polyclonal antisera

(middle and bottom panels). Apparent molecular weights are in kilodaltons (kDa).

Figure 2 FliJ binding to FlgN and FliT is required for efficient cell motility and

subunit export

A. Swimming (top panel; 0.25% soft tryptone agar) and swarming (bottom panels;

0.6% agar-tryptone with 0.5% glucose) motility of recombinant *Salmonella* strains

producing chromosomally-encoded FlgN variants (W78A, Δ76-78, or Δ90-100), the

SJW1103 wildtype, and the *Salmonella flgN* null strain (Δ*flgN*).

B. Whole cell (cell) and supernatant (secreted) proteins from late exponential-phase

cultures of *Salmonella* strains producing chromosomally-encoded FlgN variants

(W78A, Δ76-78, or Δ90-100), the SJW1103 wildtype, and the *Salmonella flgN* null

strain (Δ*flgN*) were separated by SDS (15%)-PAGE and analysed by immunoblotting

with anti-FliC, anti-FlgK or anti-FlgN polyclonal antisera. Apparent molecular weights

are in kilodaltons (kDa).

C. Swimming (top panel; 0.25% soft tryptone agar) and swarming (bottom panels;

0.6% agar-tryptone with 0.5% glucose) motility of recombinant *Salmonella* strains

producing chromosomally-encoded FliT variants (L72A and Y106A), the SJW1103

wildtype, and the *Salmonella fliT* null strain (Δ*fliT*).

D. Whole cell (cell) and supernatant (secreted) proteins from late exponential-phase

cultures of *Salmonella* strains producing chromosomally-encoded FliT variants (L72A

and Y106A), the SJW1103 wildtype, and the *Salmonella fliT* null strain (Δ*fliT*) were

separated by SDS (15%)-PAGE and analysed by immunoblotting with anti-FliC, anti-

FliD or anti-FliT polyclonal antisera. Apparent molecular weights are in kilodaltons

(kDa).

Figure 3 FlgN and FliT chaperones compete with FlhA$_C$ for binding to FliJ

A. *In vitro* competition assay to assess FlhA$_C$ binding to cobalt (Co^{2+}) bead-

immobilized (His)$_6$-FliJ in the presence of the FlgN or FliT chaperones, or chaperone

variants that cannot bind FliJ (FliJ non-binder).

B. Cell lysates of *E.coli* C41 expressing FliT94, which cannot bind FlhA$_C$ (-), or its

L72A variant, which cannot bind FlhA$_C$ or FliJ (left-hand panels), or FlgN Y112A

which cannot bind FlhA$_C$ (-) or its Δ76-78 variant, which cannot bind FlhA$_C$ or FliJ

(right-hand panels) were mixed with *E.coli* C41 lysates containing FlhAc (load) and

incubated with cobalt bead-immobilized (His)$_6$-FliJ (bound). After washing, proteins

were eluted from the cobalt beads in SDS-loading buffer, separated by SDS(15%)-

PAGE and analysed by immunoblotting with anti-FlhA, anti-FliT, anti-FlgN or anti-FliJ

polyclonal antisera. Apparent molecular weights are in kilodaltons (kDa).

C. *In vivo* competition assay to assess the effect of chaperone-FliJ binding on FlhA$_C$

export activity. Export of the unchaperoned FlgD subunit was assessed in a *flgE* null

strain. Whole cell (cell) and supernatant (secreted) proteins from late exponential-

phase cultures of *Salmonella* Δ*flgE* strains producing pTrc99a-encoded FlgN Y122A

(-) or FlgN Y122A, Δ76-78 (Δ76-78) were separated by SDS (15%)-PAGE and

analysed by immunoblotting with anti-FlgD, anti-FlgN, anti-FliJ and anti-FlhA

polyclonal antisera. Apparent molecular weights are in kilodaltons (kDa). Relative

amounts of FlgD secreted into culture supernatants were quantified using image

Studio Lite software and normalized to FlgD secreted by cells expressing the FlgN

Δ76-78 Y122A variant (right-hand bar chart). Export of non-cognate subunit FlgD is

attenuated in cells producing FlgN Y122A (-), which binds FliJ but not FlhA, but is

significantly higher in cells producing the FlgN variant defective in binding FliJ (FlgN

Δ76-78,Y122A).

Figure 4 Ribosome profiling reveals cytoplasmic ratios of flagellar subunits to

their cognate chaperones

A. Smoothed scatter plot of total translation (Ribosome profiling, measurement of

ribosome occupation) and total mRNA abundance (RNA-Seq), showing flagellar

chaperones FlgN, FliT and FliS (blue), their respective cognate subunits (red), and

the regression between total translation and mRNA abundance (dotted line).

B. Ratios of the normalised translation to gene length for the coding regions

corresponding to chaperones and their cognate subunits (FlgN:FlgK+FlgL; FliT:FliD;

FliS:FliC).

Figure 5 Chaperone regulation of the fT3SS machinery in response to subunit

cargo availability

Chaperone-subunit complexes initially dock at the FliI component of the ATPase

complex before docking at the cytoplasmic domain of FlhA (left-hand panel).

Subunits are subsequently released from their cognate chaperones and translocated

into the export channel. Unladen chaperones then bind FliJ from which they are

captured by newly synthesized cognate subunits. Under conditions where subunit

availability is limited (right-hand panel), unladen FlgN and FliT chaperones remain

bound to FliJ, blocking the interaction of FliJ with FlhA and reducing pmf-driven

export by the fT3SS machinery.

Supplementary Figure Legends

Figure S1.

A. Affinity chromatography of glutathione sepharose-bound glutathione-S-transferase

(GST) incubated with cell lysates containing FliT94. After washing, proteins were

eluted in SDS-loading buffer, separated by SDS(15%)-PAGE and stained with

Coomassie Brilliant Blue (top panel). The cell lysate load was analyzed by

immunoblotting with polyclonal anti-FliT anti-sera (bottom panel). Apparent

molecular weights are in kilodaltons (kDa).

B. Affinity chromatography of glutathione sepharose-bound glutathione-S-transferase

(GST) incubated with cell lysates containing FlgN. After washing, proteins were

eluted in SDS-loading buffer, separated by SDS(15%)-PAGE and stained with

Coomassie Brilliant Blue (top panel). The cell lysate load was analyzed by

immunoblotting with polyclonal anti-FlgN anti-sera (bottom panel). Apparent

molecular weights are in kilodaltons (kDa).

Figure S2.

A. Affinity chromatography of glutathione sepharose-bound GST-FlgL (top two

panels) or glutathione-S-transferase (GST; bottom panels) incubated with cell lysates

containing FlgN or its derivative variants FlgNΔ76-78, FlgN-W78A or FlgNΔ90-100.

After washing, proteins were eluted in SDS-loading buffer, separated by SDS(15%)-

PAGE and stained with Coomassie Brilliant Blue to visualise GST-FlgL and GST, or

analysed by immunoblotting with polyclonal anti-FlgN anti-sera. The cell lysate load

(FlgN load) was also analyzed by immunoblotting with polyclonal anti-FlgN anti-sera.

Apparent molecular weights are in kilodaltons (kDa).

B. Affinity chromatography of glutathione sepharose-bound GST-FliJ (top two

panels), GST-FliD (middle two panels) or glutathione-S-transferase (GST; bottom

panels) with cell lysates containing FliT and its mutant variants. After washing,

proteins were eluted in SDS-loading buffer, separated by SDS(15%)-PAGE and

stained with Coomassie Brilliant Blue to visualise GST-FliJ, GST-FliD and GST, or

analysed by immunoblotting with polyclonal anti-FliT anti-sera. The cell lysate load

(FliT load) was also analyzed by immunoblotting with polyclonal anti-FliT anti-sera.

Apparent molecular weights are in kilodaltons (kDa).

C. Cell lysates of *E.coli* C41 expressing FliT (top panel) or FliD (bottom panel) used

as load fractions in affinity chromatography assays with GST-FlhAc in Figure 1.

Proteins were separated by SDS(15%)-PAGE and analysed by immunoblotting with

polyclonal anti-FliT anti-sera or polyclonal anti-FliD anti-sera.

Figure S3.

a. Whole cell and supernatant proteins from late exponential-phase cultures of

Salmonella strains producing chromosomally-encoded FliT variants (L72A and

Y106A), the *Salmonella fliT* null strain (Δ*fliT*), and the SJW1103 wildtype were

separated by SDS (15%)-PAGE and analysed by immunoblotting with anti-FlgK

polyclonal antisera. Apparent molecular weights are in kilodaltons (kDa).

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

active

inactive