Neuronal firing and waveform alterations through ictal recruitment in humans

Edward M. Merricks[1], Elliot H. Smith[1,2], Ronald G. Emerson[3], Lisa M. Bateman[1], Guy M. McKhann II[4],

Robert R. Goodman[5], Sameer A. Sheth[6], Bradley Greger[7], Paul A. House[8], Andrew J. Trevelyan[9],

Catherine A. Schevon[1*]

1. Department of Neurology, Columbia University Medical Center, New York NY
2. Department of Neurosurgery, University of Utah, Salt Lake City UT
3. Department of Neurology, Weill Cornell Medical Center, New York, NY
4. Department of Neurosurgery, Columbia University Medical Center, New York NY
5. Department of Neurosurgery, Lenox Hill Hospital, New York NY
6. Department of Neurosurgery, Baylor College of Medicine, Houston TX
7. School of Biology and Health Systems Engineering, Arizona State University, Tempe AZ
8. Intermountain Healthcare, Murray UT
9. Institute of Neuroscience, Newcastle University, Newcastle upon Tyne, UK

* Corresponding author: cas2044@cumc.columbia.edu

Number of figures: 9
Number of tables: 3
Words in abstract: 250
Words in introduction: 446
Words in discussion: 1,499

Funding:

    NIH R01 NS084142; CRCNS R01 NS095368

Conflicts of interest:

    None.

ABSTRACT

Clinical analyses of neuronal activity during seizures, invariably using extracellular recordings,

is greatly hindered by various phenomena that are well established in animal studies: changes in

local ionic concentration, changes in ionic conductance, and intense, hypersynchronous firing.

The first two alter the action potential waveform, whereas the third increases the "noise"; all

three factors confound attempts to detect and classify single neurons (units). To address these

analytical difficulties, we developed a novel template-matching based spike sorting method,

which enabled identification of 1,239 single units in 27 patients with intractable focal epilepsy,

that were tracked throughout multiple seizures. These new analyses showed continued neuronal

firing through the ictal transition, which was defined as a transient period of intense tonic firing

consistent with previous descriptions of the ictal wavefront. After the ictal transition, neurons

displayed increased spike duration ($p < 0.001$) and reduced spike amplitude ($p < 0.001$), in

keeping with prior animal studies; units in non-recruited territories, by contrast, showed more

stable waveforms. All units returned to their pre-ictal waveforms after seizure termination.

Waveshape changes were stereotyped across seizures within patients. Our analyses of single

neuron firing patterns, at the ictal wavefront, showed widespread intense activation, and

commonly involving marked waveshape alteration. We conclude that the distinction between

tissue that has been recruited to the seizure versus non-recruited territories is evident at the level

of single neurons, and that increased waveform duration and decreased waveform amplitude are

hallmarks of seizure invasion that could be used as defining characteristics of local recruitment.

SIGNIFICANCE STATEMENT

Animal studies consistently show marked changes in action potential waveform during epileptic

discharges, but acquiring similar evidence in humans has proved difficult. Assessing neuronal

involvement in ictal events is pivotal to understanding seizure dynamics and in defining clinical

localization of epileptic pathology. Using a novel method to track neuronal firing, we analyzed

microelectrode array recordings of spontaneously occurring human seizures, and here report two

dichotomous activity patterns. In cortex that is recruited to the seizure, neuronal firing rates

increase and waveforms become longer in duration and shorter in amplitude, while penumbral

tissue shows stable action potentials, in keeping with the "dual territory" model of seizure

dynamics.

Introduction

A complete understanding of the mechanisms underlying seizure pathology and dynamics

depends on knowledge of the local neuronal activity, and what is driving that activity.

Comparative animal models have long been used to gain insights into the underlying neuronal

activity during seizures (Purpura *et al.*, 1972; Fariello *et al.*, 1976; Grone & Baraban, 2015), with

the paroxysmal depolarizing shift (PDS) being regarded as the intracellular correlate of ictal

discharges in animal models for more than half a century (Kandel & Spencer, 1961*a*, 1961*b*;

Matsumoto & Marsan, 1964; Traub & Wong, 1982).

More recently, early PDSs have been shown to evolve into seizures *in vivo* (Steriade &

Amzica, 1999), and PDSs have been recorded in resected human cortical tissue (Marcuccilli *et*

al., 2010; Eissa *et al.*, 2016). The PDS causes a decrease in action potential amplitude and an

increase in half width – features that should impede standard spike sorting methods – and yet this

phenomenon has not been reported in several studies of single unit activity during spontaneous

human seizures (Wyler *et al.*, 1982; Babb *et al.*, 1987; Stead *et al.*, 2010; Truccolo *et al.*, 2011,

2014; Bower *et al.*, 2012). In fact, beyond the PDS, altered action potential waveforms could be

expected following recruitment of a recording site to a seizure due to alterations to Na^+ and K^+

concentrations in the intracellular and extracellular space or the effects of burst firing (Harris *et*

al., 2000).

We have shown preliminary evidence of such potential alterations (Merricks *et al.*, 2015).

In tissue recruited to the seizure, traditional spike sorting methods can fail to cluster single units

in human ictal recordings from neocortical layers 4/5, where neuronal cell body density is

particularly high (Keller *et al.*, 2018), thereby hindering the ability to track evidence of wave

shape alterations or neuronal firing patterns during and after the ictal wavefront (Merricks *et al.*,

2015). However, whether this originated from alterations to neurons' intrinsic wave shapes or

simply from interference of action potentials from nearby, highly active cells has been unclear.

Here, we present analyses of neuronal activity in the human brain during focal seizures

using novel template matching methods in order to characterize action potential waveform

alterations and single unit firing patterns, as the ictal wavefront approaches, recruits, and passes

the local tissue. We hypothesize that, similar to observations in animal models, human focal

seizures consistently display alteration of intrinsic action potential shapes upon ictal recruitment.

Specifically, we hypothesize that the distinction between recruited and penumbral tissue is

maintained at the level of single neurons, with recruited cells displaying reduced spike amplitude,

and increased duration, an effect that is absent in penumbral sites demonstrating increased firing

rates, but lacking typical seizure hallmarks.

Materials and Methods

Human recordings

Adult patients undergoing surgical evaluation for pharmacoresistant focal epilepsy at

Columbia University Irving Medical Center (CUIMC) and University of Utah were implanted

with either a 96 channel, 4 x 4 mm "Utah"-style microelectrode array (UMA; Blackrock

Microsystems, Salt Lake City, UT) or Behnke-Fried style microwires (BF array; Ad-tech

Medical Equipment Corp, Oak Creek, WI) simultaneous to standard clinical electrocorticography

(ECoG) or stereo-electroencephalography (sEEG) respectively. UMAs were implanted into

neocortical gyri based on presurgical estimation of the ictogenic region, with electrode tips

reaching layer 4/5 (1.0 mm electrode length; layer confirmed via histology in Schevon *et al.*

(2012)), while BF arrays consisted of 8 microwires protruding ~4 mm from the tips of clinical

depth electrodes.

Neural data were recorded at a sampling rate of 30 kHz on each microelectrode with a

range of $\pm$ 8 mV at 16-bit precision, with a 0.3 Hz to 7.5 kHz bandpass filter. ECoG and sEEG

data were collected with a sampling rate of either 500 Hz or 2 kHz, with 24-bit precision and a

bandpass filter of 0.5 Hz to ¼ the sampling rate. In UMAs, the reference was either subdural or

epidural, chosen based on recording quality. In BF arrays, the reference was the ninth microwire

within the bundle.

All procedures were approved by the Institutional Review Boards of CUIMC and

University of Utah, and all patients provided informed consent prior to surgery. Clinical

determination of seizure onset zone (SOZ) and seizure spread was made by the treating

physicians and confirmed prior to analysis by two board-certified neurologists (CAS & LMB).

All analyses were performed offline using custom scripts and toolboxes written in MATLAB

(MathWorks, Natick MA). Code is available at https://github.com/edmerix.

The timing of the passage of the ictal wavefront at individual electrodes was calculated

based on the MUA firing rate. A Gaussian kernel of 500 ms duration was convolved with the

timings of all detected spikes in the MUA, and a sustained, significant increase in the resultant

instantaneous firing rate was determined as the moment of local recruitment to the seizure (Smith

et al., 2016). A sustained (> 1 s), significant increase had to be present for classification as ictal

recruitment in order to discount single discharges or herald spikes. Ictal recordings without this

signature of tonic to clonic MUA firing were determined to be penumbral. Note that unlike the

SOZ, ictal recruitment and the penumbra are spatiotemporally dynamic. As such, a single

location, unless at the true origin of the ictal activity, may receive the synaptic input of the

upstream ictal activity but remain penumbral due to feed-forward inhibition initially, prior to the

transition to the ictal state which may occur at any point during the electroclinical seizure event.

Peri-ictal single unit discrimination

Initial spike sorting was performed on the peri-ictal period as per Merricks *et al.* (2015).

Briefly, neural signals were symmetrically bandpass filtered between 300 Hz and 5 kHz (1000[th]

order FIR1) to extract multi-unit activity (MUA), from which extracellular action potential

spikes were detected using a voltage threshold of 4.5 σ, where $\sigma = median\left(\frac{|x|}{0.6745}\right)$, and x is the

MUA from that channel. This method avoids the biasing effect of large spikes on channels with

units with high firing rates (Quian Quiroga *et al.*, 2004). Ictal periods were blanked so that spike

sorting was only performed on stable spikes from the peri-ictal period.

Matrices of waveforms from each channel were created from 0.6 ms prior to 1 ms post

each detection, and principal component based semi-automatic cluster cutting was performed

using a modified version of the "UltraMegaSort2000" MATLAB toolbox (Hill *et al.*, 2011).

Artifactually large waveforms were removed by calculating the FFT on spikes up-sampled by a

factor of 4, and removing those with power > 5 SD above the mean in frequencies above 2.5 kHz

or below 500 Hz. Spikes removed in this manner were visually inspected to ensure correct

classification as artefact. Clusters that satisfied the following criteria were accepted: (i) clean

separation from all other clusters in the Fisher's linear discriminant in principal component

space; (ii) less than 1% contamination of the 2 ms absolute refractory period; (iii) no clear

outliers based on the anticipated chi-squared distribution of Mahalanobis distances; and (iv) less

than 1% of estimated false negatives as estimated by the amount of a Gaussian fit to the detected

voltages fell below the threshold for detection, as described in Hill *et al.* (2011).

Template matching through seizures

Ictal recruitment has been shown to impede standard spike sorting due to either

interference of hypersynchronous activity, intrinsic waveform alterations, or both (Merricks *et al.*,

2015; Fig. 1). We therefore developed novel methods in order to match waveforms from the ictal

period, regardless of recruitment, to their putative neuronal source based on templates derived

from the peri-ictal units. In contrast to standard spike sorting methods, these minimized false

negatives at the expense of increasing false positives so as to avoid missing potential matches.

Cluster boundaries were defined as the 3-dimensional convex hull surrounding the features in

principal component space of the previously defined units from both the pre- and post-ictal

period (Fig. 2). This method accounts for two situations: that neurons within recruited cortex

maintain their wave shape but are obscured by interference from other nearby cells; or that there

are occasional or consistent alterations to a neuron's intrinsic waveform that are minor enough to

be maintained within the convex hull of feature space. The convex hull allows for alterations to

wave shape in any dimension (any direction away from the cluster's centroid).

To avoid ictal results being biased through differing methods, template matching was

performed on spikes that were extracted from a period from 10 minutes prior to 10 minutes post

seizures, including the ictal activity that had been blanked in the original peri-ictal spike sorting.

Channels with unstable units during interictal periods were excluded. Units with no spikes in

either the preictal or ictal time period after template matching and artefact removal were

excluded from further analyses ($n = 77$).

Principal component scores were calculated on these spikes based on the previously

defined principal components, and spikes that occurred within a peri-ictal unit's convex hull

were assigned to that cell. Mahalanobis distances were calculated for all matches, between their

location in principal component space and all peri-ictal waveforms from that unit, on the first n

principal components that explained > 95% of the variance in the data set (Fig. 2C). The

expected distribution of Mahalanobis distances was calculated as the chi-squared probability

distribution with n degrees of freedom. Spikes that had < 0.1% chance of occurring in the chi-

squared distribution were excluded.

Spike metrics

The full-width at half maximum (FWHM) was calculated by up-sampling each spike by a

factor of 4, normalizing the spike voltages to between [-1, 1], and finding the difference between

the zero-crossings either side of the spike's trough. When calculating spike amplitude changes

through the ictal transition, only units whose mean voltage at detection was at least 2.5 SD away

from that channel's threshold for detection were used, to minimize the floor effect from small

units that dropped below threshold.

The probability that each spike arose from its assigned peri-ictal unit was calculated by

fitting a separate Gaussian curve (with a maximum amplitude of 1) to the distribution of voltages

at each data point in the original unit, and calculating the mean probability across all time

samples. As such, a waveform passing through the most likely voltage at each separate time

point for that unit would have a match probability of 1 (Fig. 3). Instantaneous firing rates were

calculated by convolving the spike times with a Gaussian kernel (200 ms SD) with the amplitude

scaled to that spike's probability of matching the original unit, thereby creating probabilistic

firing rates for each unit through time, by probability of when the spike occurred and probability

that the spike originated from the putative neuronal source. As such, a waveform that had an

average probability of 20% across all fitted Gaussians from each data point would contribute

only 0.2 spikes s^{-1} at its most likely time point, while an exact match to the most likely voltage at

each time point would contribute 1 spike s^{-1}. Thresholds for significant increases and decreases

in firing rate were calculated as 3 times the square root of the firing rate divided by the duration

of the epoch, which approximates 3 SD for a Poisson distribution.

Timing of FWHM alteration relative to the earliest ictal activity was determined by the

earliest timepoint during the seizure that the mean FWHM remained above the preictal mean

plus the preictal standard deviation for at least 1 second, calculated in a sliding window of 5 s

duration with a time step of 50 ms, discarding windows with fewer than 5 spikes.

All statistical tests for significance were performed using the Mann-Whitney U test

unless otherwise noted, due to the non-Gaussian distributions of data requiring non-parametric

testing. For all tests, the level for statistical significance (α) was set to 0.05, and Holm-

Bonferroni correction was applied in all instances of multiple tests.

Results

We analysed ictal recordings from 27 patients undergoing invasive EEG monitoring as

part of the presurgical evaluation for intractable focal epilepsy (Tables 1 & 2; age range = 19 to

55; 13 female, 14 male). Six patients were implanted with Utah microelectrode arrays (UMA;

Blackrock Microsystems Inc, Salt Lake City, UT), and the remaining 21 patients were implanted

with between 1 and 4 Behnke-Fried depth arrays with incorporated microwire bundles (BF

arrays; Ad-tech Medical Equipment Corp, Oak Creek, WI). A total of 41 seizures were reviewed

(10 UMA; 31 BF array), of which 27 demonstrated ictal recruitment through MUA firing rate

calculation in UMAs, or subsequent waveform alterations in BF arrays (see Methods; UMA: 6

seizures from 3 patients; BF arrays: 21 seizures from 13 patients).

Dual activity types at seizure onset

To assess the presence of physiological spike shape changes across populations of

neurons, template matching using convex hulls (Fig. 2, see Methods) was employed on each

UMA-recorded seizure (Table 1; 1,107 single units in total; range of units per seizure: 61 – 240;

mean ± SD maximum units per patient: 122 ± 63). The convex hull fits a 3-dimensional region

around the peri-ictal unit's cluster in principal component space, within which waveforms are

assigned as a putative match to that neuron. This allows for assigning unit identities without the

need for maintained cluster boundaries in the data, while accepting waveform alterations that

alter the principal component scores. In total, 938 of the 1,107 units recorded on UMAs were

successfully tracked through seizures using this method.

Individually, results from the template matching method in recordings from recruited

tissue (see Methods; Patients 3, 4 & 5) showed decreases in spike amplitude and increases in

spike full-width at half maximum (FWHM; Fig. 4, ictal waveforms in red), and were stereotyped

across seizures within patient (Fig. 5). At the population level, units in recruited cortex displayed

a significant global increase in FWHM (Fig. 6A; pre-ictal vs. ictal mean $\pm$ SD: 0.470 ± 0.137 ms

vs. 0.611 ± 0.194 ms), with 457 (81.5%) of 561 single units showing a significant ($p < 0.05$)

increase in FWHM during the seizure (Holm-Bonferroni corrected Mann-Whitney U test; range

across seizures: 79% – 97%; see Table 3). Meanwhile, units in penumbral cortex showed only a

minor increase in FWHM at the population level (Fig. 6B; pre-ictal vs. ictal mean $\pm$ SD: $0.414 \pm$

0.009 ms vs. 0.429 ± 0.099 ms) with only 9 (5.8%) of 156 single units showing a significant ($p <$

0.05) increase in FWHM during the seizure (Holm-Bonferroni corrected Mann-Whitney U test;

range across seizures: 4% – 16%; see Table 3). In a single case (Patient 6), the UMA was at the

edge of the clinically defined ictal spread, and in this patient 105 (47.5%) of 221 units showed a

significant increase in FWHM (pre-ictal vs. ictal mean $\pm$ SD: 0.408 ± 0.111 ms vs. 0.421 ± 0.152

ms). These spike shape alterations co-existed with stable spike shapes elsewhere on the UMA at

the same time (Fig. 7), and this patient was incorporated into the penumbral dataset for

population representation in the figures (Fig. 6). FWHM increases in recruited tissue were

significantly larger than those in penumbral/edge case recordings ($p < 0.001$; one-tailed Mann-

Whitney U test).

Similarly, units in recruited cortex showed a significant decrease in spike amplitude

during the seizure (Fig. 6C; pre-ictal vs. ictal mean $\pm$ SD: 48.82 ± 30.91 µV vs. 34.96 ± 19.54

µV), while penumbral recordings maintained their peri-ictal amplitude (Fig. 6D; pre-ictal vs.

ictal mean ± SD: 46.69 ± 16.59 µV vs. 45.08 ± 15.05 µV in fully penumbral cases; 47.00 ± 21.18

µV vs 45.57 ± 20.93 µV in the semi-recruited UMA). The amplitude reduction in recruited tissue

was significantly greater than in penumbral/edge case recordings (p < 0.001; one-tailed Mann-

Whitney U test), with recruited recordings showing significant ($p < 0.05$) decreases in amplitude

during the seizure in 49.3% of units versus 6.4% of units in the penumbra and 38.9% in semi-

recruited tissue (Holm-Bonferroni corrected Mann-Whitney U test).

Spike shape changes in deep structures

To assess the spatiotemporal relationship between waveform alterations and seizure

recruitment within patients, beyond the capabilities of the 4 mm^2 Utah array, we analysed BF

array recordings with the equivalent template matching, blind to the clinically-defined seizure

onset zone and areas of propagation. In BF array recordings, 120 of 132 units were successfully

tracked using these methods. Thirty of 120 single units (25.0%) showed increases beyond a cut-

off significance level of $p < 0.05$ (Holm-Bonferroni corrected Mann-Whitney U test) in FWHM

(17 seizures from 13 patients), and 30 units (25.0%; 16 seizures from 11 patients) showed

reduction in spike amplitude below the same significance cut-off ($p < 0.05$; Table 2). In 9

seizures from 6 patients, single units were simultaneously present on multiple separate BF arrays

(on different bundles of microwires, as opposed to different microwires within a single BF), of

which 7 seizures (6 patients) showed waveform alterations in at least one unit. Of these, 2

seizures (2 patients) showed significant waveform alterations in dual locations (Patients 14 and

15; Fig. 8; Table 2), while 5 seizures (4 patients) showed both activity types simultaneously

(Patients 11, 12, 16 and 21; Video 1).

We found significant waveform alterations in 13 patients. Five of these had a clinically

defined SOZ in the mesial temporal lobe, and simultaneously recorded single units in the

ipsilateral hippocampus demonstrated significant waveform alterations (Patients 8, 15, 16, 26 &

27; Table 2). In 7 further patients, significant waveform alterations were found in tissue

consistent with putative seizure spread due to proximity to the SOZ, or due to seizure

generalization. In one case (Patient 10) we found waveform alterations in the contralateral

hippocampal body, consistent with propagation of the seizure through the hippocampal

commissural fibers. Conversely, in the 8 patients showing no significant waveform alterations,

the clinically defined SOZ was anatomically distant in all cases (Patients 9, 13, 17–20, 24 & 25;

Table 2).

To assess whether the classification of ictal recruitment via waveform alterations was

consistent with the clinically defined SOZ and regions of spread, the time from earliest ictal

activity to a consistent (≥ 1 s duration) increase in FWHM ≥ 1 SD above the mean preictal level

for each unit was calculated. FWHM was used independently of amplitude for these tests to

control for any potential fluctuations in amplitude introduced by the reference electrode during

seizures. Recordings determined to be in the SOZ showed a mean ($\pm$ SD) delay of 10.23 ± 3.03 s

($n = 6$ seizures from 5 patients), while those deemed to be in regions of spread showed a mean ($\pm$

SD) delay of 22.96 ± 12.59 s ($n = 8$ seizures, 8 patients; $p < 0.05$, Mann-Whitney U test). In one

case, single unit waveforms remained stable throughout a focal to bilateral tonic-clonic seizure

(Patient 18; Table 2), further countering the possibility of transient movement artefact

introducing instability in the waveforms.

In one instance of a patient recorded with BF arrays in the hippocampal head and body, a

peculiarly discrete unit cluster due to a very large amplitude spike (mean = 354 µV; background

noise level = 25 µV) enabled us to follow its action potential through the ictal transition without

the need for template matching via convex hulls, despite marked changes to spike shape and

increases in other unit activity (Fig. 8; Video 2; Schevon *et al.*, 2019). In this example, the

seizure initiated very close to, but not at the electrode site. The ictal wavefront arrived at the

electrode approximately 8 s after seizure initiation (Patient 15; Table 2). The action potential

amplitude was stable during both the pre-ictal period and the moments after seizure initiation,

but reduced sharply upon the abrupt increase in firing rate indicating ictal recruitment (Fig. 8A &

B, magenta dashed line; preictal vs. ictal mean ± SD: 354.2 µV ± 45.7 µV vs. 265.7 µV ± 33.4

µV; $p < 0.001$, Mann-Whitney U test). The inter-spike interval to spike amplitude relationship

was similar to animal models of recruitment (c.f. Supplementary Fig. 4 in Merricks *et al.*, 2015;

0 Mg^{2+} *in vitro* cell-attached mouse slice model), and echoed the template matched results in the

Utah array (note the similarity between Fig. 4B and Fig. 8D).

Notably, a separate single unit recorded at an adjacent BF array showed spike shape

changes which developed 1 second later, consistent with seizure propagation (Fig. 8B ii; Table 2).

The distinct time course of these two units recorded in the hippocampal head and body show that

the alterations are unit-specific and not caused by local changes as a result of the seizure or by

movement artefacts (the seizure had a relatively calm semiology, without stressing the recording

device).

Simultaneously, action potential FWHM was stable prior to ictal recruitment, increasing

significantly after the passage of the ictal wavefront (preictal vs. ictal mean ± SD: 0.33 ms ± 0.04

ms vs. 0.40 ms ± 0.05 ms; $p < 0.001$, Mann-Whitney U test). The time course of this transition

followed the same progression as the spike amplitude (Fig. 8D), as can be seen in the pre-

recruitment versus post-recruitment mean (± SD) waveforms (Fig. 8C, blue and red respectively).

Template matching accuracy

Spike matches from the convex hull method were found to be significantly more likely to

arise from their assigned peri-ictal units than by chance, as calculated by comparing each

matched waveform's similarity to its peri-ictal unit's distribution of voltages through time (Fig.

3; "Spike metrics" in Methods). The "null" distribution for expected match probabilities by

chance was calculated through comparing each waveform's similarity to the peri-ictal voltage-

time distributions of all other units. Comparing intra- to inter-unit similarities in this way found a

significantly higher similarity between template matched units and their presumed peri-ictal unit

than to the "null" distribution of matches to other units ($p < 0.001$; Mann-Whitney U test; Fig.

3B).

To confirm that these findings were not a result of the template matching method

introducing an unknown variable that affects these measurements, results from the original spike

sorted data and those originating from the convex hull method on the pre-ictal period were

compared, finding little difference between the traditional cluster cutting results and the convex

hull matched results (Fig. 6 A-D i, inset cumulative histograms).

Neuronal firing rates through the ictal transition

In animal models, it is possible to isolate single units experimentally, by visually guiding

electrodes directly onto cells (Trevelyan *et al.*, 2006). This obviously is not possible in human

recordings, and consequently, there is a dearth of evidence about the firing patterns of human

neurons through seizures. Instead, previous studies have identified the ictal wavefront in terms of

multi-unit activity (Schevon *et al.*, 2012; Smith *et al.*, 2016). We therefore analysed the firing

rates of template matched unit populations throughout ictal activity and related these to the ictal

wavefront (UMA recordings showing tonic to clonic firing; Patients 3–5; see Methods). Single

unit firing rate increased during ictal recruitment in all seizures with 446 (79.5%) of 561 units

showing greater than 3 SD increase in firing rate, and only 1 unit in the entire population

showing a greater than 3 SD decrease in firing rate (range of single units with > 3 SD increase

per seizure: 70% – 96%; see Table 3). An example seizure demonstrating these trends is shown

in Fig. 9.

Discussion

The analyses presented here explored the impact of spike shape alterations in

spontaneously occurring seizures, in humans, how these alterations relate to the underlying ictal

territories, and whether the ictal wavefront – the source of seizure propagation – involves local

neuronal firing or is dominated by subthreshold or synaptic activity. We have previously shown

that traditional spike sorting methods fail upon ictal invasion of the recording site (Merricks *et*

al., 2015), however, in these recordings it was not possible to differentiate between loss of spike

sorting ability due to intrinsic waveform alterations, or due to hypersynchronous activity

(Trevelyan *et al.*, 2006, 2007; Schevon *et al.*, 2012; Weiss *et al.*, 2013). Here, we sought to

overcome this limitation through novel methods to track units through the ictal transition,

retaining the identities of putative individual neurons. We hypothesized that the temporary loss

of clusters was due in part to transient alterations to spike shapes, as opposed solely to

obfuscation of stable spike shapes by the sudden increase in activity in the MUA, and that units

at brain sites not demonstrating evidence of ictal recruitment via tonic firing in the MUA would

remain stable.

We applied this method in two types of microelectrode recordings: UMA and BF arrays,

representing neocortical and deep structure, particularly hippocampal, seizure foci. Although

there must be some expected loss of detection sensitivity due to interference from highly

synchronous firing along with the reduction of amplitude of some units below the noise threshold,

the method proved remarkably effective, enabling us to define firing rate metrics for the majority

of units throughout the seizure, highlighting a lack of neuronal quiescence during seizures.

In both types of recording, template matched units during the ictal period displayed two

types of activity: deformation of waveshapes across the population, or largely stable waveforms.

The UMA afforded the ability to detect local ictal recruitment through characteristic MUA firing,

and these types of activity corresponded to recruitment and penumbral recordings respectively

(Fig. 6). Waveform alterations recovered after seizure termination and showed a stereotyped

response across seizures, highlighting that a single neuron's wave shape change, in response to

the synaptic barrage of upstream ictal activity, is maintained across multiple seizures hours apart.

Note that the template matching method would equally favor alterations to waveform in any

dimension, and so if these changes were purely a result of the methodology, we would

reasonably expect the template to capture spikes with larger amplitude and decreased FWHM at

an equal rate. In such a case, we would anticipate a broadening of the distribution of these

features during the ictal period. Instead, we see a clear shift in the distributions to the right and

left in the FWHM and amplitude respectively, and found no evidence of increased spike

amplitude during seizures, arguing for a consistent physiological cause (Fig. 6A&C).

Detection of the ictal wavefront was not possible in BF arrays, due to a combination of

lower neuronal density in mesial structures relative to layer 4/5 neocortex (Pakkenberg &

Gundersen, 1997; Keller *et al.*, 2018), and a reduced "listening sphere" relative to UMAs due to

higher impedance (BF: 50-500 kΩ; UMA: 80-150 kΩ; Tóth *et al.*, 2016). Nonetheless, the same

waveform features were detected, and their presence correlated well with the clinical assessment

of SOZ and seizure spread (Table 2). Furthermore, BF arrays allowed for sampling of multiple

sites in a given patient, and the timing of waveform alterations correlated well with clinical

observations, with seizure spread locations showing delayed waveform changes compared to

recordings from SOZ regions (22.96 vs. 10.23 s respectively). The longest delay until

recruitment occurred after spread to the contralateral hippocampus (Patient 10; 42.7 s). The only

case with discordance between single unit and clinical data showed a statistically significant

increase in FWHM during the seizure, but at no point did the mean FWHM surpass the threshold

of 1 SD above the preictal mean (Patient 21; Table 2). In this instance, the clinical SOZ was in

the right insula and somatosensory cortex, with waveform alterations in the right hippocampus.

While recruitment of the hippocampus is plausible, this may represent a false positive as a result

of the temporally coarse statistical test.

The UMA and BF array recordings together provide evidence supporting the dual-territory

model of seizures: an ictal core with waveform changes, coexisting with a penumbral territory

with stable spike shapes. Combined, these indicate that the definitions of ictal recruitment and

penumbra are maintained at the level of single neurons, and waveshape change can be

considered a defining feature of recruitment to the seizure. In a subset of recordings, stable

action potentials were found simultaneously with waveform changes both in separate BF arrays

(5 seizures) and on the same UMA (1 seizure; Patient 6), with clinical correlation matching these

observations in all cases. In the UMA, clinical observations were consistent with location of the

UMA at the outer boundary of seizure invasion, and thus we posit this is a simultaneous

recording of both recruited and penumbral cortex (Fig. 7).

Furthermore, in 2 seizures, recruitment was found at multiple BF arrays, consistent with

clinically defined SOZ and regions of spread, along with relative delays in keeping with

anatomical distance (Fig. 8; Patients 14 & 15; mid-cingulate to hippocampus in 10.3 s,

hippocampal head to body in 1.1 s respectively). Note also, that in Patient 6 there is an increase

in waveforms of brief duration during the ictal period (data available online; population data

shown in Fig. 6B). Given the UMA's proximity to the oncoming ictal wavefront, this increase

may be explained by an increase in firing of fast-spiking interneurons, which have been shown to

exhibit action potentials of shorter duration (McCormick & Feeser, 1990; Csicsvari *et al.*, 1999;

Peyrache *et al.*, 2012) and would corroborate the penumbral feedforward inhibition model

(Trevelyan *et al.*, 2007; Cammarota *et al.*, 2013; Parrish *et al.*, 2019).

More specifically, the waveform changes found in recruited tissue are in keeping with

those observed in animal models, and are indicative of the shortening and broadening of action

potentials associated with PDS (Traub & Wong, 1982). This was especially evident in a unique

BF array recording wherein a unit was able to be tracked without need for extra methods due to

its amplitude being 14 times that of the background noise (Patient 15; Fig. 8), with feature

alterations strikingly reminiscent of the UMA population data (c.f. Figs. 4B & 8D), with timing

in keeping with recruitment, during tonic firing (c.f. Figs. 7 & 8).

Even so, the extent to which these waveforms alter is likely underrepresented in the

population data, due to changing detection sensitivity from interference between synchronous

spikes or the reduction of amplitude of some spikes below the noise threshold. The template-

matching method was designed to minimize false negatives in order to capture as much single

unit activity during seizures as possible, but it is likely that spikes undergo large enough changes

to be lost outside the convex hull, or below threshold. Indeed, even physiological bursting has

been shown to result in substantial alterations to extracellularly recorded action potentials (Harris

et al., 2000; Henze *et al.*, 2000). As such, these results are necessarily a snapshot of the total

activity of any individual neuron during the seizure, and yet still show significant changes to

waveform.

Finally, our data also demonstrate that the passage of the ictal wavefront is marked by tonic,

local neuronal firing (Fig. 9), as opposed to the wavefront being a signature of increased K^+

concentration or similar subthreshold phenomena. Studies of single unit activity recorded in

humans have rarely reported such a finding. Although placement of the electrodes likely plays a

large role, our findings in this paper suggest that extreme, rapid waveform alterations may

obscure the presence of the wavefront when standard spike sorting methods are used. Note that

these firing rate statistics are probabilistic, having been scaled by the probability each spike was

a match to its preictal unit (see Methods), and as such are conservative and thus the increase

cannot be attributed purely to increases in background spikes being matched by the convex hull

method.

This method of tracking single units across the ictal transition will enable the use of both

human and animal recordings to address open questions regarding the mechanism of seizure

spread. An immediate application, given the neuronal activity presented here, is to study how cell

types relate to the propagation of pathological activity, with considerable debate having focused

recently on the role of interneurons in seizure spread (Grasse *et al.*, 2013; Elahian *et al.*, 2018;

Magloire *et al.*, 2018; Miri *et al.*, 2018; Weiss *et al.*, 2018). All studies to date, to our knowledge,

have assessed cell-type specific activity at seizure onset in regions without waveform alterations,

and thus we suggest are recordings from penumbral territories. As such, we anticipate elucidation

of these mechanisms will come from population data confirmed to be in recruited tissue. These

methods lay important groundwork for analyses into how ictal propagation relates to the

underlying firing of local inhibitory and excitatory cells.

REFERENCES

Babb TL, Wilson CL & Isokawa-Akesson M (1987). Firing patterns of human limbic neurons
during stereoencephalography (SEEG) and clinical temporal lobe seizures.
Electroencephalogr Clin Neurophysiol 66, 467–482.

Bower MR, Stead M, Meyer FB, Marsh WR & Worrell GA (2012). Spatiotemporal neuronal
correlates of seizure generation in focal epilepsy. *Epilepsia* 53, 807–816.

Cammarota M, Losi G, Chiavegato A, Zonta M & Carmignoto G (2013). Fast spiking
interneuron control of seizure propagation in a cortical slice model of focal epilepsy. *The*
Journal of Physiology 591, 807–822.

Csicsvari J, Hirase H, Czurkó A, Mamiya A & Buzsaki G (1999). Oscillatory coupling of
hippocampal pyramidal cells and interneurons in the behaving Rat. *Journal of*
Neuroscience 19, 274–287.

Eissa TL, Tryba AK, Marcuccilli CJ, Ben-Mabrouk F, Smith EH, Lew SM, Goodman RR,
McKhann GM, Frim DM, Pesce LL, Kohrman MH, Emerson RG, Schevon CA & van
Drongelen W (2016). Multiscale Aspects of Generation of High-Gamma Activity during
Seizures in Human Neocortex. *eneuro* 3, ENEURO.0141–15.2016–17.

Elahian B, Lado NE, Mankin E, Vangala S, Misra A, Moxon K, Fried I, Sharan A, Yeasin M,
Staba R, Bragin A, Avoli M, Sperling MR, Engel J & Weiss SA (2018). Low-voltage fast
seizures in humans begin with increased interneuron firing. *Annals of neurology* 84, 588–
600.

Fariello RG, Portera A & Scheffner D (1976). Parenteral Penicillin in Rats: An Experimental
Model of Multifocal Epilepsy. *Epilepsia* 17, 217–222.

Grasse DW, Karunakaran S & Moxon KA (2013). Neuronal synchrony and the transition to
spontaneous seizures. *Experimental Neurology* 248, 72–84.

Grone BP & Baraban SC (2015). Animal models in epilepsy research: legacies and new
directions. *Nature neuroscience* 18, 339–343.

Harris KD, Henze DA, Csicsvari J, Hirase H & Buzsaki G (2000). Accuracy of tetrode spike
separation as determined by simultaneous intracellular and extracellular measurements.
Journal of Neurophysiology 84, 401–414.

Henze DA, Borhegyi Z, Csicsvari J, Mamiya A, Harris KD & Buzsaki G (2000). Intracellular
features predicted by extracellular recordings in the hippocampus in vivo. *Journal of*
Neurophysiology 84, 390–400.

Hill DN, Mehta SB & Kleinfeld D (2011). Quality metrics to accompany spike sorting of
extracellular signals. *The Journal of Neuroscience* 31, 8699–8705.

Kandel ER & Spencer WA (1961*a*). The pyramidal cell during hippocampal seizure. *Epilepsia* 2,
63–69.

Kandel ER & Spencer WA (1961*b*). Electrophysiology of hippocampal neurons. II. After-
potentials and repetitive firing. *J Neurophysiol* 24, 243–259.

Keller D, Erö C & Markram H (2018). Cell Densities in the Mouse Brain: A Systematic Review.
Frontiers in Neuroanatomy 12, 506–21.

Magloire V, Mercier MS, Kullmann DM & Pavlov I (2018). GABAergic Interneurons in
Seizures: Investigating Causality With Optogenetics. *The Neuroscientist* 35,
107385841880500–15.

Marcuccilli CJ, Tryba AK, van Drongelen W, Koch H, Viemari JC, Peña-Ortega F, Doren EL,
Pytel P, Chevalier M, Mrejeru A, Kohrman MH, Lasky RE, Lew SM, Frim DM &
Ramirez J-M (2010). Neuronal Bursting Properties in Focal and Parafocal Regions in
Pediatric Neocortical Epilepsy Stratified by Histology: *Journal of Clinical*
Neurophysiology 27, 387–397.

Matsumoto H & Marsan CA (1964). Cortical cellular phenomena in experimental epilepsy: Ictal
manifestations. *Exp Neurol* 9, 305–326.

McCormick DA & Feeser HR (1990). Functional implications of burst firing and single spike
activity in lateral geniculate relay neurons. *Neuroscience*; DOI: 10.1016/0306-
4522(90)90225-S.

Merricks EM, Smith EH, McKhann GM, Goodman RR, Bateman LM, Emerson RG, Schevon
CA & Trevelyan AJ (2015). Single unit action potentials in humans and the effect of
seizure activity. *Brain : a journal of neurology* 138, 2891–2906.

Miri ML, Vinck M, Pant R & Cardin JA (2018). Altered hippocampal interneuron activity
precedes ictal onset. *eLife* 7, 1277.

Pakkenberg B & Gundersen HJ (1997). Neocortical neuron number in humans: effect of sex and
age. *The Journal of comparative neurology* 384, 312–320.

Parrish RR, Codadu NK, Mackenzie Gray Scott C & Trevelyan AJ (2019). Feedforward
inhibition ahead of ictal wavefronts is provided by both parvalbumin- and somatostatin-
expressing interneurons. *The Journal of physiology* 597, 2297–2314.

Peyrache A, Dehghani N, Eskandar EN, Madsen JR, Anderson WS, Donoghue JA, Hochberg LR,
Halgren E, Cash SS & Destexhe A (2012). Spatiotemporal dynamics of neocortical
excitation and inhibition during human sleep. *Proceedings of the National Academy of*
Sciences of the United States of America 109, 1731–1736.

Purpura DP, Penry JK, Tower DB & Woodbury DM eds. (1972). *Experimental models of*
epilepsy: A manual for the laboratory worker. Raven Press, New York.

Quian Quiroga R, Nadasdy Z & Ben-Shaul Y (2004). Unsupervised Spike Detection and Sorting
with Wavelets and Superparamagnetic Clustering. *Neural Computation* 16, 1661–1687.

Schevon CA, Tobochnik S, Eissa T, Merricks E, Gill B, Parrish RR, Bateman LM, McKhann GM, Emerson RG & Trevelyan AJ (2019). Multiscale recordings reveal the dynamic spatial structure of human seizures. *Neurobiology of disease* 127, 303–311.

Schevon CA, Weiss SA, McKhann G, Goodman RR, Yuste R, Emerson RG & Trevelyan AJ (2012). Evidence of an inhibitory restraint of seizure activity in humans. *Nature Communications* 3, 1060–11.

Smith EH, Liou J, Davis TS, Merricks EM, Kellis SS, Weiss SA, Greger B, House PA, McKhann GM, Goodman RR, Emerson RG, Bateman LM, Trevelyan AJ & Schevon CA (2016). The ictal wavefront is the spatiotemporal source of discharges during spontaneous human seizures. *Nat Commun* 7, 11098.

Stead M, Bower M, Brinkmann BH, Lee K, Marsh WR, Meyer FB, Litt B, Van Gompel J & Worrell GA (2010). Microseizures and the spatiotemporal scales of human partial epilepsy. *Brain* 133, 2789–2797.

Steriade M & Amzica F (1999). Intracellular study of excitability in the seizure-prone neocortex in vivo. *Journal of Neurophysiology* 82, 3108–3122.

Tóth E, Fabó D, Entz L, Ulbert I & Erőss L (2016). Intracranial neuronal ensemble recordings and analysis in epilepsy. *Journal of Neuroscience Methods* 260, 261–269.

Traub RD & Wong RK (1982). Cellular mechanism of neuronal synchronization in epilepsy. *Science* 216, 745–747.

Trevelyan AJ, Sussillo D, Watson BO & Yuste R (2006). Modular propagation of epileptiform activity: evidence for an inhibitory veto in neocortex. *The Journal of Neuroscience* 26, 12447–12455.

Trevelyan AJ, Sussillo D & Yuste R (2007). Feedforward inhibition contributes to the control of epileptiform propagation speed. *The Journal of Neuroscience* 27, 3383–3387.

Truccolo W, Ahmed OJ, Harrison MT, Eskandar EN, Cosgrove GR, Madsen JR, Blum AS, Potter NS, Hochberg LR & Cash SS (2014). Neuronal ensemble synchrony during human focal seizures. *The Journal of Neuroscience* 34, 9927–9944.

Truccolo W, Donoghue JA, Hochberg LR, Eskandar EN, Madsen JR, Anderson WS, Brown EN, Halgren E & Cash SS (2011). Single-neuron dynamics in human focal epilepsy. *Nature neuroscience* 14, 635–641.

Weiss SA, Banks GP, McKhann GM, Goodman RR, Emerson RG, Trevelyan AJ & Schevon CA (2013). Ictal high frequency oscillations distinguish two types of seizure territories in humans. *Brain* 136, 3796–3808.

Weiss SA, Staba R, Bragin A, Moxon K, Sperling M, Avoli M & Engel J (2018). Interneurons and principal cell firing in human limbic areas at focal seizure onset. *Neurobiology of disease* 124, 183–188.

Wyler AR, Ojemann GA & Ward AA (1982). Neurons in human epileptic cortex: correlation
between unit and EEG activity. *Annals of neurology* 11, 301–308.

FIGURES

Fig. 1. Effects of ictal recruitment on traditional spike sorting methods

Spike sorting relies on stable waveforms from nearby neurons, but ictal activity disrupts features

used to cluster single units. **A.** Example broadband LFP from a single channel of a Utah array

implanted in the posterior temporal lobe of a patient with pharmacoresistant epilepsy (Patient 4,

seizure 1). Dashed red line denotes "global" seizure onset. **B.** Bandpass filtered signal between

300 Hz and 5 kHz of the same signal in A, showing stable single unit activity in the preictal

period. **C.** First principal component score versus time (**i**) of all detected spikes in the multi-unit

activity shown in B, with three clearly separable clusters highlighted, along with the multi-unit

cluster from background distal cells (black). (**ii**) Equivalent first versus second principal

component scores from the preictal period, and (**iii**), during the seizure. Note loss of well-defined

clusters in principal component space.

Fig. 2. Template match spike sorting via convex hulls

A. Well-isolated neurons form distinct clusters in principal component space during interictal time points (**i**), around which a convex hull can be fitted to define boundaries in 3D space within which spikes that match that unit should exist (**ii**). Despite a lack of defined clusters during ictal activity in recruited cortex, this convex hull can be used to select waveforms that are likely to correspond to the preictal unit (**iii**; black), amongst distributed noise (grey). **B(i)** Waveforms from the preictal cluster shown in A(i). **(ii)** Waveforms matched using the convex hull shown in A(iii) (black) from the large distribution of spikes during a seizure (grey). **C.** The probability of each matched waveform originating from the same neuron as its preictal counterpart is calculated by calculating their Mahalanobis distance from the preictal cluster using the first n principal components that explain $\geq$ 95% of the variance. Outliers in the chi-squared distribution for n degrees of freedom denote likely incorrect matches.

Fig. 3. Waveform match probabilities

A(i). All waveforms from a traditionally spike sorted unit in a 10 minute preictal epoch. The probability distribution of each voltage at each time point for these waveforms is shown in **(ii)**, and the spike match probabilities for ictal waveforms matching this unit are shown in **(iii)**, with the distribution of these probabilities inset. **B.** Upper panel shows the probability distribution of all matched waveforms (red), compared to the probability distribution for the original waveforms in the preictal time point (orange) for an example template-matched single unit. A bootstrap estimate of waveform matches expected by chance by comparing against waveforms from other electrodes is shown in blue. The lower panel shows the equivalent distributions across the full population. Mean ± SD is shown above the distributions.

Fig. 4. Example wave shape changes at ictal recruitment in template matched data

A. Waveforms from a convex hull-matched unit in Patient 5, seizure 2, showing reduction in amplitude and increase in FWHM during the seizure (red shaded waveforms, fading from black to red through the seizure; color maintained throughout figure). **B.** Spike amplitude versus FWHM in the unit in A, showing equivalent relationship to the high amplitude BF single unit in Fig. 8D. Wideband LFP from this channel through time (**C**), with time-locked FWHM (**D**) and amplitude (**E**) showing temporal relationship of spike shape changes through the seizure. Note the return to preictal values after seizure termination, and the lack of changes towards decreased FWHM or increased amplitude during the seizure despite the convex hull being equally permissive of alterations in any direction.

Fig. 5. Stereotypy of waveform changes within neurons across seizures

Example waveforms from a single unit in patient 5 showing stereotypy of response of extracellularly recorded action potentials in the peri-recruitment period in 2 seizures separated by 22 hours (**A** and **B** respectively). All waveforms from the 6 seconds prior to ictal recruitment are shown overlaid in **(i)**, and plotted relative to time in **(ii)**, scales maintained throughout. Saturation of color fades from black at -6 seconds, to brightest at the moment of maximal firing rate of MUA at that electrode.

Fig. 6. *Population spike shape alterations in recruited cortex versus penumbral territories*

A & B (i) Probability density plots of spike FWHM (full-width at half maximum) for every detected waveform in the preictal (blue) and ictal time periods for all seizures in recruited cortex (A; *n* = 625 units from 6 seizures in 3 patients) and penumbral tissue (B; *n* = 405 units from 4 seizures in 3 patients). Cumulative probability densities show same calculation on preictal, original data (black), **(ii)** Paired mean FWHM for each unit in the preictal (blue), ictal (red), and postictal (pink) epochs. Note return to preictal ranges after seizure termination. **(iii)** Spike

FWHM of the population through time (10 second window, sliding every 100 ms). Brighter

indicates more density, blue line shows the mean through time, purple line shows the mean value

in the preictal period, red dashed line denotes "global" seizure onset. **C & D.** Same format as A

& B, showing spike amplitude in place of FWHM, for recruited cortex and penumbra

respectively.

Fig. 7. Simultaneous recruitment and penumbral recording.

Two types of activity pattern recorded simultaneously in a patient with the Utah array at the edge

of the clinically defined seizure spread (Patient 6). **A.** LFP from the closest macro-electrode to

the UMA, with time-locked MUA (blue) and first principal component score (purple) through

time from channels 88 and 21 in **B & C** respectively. Note the stability of waveform and

principal component score throughout the seizure in channel 88, with no evidence of tonic firing,

while there is a large spike shape change at the same time in channel 21, at the moment of tonic

firing (maroon bar, C). Paired orange bars in B and C denote burst firing at the end of the seizure

in both locations. These dual activity types both occurred immediately next to the LFP in A, and

thus these patterns cannot be differentiated at the macro LFP level.

Fig. 8. Ictal recruitment in the mesial temporal lobe recorded with BF electrodes

A. (i) Broadband LFP from the closest macro contact to the microwires in a BF electrode in the mesial temporal lobe during a spontaneous seizure, and **(ii)** the bandpass filtered MUA from one of the microwires in the temporal pole. "Global" seizure onset is denoted by the red dashed line, with subsequent local recruitment to the seizure at the pink dashed line. **B.** Magnification of the region denoted by the grey bar in A, showing seizure onset through passage of the ictal wavefront in LFP and MUA (colours maintained) in the same microwire **(i)**, and a microwire from a nearby separate BF in the hippocampal body **(ii)**. Note the pre-recruitment stability in the spike amplitude, which is immediately reduced upon ictal invasion, and the simultaneous quiescence in the hippocampal body, followed by a similar, time-delayed amplitude change after recruitment. **C.** Mean ± SD of waveforms prior to ictal invasion (blue) and after recruitment during the seizure (red), showing reduction in amplitude and increase in FWHM in A(ii). **D.** Spike amplitude versus spike FWHM for the defined unit, with color maintained from A,

632   transitioning from blue to red through seizure invasion. Note the bimodal clusters that

633   correspond to pre- and post-recruitment, and the similarity to Fig. 4B.

Fig. 9. Neuronal firing through ictal recruitment in an example focal seizure

A. Local field potential (LFP) from an example channel during seizure 1 in Patient 3, with the

calculated passage of ictal wavefront marked by red dashed line, with ± 2 seconds shaded red. **B.**

Raster plot of all units found using convex hull template matching in this seizure, ordered by

firing rate, and plotted relative to the moment of ictal recruitment at that channel. **C.** Probabilistic

instantaneous firing rate of the population of single units as estimated by convolving a Gaussian

kernel (200 ms SD) over the spike times. The firing rate shown is probabilistic by scaling the

Gaussian kernel's amplitude by the likelihood of each individual spike originating from its

assigned preictal unit, as calculated by its voltage-time probabilities (see Methods & Fig. 3). As

such, the firing rate has not been biased by excessive matching of dissimilar waveforms during

the ictal activity. Note the intense, tonic firing during the seizure invasion, and sustained, above

baseline firing until seizure termination.

TABLES

Patient	Demographics	Implant location	Seizure type	Seizure onset	Seizures analyzed	# units
1	30 M	Frontal (Left frontal convexity)	FIA	Left supplementary motor area	1	79
2	39 M	Frontal (Left dorsolateral frontal lobe)	FIA	Left frontal operculum	2	[61, 61]
3	25 F	Temporal (Left inferior temporal gyrus)	FIA	Left basal/ anterior temporal	3	[111, 101, 91]
4	19 F	Temporal (Right posterior temporal)	FTBTC	Right posterior lateral temporal	1	110
5	24 M	Temporal (Left inferior temporal gyrus)	FIA	Left mesial temporal with spread to lateral temporal	2	[120, 133]
6	30 M	Temporal (Right mesial temporal gyrus)	FA	Right subtemporal	1	240

649   *Table 1. Demographics and data for patients implanted with Utah arrays*

650   Individual patient demographics for grid/Utah array implant cases, including the number of

651   isolated single units found for each seizure in each patient. FIA: focal with impaired awareness;

652   FTBTC: focal to bilateral tonic-clonic.

Patient	Demogr aphics	Implant locations with units	Seizure type	Seizure onset	Seizures analyzed	# units	Microwires with waveform alterations	Time of recruitment (s)	Clinical correlation
7	29 F	Mesial temporal (Left hippocampus)	FTBTC	Left lateral frontal	2	[2, 2]	Left hippocampus	16.3	Spread
8	29 F	Mesial temporal (Left hippocampus)	FA	Left mesial temporal	2	[2, 1]	Left hippocampal head	16.1	✓+
9	55 F	Mesial temporal (Right hippocampus)	FIA	Right lateral temporal	1	9	–	–	✓−
10	25 M	Mesial temporal (Left hippocampal body)	FIA	Right mesial temporal and right parietal (dual)	1	1	Left hippocampal body	42.7	Spread
11	19 M	Frontal (Right anterior cingulate)	FIA	Right lateral temporal	1	3	Right supplementary motor area	9.1	Spread
12	20 F	Frontal (Left anterior & mid-cingulate)	FA	Left mesial temporal	1	15	Left mid-cingulate	19.5	Spread
13	21 F	Mesial temporal (Left hippocampus)	FA	Left anterior temporal pole	1	1	–	–	✓−
14	40 M	Mesial temporal/frontal (Left hippocampus & anterior cingulate)	FIA	Left lateral frontal	2	[9, 10]	Left hippocampus; Left mid-cingulate	39.3 (HC); 29.0 (MC)	Spread
15	25 F	Mesial temporal (Right hippocampus body & head)	FIA	Right lateral temporal spreading to mesial temporal	1	11	Right hippocampal body; Right hippocampal head	9.1 (HCB); 8.0 (HCH)	✓+
16	35 F	Mesial temporal (Bilateral hippocampus)	FIA	Left mesial temporal	1	5	Left hippocampal head	8.0	✓+
17	40 M	Mesial temporal (Left hippocampus)	FIA	Left mesial parietal	1	2	–	–	✓−
18	39 M	Frontal (Right mesial cingulate)	FIA; FTBTC	Right mesial temporal and left orbitofrontal (2 types)	2	[8, 4]	–	–	✓−
19	30 M	Frontal (Right mesial cingulate)	FA; FIA	Left mesial temporal and left insula	2	[1, 1]	–	–	✓−
20	23 F	Mesial temporal (Right hippocampus)	FIA	Right posterior temporal	1	3	–	–	✓−
21	35 M	Mesial temporal (Bilateral hippocampus)	FA	Right insula and somatosensory cortex	3	[7, 8, 7]	Right hippocampal head	Sub-threshold	Spread /X+
22	44 F	Frontal (Right anterior cingulate)	FTBTC	Left cingulate	1	1	Right anterior/mid-cingulate	15.2	Spread
23	51 F	Frontal (Left anterior cingulate)	FIA	Left mesial temporal	1	4	Left anterior cingulate	12.6	Spread
24	20 M	Frontal (Left anterior cingulate)	FIA	Left posterior temporal	1	1	–	–	✓−
25	30 F	Frontal (Right anterior cingulate)	FIA	Left mesial temporal	1	1	–	–	✓−
26	20 M	Mesial temporal (Right hippocampus)	FIA	Right mesial temporal	2	[1, 2]	Right hippocampal body	10.4	✓+
27	32 M	Mesial temporal (Bilateral hippocampus)	FIA	Bilateral mesial temporal	3	[4, 3, 3]	Hippocampal body	9.8	✓+

Table 2. Demographics and results for patients implanted with Behnke-Fried arrays

Individual patient demographics for stereo-EEG/Behnke-Fried array cases, including the number

of isolated single units found for each seizure in each patient, implant locations that showed

significant waveshape alterations, the delay until waveforms surpassed > 1 SD beyond the

preictal mean, and whether clinical observations matched these findings. FTBTC: focal to

bilateral tonic-clonic; FA: focal aware; FIA: focal with impaired awareness; HC: hippocampus;

MC: mid-cingulate; "Spread": matches clinical observations of seizure spread; ✓+: true positive

match for clinical observations; ✓–: true negative match for clinical observations; ✗+: false

positive match for clinical observations.

Patient	Seizure #	Recruited?	Units from convex hull [n]	Units showing FR increase in ictal wavefront [n (%)]		Units showing significant FWHM increase [n (%)]		Units showing significant amplitude decrease [n (%)]	
1	1	✗	52	N/A		2	(4%)	3	(6%)
2	1	✗	55	N/A		9	(16%)	8	(15%)
	2	✗	49	N/A		3	(6%)	2	(4%)
3	1	✓	80	77	(96%)	63	(79%)	20	(25%)
	2	✓	80	70	(88%)	68	(85%)	38	(48%)
	3	✓	79	67	(85%)	66	(84%)	40	(51%)
4	1	✓	101	77	(76%)	93	(92%)	45	(45%)
5	1	✓	117	82	(70%)	114	(97%)	103	(88%)
	2	✓	104	73	(70%)	85	(82%)	63	(61%)
6	1	✗	221	N/A		105	(48%)	86	(39%)

Table 3. Single unit data from Utah array population analyses

Total number of units found in each patient and seizure for population analyses in Utah array

cases via the convex hull template-matching method, along with the number of units showing

significant increases in firing rate, spike full-width-at-half-maximum (FWHM), and decreases in

amplitude during the seizure.

VIDEO LEGENDS

Video 1. Location specific waveform changes during a spontaneous human seizure

Local field potential (top) and spike amplitudes for three units in three different locations

(middle; red, blue and purple), with associated spike waveforms shown to the right. Shading

shows the mean ± 2 SD for these units' preictal spike amplitudes. The locations of the BF

microwires that recorded these units are shown below, with colors maintained. Note the three

activity patterns: cessation of firing at seizure onset in the anterior temporal lobe (blue), stability

followed by loss of spike amplitude in the anterior cingulate (red), and stability throughout in the

mid-cingulate (purple).

Video 2. Single unit waveform alterations in an ictal Behnke-Fried recording

Single units undergo waveshape changes upon recruitment to the seizure, shown in real-time.

Upper trace: MUA bandpassed signal (300 Hz to 5 kHz; white), with current time shown in red,

earliest electrographic ictal activity in the patient occurs at dashed magenta line, with local

recruitment occurring at blue dashed line. Lower panels: shaded regions show mean ± 2 SD for

preictal single units in red and blue, and lower amplitude multiunit activity in yellow (left).

Waveforms are displayed in real-time, with colors matching their assigned unit, and color

saturation showing the probability of a true match to that unit. The first two principal component

scores for these waveforms are shown on the right, with colors maintained. Note the stability of

waveforms prior to local recruitment, including after seizure onset, followed by marked loss of

amplitude at the moment of recruitment with associated tonic firing.