Calcium-independent lipid release from astrocytes modulates neuronal excitability

Fushun Wang[1,2], Heather B. Bradshaw[3], Salvador Pena[4], Beata Jablonska[5], Julia Xavier[6], Sheng Gong[7], Baoman Li[1], Devin Chandler-Militello[1], Lane K. Bekar[8], and Nathan A. Smith[9,10*]

[1]Departments of Neurosurgery and Neurology, University of Rochester School of Medicine and Dentistry, Rochester, NY 14642, USA.
[2]Institute of Brain and Psychological Sciences, Sichuan Normal University, Chengdu, Sichuan 610060, China.
[3]Department of Psychological and Brain Sciences, Indiana University, Bloomington, IN 47405, USA
[4]Department of Anesthesiology, Wake Forest School of Medicine, Winston-Salem, North Carolina
[5]Center for Neuroscience Research, Children's National Research Institute, Children's National Hospital, Washington DC 20010, USA.
[6]George Washington University, Washington DC 20052, USA.
[7]Department of Physiology and Biophysics, Case Western Reserve University School of Medicine, Cleveland, OH 44106, USA
[8]Department of Anatomy, Physiology, and Pharmacology, University of Saskatchewan, Saskatoon, SK S7N 5E5.
[9]Del Monte Institute for Neuroscience, Department of Neuroscience, University of Rochester, School of Medicine and Dentistry, Rochester, USA
[10]Lead Contact

*Correspondence should be addressed to Dr. Nathan Anthony Smith,
 Email: Nathan_Smith@urmc.rochester.edu, Twitter: @SmithLab2018

ABSTRACT

An accumulating amount of data suggests that Ca^{2+}-dependent gliotransmitter release plays a key role in the modulation of neuronal networks. Here, we tested the hypothesis that in response to agonist exposure, astrocytes release lipid modulators through activation of Ca^{2+}-independent phospholipase A_2 ($iPLA_2$) activity. We found that cultured rat astrocytes treated with selective ATP and glutamatergic agonists released arachidonic acid (AA) and/or its derivatives, including the endogenous cannabinoid 2-arachidonoyl-sn-glycerol (2AG) and prostaglandin E2 (PGE_2). Surprisingly, the buffering of cytosolic Ca^{2+} resulted in a sharp increase in agonist-induced lipid release by astrocytes. In addition, the astrocytic release of PGE_2 increased miniature excitatory postsynaptic potentials (mEPSPs) by inhibiting the opening of neuronal Kv channels in brain slices. This study provides the first evidence showing that a Ca^{2+}-independent pathway regulates the release of PGE_2 from astrocytes and further demonstrates the functional role of astrocytic lipid release in the modulation of synaptic activity.

SIGNIFICANCE

Until now, most studies that implicate astrocytes in the modulation of synaptic activity have focused on Ca^{2+}-dependent release of traditional gliotransmitters such as D-serine, ATP, and glutamate. Mobilization of intracellular stores of Ca^{2+} occurs within a matter of seconds, but this novel Ca^{2+}-independent lipid pathway in astrocytes could occur on a faster time scale and thus play a role in the rapid signaling processes involved in synaptic potentiation, attention, and neurovascular coupling.

INTRODUCTION

In addition to being a major contributor to the dry weight of the adult brain, lipids are an essential component of the phospholipid bilayer and are mainly composed of long-chain polyunsaturated fatty acids (PUFAs), such as arachidonic acid (AA) and docosahexaenoic acid (DHA) (Sinclair, 1975). The liver is the major site of AA synthesis, but the brain can produce AA and DHA in situ from their precursor fatty acids, linoleic and linolenic acids (Dhopeshwarkar and Subramanian, 1976). Astrocytes play a central role in the synthesis of AA and DHA in the brain. Through their vascular endfeet, astrocytes have prime access to fatty acid precursors that arrive by crossing the blood–brain barrier (BBB) and serve as the major site for processing essential fatty acids in the central nervous system (CNS) (Moore, 1993). Astrocytes also play a key role in the macroscopic distribution of lipids in the brain parenchyma via perivascular glymphatic flux (Rangroo Thrane et al., 2013a; Plog and Nedergaard, 2018).

Lipids have gained much attention for their role as bioactive mediators in the CNS (Carta et al., 2014; Ledo et al., 2019). Numerous studies have focused on lipids in relation to functional hyperemia and synaptic activity. For instance, PGE_2 is a potent vasodilator and vasoconstrictor that regulates CNS blood flow (Zonta et al., 2003; Takano et al., 2006; Gordon et al., 2007; Dabertrand et al., 2013; MacVicar and Newman, 2015; Czigler et al., 2019) and modulates the membrane excitability of CA1 pyramidal neurons during synaptic activity (Chen and Bazan, 2005). Furthermore, AA and its derivatives are important intracellular secondary messengers that can modulate the activities of various ion channels (Piomelli, 1993; Meves, 1994; Horimoto et al., 1997; Boland and Drzewiecki, 2008; Cordero-Morales and Vasquez, 2018). In addition, PGE_2 can suppress the outwardly rectifying Kv current in sensory neurons (Nicol et al., 1997; Evans et al., 1999), whereas AA suppresses Kv channels in the soma or dendrites of pyramidal neurons and consequently broadens their presynaptic action potentials (Carta et al., 2014) and enhances EPSPs (Ramakers and Storm, 2002). However, those studies focused on the effects of neuronal lipid release on synaptic activity and paid little attention to receptor-mediated pathways by which

astrocytic lipids might influence synaptic activity. Given that the majority of AA, DHA, and other lipids present in the extracellular fluid are produced by astrocytes (Moore et al., 1991), receptor-mediated release could be a significant factor in the modulation of synaptic activity.

In culture, astrocytes can release AA via a $cPLA_2$ Ca^{2+}-dependent pathway upon the activation of metabotropic glutamate (mGluR) and P2Y purine (ATP) receptors (Bruner and Murphy, 1990; Stella et al., 1994; Stella et al., 1997; Chen and Chen, 1998). In addition, astrocytes also express the Ca^{2+}-independent PLA_2 ($iPLA_2$) enzyme (Sun et al., 2005). Both of these isoforms are activated by the G-protein βγ subunit (Jelsema and Axelrod, 1987; Murayama et al., 1990; van Tol-Steye et al., 1999), but iPLA2 does not require Ca^{2+} or PKC phosphorylation for activation (Winstead et al., 2000). Receptor-stimulated $iPLA_2$ activation can induce the release of AA and DHA by numerous types of cells (Gross et al., 1993; Portilla et al., 1994; Akiba et al., 1998; Seegers et al., 2002; Tay and Melendez, 2004), but this activation has not been fully explored in the case of astrocytes.

Previous work in our laboratory and elsewhere has demonstrated that astrocytes are capable of releasing gliotransmitters upon stimulation with mGluR or ATP receptor agonists, but the impact of this signaling pathway on synaptic activity remains controversial. Recent studies with genetically encoded calcium indicators allowed for the identification of localized Ca^{2+} signals within astrocytic fine processes and confirmed Ca^{2+}-dependent astrocytic effects on synaptic activity (Yu et al., 2018). However, the existence of Ca^{2+}-independent lipid signaling in astrocytes is still unclear. Therefore, in this study, we tested the hypothesis that astrocytes support Ca^{2+}-independent lipid signaling to modulate synaptic transmission. We used Ca^{2+} chelation to show that ATP and mGluR agonists can release lipids through $iPLA_2$ activation, resulting in the potentiation of synaptic activity. Because we currently lack tools to assess Ca^{2+}-independent signaling in astrocyte fine processes adjacent to synapses, we used Ca^{2+} inhibition on a large scale to evaluate this phenomenon (see discussion).

MATERIALS AND METHODS

Culture

Cultured neocortical astrocytes were prepared from postnatal day 1 or 2 Wistar rat pups (Taconic Farms, Inc.) of either sex, as previously described (Lin et al., 1998). In brief, cerebral cortices were dissected on ice, and the meninges were removed. The tissue was washed three times in Ca^{2+}-free Hanks' balanced salt solution, triturated, filtered through a 70 μm nylon mesh, and centrifuged. The pellet was resuspended in 10% fetal bovine serum in Dulbecco's modified Eagle's medium (DMEM)/F12 containing penicillin (100 IU ml^{-1}) and streptomycin (100 μg ml^{-1}) and transferred to culture flasks. The cells were maintained at 37 °C in an incubator with humidified air and 5% CO_2. The medium was changed after 24 hours and twice a week thereafter. More than 95% of the cells immunostained were GFAP positive. When the cells became confluent, they were rinsed two times in Ca^{2+}-free Hanks' balanced salt solution, suspended in 0.05% trypsin-containing PBS for 1 minute, resuspended in DMEM/F12, centrifuged to remove the trypsin-containing supernatant, and then plated in 24-well plates. The experiments were performed when the cells were 95% confluent.

Viral vectors and viral transduction

Viral vectors driving GFAP cyto-GCaMP6f (Baljit S. Khakh) were obtained from the University of Pennsylvania Vector Core (AAV 2/5 serotype). Secondary rat astrocytic cultures were transduced with the AAV GFAP cyto-GCaMP6f. After transduction, the cultures were incubated at 37 °C for 5 days prior to Ca^{2+} imaging experiments.

Ca^{2+} imaging

Cultured cells in 24-well plates were transduced with the AAV GFAP cyto-GCaMP6f and incubated with various pharmacological agents for 30 minutes at 37 °C. Using confocal microscopy (Olympus FV500), calcium wave activity was evaluated by adding an equal volume of medium containing 100 μM ATP to each well. Relative increases in the fluorescence signal evoked by P2Y

receptor agonist exposure were compared with baseline fluorescence (ΔF/F) as previously described (Nedergaard, 1994; Smith et al., 2018).

Radiolabeling and assessment of AA release

Confluent rat astrocytic cultures were incubated with 100 nCi [5,6,8,9,11,12,14,15-^{3}H]-AA (PerkinElmer) overnight before experiments. The cells were washed three times with serum-free medium and then allowed to recover for 20 minutes. Before stimulating P2YRs with 100 μM ATP, the cells were incubated with the appropriate inhibitors for 10 to 12 minutes. Aliquots of medium were taken 15 minutes after agonist stimulation, and the levels of ^{3}H-AA and/or metabolites were measured by liquid scintillation counting.

HPLC/MS/MS analysis of lipids

Confluent rat astrocytes were washed with serum-free media and then allowed to recover for 1 hour in serum-free media. At this time, the cells were incubated with vehicle or 20 μM cyclopiazonic acid (CPA) for 15 minutes and then challenged with 100 μM ATP or vehicle for 15 minutes, as described in the section on [^{3}H]-AA experiments. The media were then removed and put aside, and 2 ml of HPLC-grade methanol was added to the flask for 5 minutes. After removing the methanol, an additional 2 ml of HPLC-grade methanol was added, the cells were scraped from the sides of the flask, and the contents were added to the previous two retained fractions. Deuterium standards were added to a final concentration of 200 pM, and the samples were centrifuged at 19,000 × g and 24 °C. Lipids in the eluent were partially purified on 500 mg C18 Bond Elut solid phase extraction columns (Varian), concentrated into fractions of 40, 60, 85, and 100% methanol, and analyzed using HPLC/MS/MS (Shimadzu SCL10Avp-Wilmington, DE, USA; API 3000 Applied Biosystems/MDS SCIEX, Foster City, CA, USA) as previously described (Leishman et al., 2018). Over 50 lipids were targeted in these analyses, including 40 lipoamines, 3 acylglycerols, 2 free fatty acids, and 2 prostaglandins (namely, X and Y). The results of pilot studies showed that the analyte concentrations were too low for reliable detection in the medium

alone, but the extraction of medium and cells together provided sufficient lipid concentrations for reliable assays.

PGE$_2$ release assessment via PGE$_2$ immunoassay

Confluent rat astrocytic cultures were washed 3 times with serum-free medium and then allowed to recover for 20 minutes. Before stimulation with 100 µM ATP, the cells were incubated with the appropriate P2Y inhibitors for 10 to 12 minutes. Aliquots of the medium were collected 15 minutes after stimulation and PGE2 content was determined using an immunoassay kit (Cayman Chemicals) according to the manufacturer's instructions.

Western blotting

Protein in samples harvested from the 24-well plates was separated by SDS–PAGE and transferred to a nitrocellulose membrane, which was then blocked with Tris-buffered saline containing 0.05% (wt/vol) Tween 20 and 5% nonfat dry milk. The primary antibodies used were anti-iPLA2 (Sigma, St. Louis, MO) and anti-β-actin (Cell Signaling, Danvers, MA) at 1:1000 to 1:2000 dilutions in blocking buffer. Chemiluminescence from horseradish peroxidase-linked secondary antibodies was detected using the ChemiDoc™ XRS+ System and running Image Lab™ software.

Isolation of human fetal astrocytes

Human fetal forebrain tissues were obtained from second-trimester aborted fetuses at 20 weeks of gestational age. Tissues were obtained from aborted fetuses, with informed consent and tissue donation approval from the mothers using protocols approved by the Research Subjects Review Board of the University of Rochester Medical Center. No patient identifiers were made available to or known by the investigators; no samples with known karyotypic abnormalities were included. Forebrain tissue samples were collected and washed 2-3 times with sterile Hank's balanced salt solution containing Ca^{2+}/Mg^{2+} (HBSS$^{+/+}$). The cortical plate region (CTX) of the fetal forebrain was dissected and separated from the ventricular zone/subventricular zone (VZ/SVZ) portion. CTX was then dissociated from papain as previously described (Keyoung et al., 2001). The cells were

resuspended at a density of 2-4 x 10^6 cells/ml in DMEM/F12 supplemented with N2, 0.5% FBS, and 10 ng/ml bFGF and plated in suspension culture dishes. The day after dissociation, the cortical cells were recovered and subjected to magnetic activated cell sorting (MACS) to purify the astrocyte progenitor population. The recovered cells were briefly incubated with CD44 microbeads according to the manufacturer's recommendations (Miltenyi Biotech). The cells were then washed, resuspended in Miltenyi Washing buffer, and bound to a magnetic column (Miltenyi Biotech). The bound CD44+ astrocyte progenitor cells were eluted, collected, and then washed with DMEM/F12. The purified human fetal astrocyte progenitors were cultured in DMEM/F12 supplemented with N2 and 5% FBS to further differentiate them. To prepare culture dishes for PGE2 immunoassays or immunocytochemistry, fetal cortical astrocytes were dissociated with TrypLE (Invitrogen) into single cells and then plated onto poly-*L*-ornithine/laminin-coated 24-well plates (50,000 cells per well).

shRNA-mediated lentiviral knockdown of iPLA$_2$ in astrocytes

Rat astrocyte cultures were plated in a 24-well plate and grown to approximately 50% confluence. The cultures were transduced overnight by adding either group VI iPLA$_2$ shRNA (r) lentiviral particles (sc-270117-V) or control shRNA lentiviral particles-A (sc-108080) directly to culture medium containing polybrene (sc-134220) according to the manufacturer's instructions with minor modifications (all from Santa Cruz Biotechnology, Santa Cruz). At 24 hours after transfection, the culture medium was removed, and fresh culture medium without polybrene was added. Experiments and Western blot analysis were performed 7 days after transduction.

Acute hippocampal slice preparation and electrophysiology

Unless otherwise noted, 15–21-day-old C57BL/6 (Charles River, Wilmington, MA), MrgA1[+/-] transgenic, and littermate control MrgA1[-/-] pups (courtesy of Dr. Ken McCarthy) (Fiacco et al., 2007) of either sex were used to prepare hippocampal slices as previously described (Wang et al., 2012). The pups were anesthetized in a closed chamber with isoflurane (1.5%) and

decapitated. The brains were rapidly removed and immersed in an ice-cold cutting solution that contained (in mM) 230 sucrose, 2.5 KCl, 0.5 $CaCl_2$, 10 $MgCl_2$, 26 $NaHCO_3$, 1.25 NaH_2PO_4, and 10 glucose, at a pH of 7.2-7.4. Coronal slices (400 µm) were cut using a vibratome (Vibratome Company, St. Louis) and transferred to oxygenated artificial cerebrospinal fluid (aCSF) that contained (in mM) 126 NaCl, 4 KCl, 2 $CaCl_2$, 1 $MgCl_2$, 26 $NaHCO_3$, 1.25 NaH_2PO_4, and 10 glucose (pH = 7.2-7.4; osmolarity = 310 mOsm). The slices were then incubated in aCSF for 1-5 hours at room temperature before electrophysiological measurement. The experiments were performed at room temperature (21-23 °C). During the recordings, the slices were placed in a perfusion chamber and superfused with aCSF gassed with 5% CO_2 and 95% O_2 at room temperature. The cells were visualized with a 40× water-immersion objective and differential inference contrast (DIC) optics (BX51 upright microscope, Olympus Optical, New York, NY). Patch electrodes were fabricated from thin-wall glass filaments (World Precision Instruments) on a vertical puller; the resistance of the pipette was approximately 6-9 MΩ with the addition of intracellular pipette solution. The pipette solution contained 140 mM K-gluconate, 5 mM Na-phosphocreatine, 2 mM $MgCl_2$, 10 mM HEPES, 4 mM Mg-ATP, and 0.3 mM Na-GTP (pH adjusted to 7.2 with KOH). The current–voltage (I–V) curves of the voltage–gated potassium currents were recorded under a voltage clamp using an AxoPatch MultiClamp 700B amplifier (Axon Instruments, Forster City, CA). When measuring outward currents, QX314 (0.5 mM) was added to the pipette solution to block Na^+ currents. For recordings of miniature excitatory postsynaptic potentials (mEPSPs), 0.5 µM TTX was added to the aCSF. The junction potential between the patch pipette and the bath solution was zeroed before the giga seal was formed. Patches with seal resistances less than 1 GΩ were rejected. The data were low-pass filtered at 2 kHz and digitized at 10 kHz with a Digidata 1440 interface controlled by pClamp Software (Molecular Devices, Union City, CA).

Pharmacological agents used in culture and slice experiments

Adenosine 5 -triphosphate (ATP, 100 µM); cyclopiazonic acid (CPA, 20 µM); *trans*- (1S, 3R)-1-Amino-1,3-cyclopentanedicarboxylic acid (*t*-ACPD, 100 µM); (±)-α-amino-3-hydroxy-5-

methylisoxazole-4-propionic acid hydrobromide (($\pm$)-AMPA, 100 µM); Prostaglandin E_2 (PGE_2, 50 µM, Tocris); Phe-Met-Arg-Phe amide (FMRF, 15 µM); Thr-Phe-Leu-Leu-Arg-NH_2 (TFLLR-NH_2, 30 µM, Tocris); N-acetyl-Asp-Glu (NAAG, 100 µM); (1R, 4R, 5S, 6R)-4-amino-2-oxabicyclo [3.1.0] hexane-4,6-dicarboxylic acid disodium salt (LY379268, 100 µM, Tocris); calmidazolium chloride (CMZ, 2 µM, Tocris); methyl arachidonyl fluorophosphonate (MAFP, 10 µM); bromoenol lactone (Bel, 10 µM); AA (50 µM); N-(2,6-dimethylphenylcarbamoylmethyl) triethylammonium chloride (QX314, 1 mM, Tocris); 4-(4-,9-diethoxy-1,3-dihydro-1-oxo-2H-benz[f]isoindol-2-yl)-N-(phenylsulfonyl) benzeneacetamide (GW627368X 3 µM, Tocris); 6-Isopropoxy-9-xanthone-2-carboxylic acid (AH6809, 10 µM, Tocris); N-(piperidin-1-yl)-5-(4-iodophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1H-pyrazole-3-carboxamide (AM251, 5 µM, Tocris); Tetrodotoxin (TTX, 0.5 µM, Tocris); 1,2-bis(2-aminophenoxy)ethane-N,N,N',N'-tetraacetic acid (BAPTA, 50 µM, Tocris); and 1,2-Bis(2-aminophenoxy)ethane-N,N,N',N'-tetra-acetic acid tetrakis (acetoxymethylester) (BAPTA-AM, 20 µM) were utilized. All chemicals were obtained from Sigma unless otherwise noted.

Statistical analysis

Statistical significance was evaluated by one-way ANOVA and *post hoc* tests (Tukey and Dunn) using Prism software and deemed significant when $P<0.05$ for the [^{3}H]-AA and PGE_2 assay. The normality of the data was evaluated by the Shapiro–Wilk test with a = 0.05. For electrophysiology experiments, significance was determined by paired or unpaired t tests or Tukey–Kramer *post hoc* multiple comparison tests. HPLC/MS/MS lipidomic data were analyzed with ANOVA and Fisher's LSD *post hoc* test using SPSS when $P<0.05$ or $P<0.10$. All results are reported as the mean ± s.e.m.

RESULTS

Ca^{2+}-independent release of [^{3}H]-AA and its metabolites from cultured astrocytes

We first assessed the efficiency of preloading with an inhibitor of the endoplasmic reticulum (ER) Ca^{2+} pump, cyclopiazonic acid (CPA) (20 µM), or the cytosolic Ca^{2+} chelator 1,2-bis(o-

aminophenoxy) ethane-N,N,N′,N′-tetraacetic acid (acetoxymethyl ester) (BAPTA-AM) (20 µM), which was able to block ATP (100 µM)-induced increases in cytosolic Ca^{2+} in cultured rat astrocytes. Images of cytosolic Ca^{2+} (AAV GFAP cyto-GCaMP6f) showed that the purine agonist ATP (100 µM) induced a prompt increase in Ca^{2+} that was completely blocked in CPA- and BAPTA-loaded cultures, whereas 10 µM methylarachidonyl fluorophosphate (MAFP), a nonspecific inhibitor of both $cPLA_2$ and $iPLA_2$, or 10 µM bromoenol lactone (Bel) (Cornell-Bell et al.), a specific inhibitor of $iPLA_2$, did not affect ATP-induced increases in Ca^{2+} in astrocytic culture **(Figure 1A-C)**.

As a broad-based approach to AA-specific lipidomics, Ca^{2+}-independent release of AA and/or its metabolites was performed using a $[^3H]$-AA assay **(Figure 2A)**. Cultured rat astrocytes were preincubated overnight with $[^3H]$-AA with sufficient time to allow for its incorporation into multiple biosynthetic and metabolic pathways. This process includes incorporation into membrane phospholipids, which are precursors for endocannabinoids, and related lipids, which are precursors for AA release (Chen and Chen, 1998; Strokin et al., 2003). Therefore, this assay was used to determine whether any AA precursors or metabolites derived from the $[^3H]$-AA incorporated into the cell were being released; hereafter, we refer to the composite of $[^3H]$-labeled AA metabolites as $[^3H]$-AA. ATP alone failed to induce a detectable release of $[^3H]$-AA (**Figure 2B)**; however, in cultures pretreated for 10 to 12 minutes with CPA or BAPTA-AM, ATP induced a robust increase in the release of $[^3H]$-AA, whereas CPA alone had no such effect **(Figure 2B)**. Similarly, upon stimulation with a combination of the nonselective mGluR agonist tACPD (100 µM) and the ionotropic glutamate receptor agonist AMPA (100 µM), we observed significant release of $[^3H]$-AA when cytosolic Ca^{2+} was blocked with CPA but not in cultures without CPA pretreatment **(Figure 2C)**. These observations confirm that rat astrocytes can release AA derivatives, which are key precursors of bioactive eicosanoids (Strokin et al., 2003; Rosenegger et al., 2015), but that AA release is, surprisingly, inhibited by increases in cytosolic Ca^{2+}.

To explore the mechanism of Ca^{2+}-independent release of $[^3H]$-AA, we next evaluated whether inhibition of Ca^{2+}-sensitive cPLA$_2$ or Ca^{2+}-insensitive iPLA$_2$ enzymes would reduce $[^3H]$-AA release. Pretreatment of astrocytes with 10 μM MAFP significantly decreased ATP-induced release of $[^3H]$-AA in cultures exposed to CPA **(Figure 2D)**. Similarly, 10 μM BEL treatment suppressed the release of $[^3H]$-AA, thus confirming that iPLA$_2$ plays a role in Ca^{2+}-independent lipid release **(Figure 2D)**. $[^3H]$-AA release was observed only when agonist-induced increases in cytosolic Ca^{2+} were blocked **(Figure 2B-C)**, suggesting that intracellular Ca^{2+} inhibits Ca^{2+}-independent iPLA$_2$.

Calmodulin is a potent Ca^{2+}-dependent inhibitor of iPLA$_2$ (Wolf and Gross, 1996). To assess the interaction between calmodulin and iPLA$_2$, cells were treated with calmidazolium (CMZ), which is an inhibitor of Ca^{2+}/calmodulin interactions that has been shown to eliminate calmodulin blocking of iPLA$_2$ (Wolf and Gross, 1996). In the presence of CMZ (2 μM), ATP treatment led to the release of a significant amount of $[^3H]$-AA, which was comparable to the release induced by blocking increases in cytosolic Ca^{2+} by preloading astrocytes with either CPA or BAPTA **(Figure 2D)**. This observation suggested that Ca^{2+} acts primarily as a brake through calmodulin, which effectively inhibits iPLA$_2$ activity (Wolf and Gross, 1996; Wolf et al., 1997). These findings are consistent with previous studies showing that iPLA$_2$ is involved in receptor-mediated AA release from pancreatic islet cells (Gross et al., 1993), smooth muscle cells (Lehman et al., 1993), and endothelial cells (Seegers et al., 2002). Taken together, these data provide evidence of a new signaling mechanism in astrocytes involving Ca^{2+}-independent iPLA$_2$, which is the major PLA$_2$ isoform in the brain and accounts for 70% of total PLA$_2$ activity (Yang et al., 1999)

Targeted lipidomics reveals Ca^{2+}-independent lipid production in cultured astrocytes

Using lipid extraction and partial purification methods coupled to HPLC/MS/MS, we performed targeted lipidomic screening of cultured astrocytes that were preincubated with CPA and then challenged with ATP and compared the results to that of vehicle control astrocytes. Of the 50 lipids screened, 30 were present in each of the samples and were used for comparative analyses. The

lower panel in Figure 3A lists all of the concentrations as the mean ± SEM **(Figure 3A)**. **Figure 3A** summarizes the lipids that showed significant concentration differences, as well as the magnitudes of the differences. Among the 30 lipids detected, 16 (including AA and PGE_2) increased upon ATP challenge in the presence of CPA. **Figure 3B-C** shows representative chromatograms of the HPLC/MS/MS methods used to detect PGE_2. Notably, the levels of 5 of the 8 AA derivatives in the set, including docosahexaenoyl ethanolamine, were significantly increased following ATP challenge with CPA preincubation, although the most dramatic increases were observed for the prostaglandin PGE_2. **Figure 3D** shows the representative differences in concentration of PGE2, DEA, and AA.

Ca^{2+}-independent release of PGE_2 from cultured astrocytes

Given that Ca^{2+}-dependent lipid release from astrocytes has been previously implicated in vasoregulation (Zonta et al., 2003; Takano et al., 2006; Gordon et al., 2008), we were surprised that PGE_2 was released via a Ca^{2+}-independent mechanism. Our HPCL/MS/MS method requires large quantities of cells to detect PGE_2, whereas the PGE_2 ELISA can be used to accurately detect PGE_2 release in 24-well cultures. Therefore, we used ELISAs to explore the Ca^{2+} dependence of astrocytic PGE_2 release. Because $iPLA_2$ is essential for Ca^{2+}-independent liberation of AA-derived lipids, we first assessed whether knockdown of $iPLA_2$ via viral transduction would inhibit astrocytic release of PGE_2. In the presence of CPA (20 µM), ATP (100 µM) failed to induce PGE_2 release in shRNA-transduced astrocytic cultures, whereas there was a significant increase in the release of PGE_2 by ATP in the presence of CPA in control shRNA cultures **(Figure 4A)**. Knockdown of $iPLA_2$ via shRNA-mediated viral transduction was confirmed by Western blot analysis **(Figure 4B)**.

As cultured astrocytes express mGluR5 (Balazs et al., 1997; Silva et al., 1999; Gebremedhin et al., 2003), we next determined whether the agonists tACPD (100 µM) and AMPA (100 µM) evoked PGE_2 release in the presence/absence of CPA. In the absence of CPA, these agents evoked little to no release of PGE_2 **(Figure 4C)**. However, when CPA was used to block the

release of Ca^{2+} from internal stores, tACPD and AMPA significantly increased PGE_2 release **(Figure 4C)**.

The most abundant mGluRRs expressed by astrocytes are mGluR5 and mGluR3 (Petralia et al., 1996; Aronica et al., 2000; Tamaru et al., 2001). However, mGluR5 is developmentally regulated and is not expressed by astrocytes in the adult brain, whereas mGluR3 is persistently expressed at high levels throughout adulthood (Sun et al., 2013). Determining whether the activation of mGluR3 can induce Ca^{2+}-independent PGE_2 release is important because the activation of this receptor has recently been shown to induce Ca^{2+} transients in adult hippocampal astrocytes (Haustein et al., 2014; Tang et al., 2015). In the presence of CPA, the mGluR3 agonists NAAG and LY379268 (Wroblewska et al., 1997; Bond et al., 1999) significantly increased the release of PGE_2, whereas the same agonists failed to induce the release of PGE_2 in the absence of CPA **(Figure 4D)**. Notably, this result shows that activation of an astrocytic Gi-coupled receptor can induce the release of gliotransmitters via a Ca^{2+}-independent mechanism.

To further investigate human astrocytes, we performed the same experiments in primary cultured astrocytes harvested from human embryonic tissue (Windrem et al., 2004; Windrem et al., 2008; Han et al., 2013). After pharmacologically assessing $iPLA_2$ activity in rat astrocytes **(Figure 2D)**, we evaluated $iPLA_2$ expression in all culture models. Western blot analysis revealed that $iPLA_2$ expression was not limited to rat astrocytes but iPLA2 was also expressed in human and mouse astrocytes **(Figure 4E)**. In the presence of CPA, ATP evoked significant PGE_2 release from cultured human astrocytes, whereas little to no PGE2 release was observed with ATP alone **(Figure 4F)**. Similarly, coapplication of tACPD and AMPA had the same effect **(Figure 4F)**. Taken together, these findings show that ATP or mGluR3 activation leads to Ca^{2+}-independent PGE_2 release and that $iPLA_2$ is expressed in mouse, rat, and human astrocyte cultures.

Ca^{2+}-independent astrocytic lipid release enhances mEPSPs via Kv channel blockade

Thus far, our experiments demonstrated that in response to agonist exposure, cultured astrocytes can indeed release lipids via a Ca2+-independent pathway.- In fact, the in vitro analysis showed that agonist-induced Ca^{2+} increases impeded PGE_2 release, whereas CPA, BAPTA, and CMZ pretreatment potentiated PGE_2 release.

Earlier studies have shown that AA and its metabolite PGE_2 inhibit neuronal Kv currents (Horimoto et al., 1997; Nicol et al., 1997; Evans et al., 1999) and thereby enhance excitability (Sekiyama et al., 1995; Chen and Bazan, 2005; Sang et al., 2005). We therefore evaluated whether astrocytic lipid release also inhibits neuronal Kv currents and enhances neuronal excitability in acute hippocampal slices. We performed dual patch-clamp recordings of pairs of CA1 pyramidal neurons and astrocytes in acute hippocampal brain slices prepared from 12–18-day-old mice **(Figure 5A)**. To isolate transiently active potassium currents, we used 100 ms voltage ramps, which are sufficient to capture transient A-type currents (Phillips et al., 2018). Notably, the I–V ramp enables one to discern what voltage deflection in the outward current a drug affects (Jackson and Bean, 2007). Kv currents in CA1 neurons are used as an assay for astrocytic lipid release as these currents are sensitive to AA and/or its metabolites (Villarroel and Schwarz, 1996; Carta et al., 2014). We isolated the Kv currents by adding 1 mM QX314 to the patch pipette to block sodium channels (Talbot and Sayer, 1996; Kim et al., 2010) and by imposing a voltage ramp (from -100 mV to 50 mV) every 5 seconds to continuously monitor changes in the Kv current (Ji et al., 2000; Rangroo Thrane et al., 2013b; Carta et al., 2014) to assess agonist-induced astrocytic lipid release (Figure 4A). As expected, direct puffing of PGE_2 (50 µM) or AA (50 µM) from a micropipette significantly reduced the Kv current, which is consistent with the results of previous studies (Evans et al., 1999; Carta et al., 2014) **(Figure 5B-C)**.

To assess the effects of astrocytic Ca^{2+} signaling on neighboring neuronal Kv, we stimulated astrocytes with ATP or $TFLLR-NH_2$, an agonist of protease-activated receptor-1 (PAR1), which is primarily expressed by astrocytes (Shigetomi et al., 2008). ATP (100 µM) induced a comparable change in neuronal Kv when BAPTA (50 µM) was present in the astrocyte patch pipette but not

in its absence **(Figure 5B-C).** A similar transient decrease in the Kv current was observed when astrocytes were activated by TFLLR-NH$_2$ (30 µM). Again, a reduction in neuronal Kv currents was detected only when TFLLR was applied and when BAPTA was added to the astrocytic solution **(Figure 5B-C).** To selectively activate astrocytic Ca^{2+} signaling, we next generated acute hippocampal slices from MrgA1 transgenic animals (MrgA1$^{+/-}$), which express the exogenous Gq-coupled MRG receptor (MrgA1) under the control of the GFAP promoter. The MrgA1 agonist Phe-Met-Arg-Phe amide (FMRF) mobilizes intracellular astrocytic Ca^{2+} stores, thus allowing use to assess the effects of astrocytic Ca^{2+} signaling on neighboring neuronal Kv currents. Although FMRF (15 µM) induces potent and selective increases in astrocytic cytosolic Ca^{2+} in hippocampal slices (Fiacco et al., 2007; Agulhon et al., 2010; Wang et al., 2012), we observed no detectable changes in neuronal Kv current when the pipette used to patch astrocytes did not contain BAPTA **(Figure 5B-C)**. However, when BAPTA was present in the astrocyte patch pipette, there was a marked decrease in the neuronal Kv current evoked by FMRF exposure **(Figure 5B-C)**. Thus, astrocytes can modulate neuronal Kv currents via a previously undocumented Ca^{2+}-independent lipid release mechanism.

To assess whether a decrease in the neuronal Kv current is a consequence of Ca^{2+}-independent astrocytic lipid release, we employed specific lipid receptor antagonists for PGE$_2$ and endocannabinoids. As PGE$_2$ receptors 1, 2, 3, and 4 are expressed on hippocampal pyramidal neurons (Andreasson, 2010; Maingret et al., 2017), we used AH6809, a PGE$_2$ EP1, 2, and 3 antagonist (Abramovitz et al., 2000; Ganesh, 2014), and GW627368, a PGE$_2$ EP4 antagonist (Jones and Chan, 2005; Wilson et al., 2006). With AH6809 (10 µM) and GW627368X (3 µM) in perfusion solution, the TFLLR-, FMRF-, and ATP-induced Ca^{2+}-independent decrease in the neuronal Kv current caused by PGE2 was abolished **(Figure 5C)**. In contrast, the CB1 antagonist AM251 (5 µM) failed to abolish the decrease in Kv current **(Figure 5C)**. Taken together, these

data suggest that the observed decrease in the neuronal Kv current is a result of astrocytic Ca^{2+}-independent release of PGE_2.

Interestingly, the presence or absence of BAPTA in the astrocytic pipette solution affected agonist-induced changes in neuronal membrane potential **(Figure 6A)**. Without BAPTA in astrocyte pipettes, TFLLR induced hyperpolarization (-2.4 ± 0.26 mV) **(Figure 6A)**, which has been linked to a decrease in extracellular potassium (Wang et al., 2012). However, with BAPTA present in astrocyte pipettes, TFLLR induced depolarization (2.1 ± 0.25 mV) **(Figure 6A)**, an effect that could be attributed to blockage of the potassium current.

PGE_2 has been shown to enhance neuronal mEPSPs (Sekiyama et al., 1995) (Sang et al., 2005). We found that TFLLR-induced activation of astrocytes triggered a decrease in mEPSP amplitude and frequency in acute hippocampal slices from normal mice **(Figure 6B)**. The opposite effect, an increase in the amplitude and frequency of mEPSCs, was observable only when BAPTA was added to the astrocytic pipette solution **(Figure 6C)**. This observation supports the notion that astrocytic Ca^{2+}-independent lipid release may function as a signaling mechanism that can modulate proximal synaptic activity by blocking Kv, which in turn may increase the frequency and amplitude of mEPSCs. Interestingly, the opposing effects of TFLLR on mEPSP amplitude and frequency were dependent upon astrocytic Ca^{2+} levels (**Figure 6D-E**), suggesting that Ca^{2+} depletion provides a pathway that enhances neuronal excitability by astrocytic PGE2.

DISCUSSION

Lipidomics has the potential to open exciting new avenues within the field of gliotransmission. In the present study, we utilized a series of lipidomic methodologies to show that Ca^{2+} chelation followed by metabotropic glutamate or purinergic receptor stimulation promoted the formation of a variety of lipids (**Figures 2, 3**) in astrocytes and drove the release of PGE_2 in rat and human embryonic astrocyte cultures (**Figure 4**). In addition, we demonstrated that receptor-mediated

Ca^{2+}-independent PGE_2 release modulates neuronal Kv, resulting in enhanced synaptic activity in slices, as was determined by the increase in mEPSP amplitude and frequency **(Figures 5, 6)**. Multiple findings presented herein also support our hypothesis that this agonist-induced and Ca^{2+}-independent lipid release from astrocytes depends on the activation of $iPLA_2$. These observations represent, to our knowledge, the first evidence of Ca^{2+}-independent release of lipid gliotransmitters and thus add a novel dimension to our understanding of glial–neuronal communication.

Importance of the $iPLA_2$ pathway in Ca^{2+}-independent astrocytic lipid production

Although astrocytes are known to constitutively release AA and other fatty acids (Moore, 2001; Bouyakdan et al., 2015), the pathways involved in constitutive AA release are currently not known; however, inverse agonists of purinergic receptors (Ding et al., 2006) and mGluR (Carroll et al., 2001) might serve as useful pharmacological tools for investigating constitutive PLA_2 activity in relation to gliotransmission. To date, many studies have shown the multifaceted functions of PLA_2 in the CNS, but $iPLA_2$ accounts for 70% of the PLA_2 activity in the rat brain (Yang et al., 1999). Although the effects of $cPLA_2$ activation are well documented (Malaplate-Armand et al., 2006; Schaeffer and Gattaz, 2007; Kim et al., 2008), $iPLA_2$ has also been shown to participate in phospholipid remodeling (Sun et al., 2004) and regulate hippocampal AMPA receptors involved in learning and memory (Menard et al., 2005). Furthermore, $iPLA_2$ regulates store-operated Ca^{2+} entry in cerebellar astrocytes (Singaravelu et al., 2006) and provided neuroprotection in an oxygen–glucose deprivation model (Strokin et al., 2006). Astrocytes express both isoforms of PLA_2 (Sun et al., 2005), and numerous studies have reported that the release of AA and its metabolites is under the regulation of Ca^{2+}-dependent PLA_2 ($cPLA_2$) signaling in astrocytes (Bruner and Murphy, 1990; Stella et al., 1994; Stella et al., 1997; Chen and Chen, 1998). Previous studies have demonstrated astrocytic release of DHA via iPLA2 (Strokin et al., 2003, 2007). Here, we focused on determining whether the largely unexplored $iPLA_2$ lipid pathway mediates PGE2 release from astrocytes. Given the ubiquitous expression of $iPLA_2$ in astrocytes throughout the

brain, we contend that various receptor-activated pathways rely on iPLA$_2$ for the regulation of lipid release from astrocytes **(Figure 7)**.

Ca^{2+}-independent astrocytic PGE$_2$ release rapidly modulates neuronal Kv

An important aspect of astrocytic gliotransmitter release is timing. Agonist-induced increases in astrocytic Ca^{2+} occur on a slow time scale of seconds (Cornell-Bell et al., 1990; Wang et al., 2006; Srinivasan et al., 2015), which allows for the integration of many factors on that time scale. Increased intracellular Ca^{2+} levels are key in the release of gliotransmitters (Parpura et al., 1994; Bezzi et al., 1998; Kang et al., 1998), such as ATP (Coco et al., 2003; Parpura and Zorec, 2010; Illes et al., 2019) and D-serine (Mothet et al., 2000; Yang et al., 2003; Li et al., 2018; Neame et al., 2019). We have recently shown that agonist-induced astrocytic Ca^{2+} signaling can modulate synaptic activity by promoting K$^+$ uptake, resulting in a transient decrease in extracellular K$^+$ and a decrease in synaptic activity (Wang et al., 2012). More recently, we demonstrated that astrocytes can modulate rapid synaptic activity ($\geq$ 500 ms) via 2AG release upon agonism of mGluR3 receptors (Smith et al., 2019). Because receptor-mediated astrocytic Ca^{2+} drives K$^+$ uptake and the release of gliotransmitters, these processes must occur over a relatively prolonged time course (>500 ms). However, agonist-induced Ca^{2+}-independent iPLA$_2$ lipid release does not require the mobilization of intracellular Ca^{2+} stores; thus, a much faster time scale for signaling, within 10 seconds to milliseconds, is possible. We speculate that iPLA$_2$-mediated lipid release may act as a feedback system that enhances fast synaptic transmission in the short term or under conditions of minimal neuronal activity, whereas subsequent activity-mediated increases in Ca^{2+} serve as a brake in the form of calmodulin-dependent inhibition of iPLA$_2$. The slower Ca^{2+}-dependent release of gliotransmitters and stimulation of K$^+$ uptake seems more suited toward the slow and widespread modulation of brain activity (typically inhibition) that occurs in the setting of, for example, tonic activation of the locus coeruleus and consequent norepinephrine release (Bekar et al., 2008; Ding et al., 2013).

Voltage-gated potassium channels are located on the dendrites of hippocampal pyramidal neurons (Johnston et al., 2000), where they play a major role in controlling dendritic excitability by modulating the amplitude of EPSPs. A morphological study revealed that the density of Kv in the dendrites of pyramidal neurons increased 5-fold from the soma to the most distal point in apical dendrites (Hoffman et al., 1997). Inhibition of voltage-gated K^+ currents consequently enhances EPSPs, possibly explaining why PGE_2 enhances synaptic transmission and increases LTP in the hippocampus (Sang et al., 2005).

Lipidomics models as a guide for future gliotransmitter discoveries

Lipidomics is one of the fastest-growing branches in the field of metabolomics and offers the possibility of describing the enormous diversity of lipid species throughout the body, especially in the brain, which is largely composed of lipids (Sinclair, 1975). The complex and fine structure of astrocytic processes gives astrocytes a larger surface area-to-volume ratio than most other types of cells, which makes them uniquely responsive to changes in the extracellular milieu. HPLC/MS/MS lipidomics techniques have provided a means to quantify specific lipid species with an accuracy and sensitivity that is not possible with more traditional methods employing radiolabeled fatty acids and ELISA. Here, we combined these classical techniques with lipidomics to test the hypothesis that lipids are produced and released from astrocytes in a Ca^{2+}-independent manner. Although our present focus was on the release of PGE_2, we found increases in the release of 15 additional lipids. Notably, among the 30 lipids measured in the targeted HPLC/MS/MS assay, the levels of 14 did not change upon the activation of glial receptors, supporting of the idea that GPCR-stimulated, Ca^{2+}-independent lipid release is specific. Interestingly, two of the lipids released in response to agonist exposure, namely, *N*-arachidonyl taurine and *N*-palmitoyl tyrosine, were recently shown to activate TRPV4, a cation channel involved in osmotic sensitivity and mechanosensitivity (Raboune et al., 2014). Emerging evidence suggests that the TRPV4/AQP4 complex regulates the response to hypo-osmotic stress in

astrocytes (Benfenati et al., 2011). The lipidomics data presented here thus elucidated unpredicted signaling mechanisms that involve astrocytic-derived lipid modulators **(Figure 3)**.

Potential physiological relevance of calcium-independent lipid release

Astrocytes are known to respond to single experimental stimulation events with small calcium increases in their distal fine processes (Panatier et al., 2011). This effect results from the synchronous firing of as many as 50-1000 synapses in the astrocyte microdomain, as may occur when a large stimulation is applied as far as 500 μm away. However, the spontaneous activity of asynchronous synaptic events does not need to elicit calcium events in astrocytes. Metabotropic GluR- and AMPA-mediated activation of astrocytes may elicit iPLA$_2$ activation and the concerted release of AA metabolites and PGE$_2$ to affect pre- and postsynaptic potassium channels on nearby neurons. When the stimulation frequency and/or amplitude increase, increases in astrocytic calcium then serve to suppress the overactivation of synapses. In this way, astrocytes may help maintain the strength of relatively quiescent synapses in a calcium-independent fashion while employing multiple calcium-dependent mechanisms to suppress overactivation. Substantiation of this model will require rigorous testing of the spatial and temporal dynamics of astrocytic and individual dendritic activation but is limited by current methodological capabilities that do not allow for electrical recording of individual dendritic spines.

In conclusion, Ca^{2+}-independent astrocytic lipid release constitutes a largely unexplored factor in the regulation of complex neuro-glial signaling interactions. Our present analysis contributes to the understanding of agonist-induced Ca^{2+} signaling by demonstrating that the agonism of several Gq and Gi-linked astrocytic receptors can promote the release of lipid modulators and that increases in cytosolic Ca^{2+} act as a brake to prevent PGE$_2$ release.

Author contributions F.W., H.B.B., and N.A.S. performed the experiments. F.W., H.B.B., J.X., and N.A.S. analyzed the data. S.P. and B.J. performed the Western blotting. S.G. and B.L. made

the cultures. L.B. and N.A.S. planned the experiments. D.C.M. provided the human astrocytes. F.W., H.B.B., and N.A.S. wrote the paper.

Acknowledgments We thank Dr. Ken McCarthy for generously sharing the transgenic mice. This work was supported by the National Institutes of Health Grants K01NS110981 and NSFNCS-FR 1926781 and the Department of Defense Army Research Office Award W91NF2020189 to N.A.S. We thank Vittorio Gallo, Baljit Khakh, Bartosz Kula, Stefano Vicini, Alexander S. Thrane, Vinita Rangroo Thrane, Takahiro Takano, Paul Cumming, and Fernando R. Fernandez for comments and critical discussion on this manuscript. The authors declare no competing financial interests.

Figure legends

Figure 1: GPCR-evoked Ca^{2+} transients in astrocytic cultures.

A.) Cultured astrocytes were transduced with the AAV GFAP cyto-GCaMP6f, and the changes in fluorescence associated with increases in intracellular Ca^{2+} upon ATP stimulation were measured using confocal microscopy. B.) Representative individual traces of changes in GCaMP6f fluorescence ($\Delta F/F_0$) in response to ATP stimulation (upper panel) are shown. C.) BAPTA-AM (20 µM, n=5 wells) and CPA (20 µM, n=5 wells) eliminated the ATP (100 µM, n=5 wells)-induced increase in Ca2+ concentration. MAFP (10 µM, n=5 wells) and Bel (10 µM, n=5 wells) did not inhibit ATP (100 µM)-induced increases in Ca^{2+} in astrocytes. ***P≤0.0001, Tukey's *post hoc* test. All bar graphs show the mean $\pm$ s.e.m.

Figure 2: GPCR-mediated Ca^{2+}-independent release of ^{3}H-AA and/or its metabolites from astrocytic cultures.

A.) The schematic shows the ^{3}H-AA Radioactive Assay. B.) The effects of ATP (100 µM, n=45 wells), CPA (20 µM, n=9 wells), ATP/CPA (n=44 wells), and ATP/BAPTA-AM (n=24 wells) on Ca^{2+}-independent release of [^{3}H]-AA and its derivatives in astrocytic cultures were compared to those of the control and each other. ***P≤0.0001; **P≤0.001; *P≤0.05, Tukey's *post hoc* test. C.) The effects of the mGluR agonists tACPD/AMPA (100 µM, n=7), tACPD/AMPA/CPA (n=7 wells), and ATP/CPA (n=7 wells) on the Ca^{2+}-independent release of [^{3}H]-AA and its derivatives in astrocytic cultures were compared to those of the control and individual effects. ***P≤0.0001; **P≤0.001; *P≤0.05, Tukey's *post hoc* test. D.) The effects of the iPLA$_2$ inhibitor BEL (10 µM, n=12 wells), the cPLA$_2$ inhibitor MAFP (10 µM, n=12 wells), or the calmodulin/Ca^{2+} complex inhibitor CMZ (2 µM, n=20 wells) with ATP on Ca^{2+}-independent release of [^{3}H]-AA and its derivatives in astrocytic cultures were compared to ATP alone. ***P≤0.0001; **P≤0.001, Tukey's *post hoc* test. All bar graphs show the mean $\pm$ s.e.m.

Figure 3. Targeted lipidomics of rat astrocytic cultures using HPLC/MS/MS.

A.) The significant differences and the magnitude of change between the vehicle controls and the

CPA/ATP-stimulated cells for the 30 lipids that were detected across samples are summarized.

Dark green denotes significant differences ($p \leq 0.05$); light green denotes increasing trends (0.10

$\geq p \leq 0.05$). No significant difference is indicated by "*n.d.*". Arrows indicate the magnitude of

effects, as indicated in the legend. *The raw data and statistical analyses for all lipids are shown

in the figure table.* B.) An example chromatograph of a PGE2 standard (1 pmol) analyzed by the

HPLC/MS/MS method that was optimized for this standard in negative ion mode is shown. C.) An

example of an analysis of a partially purified astrocytic extract using the HPLC/MS/MS method

that was optimized for PGE2 is shown. D.) Differences in the astrocytic production of AA and two

of its derivatives, PGE2 and docosahexaenoyl ethanolamine (DEA), were expressed as the

average moles/sample; vehicle, n=3 flasks; CPA/ATP, n=4 flasks.

Figure 4: GPCR-mediated Ca^{2+}-independent release of PGE_2 from astrocytic cultures.

A.) The effects of ATP (100 µM, n=4 wells) and ATP/CPA (n=4 wells) on Ca^{2+}-independent PGE_2

release in rat astrocytic cultures virally transduced with shRNA to knockdown $iPLA_2$ was

compared to that in control shRNA cultures. *P≤0.05, Dunn posttest. B.) Western blot analysis of

$iPLA_2$ knockdown via shRNA was performed. β-actin-specific antibodies were used for the

normalization of protein loading. C.) The effects of the GPCR agonists ATP (100 µM, n=12 wells),

tACPD/AMPA (100 µM, n=7 wells), tACPD/AMPA/CPA (n=7), and ATP/CPA (n=12 wells) on Ca^{2+}-

independent PGE_2 release in rat astrocytic cultures was compared to that in control cultures.

****P≤0.0001; ***P≤0.001, Tukey's post hoc test. D. The effects of ATP/CPA (100 µM, n=4) and

the mGluR3 agonists NAAG (100 µM, n=4 wells), LY379268 (100 µM, n=4 wells), NAAG/CPA

(n=4), and LY379268/CPA (n=4 wells) on Ca^{2+}-independent PGE_2 release in rat astrocytic cultures

were compared to that in control cultures. ***P≤0.001, Tukey's post hoc test. E.) Western blot

analysis of the $iPLA_2$ protein in human, rat, and mouse cultures was performed. β-actin-specific

antibodies were used for the normalization of protein loading. F. The effects of the GPCR agonists

ATP (100 µM, n=8 wells), ATP/CPA (n=8 wells), tACPD/AMPA (100 µM, n=8 wells), and

tACPD/AMPA/CPA (n=8 wells) on Ca^{2+}-independent PGE_2 release in human astrocytic cultures were compared to that in control cultures. ***P≤0.0001; **P≤0.001; *P≤0.05, Tukey's *post hoc* test. All bar graphs show the mean ± s.e.m.

Figure 5: Ca^{2+}-independent astrocytic lipid release inhibits Kv blockade.

A.) of the image shows a patched astrocyte and a patched neuron filled with fluorescent dyes. B.) Direct application of AA (50 µM) and PGE2 (50 µM) decreased the Kv current. ATP (100 µM) and TFLLR (30 µM) together with BAPTA also decreased the Kv current, as indicated by the peak currents. FMRF (15 µM) decreased the Kv current in $MrgA1^{+/-}$ mice in the presence of BAPTA but not in the absence of BAPTA. Traces to the left are representative recordings of changes in membrane voltages in the presence of the previously described drugs. C.) The effects of different drugs on the voltage ramp-induced voltage-gated K^+ currents was quantified (**P<0.01; *P<0.05; n =5-7 mice). All bar graphs show the mean ± s.e.m.

Figure 6: Ca^{2+}-independent astrocytic lipid release increases the amplitude and frequency of mEPSP.

A.) The representative traces show changes in membrane potential (VM) upon TFLLR application in the presence or absence of BAPTA, and quantitative values are shown in the inset (**P<0.01; t test; n =5-6 mice). B.) The representative trace show hyperpolarization and the reduction in mEPSP amplitude and frequency in excitatory hippocampal neurons upon TFLLR (30 µM) application without BAPTA in neighboring astrocytes. C.) The representative trace shows depolarization and an increase in mEPSP amplitude and frequency in excitatory hippocampal neurons upon TFLLR (30 µM) application with BAPTA present in neighboring astrocytes. The lower traces are extensions of parts of the recordings in the upper traces. D.) The cumulative mEPSP amplitude and cumulative mEPSP frequency distribution recorded in B) were compared. E.) Statistical analysis of the amplitude and interevent intervals before and after TFLLR treatment

in the presence and absence of BAPTA was performed (**P<0.001; paired t test; n =5-7 mice). All bar graphs show the mean $\pm$ s.e.m.

Figure 7: Models of GPCR-mediated Ca^{2+}-independent and Ca^{2+}-dependent lipid release in astrocytes.

A.) The schematic depicts receptor-mediated Ca^{2+}-independent lipid release across the astrocytic membrane. B.) The schematic depicts receptor-mediated Ca^{2+}-dependent lipid release across the astrocytic membrane.

REFERENCES

Abramovitz M, Adam M, Boie Y, Carriere M, Denis D, Godbout C, Lamontagne S, Rochette C, Sawyer N, Tremblay NM, Belley M, Gallant M, Dufresne C, Gareau Y, Ruel R, Juteau H, Labelle M, Ouimet N, Metters KM (2000) The utilization of recombinant prostanoid receptors to determine the affinities and selectivities of prostaglandins and related analogs. Biochim Biophys Acta 1483:285-293.

Agulhon C, Fiacco TA, McCarthy KD (2010) Hippocampal short- and long-term plasticity are not modulated by astrocyte Ca2+ signaling. Science 327:1250-1254.

Akiba S, Hayama M, Sato T (1998) Inhibition of Ca2+-independent phospholipase A2 by bromoenol lactone attenuates prostaglandin generation induced by interleukin-1 beta and dibutyryl cAMP in rat mesangial cells. FEBS Lett 437:225-228.

Andreasson K (2010) Emerging roles of PGE2 receptors in models of neurological disease. Prostaglandins Other Lipid Mediat 91:104-112.

Aronica E, van Vliet EA, Mayboroda OA, Troost D, da Silva FH, Gorter JA (2000) Upregulation of metabotropic glutamate receptor subtype mGluR3 and mGluR5 in reactive astrocytes in a rat model of mesial temporal lobe epilepsy. The European journal of neuroscience 12:2333-2344.

Balazs R, Miller S, Romano C, de Vries A, Chun Y, Cotman CW (1997) Metabotropic glutamate receptor mGluR5 in astrocytes: pharmacological properties and agonist regulation. J Neurochem 69:151-163.

Bekar LK, He W, Nedergaard M (2008) Locus coeruleus alpha-adrenergic-mediated activation of cortical astrocytes in vivo. Cerebral cortex 18:2789-2795.

Benfenati V, Caprini M, Dovizio M, Mylonakou MN, Ferroni S, Ottersen OP, Amiry-Moghaddam M (2011) An aquaporin-4/transient receptor potential vanilloid 4 (AQP4/TRPV4) complex is essential for cell-volume control in astrocytes. Proc Natl Acad Sci U S A 108:2563-2568.

Bezzi P, Carmignoto G, Pasti L, Vesce S, Rossi D, Rizzini BL, Pozzan T, Volterra A (1998) Prostaglandins stimulate calcium-dependent glutamate release in astrocytes. Nature 391:281-285.

Boland LM, Drzewiecki MM (2008) Polyunsaturated fatty acid modulation of voltage-gated ion channels. Cell biochemistry and biophysics 52:59-84.

Bond A, Ragumoorthy N, Monn JA, Hicks CA, Ward MA, Lodge D, O'Neill MJ (1999) LY379268, a potent and selective Group II metabotropic glutamate receptor agonist, is neuroprotective in gerbil global, but not focal, cerebral ischaemia. Neurosci Lett 273:191-194.

Bouyakdan K, Taib B, Budry L, Zhao S, Rodaros D, Neess D, Mandrup S, Faergeman NJ, Alquier T (2015) A novel role for central ACBP/DBI as a regulator of long-chain fatty acid metabolism in astrocytes. J Neurochem 133:253-265.

Bruner G, Murphy S (1990) ATP-evoked arachidonic acid mobilization in astrocytes is via a P2Y-purinergic receptor. J Neurochem 55:1569-1575.

Carroll FY, Stolle A, Beart PM, Voerste A, Brabet I, Mauler F, Joly C, Antonicek H, Bockaert J, Muller T, Pin JP, Prezeau L (2001) BAY36-7620: a potent non-competitive mGlu1 receptor antagonist with inverse agonist activity. Mol Pharmacol 59:965-973.

Carta M, Lanore F, Rebola N, Szabo Z, Da Silva SV, Lourenco J, Verraes A, Nadler A, Schultz C, Blanchet C, Mulle C (2014) Membrane Lipids Tune Synaptic Transmission by Direct Modulation of Presynaptic Potassium Channels. Neuron.

Chen C, Bazan NG (2005) Endogenous PGE2 regulates membrane excitability and synaptic transmission in hippocampal CA1 pyramidal neurons. J Neurophysiol 93:929-941.

Chen WC, Chen CC (1998) ATP-induced arachidonic acid release in cultured astrocytes is mediated by Gi protein coupled P2Y1 and P2Y2 receptors. Glia 22:360-370.

Coco S, Calegari F, Pravettoni E, Pozzi D, Taverna E, Rosa P, Matteoli M, Verderio C (2003) Storage and release of ATP from astrocytes in culture. J Biol Chem 278:1354-1362.

Cordero-Morales JF, Vasquez V (2018) How lipids contribute to ion channel function, a fat perspective on direct and indirect interactions. Curr Opin Struct Biol 51:92-98.

Cornell-Bell AH, Finkbeiner SM, Cooper MS, Smith SJ (1990) Glutamate induces calcium waves in cultured astrocytes: long-range glial signaling. Science 247:470-473.

Czigler A, Toth L, Szarka N, Szilagyi K, Kellermayer Z, Harci A, Vecsernyes M, Ungvari Z, Szolics A, Koller A, Buki A, Toth P (2019) Prostaglandin E2, a postulated mediator of neurovascular coupling, at low concentrations dilates whereas at higher concentrations constricts human cerebral parenchymal arterioles. Prostaglandins Other Lipid Mediat 146:106389.

Dabertrand F, Hannah RM, Pearson JM, Hill-Eubanks DC, Brayden JE, Nelson MT (2013) Prostaglandin E2, a postulated astrocyte-derived neurovascular coupling agent, constricts rather than dilates parenchymal arterioles. J Cereb Blood Flow Metab 33:479-482.

Dhopeshwarkar GA, Subramanian C (1976) Biosynthesis of polyunsaturated fatty acids in the developing brain: I. Metabolic transformations of intracranially administered 1-14C linolenic acid. Lipids 11:67-71.

Ding F, O'Donnell J, Thrane AS, Zeppenfeld D, Kang H, Xie L, Wang F, Nedergaard M (2013) alpha1-Adrenergic receptors mediate coordinated Ca2+ signaling of cortical astrocytes in awake, behaving mice. Cell calcium 54:387-394.

Ding Z, Kim S, Kunapuli SP (2006) Identification of a potent inverse agonist at a constitutively active mutant of human P2Y12 receptor. Mol Pharmacol 69:338-345.

Evans AR, Vasko MR, Nicol GD (1999) The cAMP transduction cascade mediates the PGE2-induced inhibition of potassium currents in rat sensory neurones. J Physiol 516 (Pt 1):163-178.

Fiacco TA, Agulhon C, Taves SR, Petravicz J, Casper KB, Dong X, Chen J, McCarthy KD (2007) Selective stimulation of astrocyte calcium in situ does not affect neuronal excitatory synaptic activity. Neuron 54:611-626.

Ganesh T (2014) Prostanoid receptor EP2 as a therapeutic target. J Med Chem 57:4454-4465.

Gebremedhin D, Yamaura K, Zhang C, Bylund J, Koehler RC, Harder DR (2003) Metabotropic glutamate receptor activation enhances the activities of two types of Ca2+-activated k+ channels in rat hippocampal astrocytes. The Journal of neuroscience : the official journal of the Society for Neuroscience 23:1678-1687.

Gordon GR, Mulligan SJ, MacVicar BA (2007) Astrocyte control of the cerebrovasculature. Glia 55:1214-1221.

Gordon GR, Choi HB, Rungta RL, Ellis-Davies GC, MacVicar BA (2008) Brain metabolism dictates the polarity of astrocyte control over arterioles. Nature 456:745-749.

Gross RW, Ramanadham S, Kruszka KK, Han X, Turk J (1993) Rat and human pancreatic islet cells contain a calcium ion independent phospholipase A2 activity selective for hydrolysis of arachidonate which is stimulated by adenosine triphosphate and is specifically localized to islet beta-cells. Biochemistry 32:327-336.

Han X, Chen M, Wang F, Windrem M, Wang S, Shanz S, Xu Q, Oberheim NA, Bekar L, Betstadt S, Silva AJ, Takano T, Goldman SA, Nedergaard M (2013) Forebrain engraftment by human glial progenitor cells enhances synaptic plasticity and learning in adult mice. Cell Stem Cell 12:342-353.

Haustein MD, Kracun S, Lu XH, Shih T, Jackson-Weaver O, Tong X, Xu J, Yang XW, O'Dell TJ, Marvin JS, Ellisman MH, Bushong EA, Looger LL, Khakh BS (2014) Conditions and constraints for astrocyte calcium signaling in the hippocampal mossy fiber pathway. Neuron 82:413-429.

Hoffman DA, Magee JC, Colbert CM, Johnston D (1997) K+ channel regulation of signal propagation in dendrites of hippocampal pyramidal neurons. Nature 387:869-875.

Horimoto N, Nabekura J, Ogawa T (1997) Arachidonic acid activation of potassium channels in rat visual cortex neurons. Neuroscience 77:661-671.

Illes P, Burnstock G, Tang Y (2019) Astroglia-Derived ATP Modulates CNS Neuronal Circuits. Trends in neurosciences 42:885-898.

Jackson AC, Bean BP (2007) State-dependent enhancement of subthreshold A-type potassium current by 4-aminopyridine in tuberomammillary nucleus neurons. The Journal of neuroscience : the official journal of the Society for Neuroscience 27:10785-10796.

Jelsema CL, Axelrod J (1987) Stimulation of phospholipase A2 activity in bovine rod outer segments by the beta gamma subunits of transducin and its inhibition by the alpha subunit. Proc Natl Acad Sci U S A 84:3623-3627.

Ji J, Salapatek AM, Diamant NE (2000) Inwardly rectifying K(+) channels in esophageal smooth muscle. Am J Physiol Gastrointest Liver Physiol 279:G951-960.

Johnston D, Hoffman DA, Magee JC, Poolos NP, Watanabe S, Colbert CM, Migliore M (2000) Dendritic potassium channels in hippocampal pyramidal neurons. J Physiol 525 Pt 1:75-81.

Jones RL, Chan KM (2005) Investigation of the agonist activity of prostacyclin analogues on prostanoid EP4 receptors using GW 627368 and taprostene: evidence for species differences. Prostaglandins Leukot Essent Fatty Acids 72:289-299.

Kang J, Jiang L, Goldman SA, Nedergaard M (1998) Astrocyte-mediated potentiation of inhibitory synaptic transmission. Nature neuroscience 1:683-692.

Kim DH, Fitzsimmons B, Hefferan MP, Svensson CI, Wancewicz E, Monia BP, Hung G, Butler M, Marsala M, Hua XY, Yaksh TL (2008) Inhibition of spinal cytosolic phospholipase A(2) expression by an antisense oligonucleotide attenuates tissue injury-induced hyperalgesia. Neuroscience 154:1077-1087.

Kim JH, Kushmerick C, von Gersdorff H (2010) Presynaptic resurgent Na+ currents sculpt the action potential waveform and increase firing reliability at a CNS nerve terminal. The Journal of neuroscience : the official journal of the Society for Neuroscience 30:15479-15490.

Ledo A, Rocha BS, Laranjinha J (2019) Bioactive Lipids and the Gut-Brain Axis: Diet as a Modulator of Bioactivity and Diversity of Lipids in the Brain. Adv Exp Med Biol 1127:147-168.

Lehman JJ, Brown KA, Ramanadham S, Turk J, Gross RW (1993) Arachidonic acid release from aortic smooth muscle cells induced by [Arg8]vasopressin is largely mediated by calcium-independent phospholipase A2. J Biol Chem 268:20713-20716.

Leishman E, Manchanda M, Thelen R, Miller S, Mackie K, Bradshaw HB (2018) Cannabidiol's Upregulation of N-acyl Ethanolamines in the Central Nervous System Requires N-acyl

Phosphatidyl Ethanolamine-Specific Phospholipase D. Cannabis Cannabinoid Res 3:228-241.

Li S, Uno Y, Rudolph U, Cobb J, Liu J, Anderson T, Levy D, Balu DT, Coyle JT (2018) Astrocytes in primary cultures express serine racemase, synthesize d-serine and acquire A1 reactive astrocyte features. Biochem Pharmacol 151:245-251.

Lin JH, Weigel H, Cotrina ML, Liu S, Bueno E, Hansen AJ, Hansen TW, Goldman S, Nedergaard M (1998) Gap-junction-mediated propagation and amplification of cell injury. Nat Neurosci 1:494-500.

MacVicar BA, Newman EA (2015) Astrocyte regulation of blood flow in the brain. Cold Spring Harb Perspect Biol 7.

Maingret V, Barthet G, Deforges S, Jiang N, Mulle C, Amedee T (2017) PGE2-EP3 signaling pathway impairs hippocampal presynaptic long-term plasticity in a mouse model of Alzheimer's disease. Neurobiol Aging 50:13-24.

Malaplate-Armand C, Florent-Bechard S, Youssef I, Koziel V, Sponne I, Kriem B, Leininger-Muller B, Olivier JL, Oster T, Pillot T (2006) Soluble oligomers of amyloid-beta peptide induce neuronal apoptosis by activating a cPLA2-dependent sphingomyelinase-ceramide pathway. Neurobiol Dis 23:178-189.

Menard C, Valastro B, Martel MA, Chartier E, Marineau A, Baudry M, Massicotte G (2005) AMPA receptor phosphorylation is selectively regulated by constitutive phospholipase A(2) and 5-lipoxygenase activities. Hippocampus 15:370-380.

Meves H (1994) Modulation of ion channels by arachidonic acid. Prog Neurobiol 43:175-186.

Moore SA (1993) Cerebral endothelium and astrocytes cooperate in supplying docosahexaenoic acid to neurons. Adv Exp Med Biol 331:229-233.

Moore SA (2001) Polyunsaturated fatty acid synthesis and release by brain-derived cells in vitro. J Mol Neurosci 16:195-200; discussion 215-121.

Moore SA, Yoder E, Murphy S, Dutton GR, Spector AA (1991) Astrocytes, not neurons, produce docosahexaenoic acid (22:6 omega-3) and arachidonic acid (20:4 omega-6). J Neurochem 56:518-524.

Mothet J, Parent A, Wolosker H, Jr RB, Linden D, Ferris C, Rogawski M, Synder S (2000) D-serine is an endogenous ligand for the site of N-methyl D-aspartate receptor. Proc Natl Acad Sci U S A 97:4926-4931.

Murayama T, Kajiyama Y, Nomura Y (1990) Histamine-stimulated and GTP-binding proteins-mediated phospholipase A2 activation in rabbit platelets. J Biol Chem 265:4290-4295.

Neame S, Safory H, Radzishevsky I, Touitou A, Marchesani F, Marchetti M, Kellner S, Berlin S, Foltyn VN, Engelender S, Billard JM, Wolosker H (2019) The NMDA receptor activation by d-serine and glycine is controlled by an astrocytic Phgdh-dependent serine shuttle. Proceedings of the National Academy of Sciences of the United States of America 116:20736-20742.

Nedergaard M (1994) Direct signaling from astrocytes to neurons in cultures of mammalian brain cells. Science 263:1768-1771.

Nicol GD, Vasko MR, Evans AR (1997) Prostaglandins suppress an outward potassium current in embryonic rat sensory neurons. J Neurophysiol 77:167-176.

Panatier A, Vallee J, Haber M, Murai KK, Lacaille JC, Robitaille R (2011) Astrocytes are endogenous regulators of basal transmission at central synapses. Cell 146:785-798.

Parpura V, Zorec R (2010) Gliotransmission: Exocytotic release from astrocytes. Brain Res Rev 63:83-92.

Parpura V, Basarsky TA, Liu F, Jeftinija K, Jeftinija S, Haydon PG (1994) Glutamate-mediated astrocyte-neuron signalling. Nature 369:744-747.

Petralia RS, Wang YX, Niedzielski AS, Wenthold RJ (1996) The metabotropic glutamate receptors, mGluR2 and mGluR3, show unique postsynaptic, presynaptic and glial localizations. Neuroscience 71:949-976.

Phillips WS, Del Negro CA, Rekling JC (2018) Dendritic A-Current in Rhythmically Active PreBotzinger Complex Neurons in Organotypic Cultures from Newborn Mice. The Journal of neuroscience : the official journal of the Society for Neuroscience 38:3039-3049.

Piomelli D (1993) Arachidonic acid in cell signaling. Current opinion in cell biology 5:274-280.

Plog BA, Nedergaard M (2018) The Glymphatic System in Central Nervous System Health and Disease: Past, Present, and Future. Annu Rev Pathol 13:379-394.

Portilla D, Shah SV, Lehman PA, Creer MH (1994) Role of cytosolic calcium-independent plasmalogen-selective phospholipase A2 in hypoxic injury to rabbit proximal tubules. The Journal of clinical investigation 93:1609-1615.

Ramakers GM, Storm JF (2002) A postsynaptic transient K(+) current modulated by arachidonic acid regulates synaptic integration and threshold for LTP induction in hippocampal pyramidal cells. Proc Natl Acad Sci U S A 99:10144-10149.

Rangroo Thrane V, Thrane AS, Plog BA, Thiyagarajan M, Iliff JJ, Deane R, Nagelhus EA, Nedergaard M (2013a) Paravascular microcirculation facilitates rapid lipid transport and astrocyte signaling in the brain. Sci Rep 3:2582.

Rangroo Thrane V, Thrane AS, Wang F, Cotrina ML, Smith NA, Chen M, Xu Q, Kang N, Fujita T, Nagelhus EA, Nedergaard M (2013b) Ammonia triggers neuronal disinhibition and seizures by impairing astrocyte potassium buffering. Nature medicine 19:1643-1648.

Rosenegger DG, Tran CH, Wamsteeker Cusulin JI, Gordon GR (2015) Tonic Local Brain Blood Flow Control by Astrocytes Independent of Phasic Neurovascular Coupling. The Journal of neuroscience : the official journal of the Society for Neuroscience 35:13463-13474.

Sang N, Zhang J, Marcheselli V, Bazan NG, Chen C (2005) Postsynaptically synthesized prostaglandin E2 (PGE2) modulates hippocampal synaptic transmission via a presynaptic PGE2 EP2 receptor. J Neurosci 25:9858-9870.

Schaeffer EL, Gattaz WF (2007) Requirement of hippocampal phospholipase A2 activity for long-term memory retrieval in rats. J Neural Transm 114:379-385.

Seegers HC, Gross RW, Boyle WA (2002) Calcium-independent phospholipase A(2)-derived arachidonic acid is essential for endothelium-dependent relaxation by acetylcholine. J Pharmacol Exp Ther 302:918-923.

Sekiyama N, Mizuta S, Hori A, Kobayashi S (1995) Prostaglandin E2 facilitates excitatory synaptic transmission in the nucleus tractus solitarii of rats. Neurosci Lett 188:101-104.

Shigetomi E, Bowser DN, Sofroniew MV, Khakh BS (2008) Two forms of astrocyte calcium excitability have distinct effects on NMDA receptor-mediated slow inward currents in pyramidal neurons. J Neurosci 28:6659-6663.

Silva GA, Theriault E, Mills LR, Pennefather PS, Feeney CJ (1999) Group I and II metabotropic glutamate receptor expression in cultured rat spinal cord astrocytes. Neurosci Lett 263:117-120.

Sinclair AJ (1975) Long-chain polyunsaturated fatty acids in the mammalian brain. The Proceedings of the Nutrition Society 34:287-291.

Singaravelu K, Lohr C, Deitmer JW (2006) Regulation of store-operated calcium entry by calcium-independent phospholipase A2 in rat cerebellar astrocytes. J Neurosci 26:9579-9592.

Smith NA, Bekar LK, Nedergaard M (2019) Astrocytic Endocannabinoids Mediate Hippocampal Transient Heterosynaptic Depression. Neurochem Res.

Smith NA, Kress BT, Lu Y, Chandler-Militello D, Benraiss A, Nedergaard M (2018) Fluorescent Ca(2+) indicators directly inhibit the Na,K-ATPase and disrupt cellular functions. Sci Signal 11.

Srinivasan R, Huang BS, Venugopal S, Johnston AD, Chai H, Zeng H, Golshani P, Khakh BS (2015) Ca(2+) signaling in astrocytes from Ip3r2(-/-) mice in brain slices and during startle responses in vivo. Nat Neurosci 18:708-717.

Stella N, Tence M, Glowinski J, Premont J (1994) Glutamate-evoked release of arachidonic acid from mouse brain astrocytes. J Neurosci 14:568-575.

Stella N, Estelles A, Siciliano J, Tence M, Desagher S, Piomelli D, Glowinski J, Premont J (1997) Interleukin-1 enhances the ATP-evoked release of arachidonic acid from mouse astrocytes. J Neurosci 17:2939-2946.

Strokin M, Sergeeva M, Reiser G (2003) Docosahexaenoic acid and arachidonic acid release in rat brain astrocytes is mediated by two separate isoforms of phospholipase A2 and is differently regulated by cyclic AMP and Ca2+. Br J Pharmacol 139:1014-1022.

Strokin M, Sergeeva M, Reiser G (2007) Prostaglandin synthesis in rat brain astrocytes is under the control of the n-3 docosahexaenoic acid, released by group VIB calcium-independent phospholipase A2. J Neurochem 102:1771-1782.

Strokin M, Chechneva O, Reymann KG, Reiser G (2006) Neuroprotection of rat hippocampal slices exposed to oxygen-glucose deprivation by enrichment with docosahexaenoic acid and by inhibition of hydrolysis of docosahexaenoic acid-containing phospholipids by calcium independent phospholipase A2. Neuroscience 140:547-553.

Sun GY, Xu J, Jensen MD, Simonyi A (2004) Phospholipase A2 in the central nervous system: implications for neurodegenerative diseases. Journal of lipid research 45:205-213.

Sun GY, Xu J, Jensen MD, Yu S, Wood WG, Gonzalez FA, Simonyi A, Sun AY, Weisman GA (2005) Phospholipase A2 in astrocytes: responses to oxidative stress, inflammation, and G protein-coupled receptor agonists. Mol Neurobiol 31:27-41.

Sun W, McConnell E, Pare JF, Xu Q, Chen M, Peng W, Lovatt D, Han X, Smith Y, Nedergaard M (2013) Glutamate-dependent neuroglial calcium signaling differs between young and adult brain. Science 339:197-200.

Takano T, Tian GF, Peng W, Lou N, Libionka W, Han X, Nedergaard M (2006) Astrocyte-mediated control of cerebral blood flow. Nat Neurosci 9:260-267.

Talbot MJ, Sayer RJ (1996) Intracellular QX-314 inhibits calcium currents in hippocampal CA1 pyramidal neurons. Journal of neurophysiology 76:2120-2124.

Tamaru Y, Nomura S, Mizuno N, Shigemoto R (2001) Distribution of metabotropic glutamate receptor mGluR3 in the mouse CNS: differential location relative to pre- and postsynaptic sites. Neuroscience 106:481-503.

Tang W, Szokol K, Jensen V, Enger R, Trivedi CA, Hvalby O, Helm PJ, Looger LL, Sprengel R, Nagelhus EA (2015) Stimulation-evoked Ca2+ signals in astrocytic processes at hippocampal CA3-CA1 synapses of adult mice are modulated by glutamate and ATP. The Journal of neuroscience : the official journal of the Society for Neuroscience 35:3016-3021.

Tay HK, Melendez AJ (2004) Fcgamma RI-triggered generation of arachidonic acid and eicosanoids requires iPLA2 but not cPLA2 in human monocytic cells. J Biol Chem 279:22505-22513.

van Tol-Steye H, Lodder JC, Mansvelder HD, Planta RJ, van Heerikhuizen H, Kits KS (1999) Roles of G-protein beta gamma, arachidonic acid, and phosphorylation inconvergent activation of an S-like potassium conductance by dopamine, Ala-Pro-Gly-Trp-NH2, and Phe-Met-Arg-Phe-NH2. J Neurosci 19:3739-3751.

Villarroel A, Schwarz TL (1996) Inhibition of the Kv4 (Shal) family of transient K+ currents by arachidonic acid. The Journal of neuroscience : the official journal of the Society for Neuroscience 16:2522-2532.

Wang F, Smith NA, Xu Q, Fujita T, Baba A, Matsuda T, Takano T, Bekar L, Nedergaard M (2012) Astrocytes modulate neural network activity by Ca(2)+-dependent uptake of extracellular K+. Sci Signal 5:ra26.

Wang X, Lou N, Xu Q, Tian GF, Peng WG, Han X, Kang J, Takano T, Nedergaard M (2006) Astrocytic Ca2+ signaling evoked by sensory stimulation in vivo. Nature neuroscience 9:816-823.

Wilson RJ, Giblin GM, Roomans S, Rhodes SA, Cartwright KA, Shield VJ, Brown J, Wise A, Chowdhury J, Pritchard S, Coote J, Noel LS, Kenakin T, Burns-Kurtis CL, Morrison V, Gray DW, Giles H (2006) GW627368X ((N-{2-[4-(4,9-diethoxy-1-oxo-1,3-dihydro-2H-benzo[f]isoindol-2-yl)phenyl]acetyl} benzene sulphonamide): a novel, potent and selective prostanoid EP4 receptor antagonist. Br J Pharmacol 148:326-339.

Windrem MS, Nunes MC, Rashbaum WK, Schwartz TH, Goodman RA, McKhann G, 2nd, Roy NS, Goldman SA (2004) Fetal and adult human oligodendrocyte progenitor cell isolates myelinate the congenitally dysmyelinated brain. Nat Med 10:93-97.

Windrem MS, Schanz SJ, Guo M, Tian GF, Washco V, Stanwood N, Rasband M, Roy NS, Nedergaard M, Havton LA, Wang S, Goldman SA (2008) Neonatal chimerization with human glial progenitor cells can both remyelinate and rescue the otherwise lethally hypomyelinated shiverer mouse. Cell stem cell 2:553-565.

Winstead MV, Balsinde J, Dennis EA (2000) Calcium-independent phospholipase A(2): structure and function. Biochimica et biophysica acta 1488:28-39.

Wolf MJ, Gross RW (1996) The calcium-dependent association and functional coupling of calmodulin with myocardial phospholipase A2. Implications for cardiac cycle-dependent alterations in phospholipolysis. J Biol Chem 271:20989-20992.

Wolf MJ, Wang J, Turk J, Gross RW (1997) Depletion of intracellular calcium stores activates smooth muscle cell calcium-independent phospholipase A2. A novel mechanism underlying arachidonic acid mobilization. J Biol Chem 272:1522-1526.

Wroblewska B, Wroblewski JT, Pshenichkin S, Surin A, Sullivan SE, Neale JH (1997) N-acetylaspartylglutamate selectively activates mGluR3 receptors in transfected cells. J Neurochem 69:174-181.

Yang HC, Mosior M, Ni B, Dennis EA (1999) Regional distribution, ontogeny, purification, and characterization of the Ca2+-independent phospholipase A2 from rat brain. J Neurochem 73:1278-1287.

Yang Y, Ge W, Chen Y, Zhang Z, Shen W, Wu C, Poo M, Duan S (2003) Contribution of astrocytes to hippocampal long-term potentiation through release of D-serine. Proc Natl Acad Sci U S A 100:15194-15199.

Yu X, Taylor AMW, Nagai J, Golshani P, Evans CJ, Coppola G, Khakh BS (2018) Reducing Astrocyte Calcium Signaling In Vivo Alters Striatal Microcircuits and Causes Repetitive Behavior. Neuron 99:1170-1187 e1179.

Zonta M, Angulo MC, Gobbo S, Rosengarten B, Hossmann KA, Pozzan T, Carmignoto G (2003) Neuron-to-astrocyte signaling is central to the dynamic control of brain microcirculation. Nat Neurosci 6:43-50.

Figure 1

Figure 2

A

Relative differences of lipid levels in astrocytes treated with CPA and ATP compared to vehicle

Prostaglandins	
PGE$_2$	↑↑↑
Free fatty acids	
Arachidonic acid	↑
Linoleic acid	*n.d.*
2-arachidonoyl-sn-glycerol	
2-arachidonoyl-*sn*-glycerol	↑
2-linoleoyl-*sn*-glycerol	*n.d.*
2-oleoyl-*sn*-glycerol	*n.d.*
***N*-acyl ethanolamine**	
N-palmitoyl ethanolamine	↑↑
N-stearoyl ethanolamine	↑↑
N-oleoyl ethanolamine	↑↑
N-linoleoyl ethanolamine	*n.d.*
N-arachidonoyl ethanolamine	↑
N-docosahexaenoyl ethanolamine	↑↑
***N*-acyl glycine**	
N-palmitoyl glycine	*n.d.*
N-stearoyl glycine	↑
N-oleoyl glycine	↑
N-linoleoyl glycine	*n.d.*
N-arachidonoyl glycine	*n.d.*
***N*-acyl serine**	
N-palmitoyl serine	*n.d.*
N-stearoyl serine	*n.d.*
N-oleoyl serine	*n.d.*
N-arachidonoyl serine	*n.d.*
***N*-acyl taurine**	
N-arachidonoyl taurine	↑
***N*-acyl tyrosine**	
N-palmitoyl tyrosine	↑
N-stearoyl tyrosine	↑
N-oleoyl tyrosine	↑
N-arachidonoyl tyrosine	*n.d.*
N-docosahexaenoyl tyrosine	↑
***N*-acyl tryptophan**	
N-palmitoyl tryptophan	*n.d.*
N-oleoyl tryptophan	*n.d.*
N-docosahexaenoyl tryptophan	*n.d.*

Magnitude Differences	
2-2.99 times higher	↑↑↑
1.50-1.99 times higher	↑↑
1-1.49 times higher	↑
Significant Differences	
Significant increase p<0.05	
Trending increase p=0.05-0.10	
no difference between groups	*n.d.*

B

10pmol PGE2 standard

HPLC/MS/MS multiple reaction monitoring method scanning for parent ion [-H] 351 with [-H] 315 fragment ions

C

CPA/ATP stimulated astrocytic lipid extract using [-H]351/315 MRM on 40% partial purification fraction

D

Veh   CPA/ATP

Figure 3

Figure 4

A

Alexa 594
Lucifer Yellow
Patched astrocyte and neuron

Neuron

+BAPTA   Astrocytes

+TFLLR

B

BAPTA   AA

10s

BAPTA   PGE2

10s

BAPTA   TFLLR

10s

BAPTA   ATP

10s

FMRF   BAPTA   FMRF

20 s

Kv currents   50 mV

-100 mV

Ctr
AA +

Ctr
PGE2 +

Ctr
TFLLR +
BAPTA

Ctr
ATP+
BAPTA

FMRF
FMRF +
BAPTA

200 pA
10 ms

C

BAPTA

normalized Kv current

Ctr | PGE2 | AA | ATP | TFLLR | FMRF | ATP | ATP+EP blocker | ATP+CB blocker | TFLLR | TFLLR+EP blocker | TFLLR+CB blocker | FMRF | FMRF+EP blocker | FMRF+CB blocker

Figure 5

Figure 6

Ca²⁺-Independent Pathway

A

Ca²⁺-Dependent Pathway

B

Figure 7