Increased fronto-temporal connectivity

by modified melody in real music

Chan Hee Kim[1, 2], Jaeho Seol[2, 3], Seung-Hyun Jin[2], June Sic Kim[2, 4], Youn Kim[5],

Suk Won Yi[6, 7], and Chun Kee Chung[1, 2, 8, 9, *]

[1] Interdisciplinary Program in Neuroscience, Seoul National University College of Natural Science, Seoul 08826, Korea

[2] Human Brain Function Laboratory, Seoul National University, Seoul 03080, Korea

[3] W-Mind Laboratory, Wemakeprice Inc. Seoul 06173, Korea

[4] Research Institute of Basic Sciences, Seoul National University, Seoul 08826, Korea

[5] Department of Music, School of Humanities, The University of Hong Kong, Hong Kong

[6] College of Music, Seoul National University, Seoul 08826, Korea

[7] Western Music Research Institute, Seoul National University, Seoul 08826, Korea

[8] Department of Brain and Cognitive Science, Seoul National University College of Natural Science, Seoul 08826, Korea

[9] Department of Neurosurgery, Seoul National University Hospital, Seoul 03080, Korea

* Corresponding author: chungc@snu.ac.kr

Abstract

In real music, the original melody may appear intact, with little elaboration only, or significantly modified. Since a melody is most easily perceived in music, hearing significantly modified melody may change a brain connectivity. Mozart KV 265 is comprised of a theme with an original melody of "Twinkle Twinkle Little Star" and its significant variations. We studied whether effective connectivity changes with significantly modified melody, between bilateral inferior frontal gyri (IFGs) and Heschl's gyri (HGs) using magnetoencephalography (MEG). Among the 12 connectivities, the connectivity from the left IFG to the right HG was consistently increased with significantly modified melody compared to the original melody in 2 separate sets of the same rhythmic pattern with different melody ($p = 0.005$ and 0.034, Bonferroni corrected). Our findings show that the modification of an original melody in a real music changes the brain connectivity.

Keywords: Melody; Variation; Effective connectivity; Inferior frontal gyrus; Heschl's gyrus; Linearized time delayed mutual information; Magnetoencephalography

Introduction

Melody is a feature effortlessly extracted from music, which sometimes represents the music

itself. Beethoven symphony No. 9 is known as "Ode to joy", since the melody of "Ode to joy"

appears repeatedly across the 4^{th} movement of the symphony, which is easily and

prominently detected. It would be interesting to see how the brain reacts to phrases that

contain prominent information such as a well-known melody while listening to real music, or

phrases that do not. If this is something that can be detected without specific musical

expertise, it would be interesting to see how the non-expert brain will react.

Purpose of our study was to verify the changes in non-expert brain by the musics

including a well-known melody and its modifications using Mozart Variations. Mozart 12

Variations KV 265 on "Ah! vous dirai-je Maman" of a French chanson is well-known as

"Twinkle Twinkle Little Star" Variations, of an English lullaby. In some of the variations, the

original melody appears intact or with little elaboration only. In others, it is significantly

modified. We call the former variations with original melody (VOM) and the latter variations

with modified melody (VMM). The original melody of "Twinkle Twinkle Little Star" in the

Theme is modified or repeated in 12 variations. Five conditions of the *Theme*, *Variation I*,

Variation II, *Variation III*, and *Variation IV* are composed in the same key, harmony, meter,

timbre, and ternary form. Among the five conditions, the original melody is present in the

Variation II and *IV*. However, the original melody is significantly modified in the *Variation I*

and *III*. In terms of rhythmic pattern, the *Variation I* and *II* are of 16^{th} notes, while the

Variation III and *IV* are of 8^{th} notes. Hence, two sets of "*Variation I* vs. *II*" and "*Variation III*

vs. *IV*" are of the same rhythmic patterns, but of different melodies (Fig. 1).

The human brain has several interconnected key regions processing music, including

bilateral inferior frontal gyri (IFGs) and Heschl's gyri (HGs) [1-8]. Each region and

hemispheres are known to have a specific function. The right IFG is dominant in the

processing of musical syntax, compared to the left IFG [1]. The left IFG is activated by

pattern deviations of melodic contours [9], and is involved in the processing of familiarity in

music [10]. The bilateral HGs are involved in pitch and melody process, which is dominant in

right hemisphere [3, 5, 11]. The right HG is associated with the processing of auditory stream

[12], a deviance of tone [13, 14] and spectral pitch [5]. Extracting and interpreting a melody

in music would be accompanied with changes in a regional connectivity. A hemispheric or

interregional connectivity between related regions in bilateral IFGs and HGs would allow to

infer what characteristics the difference between VOM and VMM reflect.

Effective connectivity for 12 connections between the bilateral IFGs and HGs was

calculated by linearized time delayed mutual information (LTDMI) [15]. LTDMI is a measure

of functional coupling estimating information transmission between two time series based on

mutual information (MI). Directional information transmission calculated by time delayed

mutual information (TDMI) detects linear and non-linear correlation between two time series.

Therefore, we hypothesized that directional information transmission between related

regions associated with the processing of intrinsic differences implied in the VOM and VMM

would be reflected in the LTDMI between the bilateral IFGs and HGs. We further

hypothesized that the two sets of "*Variation I* vs. *II*" and "*Variation III* vs. *IV*", with the

original melody and modified melody, would consistently change effective connectivity

between the bilateral IFGs and HGs.

Fig 1. Musical stimuli. Five conditions of the *Theme*, *Variation I*, *II*, *III*, and *IV* selected in

Mozart 12 Variations on "Ah! vous dirai-je Maman"KV 265. The original melody of "C5 -

C5 - G5 - G5 - A5 (Twin- kle twin- kle lit-)" is included in the *Theme*, *Variation II*, and

Variation IV (orange boxes). The *Variation I* and *Variation III* include the modified melodies

(gray boxes). Rhythmic patterns are consistent in each set of "*Variation I* and *Variation II*"

(16th notes) and "*Variation III* and *Variation IV* " (8th notes).

Materials and Methods

Ethics Statement

70   The study was approved by the Institutional Review Board of the Clinical Research Institute,

71   Seoul National University Hospital (IRB No. C-1003-015-311). All experiments were

72   conducted in accordance with the ethical guidelines and regulations.

Participants

75   Twenty-five participants (15 females, mean age, 26.8 ± 3.4 years) were all right-handed

76   (mean Edinburgh Handedness coefficient, 95.7 ± 7.1). All participants had normal hearing,

77   and had not received any formal musical training. Prior to the experiments, all the

78   participants provided the informed consent in a written form.

Musical stimuli

Out of the theme and the 12 Variations in Mozart's Variation on "Ah! vous dirai-je Maman"

KV 265, we selected the five conditions of *Theme*, *Variation I*, *II*, *III*, and *IV* in Fig. 1,

composed in the same key (C major), harmony (tonic-tonic-subdominant-tonic…), meter

(2/4), and structure (ternary form). In magnetoencephalography (MEG) experiment, all

participants were asked to passively listen to the musical stimuli without to paying any

attention to any specific musical feature, such as melody.

Time window

The time window was selected based on the concept of the motif [16]. The ascending five

quarter notes of "C5 - C5 - G5 - G5 - A5" (2,100 ms) of the time window were in the motif of

"C5 - C5 - G5 - G5 - A5 - A5 - G5 - G5". A previous behavioral study reported that

differences in melody were detected by listening to a segment of three to six notes in a motif

of an original melody [17]. Even though it usually takes about 400 ms before the brain

response leads to a motor response after stimulus onset, 5 quarter notes were considered

appropriate time widow. Also we thought it was reasonable to focus on the original melody,

even considering that familiarity and congruence, etc., which might differ between the

conditions; i.e. responses to familiar and congruent conditions were faster [18, 19]. For the

five quarter notes of in the *Theme*, the number of 16th notes or 8th notes (in triplets)

subdivided in accompaniments was the same between the *Variation I*, *II*, *III*, and *IV*.

The present musical stimuli were recorded music, so those were presented without a

break between movements of a theme and variations. To select the proper time window, we

had to consider all the subdivided notes included in a quarter note in each time window in all

conditions. The time window of 2,100 ms was the most suitable duration because (1) 2,100

ms covered the onset of the first quarter note to the offset of the last quarter note in the time

window; (2) the number of additional notes did not exceed a quarter note in all the conditions;

and (3) fade-out times of the final note in the preceding condition did not overlap with the

onset of the time window in the following condition. The same phrases of the time window

were repeated four times in each condition. However, to rule out the effect of repetition [20],

we analyzed only the "C5 - C5 - G5 - G5 - A5" in the opening in each condition.

Data recording

The MEG signal was recorded with a sampling frequency of 1,000 Hz using a 0.1–200 Hz

band pass filter in a magnetically shielded room with a 306-channel whole-head MEG

System (Elekta Neuromag Vector View™, Helsinki, Finland). Electrooculograms (EOG) and

electrocardiograms (ECG) were also simultaneously recorded to remove ocular and cardiac

noise at a later time. We eliminated the environmental magnetic noise of raw MEG signals

with the temporal signal space separation (tSSS) algorithm in MaxFilter 2.1.13 (Elekta

Neuromag Oy, Helsinki, Finland) [21].

During the MEG recording, for about five minutes, all participants sat in a magnetically

shielded room and passively listened to music (Mozart 12 Variations on "Ah! vous dirai-je

Maman" KV 265, 2011, Brilliant classics, Dutch). Musical stimuli were presented binaurally

through plastic tubal silicone earpieces, 50 cm in length using STIM2™ (Neuroscan,

Charlotte, NC, USA) at 100 dB. Visual stimuli of a silent movie clip (Love Actually, 2003,

Universal Pictures, USA) were presented on a screen to keep the participants awake [22, 23]

while the musical stimuli were being presented. For MEG signals, EOG, ECG, muscle

artifact was removed using the independent component analysis. Epochs were determined

from -100 ms to 2,100 ms after the onset of each condition, and the baseline of each epoch

was from -100 ms to 0 ms.

Data analysis

The source coordinates for the ROIs of the bilateral HGs and IFGs were based on the

standard Talairach coordinates; Heschl's gyrus (transverse, BA 41, BA 42) and the inferior

frontal gyrus (triangular part, BA 45); the x, y, and *z in* Talairach coordinates (millimeters)

were -53.5, -30.5, and 12.6 in the left HG, 55.4, -30.5, and 12.6 in the right HG, -55.5, 11.7,

and 20.6 in the left IFG, and 53.5, 12.7and 20.6 in the right IFG, respectively (Fig. 2). In Fig.

2, The Talairach coordinates were visualized using BrainNet Viewer

(http://nitrc.org/projects/bnv/). The signal for each regional source of 4 ROIs (bilateral HGs

and IFGs) was extracted using a 14–30 Hz band-pass filter for each participant using BESA

5.1.8.10 (MEGIS Software GmbH, Gräfelfing, Germany), which was averaged at MATLAB

7.7.0.471 (Math Works Inc., Natick, MA, USA). Beta band rhythm of 14–30 Hz underlies the

top-down control [24], reflects a deviant word processing [25], and is related to the semantic

evaluation for the spoken word [26] and categorical perception [27]. Considering that our

hypothesis is focused on separating VOM and VMM, with the original and modified melody,

we chose the beta band rhythm as the target frequency.

For the twelve connections among the bilateral HGs and IFGs, effective connectivity for

twelve connections was estimated using LTDMI [15]. The MI is a measure testing the

relationship between the conditions by measuring the amount of information between two

random variables. The LTDMI is an information theoretic measure estimating the directional

information transmission of functional coupling based on MI:

$$MI = MI_{\text{AB}} = MI_{\text{BA}} = MI\big(\text{A}(n), \text{B}(n)\big) = -\sum_k p\big(\text{A}(n), \text{B}(n)\big) \log \frac{p(\text{A}(n),(\text{B}(n))}{p(\text{A}(n))(p(\text{B}(n))} \quad (1)$$

In case of $A(n) = B(n)$, the MI is maximum. When two time series are independent, the MI

is zero. The directional information transmission between the two time series estimated based

on TDMI, time delayed mutual information [15]:

$$TDMI_{AB} = TDMI\big(A(n), B(n+\tau)\big) = -\sum_k p(A(n), B(n+\tau)) \log \frac{p(A(n),(B(n+\tau))}{p(A(n))(p(B(n+\tau))}.$$

$$TDMI_{BA} = TDMI\big(B(n), A(n+\tau)\big) = -\sum_k p(B(n), A(n+\tau)) \log \frac{p(B(n),(A(n+\tau))}{p(A(n))(p(A(n+\tau))} \quad (2)$$

TDMI is asymmetric MI obtained by adding a time delay. Both of linear and non-linear

correlation between two time series can be detected by TDMI. The data length of 400 ms

epoch in our study was insufficient in reconstruction of a reliable PDF for general TDMI.

Based on TDMI, the linear correlation for the information transmission between time series A

and B is estimated by LTDMI:

$$LTDMI_{AB} = LTDMI\big(A(n), B(n+\tau)\big) = -\frac{1}{2}\log\big(1 - \rho^2_{A(n)B(n+\tau)}\big)$$

$$LTDMI_{BA} = LTDMI\big(B(n), A(n+\tau)\big) = -\frac{1}{2}\log\big(1 - \rho^2_{B(n)A(n+\tau)}\big). \qquad (3)$$

where τ is delay time, and $\rho_{A(n)B(n+\tau)}$ or $\rho_{B(n)A(n+\tau)}$ is a cross-correlation coefficient. The

LTDMI value was averaged over delay time, which was calculated for every individual

participant for each of the four conditions and twelve connections. See our previous study[3]

for the details of the LTDMI.

The mean LTDMI value for the time window of 2,100 ms in each condition was

calculated for the twelve connections between the four ROIs. Because the data set was not

normally distributed, the differences in LTDMI values of the conditions including "Twinkle

Twinkle Little Star" melody or not were tested by the Wilcoxon signed rank test using SPSS

21.0 software (IBM, Armonk, NY, USA). Multiple comparisons for the twelve connections in

each set were adjusted by the Bonferroni test ($p < 0.05$).

Results

The Wilcoxon signed-rank test revealed that the LTDMI from the left IFG to the right HG

was significantly different between "*Variation I* vs. *Variation II*" (Z = -3.512, p = 0.005,

Bonferroni corrected) as well as between "*Variation III* vs. *Variation IV*" (Z = -2.987, p =

0.034, Bonferroni corrected) (Table 1). In this fronto-temporal connection, the LTDMI values

increased in the VMM (*Variation I* and *III*) compared to the VOM (*Variation II* and *IV*) (Fig.

2).

Fig 2. Difference in effective connectivity from the left IFG to the right HG for the

modified and original melodies. Effective connectivity in both sets of "*Variation I* vs.

Variation II" and "*Variation III* vs. *Variation IV*" showed an identical pattern in the Wilcoxon

signed-rank test. Violin plots show that the LTDMI values are significantly higher for the

Variation I and *III* with the modified melodies than for the *Variation II* and *IV* with the

original melody; "*Variation I* vs. *Variation II*", $Z = -3.512$, $p = 0.005$, Bonferroni corrected;

"*Variation III* vs. *Variation IV*", $Z = -2.987$, $p = 0.034$, Bonferroni corrected (See also Table

1).

Table 1. LTDMI difference between the conditions with the modified or original melodies. For the both sets of "*Variation I* vs. *Variation*

II" and "*Variation III* vs. *Variation IV*", the LTDMI values differed in a connection from the left IFG to the right HG among twelve

connections (Wilcoxon signed-rank test, N = 25, Bonferroni corrected *$p < 0.05$ and **$p < 0.01$).

	Variation I vs. Variation II			Variation III vs. Variation IV		
	Z	*P$_{(uncorrected)}$*	*P$_{(corrected)}$*	*Z*	*P$_{(uncorrected)}$*	*P$_{(corrected)}$*
Left HG → Right HG	-0.565	0.572	1.0	-1.090	0.276	1.0
Left HG → Left IFG	-0.727	0.468	1.0	-1.843	0.065	0.784
Left HG → Right IFG	-0.148	0.882	1.0	-0.229	0.819	1.0
Right HG → Left HG	-1.493	0.135	1.0	-0.175	0.861	1.0
Right HG → Left HG	-1.286	0.199	1.0	-1.029	0.304	1.0
Right HG → Right HG	-1.305	0.192	1.0	-0.148	0.882	1.0
Left IFG → Left HG	-1.117	0.264	1.0	-1.655	0.098	1.0
Left IFG → Right HG	-3.512	0.0004	**0.005 **	-2.987	0.0028	**0.034 *
Left IFG → Right IFG	-0.875	0.382	1.0	-1.238	0.216	1.0
Right IFG → Left HG	-1.129	0.259	1.0	-0.498	0.619	1.0
Right IFG → Right HG	-1.762	0.078	0.936	-0.982	0.326	1.0
Right IFG → Left IFG	-2.193	0.028	0.340	-0.848	0.397	1.0

Discussion

Effective connectivity from the left IFG to the right HG increased when the original melody

of "Twinkle Twinkle Little Star" was significantly modified (VMM), compared with that

when the original melody was intact (VOM). It appeared consistent in two different sets with

the same rhythmic patterns but with the different melody pattern.

The present study employed real music with its variations as musical stimuli, dissected

musical elements in each condition, and devised the two comparable sets of conditions,

which have the same rhythmic pattern but different melody. Here, we exploited naturalistic

conditions in real music instead of devising artificial conditions, and successfully

demonstrated how significant variations of melody in real music change a regional

connectivity in the human brain. These are distinct from the attempts in previous studies used

to original musical pieces [10, 28-31].

An fMRI study, using a real music of "Adios Nonino", studied the effects of motif

repetitions on effective connectivity changes. Specifically they found that the hippocampal

connectivity is modulated by motif repetitions, showing strong connections with working

memory-relevant areas [31]. This study also shows that a real music and characteristics

inherent in it change the brain connectivities. However, unlike this study, our study did not

only focus on the motif of monody within the music. Instead, we identified how the motif

was intertwined with other elements of music through music analysis, and tried to present

results that did not exclude all possible elements within the music.

When hearing real music the brain regions would interact to support complex cognitive

processes [32, 33]. In the perspective of the brain networks, the change in effective

connectivity from the left IFG to the right HG could be interpreted in the syntactic process.

However, modified and original melody detection would be different processes from

syntactic process, although relatively similar cortical regions may be involved [1, 23, 34].

The left IFG is highly activated for the detection of novelty within a musical context based on

long term memory [35].

The left IFG is also involved in the retrieval of working memory [36] and episodic memory

[30], and in the processing of familiarity [10]. Also, the left prefrontal area is activated for the

recognition of pattern deviations in melody [9]. On the other hands, the right HG is dominant

for the processing of stream segregation for auditory stimuli [12], and is highly sensitive for a

deviance of tone [13, 14] and spectral pitch [5], compared to the left hemisphere. The right

superior temporal sulcus is also involved in the processing of familiarity [37]. In this regards,

increased and decreased fronto-temporal connectivity would involve memorizing melody or

streams, feeling the familiarity, recognizing patterns of melody, and detecting tone deviation

or novelty within the VMM with modified melody and the VOM with the original melody.

Also it is interesting that the change in fronto-temporal connectivity was observed in the

beta frequency band (see also the Materials and Methods section). The connectivity changes

were consistent in the two sets of "*Variation I* vs. *II*" and "*Variation III* vs. *IV*", which are

different in the rhythmic patterns of 16^{th} and 8^{th} notes sequences, respectively. Generally the

beta frequency band is associated with the processing of deviant stimuli [25] and

memory/categorical perception [27]. The connectivity in the beta frequency band would

reflect the categorization of the deviant VMM and the VOM based on the information of the

original melody acquired in real-life.

Also, beta frequency is related with beat perception. While regular auditory stimuli were

presented, beta oscillation was synchronized with the beat, while it was increased when the

beat was omitted [38, 39]. Therefore, the changes in fronto-temporal connectivity at the beta

frequency band might reflect omission of the beats on the original melody of quarter notes

sequence in each condition, regardless of changes in rhythmic patterns. It may be an

alternative explanation why the fronto-temporal connectivity increased with modified melody

conditions compared to the original melody.

In our study, the time window of five notes, "C5 - C5 - G5 - G5 - A5" (2.1 sec) was set

based on the "C5 - C5 - G5 - G5 - A5 - A5 - G5 - G5" motif of "Twinkle Twinkle Little Star"

song in the *Theme*. In a previous study, the isolation point, the point at which a melody is

correctly identified, is about 2.5 sec of 5 notes for highly familiar melodies [17]. Therefore,

the five notes in the motif, which we analyzed were long enough to detect the original melody.

Connectivity results in the two sets of conditions, with different rhythms, showed similar

pattern. The hypothesis of our study was to find a connection that connectivity pattern was

consistent for the two sets of "VOM and VMM" of the different rhythmic pattern. Therefore,

the results would be by pitch rather than rhythm. Therefore, it would be natural that rhythmic

affects are excluded from our results. Furthermore, the one that influenced the results could

be eventually be dismissed as "original melody and modified melody" that encompasses all

of familiarity, the stimuli characteristics, and the variation identification.

There are several limitations in our study. The present study was limited to 4 regions of

interest, including bilateral inferior frontal and superior temporal area. There should be

further studies covering the whole brain. For current findings targeting non-musicians,

comparative studies involving musicians are needed. Finally, in order to focus on melody

difference, we tried our best to eliminate the effects of other elements implied in naturalistic

conditions. However, it should be discovered through further studies.

In conclusion, the fronto-temporal connectivity from the left IFG to the right HG was

enhanced when the modified melody was presented, compared with the original melody of

"Twinkle Twinkle Little Star" in Mozart "Ah! vous dirai-je Maman" KV 265. Our findings

show that the modification of an original melody in a real music changes the brain

connectivity.

Acknowledgements

270 We would like to thank Ji Hyang Nam for the technical support in MEG acquisition.

References

1.   Maess B, Koelsch S, Gunter TC, Friederici AD. Musical syntax is processed in

Broca's area: an MEG study. Nature neuroscience. 2001;4(5):540-5. doi: 10.1038/87502.

PubMed PMID: 11319564.

2.   Janata P, Birk JL, Van Horn JD, Leman M, Tillmann B, Bharucha JJ. The cortical

topography of tonal structures underlying Western music. Science. 2002;298(5601):2167-70.

Epub 2002/12/14. doi: 10.1126/science.1076262. PubMed PMID: 12481131.

3.   Patterson RD, Uppenkamp S, Johnsrude IS, Griffiths TD. The processing of temporal

pitch and melody information in auditory cortex. Neuron. 2002;36(4):767-76. Epub

2002/11/21. PubMed PMID: 12441063.

4.   Patel AD, Balaban E. Human pitch perception is reflected in the timing of stimulus-

related cortical activity. Nature neuroscience. 2001;4(8):839-44. doi: 10.1038/90557.

PubMed PMID: 11477431.

5.   Schneider P, Sluming V, Roberts N, Scherg M, Goebel R, Specht HJ, et al. Structural

and functional asymmetry of lateral Heschl's gyrus reflects pitch perception preference.

Nature neuroscience. 2005;8(9):1241-7. Epub 2005/08/24. doi: 10.1038/nn1530. PubMed

PMID: 16116442.

6.   Grahn JA, Rowe JB. Feeling the beat: premotor and striatal interactions in musicians

and nonmusicians during beat perception. The Journal of neuroscience : the official journal of

the   Society   for   Neuroscience.   2009;29(23):7540-8.   Epub   2009/06/12.   doi:

10.1523/JNEUROSCI.2018-08.2009. PubMed PMID: 19515922; PubMed Central PMCID:

PMC2702750.

7.   Seol J, Oh M, Kim JS, Jin SH, Kim SI, Chung CK. Discrimination of timbre in early

auditory responses of the human brain. PloS one. 2011;6(9):e24959. doi:

10.1371/journal.pone.0024959. PubMed PMID: 21949807; PubMed Central PMCID:

PMC3174256.

8.   Kim CH, Kim JS, Choi Y, Kyong JS, Kim Y, Yi SW, et al. Change in left inferior

frontal connectivity with less unexpected harmonic cadence by musical expertise. PloS one.

2019;14(11):e0223283. doi: 10.1371/journal.pone.0223283. PubMed PMID: 31714920.

9.   Habermeyer B, Herdener M, Esposito F, Hilti CC, Klarhofer M, di Salle F, et al.

Neural correlates of pre-attentive processing of pattern deviance in professional musicians.

Human brain mapping. 2009;30(11):3736-47. doi: 10.1002/hbm.20802. PubMed PMID:

19492302.

10.   Plailly J, Tillmann B, Royet JP. The feeling of familiarity of music and odors: the

same neural signature? Cereb Cortex. 2007;17(11):2650-8. doi: 10.1093/cercor/bhl173.

PubMed PMID: 17289777.

11.   Warrier CM, Zatorre RJ. Right temporal cortex is critical for utilization of melodic

contextual cues in a pitch constancy task. Brain : a journal of neurology. 2004;127(Pt

7):1616-25. doi: 10.1093/brain/awh183. PubMed PMID: 15128620.

12.   Snyder JS, Alain C, Picton TW. Effects of attention on neuroelectric correlates of

auditory stream segregation. Journal of cognitive neuroscience. 2006;18(1):1-13. doi:

10.1162/089892906775250021. PubMed PMID: 16417678.

13.   Sabri M, Liebenthal E, Waldron EJ, Medler DA, Binder JR. Attentional modulation

in the detection of irrelevant deviance: a simultaneous ERP/fMRI study. Journal of cognitive

neuroscience. 2006;18(5):689-700. doi: 10.1162/jocn.2006.18.5.689. PubMed PMID:

16768370; PubMed Central PMCID: PMCPMC1769347.

14.   Nan Y, Friederici AD. Differential roles of right temporal cortex and Broca's area in

pitch processing: evidence from music and Mandarin. Human brain mapping.

2013;34(9):2045-54. doi: 10.1002/hbm.22046. PubMed PMID: 22431306.

15.   Jin SH, Lin P, Hallett M. Linear and nonlinear information flow based on time-

delayed mutual information method and its application to corticomuscular interaction.

Clinical neurophysiology : official journal of the International Federation of Clinical

Neurophysiology. 2010;121(3):392-401. doi: 10.1016/j.clinph.2009.09.033. PubMed PMID:

20044309; PubMed Central PMCID: PMCPMC2822094.

16.   Drabkin W. Grove Music Online. Oxford Music Online. 2014 [updated Thur. 18 Sep.

2014.].   Available   from:

http://www.oxfordmusiconline.com/subscriber/article/grove/music/19221.

17.   Bella SD, Peretz I, Aronoff N. Time course of melody recognition: a gating paradigm

study. Perception & psychophysics. 2003;65(7):1019-28. Epub 2003/12/17. PubMed PMID:

14674630.

18.   Tillmann B, Albouy P, Caclin A, Bigand E. Musical familiarity in congenital amusia:

evidence from a gating paradigm. Cortex; a journal devoted to the study of the nervous

system and behavior. 2014;59:84-94. doi: 10.1016/j.cortex.2014.07.012. PubMed PMID:

25151640.

19.   Sammler D, Novembre G, Koelsch S, Keller PE. Syntax in a pianist's hand: ERP

signatures of "embodied" syntax processing in music. Cortex; a journal devoted to the study

of the nervous system and behavior. 2013;49(5):1325-39. Epub 2012/07/27. doi:

10.1016/j.cortex.2012.06.007. PubMed PMID: 22832238.

20.   Guo S, Koelsch S. The effects of supervised learning on event-related potential

correlates of music-syntactic processing. Brain research. 2015;1626:232-46. doi:

10.1016/j.brainres.2015.01.046. PubMed PMID: 25660849.

21.   Taulu S, Hari R. Removal of magnetoencephalographic artifacts with temporal

signal-space separation: demonstration with single-trial auditory-evoked responses. Human

brain mapping. 2009;30(5):1524-34. Epub 2008/07/29. doi: 10.1002/hbm.20627. PubMed

PMID: 18661502.

22.   Koelsch S, Jentschke S. Short-term effects of processing musical syntax: an ERP

study.   Brain   research.   2008;1212:55-62.   Epub   2008/04/29.   doi:

10.1016/j.brainres.2007.10.078. PubMed PMID: 18439987.

23.   Ruiz MH, Koelsch S, Bhattacharya J. Decrease in early right alpha band phase

synchronization and late gamma band oscillations in processing syntax in music. Human

brain mapping. 2009;30(4):1207-25. Epub 2008/06/24. doi: 10.1002/hbm.20584. PubMed

PMID: 18571796.

24.   Fries P. Rhythms for Cognition: Communication through Coherence. Neuron.

2015;88(1):220-35. doi: 10.1016/j.neuron.2015.09.034. PubMed PMID: 26447583; PubMed

Central PMCID: PMCPMC4605134.

25.   Kim JS, Chung CK. Language lateralization using MEG beta frequency

desynchronization during auditory oddball stimulation with one-syllable words. NeuroImage.

2008;42(4):1499-507. doi: 10.1016/j.neuroimage.2008.06.001. PubMed PMID: 18603004.

26.   Shahin AJ, Picton TW, Miller LM. Brain oscillations during semantic evaluation of

speech. Brain and cognition. 2009;70(3):259-66. doi: 10.1016/j.bandc.2009.02.008. PubMed

PMID: 19324486; PubMed Central PMCID: PMC2683907.

27.   Weiss S, Mueller HM. "Too Many betas do not Spoil the Broth": The Role of Beta

Brain Oscillations in Language Processing. Frontiers in psychology. 2012;3:201. Epub

2012/06/28. doi: 10.3389/fpsyg.2012.00201. PubMed PMID: 22737138; PubMed Central

PMCID: PMC3382410.

28.   Wilkins RW, Hodges DA, Laurienti PJ, Steen M, Burdette JH. Network science and

the effects of music preference on functional brain connectivity: from Beethoven to Eminem.

Scientific reports. 2014;4:6130. doi: 10.1038/srep06130. PubMed PMID: 25167363.

29.   Sridharan D, Levitin DJ, Chafe CH, Berger J, Menon V. Neural dynamics of event

segmentation in music: converging evidence for dissociable ventral and dorsal networks.

Neuron. 2007;55(3):521-32. doi: 10.1016/j.neuron.2007.07.003. PubMed PMID: 17678862.

30.   Platel H, Baron JC, Desgranges B, Bernard F, Eustache F. Semantic and episodic

memory of music are subserved by distinct neural networks. NeuroImage. 2003;20(1):244-56.

PubMed PMID: 14527585.

31.   Burunat I, Alluri V, Toiviainen P, Numminen J, Brattico E. Dynamics of brain

activity underlying working memory for music in a naturalistic condition. Cortex; a journal

devoted to the study of the nervous system and behavior. 2014;57:254-69. doi:

10.1016/j.cortex.2014.04.012. PubMed PMID: 24949579.

32.   Salimpoor VN, Benovoy M, Larcher K, Dagher A, Zatorre RJ. Anatomically distinct

dopamine release during anticipation and experience of peak emotion to music. Nature

neuroscience. 2011;14(2):257-62. Epub 2011/01/11. doi: 10.1038/nn.2726. PubMed PMID:

21217764.

33.   Patel AD. Science & music: talk of the tone. Nature. 2008;453(7196):726-7. doi:

10.1038/453726a. PubMed PMID: 18528382.

34.   Sammler D, Koelsch S, Friederici AD. Are left fronto-temporal brain areas a

prerequisite for normal music-syntactic processing? Cortex; a journal devoted to the study of

the nervous system and behavior. 2011;47(6):659-73. Epub 2010/06/24. doi:

10.1016/j.cortex.2010.04.007. PubMed PMID: 20570253.

35.   Tillmann B, Janata P, Bharucha JJ. Activation of the inferior frontal cortex in

musical priming. Brain research Cognitive brain research. 2003;16(2):145-61. Epub

2003/04/02. PubMed PMID: 12668222.

36.   Watanabe T, Yagishita S, Kikyo H. Memory of music: roles of right hippocampus

and left inferior frontal gyrus. NeuroImage. 2008;39(1):483-91. Epub 2007/10/02. doi:

10.1016/j.neuroimage.2007.08.024. PubMed PMID: 17905600.

37.   Peretz I, Gosselin N, Belin P, Zatorre RJ, Plailly J, Tillmann B. Music lexical

networks: the cortical organization of music recognition. Annals of the New York Academy

of Sciences. 2009;1169:256-65. Epub 2009/08/14. doi: 10.1111/j.1749-6632.2009.04557.x.

PubMed PMID: 19673789.

38.   Fujioka T, Trainor LJ, Large EW, Ross B. Beta and gamma rhythms in human

auditory cortex during musical beat processing. Annals of the New York Academy of

Sciences. 2009;1169:89-92. Epub 2009/08/14. doi: 10.1111/j.1749-6632.2009.04779.x.

PubMed PMID: 19673759.

39.   Fujioka T, Trainor LJ, Large EW, Ross B. Internalized timing of isochronous sounds

is represented in neuromagnetic beta oscillations. The Journal of neuroscience : the official

journal of the Society for Neuroscience. 2012;32(5):1791-802. doi:

10.1523/JNEUROSCI.4107-11.2012. PubMed PMID: 22302818.