sgRNA level is a major factor affecting CRISPRi knockdown efficiency in K562 cells

Y. Wang[a], Y. Xie[b], Z. C. Dong[a], X. J. Jiang[a], P. Gong[a], J. Lu[a], and F. Wan[a, *]

[a] *College of Life Sciences, Inner Mongolia Agricultural University, Inner Mongolia, 010010 China*

[b] *College of Science, Inner Mongolia Agricultural University, Inner Mongolia, 010010 China*

*e-mail: fwan@imau.edu.cn

Abstract—To determine how nuclease deactivated Cas9 (dCas9) or sgRNA expression level affects the knockdown efficiency of CRISPRi, K562 cell clones expressing KRAB-dCas9 protein either with the inducible Tet-on system or with the constitutive SFFV promotor were created by lentiviral transduction, and single clones were selected by fluorescence-activated cell sorting (FACS) for further study. Six genes with various expression levels were targeted using lentiviral sgRNA from two libraries in four cell clones with various KRAB-dCas9 expression levels. We determined the knockdown efficiency and the expression level of the dCas9 protein /sgRNA level by flow cytometry. The cell clone with the highest KRAB-dCas9 expression level achieved effective CRISPRi knockdown, and is statistically different from other clones, indicating enough KRAB-dCas9 expression might be a prerequisite for CRISPRi. Utilizing this clone, we modified the expression level of sgRNA by adopting different multiplicity of infection（MOI）in lentiviral transduction and found that the knockdown efficiency was neither affected by the target gene expression level nor does it correlate with KRAB-dCas9 level, which remained relatively constant (CV=2.2%) across knockdown experiments. 74.72%, 72.28%, 39.08% knockdown of *mmadhc*, *rpia*, *znf148* genes were achieved, and the knockdown efficiency correlated well with the sgRNA expression level. Linear regression modeling of the data revealed that the knockdown efficiency is significantly affected by both KRAB-dCas9 and sgRNA level,

and the sgRNA level has a greater impact, based on the standardized coefficient（0.525

for KRAB-dCas9, 0.981 for sgRNA）, indicating that sgRNA level is a major factor

affecting CRISPRi efficiency.

Keywords: CRISPR interference, knockdown efficiency, inducible Tet-on system, the

multiplicity of infection, sgRNA expression level

INTRODUCTION

CRISPRi is gradually replacing the siRNA technique for mechanistic

investigations in various fields [1]. However, there is a lack of mechanistic study on the

efficiency of the CRISPRi system. How the two major players of the system, the dCas9

fusion protein, and the guide RNA affect the knockdown efficiency has not been

systematically studied.

Three protein domains were commonly used for repressing target gene

transcription: KRAB, MeCP2, dCas9 and its synthetic derivatives [2-3]. KRAB recruits

KAP1, which serves as a scaffold for various heterochromatin-inducing factors [4];

MeCP2 binds to the methylated DNA and interacts with histone deacetylase and the

co-repressor SIN3A [5]. A combination of dCas9 with KRAB and MeCP2 has been

shown to improve the knockdown efficiency [2-3], and dCas9, KRAB-dCas9,

dCas9-KRAB-MeCP2 are three commonly used combinations.

Although the dynamics of KRAB-dCas9 and sgRNA mediated CRISPRi

knockdown has not been studied thoroughly, the mechanistic study of the CRISPR

system could provide some clues due to the similarity: effective knockdown of target

gene started from the sgRNA being expressed from the transgene, formed a complex

with Cas9 protein, and roaming in the nucleus until the sgRNA hybrid to target DNA,

then the Cas9 protein cut the target gene [6]. Cas9 is essential for guide RNA stability

and that the nuclear Cas9–guide RNA complex levels limit the targeting

efficiency. Similar dynamics could exist in the CRISPRi system, and the very first step

is to determine how dCas9 fusion protein and the sgRNA expression level affect

CRISPRi efficiency.

In this study, different cell clones with various KRAB-dCas9 protein level and

subclones with different sgRNA expression level was evaluated for CRISPRi efficiency.

Linear regression was applied to determine the major factor that determines CRISPRi

efficiency.

EXPERIMENTAL

Cell culture and construction of plasmids. HEK293T and K562 cells were

obtained from the National Infrastructure of Cell Line Resource. HEK293T cells were

cultured at 37°C in Dulbecco's modified Eagle's medium (DMEM) containing 4.5 g/L

glucose (Gibco) supplemented with 10% (v/v) fetal bovine serum (FBS, Gibco). K562

cells were maintained in Iscove's modified Dulbecco's medium (IMDM, Gibco)

supplemented with 10% FBS. Cells were tested every 3 months for mycoplasma

contamination and consistently tested negative. To induce the expression of

KRAB-dCas9 protein using doxycycline, we replaced the promotor of

SFFV-KRAB-dCas9 plasmid (plasmid Addgene, #60954 provided by Jonathan S.

Weissman) with the pTRE3G promoter (plasmid Addgene # 96964 provided by Elena

Cattaneo) using classic cloning method, primers are listed in Supplementary Table 1.

Lentiviral Packaging. 2.1×10^7 HEK293T cells were seeded into T175 flasks 1d

before transfection. Cells were transfected with the packaging plasmids of pSPAX2

(11.6 ug) and pMD2.G (2.8 ug) (plasmid Addgene #12260 and #12259 provided by

Didier Trono), and the target plasmids (11.6 ug) using Polyetherimide (PEI) (1 mg/mL),

with the nitrogen to phosphorus (N/P) ratio 20 [2, 7]. Viral supernatants were collected

at 48 h and 72 h after transfection and filtered using 0.45 um filters. Subsequently, 4 mL

20% sucrose solution was added to the bottom of the 32 mL viral supernatants, and

centrifuge for 2 h at 82,700 g, 4°C [8], and the pellet was resuspended by 100 uL

pre-cooled PBS without Ca^{2+}/Mg^{2+}. The titer of the lentivirus was determined using

flow cytometry method as described [8].

Generating cell clones by lentiviral transduction. To generate single clones with constitutive KRAB-dCas9 protein expression, 1×10^5 K562 cells were transduced with the lentiviral particles for SFFV-KRAB-dCas9 using polybrene (final concentration 10 ug/mL) at MOI 20 in six-well plates. 72 h after transduction, cells were analyzed and sorted into 96-well plates using a cell sorter(BD FACS Aria II). Single clones emerged after 14 days and were expanded and cultured for further analysis.

For inducible KRAB-dCas9 expressing cells, 1×10^5 K562 cells were infected with lentiviral particles for expressing the trans-activator of the pTRE3G promotor (pLVX-Tet3G) and selected for 10 days using G418 (Final concentration 100 ug/mL). Subsequently, the cell clones were transduced with the lentiviral particles for pTRE3G-KRAB-dCas9 using polybrene, the following step with the same procedure described in the above section.

Flow cytometry analysis. The flow cytometry analysis was performed on Beckman CytoFLEX S. The data were analyzed using Flowjo 10 and CytoExpert 2.0 software.

Western blot analysis and ELISA. To detect the expression level of KRAB-dCas9 protein, K562-idCas9 #49 were induced by the doxycycline of 200 nM and 2000 nM for 48 h. Then, total protein was extracted using RIPA lysis buffer with 1 mM PMSF, and concentration was determined by the BCA protein assay kit (Solarbio, PC0020). For Western blot analysis, total protein (20 ug) were separated by 8% SDS-PAGE and transferred onto a 0.22 um nitrocellulose membrane at 150 V, 90 min. The membrane was incubated with the Cas9 polyclonal antibody (Clontech, 632607) and α-Tubulin (ABclonal, AC012), and the secondary antibodies IRDye™-800CW(BD, Lot No. C50113-06). The membrane was scanned by the Odyssey™CLx Imaging system (Li-COR Bioscience). For ELISA, the KRAB-dCas9 protein was quantified using the CRISPR/Cas9 assay ELISA kit (Epigentek, P-4060-48) according to the manufacturer's protocol.

The evaluation of the CRISPRi knockdown efficiency. Single clones were infected with the lentivirus of sgRNA (MOI 20 if not explicitly stated otherwise) with polybrene (Final concentration 10 ug/mL). After 72 h transduction, clones were selected

by puromycin (Final concentration 5 ug/mL) for 72 h. The expression level of target

genes was detected by qPCR using the relative quantification method. qPCR primers are

listed in Supplementary Table 1.

Statistical analysis. ANOVA and multiple comparisons were performed to

compare the four-cell clones. Linear regression modeling was performed to analyze the

relationship between the knockdown efficiency and the expression level of either

KRAB-dCas9 or sgRNA. All statistical analysis was performed using the SPSS

package.

RESULTS

To analyze how KRAB-dCas9 protein and/or the sgRNA expression level affects

the CRISPRi knockdown efficiency, we adopted a hierarchical design that firstly

compared cell clones with various KRAB-dCas9 expression level, then compared cell

subclones with various sgRNA expression level (Fig. 1a). For the first comparison, we

established K562 cell clones that have either constitutive or inducible expression of the

KRAB-dCas9-mCherry fusion protein.

Tet-on system was adopted to establish cell clones that express

KRAB-dCas9-mCherry fusion protein in doxycycline-inducible fashion: trans-activator

(Tet3G plasmid) of the pTRE3G promotor was first introduced into K562 cells by

lentiviral transduction, followed by G418 selection, and subsequently undergone

lentiviral transduction of the KRAB-dCas9-mCherry expressing vector with the

inducible pTRE3G promotor (Fig. 1b). After FACS selection based on mCherry

expression, sixty inducible clones (K562-idCas9 #1 to #60) were randomly chosen from

twelve 96-well plates for further expansion, and their transgene expression pattern was

analyzed with flow cytometry upon doxycycline induction (Supplemental Fig. 1-2).

Based on the fold change of transgene expression level, represented by the medium

fluorescent intensity (MFI), with or without doxycycline induction（Induced

MFI/Uninduced MFI）, the 60 clones were ranked (Fig. 1c) and five clones

(K562-idCas9 #9, #25, #49, #54, #60) with various fold changes were chosen for further

analysis.

We first confirmed the inducible expression of KRAB-dCas9-mCherry in

K562-idCas9 #49 and #60 by detecting mCherry, the tag that co-expressed with

KRAB-dCas9, using flow cytometry (Fig. 2a), then by detecting dCas9 with western

blot and ELISA (Fig. 2b-c). KRAB-dCas9 protein mainly localized in the nucleus, as

expected. MFI of mCherry exhibits a similar pattern as the ELISA signal, confirming

that mCherry represented the KRAB-dCas9 expression level. For sgRNA, two sgRNA

libraries, the hCRISPRi-v2 library [9] and the Dolcetto library [10], were obtained from

Addgene, both contain multiple sgRNAs targeting one gene and ranked them (Table 1).

Pilot CRISPRi knockdown experiments were carried out using the lentiviral particle of

highest-ranking sgRNA against UBE4A and DPH2, from the hCRISPRi-v2 library.

K562-idCas9 #9, #25, #54 were excluded from further analysis based on their

performance in the pilot experiments (Supplementary Fig. 3a-d). K562-idCas9 #49

displayed 24% knockdown efficiency in the DPH2 gene with 100 nM doxycycline

(Supplementary Fig. 3e), has relatively higher KRAB-dCas9-mCherry expression level,

and were chosen for further analysis, together with K562-idCas9 #60, which has

relatively low KRAB-dCas9-mCherry expression (Fig. 2d).

Similarly, we established constitutive KRAB-dCas9-mCherry expressing cell

clones, and chose three clones, K562-dCas9 #A, K562-dCas9 #B, and K562-dCas9 #C.

Pilot knockdown experiments revealed that K562-dCas9 #A failed to knockdown

UBE4A or DPH2 genes (Supplemental Fig. 3f), and was excluded from further analysis.

To compare the knockdown efficiency of cell clones with various KRAB-dCas9

expression levels, the constitutive K562-dCas9 #B, #C and the inducible K562-idCas9

#49, #60 were evaluated for their CRISPRi efficiency. The inducible KRAB-dCas9

expressing clones have lower expression levels compared to the constitutive clones,

indicating that inducible expression may not be able to reach the expression level of

constitutive expression (Fig. 2d). For sgRNA, we randomly choose three sgRNAs per

CRISPRi library of high rankings (Table 1). After lentiviral transduction of sgRNA into

the cell clones, puromycin selection was applied, and the knockdown efficiency was

assessed by qPCR. *mmadhc*, *rpia*, *znf148*, *lsm4*, and *lima1* genes have been knocked down in clone K562-dCas9 #B, the clone with the highest KRAB-dCas9 expression level (Fig. 3a); yet all other clones have relatively low or no knockdown efficiency or the target gene expression level increased (Fig. 3a), indicating sufficient KRAB-dCas9 expression level is a prerequisite for efficient CRISPRi.

To better understanding the difference among the four cell clones, ANOVA was performed and multiple comparisons were made. The results indicated that K562-dCas9 #B is significantly different from K562-dCas9 #C and K562-idCas9 #60, and the p-value for comparison between K562-dCas9 #B and K562-idCas9 #49 is 0.054, indicating that K562-dCas9 #B is also different from K562-idCas9 #49, corroborating our finding that only K562-dCas9 #B has effective CRISPRi (Table 2).

We also analyzed whether knockdown efficiency is affected by the target gene expression level, and found no correlation (Fig. 3b). The relative expression level of the target genes in four cell clones differed from each other, probably due to the single cloning procedure [11] (Fig. 3c). Continuous monitoring of the KRAB-dCas9 expression revealed that K562-dCas9 #C lost transgene expression over time, while other clones maintained their transgene expression (Supplementary Table 2).

Utilizing K562-dCas9 #B, we assessed the impact of sgRNA expression level on knockdown efficiency. Gene transfer rate, the rate of cells expressing a transgene in the cell population, were assessed to ensure successful transduction. BFP MFI, which is co-expressed with the sgRNA in a polycistronic transcript, was used for determining the gene transfer rate and found to correlate with MOI, fitting a polynomial curve (Fig. 4a). Based on this curve, the gene transfer rate was 34.1% at MOI 0.5, close to the previously reported gene transfer rate 28.6% at MOI 0.5 in K562 cells [12], confirming an effective lentiviral packaging and transduction.

We then assessed how MOI correlated with transgene expression and found that in positive sgRNA expression cells, based on flow cytometry detection (Fig. 4b) , BFP MFI linearly correlated with MOI (Fig. 4c). Moreover, a linear relationship was observed between the MOI and the knockdown efficiency (Fig. 4d), indicating that as sgRNA expression level correlates with knockdown efficiency. In our research, based

on the linear relationship between the MOI and the knockdown efficiency, around 45%

knockdown could be achieved for the *mmadhc* gene by MOI 0.5, the common MOI

used for CRISPRi library screening (Fig. 4d).

To systematically evaluate how dCas9 and sgRNA expression level affects the

knockdown efficiency, a linear regression model with dCas9 MFI and sgRNA as

independent variable individually was applied. Results have shown that the assumption

of equal variance was met. When dCas9 serve as the independent variable in the linear

model, R^2=0.243, F=8.367, p=0.008, indicating that dCas9 fusion gene expression

affected the knockdown efficiency significantly. The standardized coefficient β_1=0.525

indicated that when dCas9 change for one standard deviation, the knockdown efficiency

changes 0.525 standard deviations. Similarly, when sgRNA serves as the independent

variable in the linear model, R^2=0.949, F=75.84, p=0.003, and standardized coefficient

β_2=0.981, indicating that sgRNA level affected knockdown efficiency significantly, and

when sgRNA level change for one standard deviation, the knockdown efficiency

changes for 0.981 standard deviations. The standardized coefficient value has shown

that sgRNA level has a greater impact on the knockdown efficiency, which corroborated

with the correlation analysis.

DISCUSSION

As the first step to dissect the dynamics of the KRAB-dCas9 mediated CRISPRi

system, we assessed how the expression level of dCas9 fusion protein and the sgRNA

affect the knockdown efficiency, which provided the basis for further optimization of

the CRISPRi library screening.

We discovered that KRAB-dCas9 expression level significantly impact CRISPRi

efficiency, and our data, although limited in the number of clones tested, indicated that

enough dCas9 fusion protein probably is a prerequisite for the efficient knockdown. A

study on more cell clones is mandated for clarifying whether there is a threshold of

KRAB-dCas9 expression above which effective knockdown is guaranteed.

The requirement of the KRAB-dCas9 level has implications for CRISRPi library

screenings: its mandated selection of the cells after lentiviral infection, which can be

challenging due to many cells required: the number of the cell in each group shall exceed 1000× the sgRNA number in the library. A high-speed cell sorter is required, which is unfortunately not available in most laboratories. CRISPRi library screening without such an instrument might have to rely on obtaining and expanding a single-cell clone that stably expressing a high level of KRAB-dCas9. However, the lentiviral transgene promoter could be methylated and transgene expression shut down in the continuous culture of the cells, as indicated in one of the cell clone K562-dCas9 #C, so continuous monitoring of KRAB-dCas9 expression level is recommended, and more than one selection could be required to obtain such cell clone.

It's known that single cloning affected the transcription profile of the KRAB-dCas9 expressing cells, and poly-clonal cell populations are better in CRISPRi applications [11]. Effects on transcriptional profile brought by the single cloning procedure can be canceled out, though, as comparisons were made between groups using the same cell clone. Alternatively, with a more potent CRISPRi system, the high expression level of dCas9 fusion protein might not be necessary, and drug selection based on antibiotic resistance could be applied. The dCas9-KRAB-MeCP2 fusion protein is more effective in knockdown [13] and may offer such an option.

Among the six genes randomly chosen for CRISPRi, *npepps* and *rpia* has low efficiency in most or all cell clones. Gene annotation revealed that *npepps* is a puromycin responsive gene, and the puromycin selection after sgRNA transduction probably activated its expression; *rpia* has been reported to be responsive to doxycycline, with increased expression upon doxycycline treatment [14].

We observed that with enough KRAB-dCas9 expression level, CRISPRi efficiency correlated very well with the MOI, which correlated equally well with the sgRNA transgene expression, indicating that CRISPRi efficiency correlated well with sgRNA level. Although it has not been studied in the CRISPRi system, it is known that in the CRISPR system the nuclear concentration of the guide RNA is the limiting factor for efficient DNA targeting [6, 15]. We found that the guide RNA level has a greater impact on CRISPRi efficiency than dCas9. The dynamics of ribonucleoprotein

formation and DNA targeting is probably similar between CRISPR and CRISPRi system.

To improve the knockdown efficiency, it is critical to increasing the expression level of sgRNA since sgRNA is very unstable when it was not bound with Cas9 protein [6], improvement of the sgRNA stability could increase efficiency [16]. Alternatively, a change of promotor might serve to increase the sgRNA expression.

The correlation between the sgRNA expression level and the knockdown efficiency of the CRISPRi system has indications for the CRISPRi library screening, too: inducible sgRNA will provide better control of the knockdown. It has been reported that inducible sgRNA offered better control of knockdown in the CRISPR system [17], again pointing to the similarities between the two systems.

ACKNOWLEDGMENTS

We thank Professor Jonathan S. Weissman, David E. Root and John G. Doench for sharing their CRISPRi library on Addgene, and we acknowledge Dr. Luke A. Gilbert for offering advice on CRISPRi library.

FUNDING

This work was supported by the National Natural Science Foundation of China (81860652).

COMPLIANCE WITH ETHICAL STANDARDS

The authors declare that they have no conflict of interest. This article does not contain any studies involving animals or human participants performed by any of the authors.

REFERENCES

1. Evers B., Jastrzebski K., Heijmans J., Grernrum W., Beijersbergen R.L., Bernards R. 2016. CRISPR knockout screening outperforms shRNA and CRISPRi in identifying essential genes. *Nat. Biotechnol*. **34**, 631–633.

2.   Gilbert L.A., Horlbeck M.A., Adamson B., Villalta J.E., Chen Y.W., Whitehead

E.H., Guimaraes C.P., Panning B., Ploegh H.L., Bassik M.C., Qi L.S., Kampmann

M., Weissman J.S. 2014. Genome-Scale CRISPR-Mediated Control of Gene

Repression and Activation. *Cell*. **159**, 647-661.

3.   Yeo N.C., Chavez A., Lancebyrne A., Chan Y.C., Menn D.J., Milanova D., Kuo

C.C., Guo X.G., Sharma S., Tung A., Cecchi R.J., Tuttle M., Pradhan S., Lim

E.T., Davidsohn N., Ebrahimkhani M.R., Collins J.J., Lewis N.E., Kiani S., Church

G.M. 2018. An enhanced CRISPR repressor for targeted mammalian gene

regulation. *Nat. Methods*. **15**, 611-616.

4.   Peng H.Z., Ivanov A.V., Oh H.J., Lau Y.C., Rauscher F.J. 2009. Epigenetic Gene

Silencing by the SRY Protein Is Mediated by a KRAB-O Protein That Recruits the

KAP1 Co-repressor Machinery. *J. Biol. Chem*. **284**, 35670-35680.

5.   Hansen J.C., Ghosh R.P., Woodcock C.L. 2010. Binding of the Rett syndrome

protein, MeCP2, to methylated and unmethylated DNA and chromatin. *Iubmb Life*.

62, 732-738.

6.   Ma H.H., Tu L.C., Naseri A., Huisman M., Zhang S.J., Grunwald D., Pederson T.

2016. CRISPR-Cas9 nuclear dynamics and target recognition in living cells. *J. Cell*

Biol. **214**, 529-537.

7.   Yang S.Z., Shi H.J., Chu X.R., Zhou X.L., Sun P.N. 2016. A rapid and efficient

polyethyleneimine-based transfection method to prepare lentiviral or retroviral

vectors: useful for making iPS cells and transduction of primary cells. *Biotechnol*.

Lett. **38**, 1631-1641.

8.   Kutner R.H., Zhang X.Y., Reiser J. 2009. Production, concentration and titration of

pseudotyped HIV-1-based lentiviral vectors. *Nat. Protoc*. **4**, 495-505.

9.   Horlbeck M.A., Gilbert L.A., Villalta J.E., Adamson B.S., Pak R.A., Chen

Y.W., Fields A.P., Park C.Y., Corn J.E., Kampmann M., Weissman J.S. 2016.

Compact and highly active next-generation libraries for CRISPR-mediated gene

repression and activation. *Elife*. **5**, e19760.

10. Sanson K.R., Hanna R.E., Hegde M., Donovan K.F., Strand C., Sullender

M.E., Vaimberg E.W., Goodale A., Root D.E., Piccioni F., Doench J.G. 2018.
Optimized libraries for CRISPR-Cas9 genetic screens with multiple modalities. *Nat.*
Commun. **9**, 5416.

11. Stojic L., Lun A.T., Mangei J., Mascalchi P., Quarantotti V., Barr A.R., Bakal C.,
Marioni J.C., Gergely F., Odom D.T. 2018. Specificity of RNAi, LNA and
CRISPRi as loss-of-function methods in transcriptional analysis. *Nucleic Acids Res.*
46, 5950-5966.

12. Kustikova O.S., Wahlers A., Kuhlcke K., Stahle B., Zander A.R., Baum C., Fehse
B. 2003. Dose finding with retroviral vectors: correlation of retroviral vector copy
numbers in single cells with gene transfer efficiency in a cell population. *Blood.*
102, 3934-3937.

13. Martella A., Firth M., Taylor B.J., Goppert A., Cuomo E.M., Roth R.G., Dickson
A.J., Fisher D.I. 2019. Systematic Evaluation of CRISPRa and CRISPRi Modalities
Enables Development of a Multiplexed, Orthogonal Gene Activation and
Repression System. *ACS Synth. Biol.* **8**, 1998-2006.

14. Ahler E., Sullivan W.J., Cass A., Braas D., York A.G., Bensinger S.J., Graeber
T.G., Christofk H.R. 2013. Doxycycline alters metabolism and proliferation of
human cell lines. *Plos One.* **8**, e64561.

15. Yuen G., Khan F.J., Gao S.J., Stommel J.M., Batchelor E., Wu X.L., Luo J. 2017.
CRISPR/Cas9-mediated gene knockout is insensitive to target copy number but is
dependent on guide RNA potency and Cas9/sgRNA threshold expression level.
Nucleic Acids Res. **45**, 12039-12053.

16. M W., Zhang Y.P., X X.T., L L., W X.F., W H.Y. 2019. 5′ capped and 3′
polyA-tailed sgRNAs enhance the efficiency of CRISPR-Cas9 system. *Protein &*
Cell. **10**, 223-228.

17. Aubrey B.J., Kelly G.L., Kueh A.J., Brennan M.S., Oconnor L., Milla L., Wilcox
S., Tai L., Strasser A., Herold M.J. 2015. An Inducible Lentiviral Guide RNA
Platform Enables the Identification of Tumor-Essential Genes and
Tumor-Promoting Mutations In Vivo. *Cell Rep.* **10**, 1422-1432.

Table 1. List of sgRNA sequences

sgRNA	Sequences (5'-3')	Rank	Library
DPH2	GATGTTTAGCAGCCCTGCCG	2/10	hCRISPRi-v2
UBE4A	GACAGGGTCGAAGTTCAAGA	1/10	hCRISPRi-v2
MMADHC	GAAAATACGGGCAATACCGA	1/10	hCRISPRi-v2
RPIA	GGTGCACAGTCTGGGACCCG	1/10	hCRISPRi-v2
ZNF148	GCAGGCCGAGCGCTTTACCG	1/10	hCRISPRi-v2
LSM4	CAGTTGCCCTGCCCTCACCA	1/3	Dolcetto
NPEPPS	TGGCGGGCGGGCTGCCTACG	3/3	Dolcetto
LIMA1	GATGCTTTCTCCATGTGGCA	1/3	Dolcetto

357 **Table 2.** Effects of the four cell clones on knockdown efficiency and multiple
358 comparisons

Cell clone	The average of knockdown efficiency (%)	Standard deviation (%)
K562-dCas9 #B	43.16[a]	45.55
K562-idCas9 #49	-3.24[ab]	48.65
K562-dCas9 #C	-4.63[b]	36.48
K562-idCas9 #60	-15.24[b]	19.92

359 Different lowercase letters indicate a significant difference in knockdown efficiency
360 under different cell clones ($p < 0.05$).

Fig. 1. The construction and selection of inducible cell clones. (a) Experiment design. #B: K562-dCas9 #B; #C: K562-dCas9 #C; #49: K562-idCas9 #49; #60: K562-idCas9 #60. #B1, #B2, #B3, #B4, #B5 indicates K562-dCas9 #B infected with difference MOI of sgRNA. (b) Lentiviral vectors for inducible and constitutive KRAB-dCas9 expression. (c) Sixty inducible cell clones ranked based on induced MFI/uninduced MFI, the red prismatic indicated the single clones that chosen for further analysis. 2000 nM doxycycline was applied for induction for 48 h.

Fig. 2. The expression of KRAB-dCas9 in inducible and constitutive cell clones. (a) mCherry MFI in K562-idCas9 #49, K562-idCas9 #60, in response to various doses of

doxycycline. Triangle indicated K562-idCas9 #49, round indicated K562-idCas9 #60.

(b-c) Western blot and ELISA detection of nuclear and cytoplasmic KRAB-dCas9 in

K562-idCas9 #49 induced by 200 nM and 2000 nM doxycycline. (d) mCherry MFI of

two inducible cell clones induced by 100 nM doxycycline and two constitutive cell

clones. #B: K562-dCas9 #B; #C: K562-dCas9 #C; #49: K562-idCas9 #49; #60:

K562-idCas9 #60.

Fig. 3. Efficient CRISPRi knockdown requires enough KRAB-dCas9 expression. (a)

Knockdown efficiency of six genes in four clones, #B: K562-dCas9 #B; #C:

K562-dCas9 #C; #49: K562-idCas9 #49; #60: K562-idCas9 #60. (b) The relationship

between target gene expression level and CRISPRi knockdown efficiency in

K562-dCas9 #B. (c) The relative expression level of six genes in four clones, #B:

K562-dCas9 #B; #C: K562-dCas9 #C; #49: K562-idCas9 #49; #60: K562-idCas9 #60.

Fig. 4. CRISPRi knockdown efficiency correlated with sgRNA expression level. (a) Correlation between MOI of sgRNA infection of the *mmadhc* gene and gene transfer rate, determined by BFP MFI, in K562-dCas9 #B. (b) The scatter plot of K562-dCas9 #B with sgRNA of the *mmadhc* gene. The red frame indicated positive sgRNA expression cells with BFP fluorescence. (c) Correlation between BFP MFI in positive sgRNA expression cells and the MOI of sgRNA infection of the *mmadhc* gene in K562-dCas9 #B. positive sgRNA cells were defined by flow cytometry. (d) Correlation between knockdown efficiency and MOI of sgRNA infection of the *mmadhc* gene in K562-dCas9 #B.