1   **Single-cell transcriptome analysis reveals cell-cell communication and thyrocyte**

2   **diversity in the zebrafish thyroid gland.**

3   Pierre Gillotay[1], Meghna Shankar[1], Sema Elif Eski[1], Susanne Reinhardt[2], Annekathrin
4   Kränkel[2], Juliane Bläsche[2], Andreas Petzold[2], Gokul Kesavan[3], Christian Lange[3],
5   Michael Brand[3], Vincent Detours[1], Sabine Costagliola[1]*, Sumeet Pal Singh[1]*
6
[1] IRIBHM, Université Libre de Bruxelles (ULB), Route de Lennik 808, 1070 Brussels,
Belgium.
[2] DRESDEN-concept Genome Center, DFG NGS Competence Center, c/o Center for
Molecular and Cellular Bioengineering, TU Dresden, Dresden, 01307, Germany
[3] Center for Molecular and Cellular Bioengineering, TU Dresden, Dresden, 01307,
Germany.
* Corresponding authors: scostag@ulb.ac.be; sumeet.pal.singh@ulb.ac.be

Abstract

The thyroid gland regulates growth and metabolism via production of thyroid hormone in

follicles composed of thyrocytes. So far, thyrocytes have been assumed to be a

homogenous population. To uncover genetic heterogeneity in the thyrocyte population,

and molecularly characterize the non-thyrocyte cells surrounding the follicle, we

developed a single-cell transcriptome atlas of the zebrafish thyroid gland. The 6249-cell

atlas includes profiles of thyrocytes, blood vessels, lymphatic vessels, immune cells and

fibroblasts. Further, the thyrocytes could be split into two sub-populations with unique

transcriptional signature, including differential expression of the transcription factor

pax2a. To validate thyrocyte heterogeneity, we generated a CRISPR/Cas9-based

pax2a knock-in line, which demonstrated specific *pax2a* expression in the thyrocytes.

However, a population of *pax2a*-low mature thyrocytes interspersed within individual

follicles could be distinguished, corroborating heterogeneity within the thyrocyte

population. Our results identify and validate transcriptional differences within the

nominally homogenous thyrocyte population.

Keywords

Single-cell, transcriptomics, zebrafish, heterogeneity, thyroid gland, thyroid follicular

cells, connective septa, CRISPR/Cas9, knock-in

One-line summary

Single-cell analysis uncovers latent heterogeneity in thyroid follicular cells.

Graphical Abstract

Introduction

The thyroid gland produces hormones thyroxine (T4) and triiodothyronine (T3)

that regulate body metabolism, growth, and development. Thyroid dysfunction, a

disease afflicting almost 100 million people worldwide (1), is common and treatable by

hormone replacement. If left untreated, however, it may result in profound adverse

effects on the human body, including mental retardation, goiter or dwarfism.

The thyroid gland is an endocrine organ with an intricate structure enabling

production, storage and release of the thyroid hormones. It contains numerous variable-

sized spherical follicles composed of thyroid follicular epithelial cells, or thyrocytes. The

thyrocytes generate the thyroid hormones in a multi-step process. They secrete and

store thyroglobulin (TG) in the lumen of the follicles. Additionally, they intake iodide from

the blood via sodium-iodide symporter (NIS / Slc5a5). At the interface between

thyrocytes and the lumen, thyroid peroxidase (TPO) expressed by the cells catalyzes

the coupling of iodide to tyrosyl residues of thyroglobulin (TG). Iodinated TG is absorbed

back into the thyrocyte and cleaved by cysteine proteases in lysosomes to form T4 and

T3 (2). Though the machinery responsible for the production of thyroid hormones by

thyrocytes is well established, it remains unknown if all the thyrocytes resident in the

thyroid gland are equally capable of generating thyroid hormones. In other words, the

extent of molecular homogeneity between individual thyrocytes has not yet been

investigated.

Additionally, the thyroid gland contains many cell-types with potential roles in

modulating thyrocyte functionality. The gland contains an extensive distribution of blood

vessels, which carry iodide to the thyrocytes and carry thyroid hormones away from

them. The thyroid follicles are separated by a mesenchymal cell population, called

connective tissue septa, which also divides the gland into lobules. The mammalian

thyroid gland also contains parafollicular epithelial cells, or C-cells, that synthesize and

secrete the hormone calcitonin. These parafollicular epithelial cells are, however,

located outside the thyroid gland in fish and amphibians (3). Further, the presence of

immune cells and innervation has been demonstrated within the thyroid gland (4, 5).

Though we have a considerable understanding of these cell-types on a histological

level, we still lack the molecular characterization of the thyroid gland cell ensemble. This

extends to an incomplete appreciation of the impact of the diverse cell-populations on

thyroid follicular cell physiology.

To uncover the diversity within the thyrocyte population, and further characterize

the surrounding tissue at cellular resolution, we develop the first atlas of the thyroid

gland at single-cell resolution. For this, we build on the progress in single-cell

transcriptomics (6) to transcriptionally profile thousands of individual cells isolated from

the thyroid gland of adolescent and adult zebrafish. We demonstrate that these profiles

comprehensively represent the cells present in the zebrafish thyroid gland. Further, we

demonstrate the segregation of thyrocytes into two transcriptionally distinct sub-

populations. Utilizing the expression profiles of discrete cell populations, we build an

intercellular signaling network to uncover communication between thyrocytes and the

surrounding tissue. Finally, to enable easy access to the data, we have made the

zebrafish thyroid gland atlas available for online browsing.

Results

Single-cell transcriptomics of the zebrafish thyroid gland.

To generate the molecular catalogue of the thyroid gland at cellular resolution,

we sampled the organ from two ages of zebrafish: 2 month post-fertilization (mpf) and 8

mpf (Supp. Figure 1). The time points span adolescent to adult transition in zebrafish,

with animals containing fully differentiated functional organs at both stages. By 2 mpf,

the adolescent animals have completed morphogenesis, but are yet to reach sexual

maturity. The animals sampled at 2 mpf were on average 2.6 cm in length and 123.8 mg

in weight. In contrast, fish at 8 mpf are sexually mature adults, with an average length of

3.5 cm and an average weight of 294.4 mg (Supp. Figure 1). To characterize the organ

cell-types in an unbiased manner, we dissected out the entire thyroid gland (Fig. 1A, B)

from six animals at each stage, and prepared the single-cell suspension for cDNA

library preparation. To guide thyroid gland dissection, we utilized the *Tg(tg:nls-mVenus-*

T2A-NTR) zebrafish reporter line (7) that labels thyrocytes with bright yellow fluorescent

protein (Fig. 1B'). The micro-dissected tissue was dissociated using enzymatic

digestion. The single-cell suspension was stained with calcein, which specifically labels

live cells with blue fluorescence. The live cells were then enriched using FACS (Supp.

Figure 2) to limit false positive signals from dead and/or ruptured cells (8). Twelve

thousand live-cells, pooled from six animals, were collected in separate tubes according

to age and profiled using droplet-based high-throughput single-cell RNA-sequencing

provided by 10X Genomics (9, 10). The 10X Genomics pipeline uses molecule and cell-

specific barcoding allowing transcript quantification without amplification bias (11, 12).

Using the Cell Ranger bioinformatics pipelines, the resulting Next-Generation

Sequencing libraries were mapped to the zebrafish genome, de-multiplexed according

to their cellular barcodes and quantified to generate gene/cell UMI (unique molecular

identifier) count tables. Quality-based exclusion of single-cell transcriptomes was

implemented based on mean library size, percentage of mitochondrial reads and

number of genes detected per cell. On average, we detected 6012 UMIs and 1303

genes per cell (Supp. Figure 3). The process recovered in total 6249 cells, providing

single-cell transcriptomic profiles for 2986 and 3263 individual cells for 2 mpf and 8 mpf,

respectively.

Identification of cell-types present in the zebrafish thyroid gland.

To aid with visualization of the zebrafish thyroid gland single-cell RNA-Seq

(scRNA-Seq) data, we projected the cellular profiles onto t-distributed stochastic

neighbor embedding (t-SNE) plots, a non-linear dimensionality reduction technique (13)

(Fig. 1C). Using unsupervised graph-based clustering, we identified seven clusters for

the thyroid gland. Using the expression of genes involved in thyroid hormone

production, we could identify one of the clusters as thyroid follicular cells (Fig. 1D – F).

The cluster, labeled as thyrocytes, contains 267 cells.

To define the identity of the remaining cell clusters, we generated cluster-specific

marker genes by performing differential gene expression analysis (Fig. 2A) (Supp.

Table 1). For four clusters, the marker genes included one or more known cell type–

specific identifiers. This included *gpr182* for endothelial cells; *acta2* for musculature;

fcer1gl for immune cells; and *ponzr3* for cells from zebrafish gills (Fig. 2B – E). Based

on these cell identifiers, the atlas includes 233 endothelial cells, 135 muscle lineage

cells, 914 immune cells and 199 cells from zebrafish gills. Notably, the endothelial cell

cluster includes blood vessels (*flt1* and *kdrl*) and lymphatic vessels (*mrc1a*, *prox1a*, *flt4*

and *lyve1b*) (Supp. Fig. 4); while the immune cell cluster includes macrophages

(*mpeg1.1* and *mfap4*), neutrophils (*lyz*) and lymphocytes (*il4*, *il13* and *il11b*) (Supp. Fig.

5).

For the remaining two clusters (number six and seven), we identified marker

genes that hinted towards identity of the cell-type. Specifically, *col1a2* and *tp63*

enriched in cluster number six and seven respectively (Fig. 2 F – G), are known

markers of fibroblasts (14, 15) and epithelial tissue (16–18). We performed gene-

ontology (GO) enrichment analysis of the marker genes to aid with classification (Supp.

Fig. 6). Cluster six demonstrated an enrichment of 'extracellular matrix structural

constituent', 'connective tissue development' and 'extracellular space', confirming the

presence of tissue fibroblasts in this cluster. Thus, we labelled cluster six as 'Stromal'

cells. Cluster seven displayed an enrichment of 'cell motility', 'cell migration' and

'epithelium development', suggestive of epithelial cells. Hence, we labelled cluster

seven as 'Non-Follicular Epithelium (NFE)', to distinguish them from the thyroid follicular

epithelial cells. Our data contains 3670 stromal cells and 831 non-follicular epithelial

cells.

Our marker gene identification further established additional genes enriched in a

single cell-type in the thyroid gland (Fig. 2A) (Supp. Table 1). For instance, we identified

cx30.3, a connexin gene and *prdx1*, a gene involved in the antioxidant response, to be

specifically expressed in the thyrocytes. To enable further investigation of the clusters

and gene expression profiles, we have developed an interactive webtool for online

browsing (https://sumeet.shinyapps.io/zfthyroid/).

Development of autocrine and paracrine signaling networks in the thyroid gland using known ligand-receptor interactions.

Having defined the cell types of the thyroid gland, we quantified potential cell-cell interactions between thyrocytes and all cell types present in the organ (Fig. 3A) based on a reference list of approximately 3,100 literature-supported interactions containing receptors and ligands from receptor tyrosine kinase (RTK), extracellular matrix (ECM)-integrin, chemokine and cytokine families (19). Although anatomical barriers between cell types are not modeled in this analysis, expression patterns of ligand-receptor pairs revealed a dense intercellular communication network (Fig. 3B). The network consisted of 322 ligands expressed on different cell-types with a corresponding receptor expressed on the thyrocytes (Supp. Table 2). For instance, the stromal cells express the ligand *lpl* (Lipoprotein Lipase) that signals through the *lrp2a* (zebrafish homologue of Megalin) receptor (Fig. 3C). Stromal and muscle cells express *dcn* (Decorin) whose receptor *met* is expressed by thyrocytes. Further, the ligand *cyr61* is broadly expressed in the thyroid gland, with one of its receptors, *itgb5*, an integrin isoform, expressed specifically by the thyrocytes. The identified interactions also include autocrine signaling. For example, the ligand *sema3b* and its receptor *nrp2a* are both present on thyrocytes. GO-analysis for identified ligand-receptor pairs revealed genes involved in 'PI3K-Akt signaling pathway', 'MET signaling' and 'integrin binding' (Supp. Fig. 7).

Thyrocytes are composed of transcriptionally distinct sub-populations.

Next, we characterized the transcriptional differences within the thyrocyte population. For this, we bioinformatically isolated the thyrocytes, and re-performed the

clustering pipeline on the isolated cell population. With this, we could segregate the

thyrocytes into two smaller clusters (Fig. 4A), labeled as 'Cluster_Blue' and

'Cluster_Red'. The two clusters displayed differences in the expression levels of 265

genes (Fig. 4B) (Supp. Table 3). Notably, Cathepsin B (*ctsba*) is significantly

downregulated in the blue cluster (Fold change = 1.6, p-value = 1.47×10^{-9}) (Fig. 4B –

C). Cathepsin B is a cysteine protease that is involved in the processing of iodinated

thyroglobulin to T4 and T3 in the thyrocyte lysosomes (2, 20). Moreover, fusion of

Cathepsin B and EGFP has been previously used to track thyroid hormone processing

lysosomes in rat thyroid epithelial cell lines (21).

Along with higher expression of Cathepsin B, the red cluster displayed significant

downregulation of *pax2a* expression (Fold change = 1.7, p-value = 8.24×10^{-9}) (Fig. 4B-

C). *pax2a* belongs to the PAX (paired box DNA-binding) domain containing family of

transcription factors. The loss of *pax2a* expression in the red cluster is notable, as

pax2a is an important regulator of thyrocyte development (22). Zebrafish thyroid

primordium expresses *pax2a* at 24 hpf (22), which is required for specification of the

thyroid follicles (23, 24). Consequently, zebrafish lacking *pax2a* fail to develop thyroid

follicles (22), which is similar to the Pax8 knock-out phenotype in mouse (25). The low

expression of *pax2a* in the red cluster, without a difference in *tg* expression, suggests

the presence of a thyrocyte sub-population with a distinct gene expression signature.

Generation of *pax2a* knock-in reporter line.

To validate the heterogeneity among the zebrafish thyrocytes, we focused on the

expression of *pax2a* transcription factor. We generated a knock-in line by inserting

monomeric Kusabira Orange 2 (mKO2) fluorescent protein to the 3' end of the

endogenous *pax2a* genomic location (Fig. 5A). The *pax2a^pax2a-T2A-mKO2^* (abbreviated as

*pax2a^mKO2^*) reporter expression perfectly overlapped with PAX2A antibody staining at 9.5

hours post-fertilization (Fig. 5B). Moreover, the knock-in line displayed mKO2

fluorescence in the otic vesicle, mid-hindbrain boundary, optic stalk, pronephros and the

thyroid gland (Fig. 5C – F, Supp. Movie 1), mimicking known expression of *pax2a* during

zebrafish development (26). Additionally, in order to assess whether the dynamics of

mKO2 expression would follow modifications in the expression of endogenous *pax2a,* we

used CRISPR/Cas9 technology to generate F0 knock-outs (also known as Crispant (27))

of *pax2a* gene in our *pax2a^mKO2^* line. The crispants displayed defects in thyroid

morphogenesis (Fig. 8G – H), mimicking the phenotype of *pax2a* loss-of-function

mutation (22). Live imaging of crispants at 55 hpf revealed strong decrease of mKO2

expression (Fig. 8G – H), thereby corroborating the faithful recapitulation of *pax2a*

expression by the newly generated reporter line.

Segregation of thyrocyte sub-populations based on *pax2a* reporter expression.

Upon investigating the fluorescence expression of the *pax2a* reporter in the

thyroid gland of adult zebrafish, we found strong and specific expression of *pax2a*

reporter in the thyrocytes lining the thyroid follicles (Fig. 6A – D). Although a majority of

thyrocytes displayed uniform expression of *pax2a* reporter, we could identify a small

population of *pax2a^mKO2^*-Low thyrocytes (Fig. 6B – D). The *pax2a^mKO2^*-Low thyrocytes

were not segregated, but scattered throughout the gland, thereby suggesting a mixing

of the two thyrocyte sub-populations.

To quantify the proportions of *pax2a^mKO2^*-Low and -High thyrocytes, we

performed FACS analysis on *pax2a^mKO2^*; *Tg(tg:nls-EGFP)* double transgenic line (Fig.

6E – G). The *Tg(tg:nls-EGFP)* zebrafish line labels the thyrocyte population in green

fluorescence (27)*.* We restricted our analysis to the thyrocyte population by gating for

GFP+ cells in the thyroid gland (Fig. 6E). Within the thyrocyte population, the cells

displayed a normal distribution of GFP fluorescence; however, thyrocytes could be split

into two sub-populations based on the levels of *pax2a* reporter expression (Fig. 6F – G).

Specifically, 75% of thyrocytes (202 out of 268 cells) displayed *pax2a^{mKO2}*-High

fluorescence, while 25% of thyrocytes (66 out of 268 cells) displayed *pax2a^{mKO2}*-Low

fluorescence levels.

In summary, the analysis of *pax2a* knock-in line validates the identification of

thyrocyte sub-populations within our single-cell RNA-Seq. data, and clearly

demonstrates, for the first time, the presence of transcriptionally diverse sub-

populations of thyrocytes present in the thyroid gland.

Discussion

We have applied for the first time unbiased single-cell gene expression analysis

to the thyroid gland. In contrast with the mainstream view that thyrocytes constitute a

molecularly uniform population, we identify two transcriptionally distinct sub-populations

of thyrocytes. The two sub-populations differed, among other genes (Supp. Table 3), in

the expression levels of a transcription factor *pax2a* and a cysteine protease Cathepsin

B (*ctsba*) (Fig. 4B – C). Cathepsin B is particularly notable as it enables the liberation of

thyroid hormone from thyrocytes by proteolytic processing of thyroglobulin (2, 20).

We validate the heterogeneity among the thyrocytes using a newly generated

knock-in reporter line for *pax2a* gene (Fig. 5). The knock-in reporter line was generated

using CRISPR/Cas9-based insertion of mKO2 fluorescent protein in the endogenous

pax2a genomic location. The *pax2a* knock-in line faithfully recapitulates the embryonic

expression of *pax2a* gene (Supp. Movie 1, Fig. 5B – F). Using the *pax2a* reporter line to

characterize the adult thyroid gland, we demonstrate the presence of *pax2a*^{mKO2}-Low

thyrocytes in the follicles (Fig. 6). Notably, *pax2a*^{mKO2}-Low and *pax2a*^{mKO2}-High

thyrocytes are present in the same follicle (Fig. 6C – D), raising the possibility of

contact-mediated interactions between the two sub-populations. It would be of interest

to build on this study and investigate the functional and replicative differences among

the two sub-populations of thyrocytes.

Our single-cell transcriptomics atlas provides a comprehensive genomics

resource to study the zebrafish thyroid gland in unprecedented detail. We performed

unbiased profiling of the thyroid gland, without enrichment for a specific cell-type. This

allowed us to capture yet poorly characterized cell-populations within the thyroid gland.

Specifically, we provide the molecular characteristics of the stromal tissue present in the

zebrafish thyroid gland. The stromal cells display enrichment of extra-cellular matrix

(ECM) related genes (Supp. Table 1) and are possibly homologous to the mesenchymal

connective septa found in the mammalian thyroid gland. The connective septa helps

cluster the thyroid follicles into lobules. Notably, the expression of fgf ligands from the

mesenchymal septa cells has been implicated in lobe formation during mouse thyroid

gland development (28). It would be of interest to test if similar morphological clustering

of the thyroid follicles exists in zebrafish, and the role the stromal cells play during

development and growth of the thyrocytes.

Our atlas further identifies a non-follicular epithelial (NFE) cell-population present

in the zebrafish thyroid gland. It is interesting to note that epithelial cells apart from

follicular and parafollicular cells have been observed in the mammalian thyroid gland. In

a report from E. Baber published in 1876 (29), histological examination of the dog

thyroid gland displayed the presence of cells "beside the stroma, lymphatics, blood

vessels, & cells between the vesicles". Dr. Baber labeled the cells as 'parenchyma', and

noted the existence of "numerous cells differing markedly in size and shape from the

epithelial cells amongst which they lie" (29). Recently, epithelial cells have been

reported in a structure called the Solid Cell Nests (SCN) of the thyroid (30). SCN are

lumen containing irregular structures located between the thyroid lobes in mammals.

SCN contain two types of epithelial cells: main cells and C-cells, expressing TP63 and

calcitonin respectively (31). Notably, the NFE cells we identified in the zebrafish thyroid

gland are marked with TP63 expression (Fig. 2G), raising the possibility of their

homology with the main cells of the SCN. C-cells, however, exist outside the thyroid

gland in zebrafish (3), and thus it is unlikely that the NFE cells we have identified would

be related to cells of parafollicular origin. Currently, the developmental origin of NFE

cells and their role in thyroid gland is unclear. To study the cell-population, transgenic

zebrafish reagents driving expression using the *tp63* regulatory region (32) could be

utilized in future.

To survey the communication between thyrocytes, the functional unit of the

thyroid gland, and the other cell-types present in the thyroid gland, we constructed a

cellular interaction network (Fig. 3B). The network was built by matching the expression

of ligands in the diverse cell-types with the expression of receptor in the thyrocytes

(Supp. Table 2) (19). Based on literature survey, we manually identified multiple

interacting genes that have been implicated in thyroid diseases. For instance, the ligand

Decorin (*dcn*) is expressed by the stromal cells and musculature, with its receptor, MET,

present on thyrocytes (Fig. 3C). Decorin, a secreted proteoglycan, is considered a

"guardian from the matrix" (33), as it is an antagonist of growth factor signaling.

Importantly, Decorin expression has been reported to be downregulated in thyroid

cancer samples (34). Thus, stromal cells could modulate Decorin to control thyrocytes

growth. Further, interactions for CYR61 (associated with Graves' Disease (35)), LRP2 /

Megalin (involved in thyroglobulin processing (36)) and NRP2 (associated thyroid

cancer metastasis (37)) were identified (Fig. 3C). The hypothesis generated by the

theoretical ligand-receptor interaction network can be tested *in vivo* in zebrafish or *in*

vitro by manipulation of thyrocytes in thyroid organoid models (38) to gain valuable

insight into thyroid gland homeostasis.

The current atlas is restricted to healthy adolescent and adult thyroid gland. In

future, it would be of interest to extend the atlas by including single-cell transcriptomics

from embryonic and old fish, providing a comprehensive resource for development,

homeostasis and aging of the thyroid gland. It would be of further interest to profile

zebrafish models of thyroid disorder (39, 40) to understand the cellular and molecular

changes underlying organ dysfunction. Combined with the power of CRISPR/Cas9

based screen that we have established for the thyroid gland (27), this resource will

provide a roadmap for the functional elucidation of cell type specific programs during

thyroid gland growth and homeostasis.

In summary, our work provides the first molecular map of the zebrafish thyroid

gland at cellular resolution. The atlas contains the molecular characterization of the

thyroid gland stromal population, identification of non-follicular epithelial cells, and

demonstrate the transcriptional heterogeneity among zebrafish thyrocytes. Further, by

constructing cell-cell communication network, the atlas provides clues into tissue

dynamics present within the organ. Finally, the dataset has been made available for

browsing via an interactive webtool (https://sumeet.shinyapps.io/zfthyroid/). We hope

that our efforts will expand the understanding of thyrocytes beyond a nominally

homogenous endocrine cell population; providing a complex picture of the diversity in

thyrocyte identity and function.

Methods

Zebrafish strains and husbandry

Wild-type or transgenic zebrafish of the outbred AB, WIK, or a hybrid WIK/AB strain

were used in all experiments. Zebrafish were raised under standard conditions at 28 °C.

Animals were chosen at random for all experiments. Published transgenic strains used

in this study were *Tg(tg:nls-mVenus-T2A-NTR)* (7) and *Tg(tg:nls-GFP)* (27).

Experiments with *Tg(tg:nls-mVenus-T2A-NTR)* were conducted in accordance with the

Animal Welfare Act and with permission of the Landesdirektion Sachsen, Germany

(DD24-5131/346/11, DD24-5131/346/12, DD24.1-5131/476/2, TVV21/2018 and all

corresponding amendments). Zebrafish husbandry and experiments with *Tg(tg:nls-*

GFP) and *pax2a* knock-in line were performed under standard conditions in accordance

with institutional (Université Libre de Bruxelles (ULB)) and national ethical and animal

welfare guidelines and regulation, which were approved by the ethical committee for

animal welfare (CEBEA) from the Université Libre de Bruxelles (protocols 578N-579N).

Single cell suspension of zebrafish thyroid gland

Single cell suspension of zebrafish thyroid gland was performed by adapting the cell

dissociation protocol outlined in Singh et al., Scientific Reports, 2018 (41). In brief, the

thyroid gland was collected and dissociated into single cells by incubation in TrypLE

(ThermoFisher, 12563029) with 0.1% Pluronic F-68 (ThermoFisher, 24040032) at 37 °C

in a benchtop shaker set at 450 rpm for 45 min. Following dissociation, TrypLE was

inactivated with 10% FBS, and the cells pelleted by centrifugation at 500g for 10min at 4

°C. The supernatant was carefully discarded and the pellet re-suspended in 500 uL of

HBSS (without Ca, Mg) + 0.1% Pluronic F-68. To remove debris, the solution was

passed over a 30 µm cell filter (Miltenyi Biotec, 130-041-407). To remove dead cells,

calcein violet (ThermoFisher, C34858) was added at a final concentration of 1 µM and

the cell suspension incubated at room temperature for 20 minutes. The single cell

preparation was sorted with the appropriate gates, including excitation with UV (405 nm)

laser for identification of alive cells (calcein+). FACS was performed through 100 µm

nozzle.

Single-cell profiling of the zebrafish thyoid gland

For single-cell RNA-seq of the zebrafish thyroid gland using the 10X Genomics

platform, cell suspension was prepared as mentioned above from the thyroid glands of

six 2 month post fertilization and six 8 month post-fertilization *Tg(tg:nls-mVenus-T2A-*

NTR) animals. The cell suspension was adjusted with Hanks' Balanced Salt Solution

(without calcium and magnesium) to a density of 800cells/µl, and diluted with nuclease-

free water according to the manufacturer's instructions to yield 12,000 cells.

Subsequently, the cells were carefully mixed with reverse transcription mix before

loading the cells on the 10X Genomics Chromium system (10). After the gel emulsion

bead suspension underwent the reverse transcription reaction, emulsion was broken

and DNA purified using Silane beads. The complementary DNA was amplified with 10

cycles, following the guidelines of the 10x Genomics user manual. The 10X Genomics

single cell RNA-seq library preparation—involving fragmentation, dA tailing, adapter

ligation and indexing PCR—was performed based on the manufacturer's protocol. After

quantification, the libraries were sequenced on an Illumina NextSeq 550 machine using

a HighOutput flowcell in paired-end mode (R1: 26 cycles; I1: 8 cycles; R2: 57 cycles),

thus generating ~45mio fragments. The raw sequencing data were then processed with

the 'count' command of the Cell Ranger software (v.2.1.0) provided by 10x Genomics

with the option '–expect-cells' set to 10,000 (all other options were used as per default).

To build the reference for Cell Ranger, zebrafish genome (GRCz10) as well as gene

annotation (Ensembl 87) were downloaded from Ensembl and the annotation was

filtered with the 'mkgtf' command of Cell Ranger (options: '–

attribute=gene_biotype:protein_coding– attribute=gene_biotype:lincRNA –

attribute=gene_biotype:antisense'). Genome sequence and filtered annotation were

then used as input to the 'mkref' command of Cell Ranger to build the appropriate Cell

Ranger Reference.

Analysis of single-cell RNA-Seq. of the zebrafish thyroid gland

The raw data generated from 10x Chromium pipeline was clustered using Seurat 2.3.4

(42) using the recommended analysis pipeline. Briefly, the raw data as UMI-counts was

log-normalized, regressed to remove the effect of library size and mitochondrial counts,

and scaled. Highly variable genes were identified for PCA analysis and graph-based

clustering. Marker genes identified for each cluster were used to classify the cell-type.

The thyrocyte cluster was isolated and sub-clustered to identify and label sub-

populations.

Data Availability

The raw 10X data, along with tabulated count data are available publicly from GEO

under accession number GSE133466. The atlas for online browsing is available at

https://sumeet.shinyapps.io/zfthyroid/.

Generation of knock-in *pax2a^pax2a-T2A-mKO2*^ zebrafish line

For generation of *pax2a* reporter line, we designed a single-guide RNA (sgRNA) targeting

the STOP codon of the *pax2a* coding sequence (GCTGCGATGGTAACTAGTGG). We

then generated a donor construct in which the sequence encoding for the monomeric

Kusabira orange (mKO2) protein was fused to a viral 2A peptide linker. This reporter

cassette was flanked by left (1000bp) and right (2000bp) homology arms of the *pax2a*

genomic DNA region around the stop codon therefore preventing the sgRNA from cutting

the donor construct. sgRNA design, production and validation were done as previously

described (27, 43). Wild-type embryos were injected with 3 nL of the injection mix

containing the sgRNA (final concentration 80 ng/µL), the donor construct (final

concentration 7.5 ng/µL), the protein Cas9 (recombinant cas protein from *S. pyogenes*

PNA Bio CP01, final concentration 100 ng/µL) and KCL (final concentration 200 mM).

Upon homologous recombination of this reporter construct in the endogenous locus,

pax2a-expressing cells were fluorescently labelled by mKO2. This *pax2a^pax2a-T2A-mKO2*^ line

is referenced as *pax2a*^mKO2^ in the text.

Generation of *pax2a* crispants

400 Somatic mutagenesis of pax2a gene was carried out exactly as mentioned in Trubiroha

401 et al., Scientific Reports, 2018 (27). Briefly, sgRNA targeting the exon 2 of *pax2a* was

402 generated as described in the publication. Following the strategy described in the

403 publication, Cas9 protein along with sgRNA was injected in one-cell stage of zebrafish

404 embryos for disruption of *pax2a* gene. Non-injected animals were used as controls.

Tissue collection

To facilitate confocal imaging of the thyroid gland, the organ was dissected from fish after fixation. Fish were killed in Tricaine prior to either direct fixation or dissection of the gland, and the samples immersed in 4% paraformaldehyde + 1% Triton-X overnight at 4 °C. The gland was then manually dissected and washed multiple times in PBS.

Immunofluorescence and image acquisition

Whole-mount immunofluorescence was performed on thyroid gland collected as described above. The collected samples were permeabilized in 1% PBT (Triton-X-100) and blocked in 4% PBTB (BSA). Primary and secondary antibody stainings were performed overnight at 4 °C. Primary antibodies used in this study were anti-PAX2A (rabbit, Genetex GTX128127) at 1:250, anti-EGFP (chicken, Abcam ab13970) at 1:1000, anti-E-Cadherin (mouse, BD bioscience cat 610181) at 1:200, anti-monomeric Kusabira-Orange 2 (mouse, MBL amalgaam M-168-3M) at 1:200 and anti-monomeric Kusabira-Orange 2 (rabbit, MBL amalgam PM051M) at 1:250. Secondary antibodies at 1:250 dilutions used in this study were Alexa Fluor 488 anti-chicken (Jackson ImmunoResearch laboratories 703-545-155), Alexa Fluor 647 anti-rabbit (Jackson ImmunoResearch laboratories 711-605-152), Alexa Fluor 647 anti-mouse (Jackson ImmunoResearch laboratories 715-605-150), Cy™3-conjugated anti-rabbit (Jackson ImmunoResearch laboratories 711-165-152) and Cy™3-conjugated anti-mouse (Jackson ImmunoResearch laboratories 715-165-150). When needed nuclei were staining using DAPI at a 1:1000 dilution. Samples were mounted in NuSieve™ GTG™ Agarose (Lonza cat50080) and imaged on a glass bottom FluoroDish™ (WPI FD3510-100) using a Zeiss LSM 780

confocal microscope or Leica DMI 6000b microscope. ImageJ was used to add scale bars

and PowerPoint was used for adding arrows and labels.

FACS-based reporter analysis

For analysing the levels of $pax2a^{mKO2}$ by FACS, single-cell suspension from the thyroid

gland of 5 mpf $Tg(tg:nls-GFP)$; $pax2a^{mKO2}$ animals was prepared as described earlier

and stained with 1 µM calcein violet (ThermoFisher, C34858). Cells were sorted and

analyzed using FACS-Aria II (BD Bioscience). Thyrocytes were selected by gating for

calcein+ GFP+ population, and mKO2 expression level recorded for analysis.

Gene Ontology (GO) Analysis

Gene ontology (GO) analysis was performed using DAVID (44). The list of genes was

uploaded on the web browser of DAVID and statistically significant (p-value < 0.05) GO

terms were identified using default parameters.

Statistical analysis

Statistical analysis was performed using R. No animals were excluded from analysis.

Blinding was not performed during analysis. Analysis of normal distribution was not

performed.

References

References

1.   Taylor PN, Albrecht D, Scholz A, Gutierrez-Buey G, Lazarus JH, Dayan CM,

Okosieme OE 2018 Global epidemiology of hyperthyroidism and hypothyroidism.

Nat Rev Endocrinol **14**:301–316.

2.   Brix K, Lemansky P, Herzog V 1996 Evidence for extracellularly acting cathepsins

mediating thyroid hormone liberation in thyroid epithelial cells. Endocrinology

137:1963–74.

3.   Alt B, Reibe S, Feitosa NM, Elsalini OA, Wendl T, Rohr KB 2006 Analysis of origin

and growth of the thyroid gland in zebrafish. Dev Dyn **235**:1872–1883.

4.   Linehan SA, Martínez-Pomares L, da Silva RP, Gordon S 2001 Endogenous

ligands of carbohydrate recognition domains of the mannose receptor in murine

macrophages, endothelial cells and secretory cells; potential relevance to

inflammation and immunity. Eur J Immunol **31**:1857–66.

5.   Nonidez JF 1931 Innervation of the thyroid gland. II. Origin and course of the

thyroid nerves in the dog. Am J Anat **48**:299–329.

6.   Svensson V, Vento-Tormo R, Teichmann SA 2018 Exponential scaling of single-

cell RNA-seq in the past decade. Nat Protoc **13**:599–604.

7.   McMenamin SK, Bain EJ, McCann AE, Patterson LB, Eom DS, Waller ZP, Hamill

JC, Kuhlman JA, Eisen JS, Parichy DM 2014 Thyroid hormone-dependent adult

pigment cell lineage and pattern in zebrafish. Science (80-) **345**:1358–1361.

8.   AlJanahi AA, Danielsen M, Dunbar CE 2018 An Introduction to the Analysis of

Single-Cell RNA-Sequencing Data. Mol Ther Methods Clin Dev **10**:189–196.

9.   Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I,

Bialas AR, Kamitaki N, Martersteck EM, Trombetta JJ, Weitz DA, Sanes JR,

Shalek AK, Regev A, McCarroll SA 2015 Highly Parallel Genome-wide

Expression Profiling of Individual Cells Using Nanoliter Droplets. Cell **161**:1202–

1214.

10.   Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, Ziraldo SB,

Wheeler TD, McDermott GP, Zhu J, Gregory MT, Shuga J, Montesclaros L,

Underwood JG, Masquelier DA, Nishimura SY, Schnall-Levin M, Wyatt PW,

Hindson CM, Bharadwaj R, Wong A, Ness KD, Beppu LW, Deeg HJ, McFarland

C, Loeb KR, Valente WJ, Ericson NG, Stevens EA, Radich JP, Mikkelsen TS,

Hindson BJ, Bielas JH 2017 Massively parallel digital transcriptional profiling of

single cells. Nat Commun **8**:14049.

11.   Kivioja T, Vähärautio A, Karlsson K, Bonke M, Enge M, Linnarsson S, Taipale J

2011 Counting absolute numbers of molecules using unique molecular identifiers.

Nat Methods **9**:72–4.

12.   Islam S, Zeisel A, Joost S, La Manno G, Zajac P, Kasper M, Lönnerberg P,

Linnarsson S 2014 Quantitative single-cell RNA-seq with unique molecular

identifiers. Nat Methods **11**:163–6.

13.   Maaten L van der, Hinton G 2008 Visualizing data using t-SNE. J Mach Learn

Res **9**:2579–2605.

14.   Sánchez-Iranzo H, Galardi-Castilla M, Sanz-Morejón A, González-Rosa JM,

Costa R, Ernst A, Sainz de Aja J, Langa X, Mercader N 2018 Transient fibrosis

resolves via fibroblast inactivation in the regenerating zebrafish heart. Proc Natl

Acad Sci U S A **115**:4188–4193.

15.   Denton CP, Zheng B, Shiwen X, Zhang Z, Bou-Gharios G, Eberspaecher H, Black

CM, de Crombrugghe B 2001 Activation of a fibroblast-specific enhancer of the

proalpha2(I) collagen gene in tight-skin mice. Arthritis Rheum **44**:712–22.

16.   Lisse TS, Middleton LJ, Pellegrini AD, Martin PB, Spaulding EL, Lopes O, Brochu

EA, Carter E V, Waldron A, Rieger S 2016 Paclitaxel-induced epithelial damage

and ectopic MMP-13 expression promotes neurotoxicity in zebrafish. Proc Natl

Acad Sci U S A **113**:E2189-98.

17.   Reischauer S, Levesque MP, Nüsslein-Volhard C, Sonawane M 2009 Lgl2

executes its function as a tumor suppressor by regulating ErbB signaling in the

zebrafish epidermis. PLoS Genet **5**:e1000720.

18.   Barbieri CE, Pietenpol JA 2006 p63 and epithelial biology. Exp Cell Res **312**:695–

706.

19.   Ramilowski JA, Goldberg T, Harshbarger J, Kloppman E, Lizio M, Satagopam VP,

Itoh M, Kawaji H, Carninci P, Rost B, Forrest ARR 2015 A draft network of ligand-

receptor-mediated multicellular signalling in human. Nat Commun **6**:7866.

20.   Brix K, Linke M, Tepel C, Herzog V 2001 Cysteine proteinases mediate

extracellular prohormone processing in the thyroid. Biol Chem **382**:717–25.

21.   Linke M 2002 Trafficking of lysosomal cathepsin B--green fluorescent protein to

the surface of thyroid epithelial cells involves the endosomal/lysosomal

compartment. J Cell Sci **115**:4877–4889.

22.   Wendl T, Lun K, Mione M, Favor J, Brand M, Wilson SW, Rohr KB 2002 Pax2.1 is

required for the development of thyroid follicles in zebrafish. Development

129:3751–60.

23.   Porazzi P, Calebiro D, Benato F, Tiso N, Persani L 2009 Thyroid gland

514   development and function in the zebrafish model. Mol Cell Endocrinol **312**:14–23.

515   24.   Pfeffer PL, Gerster T, Lun K, Brand M, Busslinger M 1998 Characterization of

516   three novel members of the zebrafish Pax2/5/8 family: dependency of Pax5 and

517   Pax8 expression on the Pax2.1 (noi) function. Development **125**:3063–74.

518   25.   Mansouri A, Chowdhury K, Gruss P 1998 Follicular cells of the thyroid gland

519   require Pax8 gene function. Nat Genet **19**:87–90.

520   26.   Kesavan G, Chekuru A, Machate A, Brand M 2017 CRISPR/Cas9-Mediated

521   Zebrafish Knock-in as a Novel Strategy to Study Midbrain-Hindbrain Boundary

522   Development. Front Neuroanat **11**:52.

523   27.   Trubiroha A, Gillotay P, Giusti N, Gacquer D, Libert F, Lefort A, Haerlingen B, De

524   Deken X, Opitz R, Costagliola S 2018 A Rapid CRISPR/Cas-based Mutagenesis

525   Assay in Zebrafish for Identification of Genes Involved in Thyroid Morphogenesis

526   and Function. Sci Rep **8**:5647.

527   28.   Liang S, Johansson E, Barila G, Altschuler DL, Fagman H, Nilsson M 2018 A

528   branching morphogenesis program governs embryonic growth of the thyroid

529   gland. Development **145**:dev146829.

530   29.   Baber EC 1876 XXI. Contributions to the minute anatomy of the thyroid gland of

531   the dog. Philos Trans R Soc London **166**:557–568.

532   30.   Harach HR 1988 Solid cell nests of the thyroid. J Pathol **155**:191–200.

533   31.   Ríos Moreno MJ, Galera-Ruiz H, De Miguel M, López MIC, Illanes M, Galera-

534   Davidson H 2011 Inmunohistochemical profile of solid cell nest of thyroid gland.

535   Endocr Pathol **22**:35–9.

536   32.   Rasmussen JP, Sack GS, Martin SM, Sagasti A 2015 Vertebrate epidermal cells

537   are broad-specificity phagocytes that clear sensory axon debris. J Neurosci

538   **35**:559–70.

539   33.   Neill T, Schaefer L, Iozzo R V 2012 Decorin: a guardian from the matrix. Am J

540   Pathol **181**:380–7.

541   34.   Martínez-Aguilar J, Clifton-Bligh R, Molloy MP 2016 Proteomics of thyroid

542   tumours provides new insights into their molecular composition and changes

543   associated with malignancy. Sci Rep **6**:23660.

544   35.   Planck T, Shahida B, Sjögren M, Groop L, Hallengren B, Lantz M 2014

545   Association of BTG2, CYR61, ZFP36, and SCD gene polymorphisms with

546   Graves' disease and ophthalmopathy. Thyroid **24**:1156–61.

547   36.   Marinò M, McCluskey RT 2000 Megalin-mediated transcytosis of thyroglobulin by

548   thyroid cells is a calmodulin-dependent process. Thyroid **10**:461–9.

549   37.   Tu D-G, Chang W-W, Jan M-S, Tu C-W, Lu Y-C, Tai C-K 2016 Promotion of

550   metastasis of thyroid cancer cells via NRP-2-mediated induction. Oncol Lett

551   **12**:4224–4230.

552   38.   Antonica F, Kasprzyk DF, Opitz R, Iacovino M, Liao X-H, Dumitrescu AM, Refetoff

553   S, Peremans K, Manto M, Kyba M, Costagliola S 2012 Generation of functional

554   thyroid from embryonic stem cells. Nature **491**:66–71.

555   39.   Chopra K, Ishibashi S, Amaya E 2019 Zebrafish duox mutations provide a model

556   for human congenital hypothyroidism. Biol Open **8**.

557   40.   Anelli V, Villefranc JA, Chhangawala S, Martinez-McFaline R, Riva E, Nguyen A,

558   Verma A, Bareja R, Chen Z, Scognamiglio T, Elemento O, Houvras Y 2017

Oncogenic BRAF disrupts thyroid morphogenesis and function via twist

expression. Elife **6**.

41.   Singh SP, Janjuha S, Chaudhuri S, Reinhardt S, Kränkel A, Dietz S, Eugster A,

Bilgin H, Korkmaz S, Zararsız G, Ninov N, Reid JE 2018 Machine learning based

classification of cells into chronological stages using single-cell transcriptomics.

Sci Rep **8**:17156.

42.   Butler A, Hoffman P, Smibert P, Papalexi E, Satija R 2018 Integrating single-cell

transcriptomic data across different conditions, technologies, and species. Nat

Biotechnol doi: 10.1101/164889.

43.   Varshney GK, Pei W, LaFave MC, Idol J, Xu L, Gallardo V, Carrington B, Bishop

K, Jones M, Li M, Harper U, Huang SC, Prakash A, Chen W, Sood R, Ledin J,

Burgess SM 2015 High-throughput gene targeting and phenotyping in zebrafish

using CRISPR/Cas9. Genome Res **25**:1030–42.

44.   Huang D, Sherman BT, Tan Q, Collins JR, Alvord WG, Roayaei J, Stephens R,

Baseler MW, Lane HC, Lempicki RA 2007 The DAVID Gene Functional

Classification Tool: a novel biological module-centric algorithm to functionally

analyze large gene lists. Genome Biol **8**:R183.

Acknowledgements

We thank members of the Costagliola and Singh lab for comments on the manuscript,

members of Center for Regenerative Therapies Dresden (CRTD) fish, FACS and

sequencing facility, and members of IRIBHM fish facility for technical assistance. We

thank J.-M. Vanderwinden from the Light Microscopy Facility and Christine Dubois from

the FACS facility for technical assistance at ULB. We are grateful to Priyanka Oberoi for

illustrations. P.G is Fund for Research in the Industry and the Agriculture (FRIA)

Research fellow; M.S. is FNRS Research Fellow (34985615 - THYSCEFA); S.C. is

FNRS Senior Research Associate. V.D. acknowledges grants from the Fond Naets

(J1813300), the Fondation Contre le Cancer (2016-093) and FNRS (EQP/OL

U.N019.19, J006120F). Work by M.B., C.L. and G.K. was supported by grants to M.B.

from the Deutsche Forschungsgemeinschaft and European Union (European Research

Council AdG Zf-BrainReg). Work by S.P.S. was supported by MISU funding from the

FNRS (34772792 – SCHISM). This work was supported by grants from the Belgian

National Fund for Scientific Research (FNRS) (FRSM 3-4598-12; CDR-J.0145.16, GEQ

U.G030.19), the Fonds d'Encouragement à la Recherche de l'Université Libre de

Bruxelles (FER-ULB).

Author contribution

S.P.S. conceptualized the project. G.K., C.L. and M.B. provided animals for single-cell

RNA-Sequencing. S.P.S., S.R., A.K., J.B., and A.P. performed the single-cell RNA-

Sequencing. S.P.S., S.E.E., V.D., S.C. analysed and interpreted the data. S.P.S.

developed the online browser. P.G. generated the *pax2a* knock-in line. P.G. and M.S.

analysed the *pax2a* reporter line. S.P.S. wrote the first draft and P.G. edited the

manuscript. S.P.S. acquired funding for the project. All authors read and approved the

final manuscript.

Conflict of interest

The authors declare no competing interests.

 Figure Legends

Figure 1: Single-cell RNA-Seq. of the zebrafish thyroid gland

(A – B) A brightfield image showing the zebrafish thyroid gland along with surrounding

tissue. The thyroid gland surrounds the ventral aorta, which extends from the outflow

tract of the heart into the gills towards the lower jaw. The thyroid follicular cells, or

thyrocytes, are labeled in green in the *Tg(tg:nls-mVenus-NTR)* transgenic line (B'). **(C)**

A t-SNE plot displaying the 6249 single-cells profiled in the zebrafish thyroid gland atlas.

The colors represent cell clusters denoting a specific cell-type. **(D – F)** Cluster #1

represents the thyrocytes that express *tg*, *slc5a5* (NIS) and *tpo*.

Figure 2: Gene expression signature of the different cell-types in the zebrafish

thyroid gland

(A) Heatmap depicting five genes specifically expressed in each one of the seven

clusters of the zebrafish thyroid gland atlas. **(B – G)** t-SNE plots overlaid with the

expression of a gene specific to each of the cluster. The endothelium cluster (cluster #2)

is a mix of blood vessels and lymphatic vessels (see Supp. Fig. 4), while the immune

cell cluster (cluster #4) is a mix of macrophages, neutrophils and lymphocytes (see

Supp. Fig. 5).

Figure 3: Connectome of the zebrafish thyroid gland identifies a dense

intercellular signaling network

(A) To build a connectome for the atlas, the ligands expressed specifically in each cell-

type were matched with their corresponding receptors in the thyrocytes. **(B)** A highly

connected intercellular interaction network is identified by the connectome. The number

of ligand-receptor pairs identified between two cell-types is denoted alongside the

arrows. **(C)** A dotplot depicting examples of paracrine and autocrine signaling in the

thyroid gland. The dots represent expression level in the different cell-types of the atlas.

Figure 4: Thyrocytes can be subdivided into two transcriptionally distinct sub-

populations

(A) Unsupervised clustering of the thyrocyte population identifies two sub-populations.

(B) Heatmap depicting the top ten most differentially expressed genes between the two

sub-populations. **(C)** Violin plots depicting the expression levels of three specific genes

in each sub-populations. Y-axis represents scaled data.

Figure 5: *pax2a^mKO2* knock-in line faithfully reports *pax2a* expression and knock-

down

(A) Schematic of the knock-in strategy used to generate the *pax2a^mKO2* line. Double

strand break was induced between the penultimate codon and the STOP codon of *pax2a*

gene using CRISPR/Cas9. DNA repair integrates the donor construct at the site of double

strand break, resulting in a *pax2a* reporter line. The donor construct contains T2A-mKO2

reporter cassette flanked by left homology (LH) and right homology (RH) arms. **(B)** Whole

mount immunofluorescence of 9.5hpf *pax2a*^mKO2 embryos stained with anti-mKO2

antibody (red) and anti-PAX2A antibody (green). Anterior is to the left, and dorsal side is

to the top. **(C)** Whole mount immunofluorescence of 55 hpf *pax2a*^mKO2; *Tg(tg:nls-EGFP)*

stained with PAX2A antibody (PAX2A-Ab) displays an overlap of mKO2 and PAX2A-Ab

signal. The otic vesicle (OV), mid-hindbrain barrier (MHB), interneurons (IN) and thyroid

gland (THY) is labelled. **(D – F)** Confocal microscopy imaging of a sagittal section of a 55

hpf *pax2a^mKO2*; *Tg(tg:nls-EGFP)* embryos showing co-localization of mKO2 and *pax2a* in

the pronephros (D), thyroid gland (E) and mid-hindbrain barrier (F). In the thyroid gland,

mKO2, PAX2A-Ab and thyrocyte-specific GFP (green) show co-localization. Scale bars:

100µm (C) and 50µm (D – F). Anterior to the right, white dashed line represents the

outline of the embryo. **(G – H)** Snapshots from live imaging of 55 hpf *pax2a^mKO2*; *Tg(tg:nls-*

EGFP) embryos injected with sgRNA targeting *pax2a* coding sequence. The anterior part

of a representative control embryo (G) is shown alongside a representative crispant (H).

Crispants display a strong reduction of mKO2 fluorescence, as well as an absence of

GFP signal suggesting absence of thyroid (THY) tissue.

Figure 6: A *pax2a* knock-in line validates the presence of thyrocyte sub-

populations

(A-D) Analysis of 3 mpf thyroid gland from *pax2a^mKO2* zebrafish shows heterogeneity in

pax2a reporter expression. **(A)** Whole mount confocal imaging of a 3 mpf *pax2a^mKO2*

thyroid labelled with mKO2 (red), E-cadherin (cyan, not shown in 'A' for clarity reasons)

and DAPI (dark blue) for nuclear localisation. **(B – D)** Optical sections of three follicles,

with mKO2-Low cells labelled with arrows. E-cadherin (B' – D') and DAPI (B'' – D'')

staining shows that absence of mKO2 signal does not correspond to an absence of cells.

Scale bars: 250 µm (A), 50 µm (B – D). Anterior to the bottom of the pictures. **(E – G)**

Cells from the thyroid gland of 5 mpf *Tg(tg:nls-GFP); pax2a^mKO2* animals were labelled

with calcein (live cell marker) and analysed using FACS. (E) A FACS plot showing calcein

on X-axis and GFP on Y-axis. The box encompassing the GFP+ cells represents the

thyrocyte population, which was gated for further analysis. (F) Histogram showing the

distribution of GFP intensity in thyrocytes. (G) Histogram showing the distribution of

mKO2 intensity in thyrocytes. Thyrocytes were selected by gating for GFP+ population.

Horizontal lines indicate the mKO2-Low and mKO2-High expression level, with

percentage values representing proportion of thyrocytes with mKO2-Low and mKO2-High

expression.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

5'UTR | *pax2a* | 3'UTR   + Cas9 +   LH | T2A | mKO2 | RH

STOP

Donor construct

Double strand break
and
Homologous recombination

5'UTR | *pax2a* | T2A | mKO2 | 3'UTR

STOP

B

*pax2a^mKO2^*

PAX2A-Ab

Merge

Merge Brightfield

C

Tg(tg:nls-EGFP)
*pax2a^mKO2^*
PAX2A-Ab

MHB
IN
OV
Eye
Yolk
THY
Heart

D Pronephros

E Thyroid Gland

F Mid-hindbrain boundary

G

Tg(tg:nls-EGFP)
*pax2a^mKO2^*

Yolk
THY
Eye
Heart
Control

H

Yolk
THY
Eye
Heart
pax2a Crispant

Figure 6