1

Function and evolution of B-Raf loop dynamics relevant to cancer recurrence under drug inhibition

Gregory A. Babbitt[1]*, Miranda L. Lynch[2], Matthew McCoy[3], Ernest P. Fokoue[4], and André O. Hudson[1]

[1]Thomas H. Gosnell School of Life Sciences, Rochester Institute of Technology, Rochester NY, USA 14623
[2]Hauptmann-Woodward Medical Research Institute, Buffalo NY, USA 14203
[3]Innovation Center for Biomedical Informatics, Georgetown University Medical Center, Washington DC, USA 20007
[4]School of Mathematical Sciences, Rochester Institute of Technology, Rochester NY, USA 14623

Contact Information –Gregory A. Babbitt (gabsbi@rit.edu)

Short title – cancer recurrence and protein dynamics

Abstract

Oncogenic mutations in the kinase domain of the B-Raf protein have long been associated with cancers involving the RAF-MEK-ERK pathway. One constitutive ERK activating mutation in B-Raf, the V600E (valine to glutamate) replacement occurring adjacent to a site of threonine phosphorylation (T599) occurs in many types of cancer, and in a large percentage of certain cancers, such as melanoma. Because natural ATP binding activity and the V600E mutation are both known to alter the physical behavior of the activation loop in the ATP binding domain, this system is especially amenable to comparative functional analyses using molecular dynamics of drug classes and genetic variants at all-atom resolution. Here, we employ machine learning enabled identification of functionally conserved protein dynamics to compare how disease pertinent molecular variations impact conserved loop dynamics during the binding of four ATP competitive B-Raf inhibitors (sorafenib, vemurafenib, dabrafenib, and PLX7904) that differ in their effectiveness in preventing cancer recurrence. We demonstrate that drug development targeting B-Raf has progressively moved towards ATP competitive inhibitors that demonstrate less tendency to mimic the functional dynamic shifts associated with ATP binding in this domain. We argue this functional dynamic mimicry in first generation B-Raf inhibitors increases the side effect of hyperactivation (i.e. inducing MAPK activation in non-tumorous cells in the absence of secondary mutation). Within the context of the binding interaction of each inhibitor, we compare the functional dynamic impacts of V600E and other sensitizing and drug resistance causing mutations in the activation and P-loops, confirming sites of low mutational tolerance in these two functional regions. Lastly, we investigate V600E sensitivity of B-Raf loop dynamics in an evolutionary context, demonstrating that while it probably shares partial origin with its early evolution in primitive eukaryotes, the functional sensitivity to V600E was secondarily increased during early jawed vertebrate evolution.

Author Summary

Our lab has recently developed a combined simulation-based and machine learning method and user-friendly software for identifying dynamic motions in proteins that are preserved over evolutionary time scales (i.e. are functionally conserved). Our method also can assess how genetic or chemical variation affects these functionally conserved motions. Here, we utilize this approach to computationally demonstrate how one of the most common mutations that arise in tumors, the V600E B-Raf protein mutation, disrupts the functioning of protein loop dynamics responsible for normal regulation of the oncogenic B-Raf protein and its growth signaling pathway. We demonstrate that cancer recurrence in the absence of new mutations may be caused by drug interactions impacting these dynamics. We propose that mimicry of the functional dynamics of natural chemical activators (i.e. agonists) by many cancer targeting drugs may be a potent causal factor in recurrence of cancer in the absence of secondary mutation. By combining comparative modeling of molecular dynamics and analyses of molecular evolution, we also demonstrate that the unusual sensitivity of B-Raf protein regulatory loop dynamics to this mutation has an ancient evolutionary origin.

Keywords – oncogene, mutation, B-Raf, inhibitor, molecular dynamics, molecular evolution, machine learning, dynamic mimicry

Introduction

From both a patient's and doctor's perspective, perhaps the most dreaded aspect of many cancers are their ability to undergo recurrence following therapeutic intervention. The recurrence of cancer can have obvious negative impact on patient psychology [1,2], doctor-patient relations [3] and questions the very concept of a 'cure' for cancer. The molecular mechanisms and somatic evolutionary processes that underlie the phenomena of recurrence are complex and do not act singularly within any given tumor, thereby greatly exacerbating the clinical difficulty of treatment and post-treatment monitoring. There are currently three molecular mechanistic frameworks for potentially explaining cancer recurrence. The traditional view of 'mutational deregulation' invokes a progressive molecular evolutionary Fisherian or 'runaway' process whereby tumor progression begins with one or several 'driver mutations' that start a cascade of secondary mutation and subsequent deregulation of growth and proliferation followed by eventual chromosomal disruption through aneuploidy and chromothrypsis [4]. Under this conceptual framework, cancer progression is largely due to a mutational process that progressively deregulates cell proliferation and frames cancer recurrence as a problem analogous to a 'whack a mole' game that gets more difficult as the game progresses. A second molecular evolutionary perspective that has gained recent popularity invokes Darwinian natural selection at the somatic level of the tumor microenvironment [5,6]. Under this view, the cancer treatment regime, in creating a local tumor microenvironment that is selectively detrimental to the survival and proliferation of cancer cells, is thought to create a selective microenvironment that favors more robust and treatment-resistant cancers that can recur if therapy is not modulated accordingly. Cancer recurrence under this new paradigm is framed as a game of strategy like chess, where the competitive environment changes fluidly with the progression of the game. Both of these conceptual frameworks for our understanding of cancer recurrence rely upon disruption of mutation-selection balance at a tissue level [7,8] acting through the complementary mechanisms of somatic mutation and apoptosis [9].

A third and less common cause of cancer recurrence, known as the 'hyperactivation paradox' does not rely upon disruption of selection-mutation balance and can be invoked to explain cancer recurrence in the absence of secondary mutations. Hyperactivation is observed when some cancer inhibitors have opposing effects in tumor and normal wildtype cells, repressing growth pathways in tumor tissues while paradoxically activating them in normal healthy tissue [10–12]. The mechanism of hyperactivation may be less of a paradox when one considers that small molecule inhibitors are often targeting the binding sites of natural agonists (i.e. signal pathway activators) in both tumor and wildtype cells. In growth activated tumor cells, these drugs will competitively bind the target protein, thus blocking the functional role of the natural agonist in the pathway. However, in growth deactivated cells, these drugs might mimic certain functional aspects of the agonist, allowing partial activation of growth pathways in the presence of the drugs. While not a 'game' that most would choose to play, this effect of functional mimicry in some drugs might be analogous to 'poking a sleeping bear', where cancer recurrence can be caused in the absence of secondary mutation by a drug's unintended signal activation of the growth pathway in nearby normal cells.

Proteins involved in growth signaling pathways, such as mitogen-activeted protein kinases (MAPK), are often very labile in structure, involving highly dynamic regions of intrinsic disorder (e.g. activation loops) that facilitate complex dynamic shifts between functional states that ultimately govern interactions with other proteins on the pathway [13,14]. Mutations that potentially alter function in

highly dynamic loop regions are often implicated in many cancers. A well-studied oncogenic system
exists in the ATP binding domain of the B-Raf protein, a serine/threonine-protein kinase that activates
the MAPK (i.e. RAF-MEK-ERK) growth pathway (Figure 1). B-Raf is a member of the Raf family of
serine/threonine kinases, along with A-Raf and C-Raf, and is the member with the highest basal activity
of the three [15]. B-Raf is a 766 amino acid protein with gene on chromosome 7 (7q34). There are 3
primary conserved domains for B-Raf, CR1, 2, and 3. CR3 (residues 443-735) is the domain that contains
the enzymatic ATP kinase domain (KD, residues 457-714). B-Raf regulation functions primarily by the
changes in the position and dynamics in several loop regions upon interaction of ATP with the
hydrophobic binding pocket formed by V471, A481, L514, T529, W531, and C532. Within the KD, the
glycine-rich P-loop (ATP-phosphate stabilizing loop) spans residues 464-471 and functions to anchor ATP
during enzymatic activity. The activation segment or activation loop (residues 594-623) within the KD is
partially disordered and contains the serine and threonine phosphorylation sites (S602 and T599). Thus,
the loop regions primarily affected by ATP binding interaction are the ATP phosphate stabilizing P-loop
at residues 464-471, the catalytic loop at residues 574-581 that captures and transfers the ATP γ-
phosphate, and the activation loop segment and DFG motif at residues 594-623 which breaks
hydrophobic interactions with the P-loop (i.e. activating the KD) with negative charge during
phosphorylation of T599. Activating mutations in the KD can often lead to disrupted conformation and
constitutive ERK activation and many mutations in this region are associated with loss of regulatory
function and/or abnormal dimerization [16]. In particular, a V600E mutation in the activation loop
causing the constitutive activation of the B-Raf signaling is one of the most common mutations in many
type of cancer, as well as a high percentage of certain cancers like melanoma. B-Raf mutants appear in
multiple cancers, notably in melanoma, and colorectal, but also less frequently in non-small cell lung
cancer, breast, ovarian, and several others [14,17,18]. Greater than 50% melanomas harbor B-Raf
activating mutations, and of these 90% involve the three bases of codon 600, and of these 90% are the
single nucleotide missense mutation at position 1799 resulting in V600E substitution (although V600 K,
D, R have also been less commonly observed [19]).

Fig 1. Overview of role of BRAF in mitogen-activated protein kinase pathway (MAPK) pathway.
Growth factor/s binds to the receptor (EGFR) on the cell surface which leads to phosphorylation and
subsequent activation. The scaffold proteins (GRB2 and SOS) facilitate the removal of GDP from RAS. RAS
then binds GTP which leads to activation. Activated RAS undergoes a conformational change which leads
to the phosphorylation and activation of BRAF, MEK and ERK. Activated ERK is shuttled to the nucleus
where it is involved in recruiting transcription factors involved in the up regulation of genes involved in
cell proliferation, cell survival, cell motility and cell invasion. The BRAF V600E mutation allows for
constitutive activation of BRAF to facilitate downstream signaling through MEK and ERK circumventing
upstream activation/regulation.

Several first generation B-Raf inhibitors designed to target V600E mutants cells have been
recently discovered to induce hyperactivation in normal wildtype cells in the absence of V600E [10–12].
Now, a new generation of B-Raf inhibitors that break the hyperactivation paradox are currently under
development and clinical trial [20]. Raf inhibitors are generally designed to occupy the ATP binding
pocket preventing phosphorylation. Why this region suffers from such low mutational tolerance, and
why some Raf inhibitors prove vulnerable to hyperactivation while others do not, are currently open
questions. A comparative analysis of the all-atom molecular dynamics (MD) of the B-Raf KD in both its
wildtype and various mutant states during interaction with small molecule variants of this drug class

could prove illuminating with regard to the problems of B-Raf hyperactivation and mutational tolerance
more generally. Three B-Raf inhibitors are currently in use. These include sorafenib, a potent
hyperactivation inducer in normal cells, [20](Zhang et al., 2015)[20][20][20][20][20] (Zhang et al.,
2015)[20][20][20](Zhang et al., 2015)as well as dabrafenib and vemurafenib, two more modern
inhibitors that also show potential problems with hyperactivation [14,20,21]. PLX7904, a modern
hyperactivation paradox breaker is currently in preclinical development [20] while its analogue PLX8394
is in clinical trials. Here, we employ machine learning enhanced comparative analyses of MD simulation
[22,23] to identify functionally conserved loop dynamics connected to normal ATP binding. We examine
how the V600E mutation breaks the natural regulatory loop dynamics of the B-Raf protein. We compare
the ability of each of these four B-Raf inhibitors (see Appendix A) to mimic functionally conserved loop
dynamics (i.e. creating functional dynamic mimicry (FDM) in their B-Raf targets), and we quantify the
impacts of known sensitizing and drug resistant mutations on these dynamics. We also examine the
V600E sensitivity of loop dynamics in two ancient phylogenetically reconstructed protein models, so as
to determine what evolutionary process may have led to unusual functional vulnerability of this
important signaling protein to the V600E mutation.

172   Results

Functionally conserved B-Raf loop dynamics upon ATP binding in wildtype background

The comparative analysis of wildtype B-Raf dynamics in both its ATP-bound and unbound forms (via
DROIDS 3.0) revealed upon binding, a significant dampening of atom fluctuation (i.e. decreased rmsf or
negative dFLUX defined by the KL divergence in rmsf) in the ATP binding pocket associated with highly
pronounced increase in fluctuation in the region of the activation loop as well as upstream toward the C
terminal approximately affecting residues 599-625 (Figure 2A-C). Shifts in dynamics were significant
according to multiple test corrected Kolmogorov-Smirnov tests at four major regions (Figure 2D)
including the binding pocket, the catalytic + activation loops and at two regions centered at residue 550
and 660. When color mapped to the ATP bound structure, these dynamics exhibit large separation
between the activation loop and P-loop (i.e. open active conformation for the protein) (Figure 2C).
When trained to recognize binding interaction dynamics and later deployed upon MD validation runs for
ATP bound wildtype B-Raf, our maxDemon 1.0 stacked machine learning model identified strong
canonical correlations in learning performance indicating conserved binding dynamics across the regions
connecting the ATP binding pocket to both the tip of the P-loop and a large region surrounding the
catalytic and activation loops (Figure 2E-F). High R values in these regions indicate that the learning
algorithms identified dynamics characteristic of B-Raf's interaction with ATP over 65% of the total time
of simulation, in spite of the thermal fluctuation that is always present. The plots of local learning
performance of each algorithm in each of the two validation runs (Supplemental Figure 1) also indicates
that characteristic functional B-Raf binding dynamics corresponded well with the known functional
regions of the protein as well as the C terminal region approximately from residue 625-675.

Fig 2. Comparative molecular dynamic analysis of ATP binding interaction and subsequent machine
learning derived Raf inhibitor impacts in normal wild type human B-Raf kinase domain (PDB ID =
1uwh). (A) Average atom fluctuation over time (i.e. RMSF or FLUX) for two ensembles of 300 x 0.5ns
molecular dynamic simulations of inhibitor bound and unbound structures are shown and statistically
compared. (B) Signed symmetric Kullback-Leibler (KL) divergences in local RMSF distributions (i.e. dFLUX)
from each amino acid backbone are plotted and (C) shown color mapped on the protein structure with

red indicating amplified fluctuation and blue indicating stabilized regions. Colored bars in B and circled
regions in D indicate the P-loop (red), ATP binding pocket (blue), catalytic loop (green) and a highly
flexible activation loop (yellow). Yellow arrows also indicate location of the activation loop in B, D and E
highlighting a highly pronounced increase in flexibility of this region during ATP binding. P values from a
Benjamini-Hochberg corrected Kolmogorov-Smirnov (KS) test are also shown in (D) indicating significant
differences in dynamics between bound and unbound structures. (E-F) The machine learning model
identifies functionally conserved (i.e. correlated) molecular dynamics that clearly connect the ATP
binding pocket (blue arrow; ATP structure is tan) to the activation loop (yellow arrow), p loop (red arrow)
and surrounding structures (E-F). Molecular dynamic impacts of the binding interaction of 4 different Raf
inhibitors (sorafenib, vemurafenib, dabrafenib and PLX7904) are summarized over both (G) the entire
protein and functionally conserved regions (supplemental figure 1). The local impacts on dynamics and
learning performance profiles (supplemental figure 2) upon which the canonical correlation between the
two validation sets is derived are also given (supplemental figure 3).

Supplemental Figure 1. *Raw local machine learning performance used to derive canonical correlations*
for detecting functionally conserved dynamics are shown for comparison in both (A) wildtype and (B)
V600E mutant backgrounds.   Two validation runs are plotted (red and blue) using all seven machine
learners. The greater correlation in A compared to B can be seen, indicating the effect of V600E in
destroying the functional dynamics of B-raf that is normally responsive to ATP binding.

Effects of Raf inhibitors on conserved B-Raf loop dynamics in wildtype background

When we analyzed the impacts (i.e. relative entropy of the canonical correlations indicating conserved
dynamics) of binding for four Raf inhibitors relative to the normal interaction of ATP with wildtype B-Raf
(Figure 2G), we find sorafenib more closely mimics ATP in the magnitude of its dynamic impacts on
conserved regions (Supplemental Figure 2-3, globally, locally resp.) and total protein (Figure 1G) than
more modern inhibitors, vemurafenib, dabrafenib and PLX7904. The highest dynamic impact was in
PLX7904, a potent 'breaker' of the 'paradox of hyperactivation', suggestive that unlike sorafenib, its
binding to wildtype B-Raf may render B-Raf unrecognizable to the MAPK pathway. Comparative dynamic
analysis of B-Raf bound to each of the four inhibitors within wildtype backgrounds (dFLUX profiles in
Supplemental Figure 4) also indicates that sorafenib most closely mimics ATP via its interaction with the
binding pocket and subsequent amplifications of dynamics of the activation loop (i.e. see similarity
between Figure 1B and Supplemental Figure 4A). While the modern inhibitors have dFLUX profiles
somewhat similar to that of ATP, they all differ in key respects. Vemurafenib and dabrafenib also seem
to affect P loop dynamics in ways that are very different than ATP (position 460-468 in Supplemental
Figure 4B and 4C), while vemurafenib and PLX7904 exhibits more global disruption and dampening of
dynamics across the whole B-Raf binding domain (indicated by widespread negative dFLUX in
Supplemental Figure 4B and 4D).

Supplemental Figure 2. *Impacts of four Raf inhibitor drug class variants on functionally conserved ATP*
binding dynamics in wildtype B-raf protein. Additional runs with just ATP binding impact is also shown
for validation.

Supplemental Figure 3*. Local impacts of four Raf inhibitor drug class variants on functionally conserved*
ATP binding dynamics in B-raf protein compared in both (A) wildtype and (B) V600E mutant
backgrounds. Additional runs with just ATP binding impact is also shown for validation.

Supplemental Figure 4. *Comparative molecular dynamic analysis of Raf inhibitor binding interactions in*
normal wild type human B-Raf kinase domain. Inhibitors are (A) sorafenib, (B) vemurafenib, (C)
dabrafenib and (D) PLX7904 (PDB ID = 1uwh, 4rzv, 4xv2 and 4xv1). Signed symmetric Kullback-Leibler
(KL) divergences in local RMSF distributions (i.e. dFLUX) from each amino acid backbone are plotted (left)
shown color mapped on the protein structure (right) with red indicating amplified fluctuation and blue
indicating stabilized regions.

Malfunction of conserved B-Raf loop dynamics during ATP binding in V600E mutant

Comparative dynamics analysis of ATP binding interaction in the presence of the V600E mutation
exhibited clear and striking differences in dynamics compared to the wildtype background. ATP binding
not only stabilized the regions around the P-loop and ATP binding pocket, but also massively and
significantly stabilized the dynamics of the region of the catalytic + activation loop and a large region
from residue 650-680 (Figure 3 A-D) serving to freeze the activation loop into an open configuration
characteristic of the V600E mutant (Figure 3C). In this genetic background, our machine learning model
fails to identify any functionally conserved dynamics of ATP binding, except in a very small part of the
region (650-680) that freezes to hold the activation loop (Figure 3E-F). The impacts of the four B-Raf
inhibitors on protein dynamics were roughly equivalent to that of ATP in the mutant background (Figure
3G), but generally stronger than the impact of ATP and sorafenib in the wildtype background (Figure
2G). The local machine learning performance was generally poorer across the whole domain (i.e. closer
to 0.5 in Supplemental Figure 1B) indicating that dynamics played a far less important functional role in
the V600E mutant background than in wildtype. This is indicative of how V600E destroys the functional
dynamics of ATP binding in B-Raf.

Fig 3. Comparative molecular dynamic analysis of ATP binding interaction and subsequent machine
learning derived Raf inhibitor impacts in mutant V600E human B-Raf kinase domain (PDB ID = 1uwh).
(A) Average atom fluctuation over time (i.e. RMSF or FLUX) for two ensembles of 300 x 0.5ns molecular
dynamic simulations of inhibitor bound and unbound structures are shown and statistically compared.
(B) Signed symmetric Kullback-Leibler (KL) divergences in local RMSF distributions (i.e. dFLUX) from each
amino acid backbone are plotted and (C) shown color mapped on the protein structure with red
indicating amplified fluctuation and blue indicating stabilized regions. Colored bars in B and circled
regions in D indicate the P-loop (red), ATP binding pocket (blue), catalytic loop (green) and a highly
flexible activation loop (yellow). Yellow arrows also indicate location of the activation loop in B, D and E
highlighting a highly pronounced increase in flexibility of this region during ATP binding. P values from a
Benjamini-Hochberg corrected Kolmogorov-Smirnov (KS) test are also shown in (D) indicating significant
differences in dynamics between bound and unbound structures. (E-F) The machine learning model fails
to identify functionally conserved (i.e. correlated) molecular dynamics connecting the ATP binding pocket
(blue arrow; ATP structure is tan) to the activation loop (yellow arrow), p loop (red arrow) and
surrounding structures (E-F). Molecular dynamic impacts of the binding interaction of 4 different Raf
inhibitors (sorafenib, vemurafenib, dabrafenib and PLX7904) are summarized over (G) the entire protein.

Effects of Raf inhibitors on mutant B-Raf loop dynamics

Comparative dynamics analysis of the four B-Raf inhibitors in the V600E mutants (Supplemental Figure 5) indicated that sorafenib globally destabilized B-Raf upon interaction with the ATP binding pocket (Supplemental Figure 5A). This effect was larger toward the C terminal regions of the protein. In the mutant background, Vemurafenib appeared to mimic the functional effect of ATP binding similar to wildtype, but too a much larger degree, confirming a potential indication of its well-known selectivity to mutant B-Raf (Supplemental Figure 5B). Upon its interaction with the binding pocket, Dabrafenib induced strong stabilization in the area of the catalytic and activation loops, locking B-Raf into its open confirmation and perhaps enhancing its well-known effect as an ATP competitor (Supplemental Figure 5C). Alternatively, upon binding the mutant, PLX7904 exhibited large destabilizations of the P-loop, ATP binding pocket and the regions extending from the activation loop and the region of residue 650-680 that freezes the open confirmation in the absence of the drug (Supplemental Figure 5D). Close-up images of the impacts on B-Raf inhibitor binding on conserved protein dynamics in the loop regions are shown in Supplemental Figure 6. We note that PLX7904, as the paradox breaking inhibitor, shows substantially less activity near the V600 region and activating segment in general, as it was constructed to disrupt overall B-Raf structure and affect dimerization, particularly targeting the L505 position, in order to affect the helix there. In our study, we find that the region near 490-505 appear to be selectively affected by disruptions to dynamics, suggesting a dynamic mechanism for its previously observed hyperactivation paradox breaking behavior.

Supplemental Figure 5. Comparative molecular dynamic analysis of Raf inhibitor binding interactions in mutant V600E human B-Raf kinase domain. Inhibitors are (A) sorafenib, (B) vemurafenib, (C) dabrafenib and (D) PLX7904 (PDB ID = 1uwh, 4rzv, 4xv2 and 4xv1). Signed symmetric Kullback-Leibler (KL) divergences in local RMSF distributions (i.e. dFLUX) from each amino acid backbone are plotted (left) shown color mapped on the protein structure (right) with red indicating amplified fluctuation and blue indicating stabilized regions.

Supplemental Figure 6. Close-up images of the mapping of conserved protein dynamics (darker regions) affected under binding of four raf inhibitors; (A) sorafenib, (B) PLX7904, (C) vemurafenib, and (D) dabrafenib.   Arrow indicate ATP binding pocket (blue), activation loop (yellow) and P loop (red).

Biphasic behavior in dynamics of B-Raf activation loop

A series of shorter comparative MD production runs were averaged to collect 'snapshots' in time and to investigate the presence of quasi-stable dynamic states. In most cases, the dFLUX (i.e. KL divergence) profiles in these runs matched that of longer runs. However, we did occasionally encounter a less common dFLUX profile around the V600 site, indicating strongly biphasic dynamic behavior in this region. In this situation, the typical combination of dampened and normal amplification of atom fluctuation (Supplemental Figure 7A) is replaced by only strong amplification at the same site (Supplemental Figure 7B). This potentially indicated the presences of very complex biphasic, or 'switch-like' behavior of the dynamics in the region where V600E sensitivity occurs.

Supplemental Figure 7. Natural biphasic behavior of the activation loop captured by comparative molecular dynamic analysis of PLX7904 binding interaction in mutant V600E B-Raf kinase domain (PDB

ID = 4xv1). Colored circled regions indicate the P-loop (red), ATP binding pocket (blue), catalytic loop
(green) and a highly flexible activation loop (yellow). The activation loop is shown in two stable states,
(A) open conformation with and highly flexible activation loop, and (B) closed conformation with highly
dampened activation loop. NOTE: simulation runs were averaged over only 20x0.1ns production runs (i.e.
fewer sampling runs taken over shorter time periods than training sets in Figure 2 and 3).

Comparison of impacts of drug sensitivity and resistant mutations on B-Raf loop dynamics

To further investigate the effect of both genetic and drug class variation on functional B-raf loop
dynamics, we compared the dynamic impacts on B-Raf for a variety of drug sensitizing and drug
resistance linked mutations for each of the four Raf inhibitors, sorafenib, vemurafenib, dabrafenib, and
PLX7904 (Supplemental Table 1, Figures 4-7). During the interaction of hyperactivation-inducing
sorafenib with the ATP binding pocket, dynamic shifts in the activation loop are clearly evident (as large
positive and negative dFLUX values in Figure 4A) which are quite similar in effect to changes in this
region caused by ATP binding (Figure 2A). Functional dynamic shifts identified by our machine learning
model are also quite evident upon sorafenib binding (Figure 4B-C) and similarly connect the ATP binding
pocket to the outer bend of the activation loop when ATP binds (Figure 2B). The V600E mutation which
sensitizes sofafenib targeting to B-Raf, was found to greatly impact loop dynamics when compared to
validation runs with no mutation (Figure 4D) and corroborating its large impact on functional dynamics
in general (Figure 3). The three remaining drug resistant mutations (G464V, G469E, G469V) show less
impact, but are collectively larger than validation, suggesting that mutations disrupting dynamics of the
P loop are key in creating drug resistance to sorafenib (Figure 4D). In stark contrast to the sorafenib
results, the hyperactivation-breaking compound, PLX7904, while also causing similar dynamics shifts to
the activation loop (Figure 5A), fails to produce hardly any dynamic changes that are identified as
functional by our machine learning classification model (Figure 5B-C). Thus, it appears that while
PLX7904 is a potent competitive binder to mutant B-Raf, it also has the benefit of potentially disrupting
any functional dynamics that could allow stable interactions with other components of the MAPK
pathway. Our analysis of mutational impacts on functional dynamics (Figure 4D) also support this, and
suggest that changes in PLX7904 efficacy caused by mutation are likely independent of the normal
evolutionary conserved function of B-Raf. Analysis of vemurafenib (Figure 6) shows marked similarity to
sorafenib, suggesting that it is also prone to mimic the functional dynamics of ATP binding (Figure 6B-C)
but with softening of the P-loop as well (Figure 6A). A survey of 18 drug impacting mutations does not
show any clear trend except that mutational impacts are largest in regions affecting either the P-loop or
the activation loop (Figure 6D). Similar analysis of dabrafenib indicates a hardening of P-loop dynamics
(Figure 7A) and a pattern of less coordinated impact on conserved dynamics that fails to connect the
ATP binding pocket to the activation loop (Figure 7B-C), supporting our hypothesis that more modern
Raf inhibitors are less like ATP in their overall effect on functional shifts in B-Raf dynamics during
binding. Mutations affecting dabrafenib function also appear to be largely confined to P-Loop and
activation loop, with the exception of L514V, a site whose side-chain actively engages ATP during normal
B-Raf function.

Table 1. *Genetic variants affecting drug activity of four Raf inhibitors (sorafenib, vemurafenib,*
dabrafenib, PLX7904).

Fig 4. Comparative molecular dynamic analysis of hyperactivation-inducing sorafenib binding in
wildtype BRAF and subsequent mutational impacts on conserved BRAF dynamics. *Images represent*
the (A) signed symmetric Kullback-Leibler (KL) divergences in local RMSF distributions (i.e. dFLUX) color-
mapped to the BRAF structure with red indicating amplified fluctuation and blue indicating stabilized
regions and (B) color-mapped functionally conserved dynamic regions (dark gray) identified by (C)
significant canonical correlation of performance in machine learning validation runs. (D) The impacts of
four known drug sensitizing mutations (V600E, G469V, G469E, G464V) are also compared. NOTE: blue,
yellow and red arrows indicate drug binding site, activation loop and p loop (resp.)

Fig 5. Comparative molecular dynamic analysis of hyperactivation-breaking PLX7904 binding in
wildtype BRAF and subsequent mutational impacts on conserved BRAF dynamics. *Images represent the*
(A) signed symmetric Kullback-Leibler (KL) divergences in local RMSF distributions (i.e. dFLUX) color-
mapped to the BRAF structure with red indicating amplified fluctuation and blue indicating stabilized
regions and (B) color-mapped functionally conserved dynamic regions (dark gray) identified by (C)
significant canonical correlation of performance in machine learning validation runs. (D) The impacts of
three known drug sensitizing mutations (V600E, V600R, V600D) and one drug resistant mutation (G466E)
are also compared. NOTE: blue, yellow and red arrows indicate drug binding site, activation loop and p
loop (resp.)

Fig 6. Comparative molecular dynamic analysis of vemurafenib binding in wildtype BRAF and
subsequent mutational impacts on conserved BRAF dynamics. *Images represent the (A) signed*
symmetric Kullback-Leibler (KL) divergences in local RMSF distributions (i.e. dFLUX) color-mapped to the
BRAF structure with red indicating amplified fluctuation and blue indicating stabilized regions and (B)
color-mapped functionally conserved dynamic regions (dark gray) identified by (C) significant canonical
correlation of performance in machine learning validation runs. (D) The impacts of 18 known drug
sensitizing and resistant mutations are also compared. See Table 1 for classifications. NOTE: blue, yellow
and red arrows indicate drug binding site, activation loop and p loop (resp.)

Fig 7. Comparative molecular dynamic analysis of dabrafenib binding in wildtype BRAF and
subsequent mutational impacts on conserved BRAF dynamics. *Images represent the (A) signed*
symmetric Kullback-Leibler (KL) divergences in local RMSF distributions (i.e. dFLUX) color-mapped to the
BRAF structure with red indicating amplified fluctuation and blue indicating stabilized regions and (B)
color-mapped functionally conserved dynamic regions (dark gray) identified by (C) significant canonical
correlation of performance in machine learning validation runs. (D) The impacts of 10 known drug
sensitizing and resistant mutations are also compared. See Table 1 for classifications. NOTE: blue, yellow
and red arrows indicate drug binding site, activation loop and p loop (resp.)

Phylogenetic analysis of V600E sensitivity regarding B-Raf loop dynamics

A minimum evolution tree (Figure 8A) and maximum likelihood tree (supplemental Figure 8) was used to
identify protein level variants for tetrapod (= human), ray-finned fish, and insect. The wildtype genetic
backgrounds incorporating 9 amino acid replacements (fish) and 95 replacements (insect) show no
elevated dynamic impacts over the validation run (wildtype tetrapod). However, V600E/A600E mutants
both demonstrate elevated disruption of conserved ATP binding dynamics (Figure 8B) indicating that B-
Raf sensitivity to this mutation has an ancient origin. The very large increase in dynamic impact of V600E

in ray-finned fish over that of A600E in insects indicates a large increase in V600E sensitivity in early
jawed vertebrate evolution as well (Figure 8B).   Local mapping of these impacts (Figure 8C) indicates
involvement of all three functional regions (P-loop at position 13-25, ATP binding pocket at position 60-
100, and activation loop at position 153-190) in the functional evolution of this sensitivity. The dynamic
impacts of V600E and A600E in these ancient backgrounds are also shown color-mapped to the B-Raf
structure (Figure 8D and 8E resp.).

Fig 8. Comparative molecular dynamic analysis of ATP binding in wildtype BRAF orthologs. Orthology
was determined via phylogenetic analysis to derive a (A) minimum evolution tree bootstrapped 500
times using the Kimura 2 parameter model with Gamma correction for multiple hits and invariant sites.
This best model was determined via multi-model inference (BIC, AICc).   (B-C) With a goal of determining
when V600E sensitivity first arose in this phylogeny, we compare the conserved region impacts of ATP
binding on BRAF dynamics in wildtype and V600E mutated genetic backgrounds of ray-finned fish and
insect ortholog models after training on the functional ATP binding in wildtype tetrapod (i.e. human)
background. These impacts are shown both (B) globally and (C) locally, where regions of low mutational
tolerance are observed corresponding to the P-loop, ATP binding pocket, and activation loop and
adjacent downstream residues (shaded yellow). The impacts of the genetic variation of each variant is
color mapped to the wildtype and mutant backgrounds in both (D) fish and (E) insect. NOTE: in the insect
ortholog, this mutation is A600E rather than V600E. A maximum likelihood tree is also shown in
Supplemental Figure 7.

Supplemental Figure 8. A maximum likelihood tree bootstrapped 500 times using the Kimura 2
parameter model with Gamma correction for multiple hits and invariant sites. This best model was
determined via multi-model inference (BIC, AICc).   Insect outgroup is not shown to enhance readability
of the bootstrap values.

426 Discussion

Given their involvement in a variety of cellular processes, protein kinases, specifically serine/threonine
kinases play a role in all of the major cancers. Many serine/threonine kinases have altered expression in
tumors [24]. Given that this class of enzyme is a proven putative and attractive target for the
development of cancer therapies, our study of the function and evolution of the molecular dynamics of
the serine/threonine kinase B-Raf has the potential to guide future studies and experiments aimed at
elucidating efficacy and rationale design of therapeutics. Through our novel high performance
computing approach to machine learning applied to comparative functional molecular dynamics
simulations, we have demonstrated that ATP binding to the CR3 domain of B-Raf protein induces
evolutionarily functionally conserved changes in dynamics that physically connect ATP binding pocket
with both the P-loop and activation segment. The region just downstream of the mutation intolerant
valine codon 600 is especially affected in its dynamics by ATP binding remotely on the structure of the
protein KD. The V600E mutation massively disrupts these evolutionary conserved shifts in dynamics that
are obviously a key complex functional feature of BRAF signaling and MAPK regulation. Hyperactivation
inducing drugs, especially the first generation drug sorafenib, appears to mimic functionally conserved
dynamics of natural ATP binding function in the CR3 domain, inducing almost the same spatial pattern of
functionally conserved protein dynamics in our simulations as ATP itself. Vemurafenib and dabrafenib

show less of this tendency, but still induce functional dynamics in the region joining the ATP binding
pocket to the P-loop and activation loop segment of B-Raf. In contrast, the hyperactivation breaking
drug PLX7904 shows almost no such propensity for mimicry of dynamics of the ATP agonist. See close-up
comparison of the functional dynamic mimicry (FDM) of ATP caused by the four Raf inhibitors in
Supplemental Figure 6. Our survey of single amino acid replacements most affecting patient drug
sensitivity or resistance to the four inhibitors indicates that mutations impacting the three
hyperactivation inducing drugs (i.e. sorefenib, vemurafenib and dabrafenib) appear to target
functionally conserved aspects of P-loop and activation loop dynamics. This might be expected under a
somatic mutation-selection regime induced by therapeutic treatments of tumors where functional B-Raf
dynamics are still potentially in play, albeit likely subverted into a malfunctional state from the
standpoint of the patient. However, PLX7904 currently has very few known sensitizing or resistant
mutations, an observation that may reflect lack of experimentation with the new compound, but may
also prove consistent with our demonstration that it largely avoids inducing the same sort of dynamic
changes as ATP upon binding the same location. Finally, our molecular evolutionary study, indicates that
B-Raf sensitivity to V600E has a likely ancient origin, perhaps as old as the Raf kinase protein family
itself. However, the sensitivity of functionally conserved B-Raf dynamics to V600E, appears to have
greatly increased sometime during early jawed vertebrate evolution, as represented by extant
sequences across all modern ray-finned fishes. As a modulator of multicellular growth, B-Raf likely has
origins in early eukaryotes, however poor alignment quality to putative orthologs in *C. elegans* and *S.*
cerevisiae prevented us from exploring this possibility further.

Conclusion

We propose that functional dynamic mimicry (FDM), as observed here in B-Raf inhibitors, is an
important risk factor to consider in future rational drug design. This an especially important
consideration when targeting sites of natural agonists involved in signal activation in pathways like
MAPK which are typically downregulated after early embryonic development, and a subsequently
problematic in all cancers. Pharmaceuticals that avoid FDM while effectively targeting and binding active
sites of proteins in these pathways, are most likely disrupting the functional dynamics of protein
domains to the extent that they are no longer recognized by the pathways in which they normally
participate. In the case of pathways downregulated after early development, this affect is ideal, as it
destroys signal transduction. Our DROIDS 3.0 + maxDemon 1.0 computational software [22,23] can
allow for inexpensive screening against unintended effects of FDM, potentially producing better patient
products while saving pharmaceutical industry wasted time, effort and money in preclinical
development and clinical trial. We also conclude that a more evolutionary and biophysical perspective
around complex problems like FDM that are potentially linked to naturally evolved tolerances to protein
mutation are greatly needed in future biomedical research. This cannot be easily addressed by the
combination of DNA sequence-based bioinformatics and cell/animal-based experimental research alone.
Proteins are nanometer sized machines operating in the complex context of soft matter dynamics. They
are often imperfect in their interactions with other proteins and incompletely optimized in terms of
their functional evolution. The development of modern drug therapies could benefit by embracing a
more functional evolutionary perspective that is grounded in computational biophysical method.

Appendix A

Supplementary information on the four inhibitors: The four B-Raf inhibitors (BRAFi) discussed in this manuscript are all small molecule compounds that are FDA approved for use: 1. in mutated BRAF in metastatic melanoma (Vemurafinib, Dabrafenib) or 2. in other carcinomas (Sorafenib, for advanced renal cell, thyroid, and hepatocellular carcinomas) as a more general kinase inhibitor (active against BRAF, VEGFR, etc).   PLX7904 is in preclinical development.

Sorafenib (also, Nexavar, BAY 43-9006, MW: 464.8 g/mol): A member of the class of phenylureas that acts as a kinase inhibitor and angiogenesis inhibitor.   It is nonspecific, and does not distinguish between WT and mutant B-Raf forms, and has higher specificity for C-Raf compared to B-Raf.   It is known to potently induce hyperactivation, and leads to rapid development of drug resistance.

Vemurafinib (also, PLX4032; Zelboraf; MW: 489.9 g/mol): An ATP-competitive kinase inhibitor selective for B-Raf V600E.   It functions to decrease signaling through the MAPK pathway by inhibiting the kinase domain of mutant BRAF, but is known to also induce hyperactivation in WT BRAF.   Shows increased specificity for B-Raf V600E relative to B-Raf WT or C-Raf.

Dabrafenib (also, Tafinlar, GSK2118436; 519.6 g/mol): An ATP-competitive kinase inhibitor selective for B-Raf V600 (E/K/R) mutations, which acts to induce inhibition of phosphorylated extracellular signal-regulated kinase (ERK). Often given in combination with MEK inhibitor Trametinib to achieve a more favorable resistance profile.

PLX7904 (also PB04; MW: 512.5 g/mol): A selective B-Raf kinase inhibitor known to be a potent paradox-breaker. able to efficiently inhibit activation of ERK1/2 in mutant BRAF melanoma cells but does not hyperactivate ERK1/2 in mutant RAS-expressing cells.   Was designed to interact with position 505 in B-Raf, shifting the Raf dimer interface, and cause inhibition via long-range structural alterations.   Along with analogue PLX8394, is in development as BRAFi capable of uncoupling RAF blocking and MAPK pathway activation.

Models

PDB structure preparation and ligand force field modification

Structures of the ATP binding domain of B-Raf bound to various Raf inhibitors were obtained from the Protein Data Bank (PDB). These were PDB ID 1uwh (wildtype B-Raf bound to sorafenib), 4rzv (vemurafenib bound to V600E mutant), 4xv1 (PLX7904 bound to V600E mutant), 4xv2 (dabrafenib bound to V600E mutant). Loop modeling and refinement were conducted on the activation loop where needed using Modeller in UCSF Chimera [25]. Complementary mutant (V600E in 1uwh) and wildtype structures (E600V in 4rzv, 4xv1,4xv2) were computationally induced in UCSF Chimera 1.13 [25] by first swapping amino acids using optimal configurations in the Dunbrack rotamer library, then using 2000 steepest gradient descent steps of energy minimization to relax the regions around the amino acid replacement. To obtain ATP bound versions of wildtype B-raf and the V600E mutant, the sorafenib inhibitor was deleted from 1uwh and 1uwj, then ATP was docked using the AutoDock Vina program [26] after reducing both structures (adding H). ATP docking positions most closely overlapping sorafenib were chosen. Unbound forms of all the structures were obtained by deleting the ligands in the bound structure and performing energy minimization for 5000 steps. Prior to molecular dynamic simulation in

Amber18, the GAFF2 force field modifications for ATP and the small molecule ligands were generated

using sqm, a semiempirical and DFTB quantum chemistry program employed by the Antechamber

(Ambertools18) software suite [27,28].

MD simulation protocols

Large ensembles of graphic processing unit (GPU) accelerated molecular dynamic simulations were

prepared and conducted using the particle mesh Ewald method employed by pmemd.cuda in Amber18

[29,30] via the DROIDS v3.0 interface (Detecting Relative Outlier Impacts in Dynamic Simulation) [22,23].

Simulations were run on a Linux Mint 19 operating system mounting two Nvidia Titan Xp graphics

processors. Explicitly solvated protein systems were prepared using tLeAP (Ambertools18) using the

ff14SB protein force field [31] in conjunction with modified GAFF2 small molecule force field [28].

Solvation was generated using the Tip3P water model [32] in a 12nm octahedral water box and

subsequent charge neutralization with Na+ and Cl- ions. After energy minimization, heating to 300K, and

10ns equilibration, an ensemble of 200 MD production runs each lasting 0.5 ns of time were created for

both ligand bound and unbound B-Raf. Each MD production run was preceded by a single random

length short spacing run selected from a range of 0 to 0.25ns to mitigate the effect of chaos (i.e.

sensitively to initial conditions) in the driving ensemble differences in the MD production runs [22]. All

MD was conducted using an Andersen thermostat [33] under constant pressure. Root mean square

atom fluctuations (rmsf) were calculated using the atomicfluct function in CPPTRAJ [34].

Comparative dynamics of bound and unbound functional states

The symmetric Kullback-Leibler (KL) divergence or relative entropy between the distributions of atom

fluctuation (i.e. root mean square fluctuation or rmsf taken from 0.01 ns time slices of total MD

simulation time) on ligand bound and unbound B-Raf were computed using DROIDS and color mapped

to the protein backbone with individual amino acid resolution to the bound structures using a

temperature scale (i.e. where red is hot or amplified fluctuation and blue is cold or dampened

fluctuation). Details of this calculation and representation can be found in our previous publication [22].

The reference state of the protein is unbound while the query state is bound. Therefore, this pairwise

comparison is used to represent the functional impact of ligand binding on the protein's normal

unbound motion, where it is expected that ligand contacts would typically dampen the fluctuation of

atoms around the ATP binding pocket to some degree. As ATP binding acts to destabilize the activation

loop and P-loop regions of B-Raf, this analysis will allow quantification of dynamic impacts that amplify

loop motions as well.

Machine learning performance and detection of conserved dynamic function

We applied machine learning analysis to examine normal ATP binding in wildtype B-Raf. We then

compare these results to a similar analysis of ATP binding in V600E mutants to ascertain how this

mutation alters functionally conserved dynamics in the B-Raf protein system. The detection of

functionally conserved protein dynamics in wildtype B-Raf followed the machine learning-based method

outlined in our previous DROIDS 3.0 + maxDemon 1.0 software papers [22,23]. In summary, a stacked

machine learning model implementing seven classification methods (K nearest neighbors, naïve Bayes,

linear/quadratic discriminant analysis, support vector machine, random forest and adaptive boosting)

was individually trained upon each amino acid's fluctuation data that was pre-classified to the ligand

bound and the unbound dynamic states. These learners were deployed upon the identical amino acids

within two new 5ns MD simulations of the ligand bound structure and classifications of atom fluctuation
(i.e. rmsf calculated for 500 x 0.01 ns time slices) where 0 = unbound and 1 = bound states were
averaged to obtain a learning performance value for each amino acid. Average learning performance
profiles (i.e. accuracy and precision) are generated along the protein from N to C terminus. Note that a
value of 0.5 at a given position would indicate that the learning model could not differentiate between
bound and unbound dynamics at that residue. Local significant canonical correlations in learning
performance between the two validation runs on the ligand bound structure indicate where any
position-dependent molecular dynamics that is functionally conserved with regards to the binding
interaction has occurred. The correlation was calculated within a 20 residue sliding window along the
learning profile.

$$conserved_{dynamics} = significant\ (CC_{self})$$

The central concept here is that if any atom motion is specific to a given functional state (i.e. in
this case protein bound to ligand, vs unbound protein), then after training, the machine learners will be
able to successfully classify this motion when it is observed in subsequent identical validation MD runs.
When the learners are able to repeatedly identify functional state dynamics that locally associate with a
specific position on the protein, this will create correlated profiles in learning performance. As our
machine learning model is a stacked model, incorporating seven learning algorithms, it is more robust to
artifacts related to any single method, and the resulting correlation is canonical as well (see
Supplemental Figure 1 for an example of correlated and uncorrelated sets of learning profiles). We can
label the dynamics as 'functionally conserved' if the machine learning model can consistently identify
the functional dynamic states upon which it trained across multiple and independent validation runs.
Two or more identically prepared validation runs are used to examine whether statistically significant
canonical correlations (i.e. conserved dynamics) are locally identifiable by a significant local Wilk's
lambda value for a local canonical correlation coefficient.

Method of genetic and drug variant impact assessment

The local impacts of mutations and drug class variants were obtained by first deriving the canonical
correlation of the learner performance profile of a given variant to the wildtype B-Raf in its normal
functionally bound state (= $CC_{variant}$). This also uses a 20 residue sliding window. The impact metric is
generated by comparing canonical correlation of the variant ($CC_{variant}$) to the self-similar dynamic
correlation (CC_{self}) defining functionally conserved dynamics above, using the common definition of
relative entropy below

$$variant_{impact} = CC_{self} * log\frac{CC_{variant}}{CC_{self}}$$

Effectively, this impact metric captures the difference in conserved dynamics between the wildtype
ligand bound state (here ATP or one of the four Raf inhibitors) and the variant states of binding created
by genetic mutation or different drug ligand molecules. This occurs on the same scale as that used to
differentiate bound from unbound states during the earlier training of the stacked learning model. More
details regarding our methodological approach to identifying functionally conserved dynamics and
subsequent impacts of genetic and drug class variants can be obtained in our prior publication [22,23].

The relative impacts of four B-Raf inhibitors, sorafenib, dabrafenib, vemurafenib and PLX7904
were assessed after initially training the stacked machine learning model on ATP bound and unbound B-

Raf within both the wildtype and the V600E mutant genetic background. Several drug affecting genetic
mutants were also examined after training the learning model on normal functional binding interaction
of each drug in the wildtype genetic background. These drug resistant and drug sensitizing mutations
were identified using the Variant Interpretation for Cancer Consortium (VICC) Meta-Knowledgebase
[35]. Variants were selected that contained literature support for the functional impact on at least one
of the four B-Raf inhibitors (Supplemental Table 1).

Phylogenetic analysis of B-Raf sensitivity to V600E

A set of 27 BRAF ortholog sequences were obtained using the National Center for Biotechnology
Information (NCBI) ortholog finder tool. These sequences were aligned using MEGA 7.0 [36] Clustal
multiple sequence alignment tool and then manually trimmed to contain only the ATP binding domain.
Phylogenetic model selection was also conducted in MEGA to identify the most likely (best fitting)
evolutionary substitution model. This was identified via both Bayesian and Akaike Information Criterion
to be the Kimura 2-parameter model with extra terms for both gamma correction for multiple same site
substitution and correction for invariants sites (i.e. K2+G+I). A minimum evolution tree and a maximum
likelihood tree bootstrapped 500 times confirmed the same tree topology with high confidence. Amino
acid replacements distributed on this topology identified three protein variants, one each for tetrapods,
ray-finned fish and insects. Amino acids differences from tetrapods identified 9 amino acid replacements
since a common ancestor with ray-finned fish and 95 replacements since a common ancestor with
insects. Structural models for the two ancestral protein variants were manually generated in UCSF
Chimera using swapaa commands to find non-clashing Dunbrack library rotamer configurations. These
models were then subsequently relaxed with 2000 steps of energy minimization via steepest gradient
descent. A V600E variant was subsequently created from the wildtype ray-finned fish model and an
A600E variant was created from the wildtype insect model. DROIDS software was used to train upon
ATP binding interaction in the wildtype tetrapod model (comparing ATP bound and unbound dynamics
as described above). DROIDS 3.0 and maxDemon 1.0 machine learning analysis (described above) was
then used to identify functionally conserved dynamics in tetrapod B-Raf and subsequently compare
mutational impacts on B-Raf dynamics for both the ancestral wildtype and mutant backgrounds in ray-
finned fish and insects.

The author's software can be accessed at the promotional website and GitHub repository links below

https://people.rit.edu/gabsbi/

https://github.com/gbabbitt/DROIDS-3.0-comparative-protein-dynamics

Acknowledgements

636 We acknowledge Nvidia corporation for a GPU hardware grant, and the College of Science at the
637 Rochester Institute of Technology for internal grant support.

List of abbreviations

FDM - functional dynamic mimicry

MD - molecular dynamic (simulation)

GPU – graphic processor unit

PDB – Protein Data Bank

NCBI – National Center for Biotechnology Information

BRAF – B-Raf gene sequence

B-Raf – protein product of BRAF gene

ATP – adenosine triphosphate

DNA – deoxyribonucleic acid

MAPK – mitogen-activated protein kinase

BRAFi – B-Raf inhibitor

V600E – valine to glutamic acid mutation and position 600

CR1 CR2 CR3 – conserved region 1, 2, or 3

KD – kinase domain

P-loop – ATP stabilizing loop

KL divergence – Kullback-Leibler divergence

dFLUX – change in atom fluctuation or rmsf

rmsf – root mean square fluctuation

CC – canonical correlation

DROIDS – software acronym for 'detecting outlier impacts in dynamic simulation', our software package
for comparative protein dynamics simulation

maxDemon – software acronym for 'Maxwell's demon', our machine learning application for DROIDS

References

664   1.   Andersen BL, Shapiro CL, Farrar WB, Crespin T, Wells-DiGregorio S. Psychological Responses to
665   Cancer Recurrence. Cancer. 2005;104: 1540–1547. doi:10.1002/cncr.21309

666   2.   Yang H-C, Thornton LM, Shapiro CL, Andersen BL. Surviving Recurrence: Psychological and Quality-
667   of-life Recovery. Cancer. 2008;112: 1178–1187. doi:10.1002/cncr.23272

668   3.   Janz NK, Li Y, Zikmund-Fisher BJ, Jagsi R, Kurian AW, An LC, et al. The Impact of Doctor-Patient
669   Communication on Patients' Perceptions of their Risk of Breast Cancer Recurrence. Breast Cancer
670   Res Treat. 2017;161: 525–535. doi:10.1007/s10549-016-4076-5

4.   Brown A-L, Li M, Goncearenco A, Panchenko AR. Finding driver mutations in cancer: Elucidating
the role of background mutational processes. PLOS Comput Biol. 2019;15: e1006981.
doi:10.1371/journal.pcbi.1006981

5.   Gatenby RA, Zhang J, Brown JS. First Strike–Second Strike Strategies in Metastatic Cancer: Lessons
from the Evolutionary Dynamics of Extinction. Cancer Res. 2019 [cited 4 Sep 2019].
doi:10.1158/0008-5472.CAN-19-0807

6.   Zhang J, Cunningham JJ, Brown JS, Gatenby RA. Integrating evolutionary dynamics into treatment
of metastatic castrate-resistant prostate cancer. Nat Commun. 2017;8: 1816. doi:10.1038/s41467-
017-01968-5

7.   Bürger R. Mutation-Selection Models in Population Genetics and Evolutionary Game Theory. In:
Kurzhanski AB, Sigmund K, editors. Evolution and Control in Biological Systems. Springer
Netherlands; 1989. pp. 75–89.

8.   Hartl DL. Mutation-selection balance with stochastic selection. Genetics. 1977;86: 687–696.

9.   Nunney L. The population genetics of multistage carcinogenesis. Proc R Soc B Biol Sci. 2003;270:
1183–1191. doi:10.1098/rspb.2003.2351

10.   Hatzivassiliou G, Song K, Yen I, Brandhuber BJ, Anderson DJ, Alvarado R, et al. RAF inhibitors prime
wild-type RAF to activate the MAPK pathway and enhance growth. Nature. 2010;464: 431–435.
doi:10.1038/nature08833

11.   Poulikakos PI, Zhang C, Bollag G, Shokat KM, Rosen N. RAF inhibitors transactivate RAF dimers and
ERK signaling in cells with wild-type BRAF. Nature. 2010;464: 427–430. doi:10.1038/nature08902

12.   Villanueva J, Vultur A, Herlyn M. Resistance to BRAF inhibitors: Unraveling mechanisms and future
treatment options. Cancer Res. 2011;71: 7137–7140. doi:10.1158/0008-5472.CAN-11-1243

13.   Montagut C, Settleman J. Targeting the RAF-MEK-ERK pathway in cancer therapy. Cancer Lett.
2009;283: 125–134. doi:10.1016/j.canlet.2009.01.022

14.   Wan PTC, Garnett MJ, Roe SM, Lee S, Niculescu-Duvaz D, Good VM, et al. Mechanism of activation
of the RAF-ERK signaling pathway by oncogenic mutations of B-RAF. Cell. 2004;116: 855–867.

15.   Roskoski R. RAF protein-serine/threonine kinases: structure and regulation. Biochem Biophys Res
Commun. 2010;399: 313–317. doi:10.1016/j.bbrc.2010.07.092

16.   Durrant DE, Morrison DK. Targeting the Raf kinases in human cancer: the Raf dimer dilemma. Br J
Cancer. 2018;118: 3–8. doi:10.1038/bjc.2017.399

17.   Davies H, Bignell GR, Cox C, Stephens P, Edkins S, Clegg S, et al. Mutations of the BRAF gene in
human cancer. Nature. 2002;417: 949–954. doi:10.1038/nature00766

18.   Rezaei Adariani S, Buchholzer M, Akbarzadeh M, Nakhaei-Rad S, Dvorsky R, Ahmadian MR.
Structural snapshots of RAF kinase interactions. Biochem Soc Trans. 2018;46: 1393–1406.
doi:10.1042/BST20170528

19. Ascierto PA, Kirkwood JM, Grob J-J, Simeone E, Grimaldi AM, Maio M, et al. The role of BRAF V600
mutation in melanoma. J Transl Med. 2012;10: 85. doi:10.1186/1479-5876-10-85

20. Zhang C, Spevak W, Zhang Y, Burton EA, Ma Y, Habets G, et al. RAF inhibitors that evade
paradoxical MAPK pathway activation. Nature. 2015;526: 583–586. doi:10.1038/nature14982

21. Karoulia Z, Wu Y, Ahmed TA, Xin Q, Bollard J, Krepler C, et al. An Integrated Model of RAF Inhibitor
Action Predicts Inhibitor Activity against Oncogenic BRAF Signaling. Cancer Cell. 2016;30: 485–498.
doi:10.1016/j.ccell.2016.06.024

22. Babbitt GA, Mortensen JS, Coppola EE, Adams LE, Liao JK. DROIDS 1.20: A GUI-Based Pipeline
for GPU-Accelerated Comparative Protein Dynamics. Biophys J. 2018;114: 1009–1017.
doi:10.1016/j.bpj.2018.01.020

23. Babbitt GA, Fokoue EP, Evans JR, Diller KI, Adams LE. DROIDS 3.0 - detecting genetic and drug class
variant impact on conserved protein binding dynamics. Biophys J. 2019;0.
doi:10.1016/j.bpj.2019.12.008

24. Capra M, Nuciforo PG, Confalonieri S, Quarto M, Bianchi M, Nebuloni M, et al. Frequent alterations
in the expression of serine/threonine kinases in human cancers. Cancer Res. 2006;66: 8147–8154.
doi:10.1158/0008-5472.CAN-05-3489

25. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, et al. UCSF Chimera--a
visualization system for exploratory research and analysis. J Comput Chem. 2004;25: 1605–1612.
doi:10.1002/jcc.20084

26. Trott O, Olson AJ. AutoDock Vina: improving the speed and accuracy of docking with a new scoring
function, efficient optimization and multithreading. J Comput Chem. 2010;31: 455–461.
doi:10.1002/jcc.21334

27. Wang J, Wang W, Kollman PA, Case DA. Automatic atom type and bond type perception in
molecular mechanical calculations. J Mol Graph Model. 2006;25: 247–260.
doi:10.1016/j.jmgm.2005.12.005

28. Wang J, Wolf RM, Caldwell JW, Kollman PA, Case DA. Development and testing of a general amber
force field. J Comput Chem. 2004;25: 1157–1174. doi:10.1002/jcc.20035

29. Case DA, Cheatham TE, Darden T, Gohlke H, Luo R, Merz KM, et al. The Amber biomolecular
simulation programs. J Comput Chem. 2005;26: 1668–1688. doi:10.1002/jcc.20290

30. Salomon-Ferrer R, Götz AW, Poole D, Le Grand S, Walker RC. Routine Microsecond Molecular
Dynamics Simulations with AMBER on GPUs. 2. Explicit Solvent Particle Mesh Ewald. J Chem
Theory Comput. 2013;9: 3878–3888. doi:10.1021/ct400314y

31. Maier JA, Martinez C, Kasavajhala K, Wickstrom L, Hauser KE, Simmerling C. ff14SB: Improving the
Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. J Chem Theory Comput.
2015;11: 3696–3713. doi:10.1021/acs.jctc.5b00255

32.   Mao Y, Zhang Y. Thermal conductivity, shear viscosity and specific heat of rigid water models.
742   Chem Phys Lett. 2012;542: 37–41. doi:10.1016/j.cplett.2012.05.044

743   33.   Andersen HC. Molecular dynamics simulations at constant pressure and/or temperature. J Chem
744   Phys. 1980;72: 2384–2393. doi:10.1063/1.439486

745   34.   Roe DR, Cheatham TE. PTRAJ and CPPTRAJ: Software for Processing and Analysis of Molecular
746   Dynamics Trajectory Data. J Chem Theory Comput. 2013;9: 3084–3095. doi:10.1021/ct400341p

747   35.   Wagner AH, Walsh B, Mayfield G, Tamborero D, Sonkin D, Krysiak K, et al. A harmonized meta-
748   knowledgebase of clinical interpretations of cancer genomic variants. bioRxiv. 2018; 366856.
749   doi:10.1101/366856

750   36.   Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for
751   Bigger Datasets. Mol Biol Evol. 2016;33: 1870–1874. doi:10.1093/molbev/msw054

Fig 1

Fig 2 - top

cancor for 20 AA window + conserved dynamics zones −Wilks lamda

total mutational impact (signif + ns)

Fig 2 - bottom

Fig 3 - top

cancor for 20 AA window + conserved dynamics zones −Wilks lamda

E

local R value for cancor / position (residue number)

atp_validate_V600E
conserved dynamics

F

total mutational impact (signif + ns)

G

sum impact / variant

atp_validate_V600E, dabrafenib_V600E, PLX7904_V600E, sorafenib_V600E, vemurafenib_V600

Fig 3 - bottom

cancor for 20 AA window + conserved dynamics zones −Wilks lamda

Fig 4

Fig 5

A

B

C

cancor for 20 AA window + conserved dynamics zones −Wilks lamda

local R value for cancor

position (residue number)

conserved dynamics

validation_run

D

sum impact

G464E G464R G466E G469L G469V validation_run

G596R GF595L L597Q L597R L597S L597V validation_ru'

G606E K601N K601T V600D V600E V600K V600R validation_r'

Fig 6

cancor for 21 AA window + conserved dynamics zones −Wilks lamda

- conserved dynamics
- validation_run

position (residue number)

local R value for cancor

sum impact

G466E G469A G469V L514V L597S validation_run

V600D V600E V600K V600M V600R validation_run

Fig 7

A

BRAF orthologs

tetrapod

0 replacements = training set

9 amino acid replacements

D453E
V458L
Y471Y or Y471F
S520T
M626L
T650S
Q652A
E694D
D701E

ray-finned fish

insect

95 amino acid replacements

0.050

Fig 8 - top

Fig 8 - bottom