Recent fluctuations in Mexican American genomes have altered the genetic architecture

of biomedical traits

Authors: Melissa L. Spear[1,2], Alex Diaz-Papkovich[3,4], Elad Ziv[5,6,7,8], Simon Gravel[4,9], Dara G.

Torgerson[4,9], Ryan D. Hernandez[2,4,7,9,10,11*]

1. Biomedical Sciences Graduate Program, University of California, San Francisco, San

Francisco, CA, USA

2. Department of Bioengineering and Therapeutic Sciences, University of California, San

Francisco, San Francisco, CA, USA

3. Quantitative Life Sciences Program, McGill University, Montreal, QC, Canada

4. McGill University and Genome Quebec Innovation Center, McGill University, Montreal, QC,

Canada

5. Division of General Internal Medicine, University of California, San Francisco, San Francisco,

CA, USA

6. Department of Medicine, University of California, San Francisco, San Francisco, CA, USA

7. Institute of Human Genetics, University of California, San Francisco, San Francisco, CA, USA

8. Helen Diller Family Comprehensive Cancer Center, University of California, San Francisco,

San Francisco, CA, USA

9. Department of Human Genetics, McGill University, Montreal, QC, Canada

10. Quantitative Biosciences Institute, University of California, San Francisco, San Francisco,

CA, USA

11. Institute for Human Genetics, University of California, San Francisco, CA, USA

*Corresponding author: ryan.hernandez@me.com

Abstract

Hispanics/Latinos are a diverse group of admixed populations with African, European, and Native American ancestries. They remain understudied, and thus little is known about the genetic architecture of phenotypic variation in these populations. Using genome-wide genotype data from the Hispanic Community Health Study/Study of Latinos, we find that Native American ancestry has increased over time across Hispanic/Latino populations, particularly in Mexican Americans where Native American ancestry increased by an average of ~20% over the 50-year period spanning 1940s-1990s. We find similar patterns across American cities, and replicate our observations in an independent sample of Mexican Americans. These dynamic ancestry patterns are a result of a complex interaction of several population and cultural factors, including strong ancestry-related assortative mating and subtle shifts in migration with differences in subcontinental Native American ancestry over time. These factors have shaped patterns of genetic variation, including an increase in runs of homozygosity in Native American ancestral tracts, and also influenced the genetic architecture of complex traits within the Mexican American population. We show for height, a trait correlated with ancestry, polygenic risk scores based on summary statistics from a European-based genome-wide association study perform poorly in Mexican Americans. Our findings reveal temporal changes in population structure within Hispanics/Latinos that may influence biomedical traits, demonstrating a crucial need to improve our understanding of the genetic diversity of admixed populations.

Introduction

Hispanics/Latinos living in the United States are culturally, phenotypically, and genetically diverse populations. Individuals who identify as Hispanic/Latino have varying proportions of Native American, African, and European genetic ancestry, each with its own unique continental demographic history. Demographic forces such as population bottlenecks and expansions, migration and adaptation to novel environments resulted in observable differences in continental

patterns of genetic variation (1-3). These differing patterns were shaped by many historical events

of migration which included the founding of the Americas by Native American populations, the

colonization by Europeans, and the African slave trade (4-8). These large-scale migrations and

additional demographic events shaped the genetic diversity of individuals living today within the

United States (9-13).

Demographic history has shaped the genetic architecture of modern human phenotypic

variation (14-19), and is critical to consider in the search for the genetic basis of complex diseases.

The demography of the United States has changed drastically over the 20th century, and by 2044

is predicted to become a 'minority-majority' country whereby no one racial/ethnic group comprises

more than 50% of the population (20). By 2060 Hispanics/Latinos are projected to make up the

largest of that share at 29% or 119 million individuals (20). However, to date, population-based

medical genomics research [and its subsequent benefits, including polygenic risk score (PRS)

profiling] have been disproportionately focused on individuals of European descent, with the

findings primarily benefiting European populations (21, 22). Despite the increases in sample

sizes, rates of discovery, and traits studied, Hispanic or Latin American participation in genome-

wide association studies (GWAS) has continued to hover around 1% (23, 24). This shows a

continuing trend of leaving this population particularly vulnerable to falling behind in receiving the

benefits of the precision medicine revolution (22, 23).

In this study we utilize the largest genetic study of Hispanics/Latinos in the U.S. to date --

the Hispanic Community Health Study/Study of Latinos (HCHS/SOL) (10) -- to understand how

patterns of genetic variation in Hispanic/Latino populations in the United States have changed

over the last century, and evaluate the impact such changes may be having on complex traits.

Results

Global ancestry proportions among HCHS/SOL Hispanic/Latino Populations

Using the subset of sites that overlapped with our African, European, and Native American

reference panels, we called 3-way global ancestry estimates for 10,268 unrelated HCHS/SOL

individuals (see methods). Figure 1A summarizes the global ancestry proportions shaded by

admixture estimates in a ternary plot, recapitulating the original HCHS/SOL analysis of continental

ancestry (10). However, while several population groups appear to have overlapping ancestry

proportions, this analysis masks more subtle structure in subcontinental ancestry. To investigate

subtle population structure across these self-identified population groups, we performed UMAP

on the top 3 principal components (see methods), and find substantial structure across self-

identified groups (Figure 1B and Supplementary Figure 1B). We find that Dominicans, who have

the highest average proportions of African ancestry, are in the middle, with Puerto Ricans and

Cubans, diverging in opposite directions (Supplementary Figure 1B) with clines of increasing

European ancestry proportions (Figure 1B). Further, while self-identified Mexican, Central, and

South American groups appear to have overlapping ancestry proportions in Figure 1A, UMAP

represents the Mexican Americans and Central/South American groups as large, separate wings

that diverge from self-identified Cubans and Dominicans, with both clusters diverging with clines

of increasing ancestry toward different Native American populations (Figure 1B and

Supplementary Figure 1B and 1C).

Dynamic Global Ancestry Proportions in Mexican Americans

For each of the HCHS/SOL Latino populations, we evaluated differences in global

ancestry estimates over time while accounting for the sampling method (referred to as "sampling

weight", see methods) used for the design of the HCHS/SOL study (25). We found that in all

populations, the effect size for Native American ancestry on birth year is positive, though only

statistically significant after multiple testing in the Mexican American (β =0.0023; P=3.58E-22;

Figure 1C) and Central American (β=0.0013; P=0.0013) cohorts (Supplementary Table 1). Due

to the larger sample size, magnitude of the effect, and statistical significance, we shift our focus

to Mexican Americans. In Mexican Americans, the increase in Native American global ancestry

over time was consistent across multiple data stratifications including recruitment region, US born

or not US born, educational attainment, and sex (Table 1), and was robust to alternative methods

for estimating global ancestry proportions (e.g. based on the summation of RFMix local ancestry

estimates; Supplementary Figures 2 and 3). We performed bootstrap resampling (n=1000) of

global Native American ancestry for the Mexican Americans and observed a consistent increase

in Native American ancestry with fitted LOESS smoothing (Figure 1D) and when individuals were

binned by birth year decades (Supplementary Figure 4). On average, global Native American

ancestry has increased ~20% over the last 50 years in Mexican Americans.

We replicated our original findings of the increase in global Native American ancestry over

time in a smaller, independent cohort of self-identified Mexican Americans (n=705) from the

Health and Retirement Study (HRS) [34]. The HRS Mexican Americans in this study are older

compared to the HCHS/SOL Mexican Americans (birth year distribution: 1915-1981;

mean=1943,median:1942) and have lower levels of global Native American ancestry on average

(mean=0.29), but we still observed an increase in global Native American ancestry over time

(β=0.00082; P=0.02; SE=0.0003673) (Supplementary Figure 5A). We performed 1000 bootstrap

resampling iterations of the linear regression model (global Native American ancestry ~ birth year)

fitted to the data. From these resampling iterations, 98.2% of the tests had a slope > 0 (average

β=0.00083) and 61.5% of the regression p-values were less than 0.05 as illustrated in

(Supplementary Figures 5B-5D).

A previous study (12) identified ancestry biased migration in African Americans where

individuals with higher proportions of European ancestry migrated first out of the South during the

Great Migration followed by individuals with higher proportions African ancestry. We hypothesized

that earlier immigrants to the US had higher proportions of European ancestry followed by recent

immigrants having higher proportions of global Native American ancestry. In our non-US born individuals (N=2987), we evaluated differences in ancestry estimates over time while accounting for years in the US and sampling weight and identified a significant effect of years in the US (β=-0.0009; P=0.0006; SE=0.0003). However, this did not change the effect of birth year on the proportion of global Native American ancestry (β =0.0028; P<2e-16, SE=0.0003).

For US born individuals we assessed whether parental birth place could explain the increases in global Native American ancestry. Of the 634 US born individuals, 385 had parents both born outside of the US, 149 had one parent born outside of the US. 97 had both parents born within the US. For the 385 individuals, we identified a significant association between birth year and global Native American ancestry (β=0.004, P=2.34e-10, SE=0.0006). For the remaining individuals we were unable to identify a significant effect of birth year on global Native American ancestry, possibly due to a small sample size.

Individual loci are not driving global ancestry proportions

We used local ancestry estimates generated across the genome to perform admixture mapping in HCHS/SOL Mexican Americans to determine if younger individuals harbored excess Native American ancestry in certain regions of the genome. Although we tried two different models (see methods), we did not find any loci to be significantly associated with birth year across the genome (Supplementary Figure 6).

Little evidence for subcontinental population structure

It is possible that the increase in global Native American ancestry over time could be biased by changes in the specific subcontinental Native American ancestries over time (though such an effect is not visible in our UMAP analysis, Figure 1B). If it were the case, then we would expect subtle signals of genetic divergence in Native American ancestry tracts over time. To

investigate this, we calculated F_{ST} between all pairs of birth-decades (see methods). Figure 2A

shows all pairwise comparisons among birth-decades, and demonstrates that while the estimates

of F_{ST} are negligible (with many estimates below 0), there is a subtle trend of increasing F_{ST} as

birth-decade differences increase (though individuals born in the 80s and 90s show a conflicting

pattern). We further investigated this pattern using genetic diversity, π, in Native American

ancestry tracts for each birth-decade (see methods). We hypothesized that if there were

increased migration from multiple Native American source populations (coupled with rapid

population growth in Mexican American communities), then genetic diversity should be increasing

over time. We found the opposite: Supplementary Figure 7 shows a subtle decrease in genetic

diversity (π) over time from the 1930s to the 1980s in non-US born Mexican Americans, and a

subtle decrease in US born Mexican Americans from the 70s to the 90s (while remaining roughly

constant from the 30s to the 70s).

Little evidence that Native American ancestry tract lengths have changed

We next sought to test whether differences at the local ancestry level could explain the

shift in global Native American ancestry over time in the Mexican Americans. We calculated the

length of each RFMix inferred local ancestry tract in each Mexican American individual, and tested

for differences in the distribution of tract lengths across birth-decades using a multiple linear

regression model (see methods). We found no significant associations between the decade bin

and the proportion of Native American ancestral tracts at various lengths (Figure 2B), even when

testing for violations of model assumptions (e.g. normalizing the tracts per bin by the number of

individuals, or excluding the 1930s and/or 1990s individuals due to the small sample size in each

bin).

Increased runs of homozygosity over time

Since genetic diversity has decreased over time in the Native American ancestry tracts of Mexican Americans (despite rapid growth of the census population size), it is possible that this population has also experienced increased haplotype homozygosity over time. We investigated this possibility by exploring runs of homozygosity (ROH) in Native American ancestry tracts in each of the 3622 Mexican Americans. We classified ROH into three categories: short, medium, and long, based on the length distribution in the population. Generally, short ROH are tens of kilobases in length and likely reflect the homozygosity of old haplotypes; medium ROH are hundreds of kilobases in length and likely reflect background relatedness in the population; and long ROH are hundreds of kilobases to several megabases in length and are likely the result of recent parental relatedness. Figure 2C shows a fitted loess curve to the log of the total length of ROH summed across each Mexican American's genome as a function of their birth year, broken down by ROH size class (as well as the total of each size class that overlaps all ancestry tracts (Supplementary Figure 8A) and Native American ancestry tracts (Supplementary Figure 8B)). Overall, we find a significant positive correlation between birth year and the total summed ROH across size classes (τ=0.0449, P=6.12e-5, Kendall's rank correlation), but this becomes more significant when we restrict our analysis to ROH calls that overlap Native American ancestry tracts (τ=0.0873, P=9.46e-15). When stratified by size class, the associations (all Kendall's rank correlation) in ROH were primarily driven by the short (τ=0.0833, P=9.45e-14), and medium (τ=0.0718, P=1.46e-10) size classes, and are again strongest when ROH overlap Native American ancestry tracts (short τ=0.107, P<2.2e-16, medium τ=0.1003, P<2.2e-16). The long ROH had a negative correlation with birth year, but was insignificant after multiple testing (τ=-0.0291, P=0.01499; note that 1694 individuals did not have any long ROH calls in their genome).

Strong ancestry-related assortative mating in HCHS/SOL Mexicans

Given that short and medium length ROH have increased over time, it appears that background relatedness within Native American ancestry in Mexican Americans has increased over time (but not an increase in recent parental relatedness). One way for this to occur is if individuals with similar ancestry patterns tend to mate with one another more often than expected under a model of random mating (i.e. assortative mating). To measure assortative mating, we estimated the ancestral proportions of the biological parents of each HCHS/SOL Mexican American (see methods). With individuals from all decades pooled together, we found the inferred biological parental Native American ancestries to be significantly correlated (Figure 2D, r=0.708, 95% CI:0.69-0.72, P<2.2e-16, Pearson correlation). When stratified by decade, the correlation in inferred parental Native American global ancestry ranged from 0.65 to 0.74 (Supplementary Figure 9), but were not statistically different from each other. This shows that there was a strong parental ancestry correlation among Mexican Americans over different generations. This signature of assortative mating is not due to recent parental relatedness, because there is no trend in long ROH with birth year (and an overall low rate of long ROH among Mexican Americans).

Genetic correlation of global Native American ancestry with biomedical traits

We have shown that genetic variation in the Mexican American population is dynamic, with Native American ancestry increasing over a short period of time (combined with decreased genetic diversity and increased short and medium length ROH within Native American ancestry tracts). These features may have implications for the genetic architecture of complex traits within Mexican Americans, a topic that is understudied and poorly understood. To further our understanding of the genetic architecture of complex traits in Mexican Americans, we investigated the relationship between Native American ancestry and various complex traits that may be relevant to biomedical phenotypes. Specifically, we tested for a correlation between 66 complex traits from the

HCHS/SOL phenotypic dataset and global Native American ancestry (Kendall's tau). As illustrated

in Figure 3, many of these traits are significantly correlated after Bonferroni correction

(P<0.00076). We found that the effect of global Native American ancestry on many of these

phenotypes persisted when using multiple regression to account for age, sex, center, and the

sampling weight (Supplementary Table 2), highlighting the need for increased investigation into

the role of Native American genetic ancestry in admixed populations such as Mexican Americans.

Assessing the genetic contribution of Native American ancestry to height

Among the traits we tested for a correlation with global Native American ancestry, height

had the strongest negative correlation, and our regression model indicated that height also had a

strong positive relationship with birth year (Figure 4A and Supplementary Table 3). Globally,

populations have grown taller over time due to a variety of non-genetic, environmental factors

(26). We find a similar trend in the HCHS/SOL Mexican Americans (Figure 4A). Indeed, when we

stratified individuals by quartiles of global Native American ancestry, we see that all quartiles have

increased in height by a similar amount over the period investigated (though individuals with lower

Native American ancestry were taller on average).

Height is one of the most highly studied complex traits, with GWAS sample sizes

numbering in the hundreds of thousands (27). Results for many of these studies have been made

readily available on public databases as summary association statistics that can be leveraged to

build genetic predictions through polygenic risk scores (PRS) (28). In Europeans, PRS have been

shown to have great predictive power for several traits, including breast cancer, prostate cancer,

and type 1 diabetes (22, 29-31). PRS are most effective in populations of European descent as

GWAS studies have been primarily performed in these populations (21-23) and are expected to

be biased when applied to other populations due to differences in the genetic architecture of traits

across diverse populations (32). Since Mexican Americans have some fraction of European

ancestry, we sought to determine whether PRS calculated utilizing GWAS summary statistics

from European populations could still provide useful insight.

To evaluate the effectiveness of PRS for height (i.e. the polygenic height score, or PHS,

see methods), we first tested whether there was an association between the observed height and

the predicted height estimates while controlling for sampling weight, sex, and recruitment center

(see methods). We identified a significant association between observed height and predicted

height for the population as a whole (β=0.0044881, P=2.19e-12; Figure 4B, Supplementary Table

4). However, when we stratified by quartiles of Native American global ancestry, the association

only remained for the individuals in the lower two quartiles of global Native American ancestry

(NAM<0.37: β=0.004, P=0.0008 and 0.36<NAM<0.46: β=0.004, P=0.003, Supplementary Table

4). The association between predicted height and observed height was no longer significant for

individuals in the upper two quartiles of global Native American ancestry proportions

(0.46<NAM<0.58: β=0.0011, P=0.39 and 0.58<NAM: β=0.0022, P=0.08, Supplementary Table

4).

As we had found global Native American ancestry to be increasing over time, we

hypothesized that there would be a change in PHS over time as well. However, we find little

evidence supporting this hypothesis. While individuals born earlier than 1950 or in the 1950s have

a stronger correlation between their PHS and observed height (β=0.034 and 0.039; p=5.6e-4 and

2.7e-7 respectively) than individuals born in the 1960s, 1970s, or 1980s (β=0.016, 0.029, and

0.029; p=0.044, 0.0066, and 7.8e-5 respectively), there is no clear trend and we did not find a

significant effect of birth year on PHS (P=0.09) even when we stratified by the quartiles of global

Native American ancestry.

Discussion

The United States is a dynamic, rapidly changing population, and this will continue to occur

as the population size grows (20). Hispanics/Latinos are the largest and fastest growing minority

group, and are projected to comprise over 25% of the US population by 2060. They are a

genetically and phenotypically diverse population as a result of extensive admixture between

Native Americans and immigrants from multiple geographic locations around the world. In this

study, we identified additional population substructure complexities that may contribute to

phenotypic variation within Hispanics/Latinos.

Specifically, we demonstrated how the admixture dynamics of Mexican Americans have

changed over time, resulting in an increase of ~20% Native American ancestry on average over

the 50-year period studied. This change in ancestry is equivalent to a mean increase in Native

American ancestry of ~0.4% per year. While the effect sizes vary to some extent, we replicate the

underlying pattern across multiple data stratifications (two metropolitan cities, US born and non-

US born) and also replicate this feature in an independent cohort of Mexican Americans. Further,

we find that a similar trend holds across multiple self-identified Hispanic/Latino populations in the

US (and is statistically significant in Central Americans). This effect does not appear to have a

simple explanation: we do not see any statistically significant increases at individual loci, we do

not see more than a negligible degree of population differentiation over time, and this increase

cannot be entirely explained by very recent migration. We do, however, find that as Native

American ancestry has increased, genetic diversity within Native American ancestry tracts across

Mexican Americans has decreased over time, and is associated with increased short and medium

length ROH over time. This suggested that there could be increased relatedness within Native

American ancestries within Mexican Americans, and we confirmed that there is a very high degree

of ancestry-based assortative mating within the Mexican American population.

What could be driving the increased Native American ancestry in Mexican Americans

(12)? Population genetic theory suggests that while assortative mating could result in increased

ROH and decreased genetic diversity, ancestry-based assortative mating alone should not result

in mean changes in global ancestry proportions. Regardless of the underlying mechanisms driving

increased Native American ancestry in Mexican Americans, this additional source of temporal

substructure within this population has substantial consequences for phenotypic variation in

biomedical traits. We identify several biomedical traits that are correlated with Native American

ancestry, and show that in the case of height, there are both ancestry and temporal effects.

Further study is necessary to understand whether other biomedical traits are also changing over

time as the genomic ancestry proportions change in this population.

In our study, we bring specific attention to the biases that continue to exist with using

European GWAS summary statistics to calculate polygenic risk scores in admixed populations

such as Mexican Americans that are comprised of European, Native American, and African

genetic ancestries. In particular, in the case of height, we found that the polygenic height score

(PHS) correlated with observed height only in the subset of individuals with the lowest levels of

Native American ancestry (i.e. the subset of individuals with highest European ancestry). As the

population dynamics of the US continue to change, it is imperative that we study diverse

populations, or we risk exacerbating the health disparities that currently exist. To date, population-

based medical genomics research (and its subsequent benefits) have been disproportionately

focused on populations of European ancestry. In order to improve the design and implementation

of medical genetics studies for the ethnically diverse U.S. population, we need detailed insights

into the population history of diverse U.S. populations. This includes characterizing the admixture

dynamics of Hispanic/Latino populations, as well as the evolutionary forces that shaped patterns

of genetic variation of the ancestral populations that contributed to modern day Hispanic/Latino

populations.

The genetic variation of the Hispanic community in the United States belies

categorization under a single label (10). The events that have shaped and continue to shape

this genetic diversity are complex, numerous, and nuanced, and the social history of such a

diverse population is intrinsic to any genetic study. Mexico's society was largely defined by an

established social caste system based on ancestry, which disappeared after Mexico's

independence in 1821 (33). Even so, social inequalities persist today with skin colour having a

significant effect on wealth and education (34). A multitude of factors within and outside Mexico

— whether related to trade, immigration policies, or armed conflicts — acted to influence who

immigrated to the United States, and the impact of each of these fluctuates over time (35-37).

These changes shift the demographics of immigration, which is inherently related to the genetic

ancestry of the population.

Consequently, this shapes the genetic architecture of complex traits. Diverse

populations are at risk not only from underrepresentation in research, but because of poor

understanding of the temporal and spatial dynamics at play in genetic variation. The promise of

equitable precision medicine — one of the ultimate goals of medical genomics — cannot be

kept without understanding this interplay. Health disparities in the United States are fed by

structural inequalities. For example, studies that use modern Artificial Intelligence (AI)

techniques have already been shown to inflate existing disparities between Black Americans

and White Americans (38). Such biases, whether from algorithms, study designs, or

misunderstandings of subtleties in data, feed into the larger systemic pressures faced by

minority populations in the United States.

While we have shown a dramatic shift in ancestry proportions in US Hispanic/Latinos, one

of the caveats of this study is that the HCHS/SOL cohort is not representative of all US

Hispanics/Latinos. HCHS/SOL participants were recruited at four primary centers: Bronx,

Chicago, Miami, and San Diego. There may be additional genetic diversity that has not been

captured by this dataset and trends exhibited in this dataset may not translate to Hispanic/Latino

populations living in other regions of the US (though the temporal increase in Native American

ancestry was replicated in an independent sample of Mexican Americans). With better population

genetic modeling and a deeper understanding of the social and historical aspects of

Hispanic/Latino populations, we will be able to improve our understanding of the genetic and

phenotypic diversity across these populations, and subsequently improve our ability to understand

genetic contributions to complex traits and disease. These insights will lead to optimization of

population sampling for the design of future medical genetic studies, the identification of disease

risk variants, and ultimately, precision medicine for all.

Methods

Study dataset and initial quality control

The HCHS/SOL study is a community-based cohort study of self-identified Hispanic/Latino

individuals from four US metropolitan areas with the general goal of identifying risk and protective

factors for various medical conditions including cardiovascular disease, diabetes, pulmonary

disease, and sleep disorders (25). 12,434 participants with birth year estimates between 1934-

1993 who self-identified as being of Cuban, Dominican, Puerto Rican, Mexican, Central American,

or South American background consented to genetics studies and posting of their genetic and

phenotype data on the publicly available Database of Genotypes and Phenotypes (dbGaP)

through Study Accession phs000810.v1.p1. Samples were genotyped on an Illumina custom

array, SoL HCHS Custom 15041502 array (annotation B3, genome build 37), consisting of the

Illumina Omni 2.5M array and 148,353 custom single nucleotide polymorphisms (SNPs) (10).

Data posted to dbGaP had passed initial sample quality control filters, including removing samples

with differences in reported vs. genetic sex, call rates > 95%, and evidence for sample

contamination (e.g. heterozygosity and sample call rates). For initial SNP quality control, we

filtered out SNPs that were monomorphic, positional duplicates, or Illumina technical failures, as

well as SNPs that had cluster separation <= 0.3, call rate <=2%, >2 disconcordant calls in 291

duplicate samples, >3 Mendelian errors in parent-offspring pairs/trios, Hardy-Weinberg

Equilibrium combined P-value $<10^{-5}$, and sex differences in allele frequency ≥0.2. Our filtering

resulted in 1,763,935 genotyped SNPs with minor allele frequency (MAF) >0.01.

Additional sample quality control performed in the HCHS/SOL dataset included filtering

out samples with 1) large chromosomal anomalies, 2) substantial Asian ancestry as previously

identified in HCHS/SOL (*12*) and 3) individuals with up to third degree genetic relatedness in the

dataset as inferred by REAP (39). For genetic relatedness filtering, individuals from pairs were

kept to maximize representation of the birth year distribution, which resulted in 10,268 unrelated

remaining individuals.

From the original HCHS/SOL analysis, individuals were classified into genetic-analysis

groups, similar to self-identified background groups in that they share cultural and environmental

characteristics, but are also more genetically homogenous (10).

Birth year for all individuals was estimated by subtracting the difference between date of

first clinic visit for the baseline examination (25) and age. Year of arrival was estimated by

subtracting the difference between date of first clinic visit for the baseline examination and years

in the US.

Global, local, and parental ancestry inference

All ancestry analyses were restricted to the 211,152 autosomal SNP markers that overlapped

between the study and reference panel genotyping array. For the HCHS/SOL dataset, global

African, European, and Native American ancestries were inferred with ADMIXTURE, in an

unsupervised manner, with K=3. For some analyses, HCHS/SOL individuals with greater than

95% of either African, European, or Native American ancestry were filtered out resulting in 9,913

individuals: 1,099 Central American, 1,536 Cuban, 954 Dominican, 3,622 Mexican, 1,783 Puerto

Rican, 652 South American and 267 "Other" individuals.

Ancestral tracts, known as 'local' ancestry, along the genome for all HCHS/SOL

individuals were inferred using RFMix (40) and a three population reference panel, comprised of

315 individuals: 104 HapMap phase 3 CEU (European) and 107 YRI (African) individuals (41)

and 112 Native American individuals from throughout Latin America (8). The reference panel was

limited to individuals with 99% continental ancestry as inferred by unsupervised ADMIXTURE

(42). Prior to local ancestry inference, HCHS/SOL individuals were merged with the reference

panels and then phased using SHAPEIT2 (43). For all HCHS/SOL Mexican American individuals,

parental genomic ancestry was inferred with ANCESTOR (44) using the local ancestry estimates

generated by RFMix.

Uniform Manifold Approximation and Projection (UMAP)

Principal components for HCHS/SOL and the reference panel were computed using smartPCA

(45). UMAP (version 0.3.8) was run using the Python script freely available at

https://github.com/diazale/gt-dimred with parameter specification set at 15 nearest neighbours

and a minimum distance between points of 0.5.

Admixture mapping

Local ancestry estimates for 211,151 SNPs across the genome were used to perform admixture

mapping in HCHS/SOL Mexican Americans to determine if younger individuals harbored excess

Native American ancestry in certain regions of the genome. Admixture mapping was performed

applying two different models: 1) a linear regression model with age as the dependent variable

adjusting for global Native American ancestry, sampling weight and center and 2) a logistic

regression model dividing the HCHS/SOL Mexican cohort in to an older vs younger generation

with 1965 set as the dividing point while also adjusting for global Native American ancestry,

sampling weight, and center. The threshold for genome-wide significance, 1.38×10^{-4} was

calculated using the empirical autoregression framework with the package *coda* in R to estimate

the total number of ancestral blocks (46, 47).

Tract Lengths

The multiple regression model: $\log(f) = \beta_0 + \beta_1\ T + \beta_2\ A + \beta_3\ TA + \varepsilon$, where f is a matrix

containing the proportion of lengths of all ancestral tracts across the genome for all 3622 Mexican

American individuals, T the tract length bin and A decade of birth year bin, was used to test for an

effect of birth decade on the proportion of Native American ancestral tract lengths. For

assessment between the fraction of ancestry tracts in an individual's genome and birth year, long

tract cutoffs were chosen based on tract separation between the birth year decades in Figure 2B.

Diversity Calculations

Subcontinental ancestry was assessed using the diversity measurements π and F_{ST}. π was

calculated as the average number of pairwise genetic differences among all pairs of overlapping

Native American ancestry tracts across individuals. F_{ST} was calculated as:

$$F_{ST} = (H_T - H_S)/H_T$$

where H_T is the average heterozygosity when all individuals are pooled across decades and H_S is

the average heterozygosity within each decade of individuals.

Inference of Runs of Homozygosity

Runs of homozygosity (ROH) were called using the program GARLIC v1.1.4 (48) on 211,152

sites for the Mexican American individuals. An analysis window size of 50 SNPs and an

overlap fraction of 0.25 were both chosen using GARLIC's rule of thumb parameter

estimation. GARLIC chose a LOD score cutoff of 0. Using a three-component Gaussian

mixture, GARLIC determined class A/B (short/medium) and class B/C (medium/long) size

boundaries as 845,097 bp and 2,501,750 bp, respectively.

Imputation

Imputation for HCHS/SOL was performed locally using IMPUTE2 with the 1000 Genomes Project Phase 3 haplotypes used as a reference panel. After filtering on an info score cutoff of 0.3, this resulted in 33,041,084 SNPs.

Polygenic Risk Score Calculations

Polygenic risk scores for height were calculated using the publicly available UK Biobank (UKBB) GWAS Round 2 Summary Statistics retrieved from http://www.nealelab.is/uk-biobank. Briefly, for sample quality control, sample inclusion was limited to unrelated samples who passed the sex chromosome aneuploidy filter. British ancestry was determined using the 1^{st} 6 PCs; individuals more than 7 standard deviations away from the 1^{st} 6 PCs were excluded. Further filtering included limiting to self -reported 'white-British' / 'Irish' / 'White' resulting in a QCed sample count of 361,194 individuals https://github.com/Nealelab/UK_Biobank_GWAS#imputed-v3-sample-qc. An imputation panel of ~90 million SNPs from HRC, UK10K and 1KG were used to impute genotypes. 13.7 million autosomal and X-chromosome SNPs passed quality control thresholds including Info score>0.8, MAF>0.0001, and HWE p-value>1e-10. For the phenotype, a linear regression model in Hail was run for all individuals (both sexes) adjusting by the first 20 PCs + sex + age + age^2 + (sex*age) + $(sex*age)^2$. For height, there was complete phenotype information for 360,388 individuals.

Risk scores were calculated by extracting the overlapping genome-wide significant hits initially discovered in the UKBB GWASs of height and selecting SNPs with the lowest p-value in each 1Mb window across the genome. For height this resulted in a dataset of 1,103 overlapping SNPs that were present in our dataset of genotyped and imputed SNPs.

Health and Retirement Study (HRS)

For replication, we used genotype data from 705 self-identified Mexican-Americans from the

Health and Retirement Study (HRS) (49), genotyped on the Illumina Human Omni 2.5M platform.

HRS data was made available under IRB Study No. A11-E91-13B - The apportionment of genetic

diversity within the United States. Estimated global ancestry proportions for the Mexican American

population in the HRS were calculated as in Baharian et al. (12), which used an alternative

reference panel and alternative ancestry inference approach. Briefly, RFMix was used to infer

local ancestry estimates across the genome utilizing CHS, YRI, and CEU individuals from the

1000 Genomes Project as reference populations for Native American/Asian, African, and

European ancestries, respectively. Global ancestry estimates were calculated using the summed

RFMix calls.

Statistical Analyses and Plots

Statistical analyses and plot generation were performed within Rstudio using Version 1.1.463 and

R version 3.5.3. ternary and ggridges/ggplot2 packages were used to create the simplex and

ridgeline plots.

Acknowledgements

MLS was supported through the National Human Genome Research Institute (NHGRI) of the

National Institutes of Health (NIH) under Award Number F31HG010104. ADP and SG were

supported, in part, thanks to funding from the Canada Research Chairs program and CIHR

grant MOP-136855. RDH was supported, in part, by NHGRI grant R01HG007644 and the

Canadian Research Chairs program.

Declaration of Interests

The authors declare no competing interests.

Figure 1. Recent dynamics continually shape the continuum of continental ancestry Hispanic/Latino populations. A. Ternary plot of HCHS/SOL (n=10,268) colored by admixture proportions. B. Uniform Manifold Approximation and Projection (UMAP) plot depicting the genetic diversity of HCHS/SOL and the reference panel (n=10,591) using 3 principal components, colored by admixture proportions (see Supplemental Fig 1 for population labels). Within the legend, AFR, EUR, and NAM refer to African, European, and Native American global ancestries, respectively. C. Global Native American ancestry proportions plotted by birth year for Mexican Americans (n=3,622). Fitted line is multiple regression of Native American ~ birth year + sampling weight. Bars represent 95% confidence intervals for individuals grouped by decade. D. Bootstrap resampling (n=1000 iterations) of Native American global ancestry for the Mexican American individuals with a fitted LOESS regression line for each iteration. Dashed lines represent the 95% confidence interval and the blue line represents the fitted regression line from Figure 1C.

Figure 2. Diversity of and within Native American ancestral tracts. A) F_{ST} estimates calculated

between each decade group. Bars represent the 95% CI. B) Proportion of total Native American

ancestral tracts in the HCHS/SOL Mexican American population by decade. C) Loess regression

of the log of the sum of total ROH and ROH overlapping Native American ancestral tracts

separated by ROH class. Total long ROH is not represented as an individual group due to the

high number of individuals missing long ROH (1694 for long ROH across ancestries and 1987 for

long NAM ROH) but was included in the sum of "All ROH" and "All NAM ROH". D) Correlation of

parent's inferred global Native American ancestries using ANCESTOR.

Figure 3. Correlation of 66 quantitative traits with global Native American ancestry.

Significance level was determined using Bonferroni correction adjusting by the number of

quantitative traits tested (0.05/66=0.00075).

Figure 4: Height and global Native American ancestry in HCHS/SOL Mexican Americans.

Each plot illustrates the relationship between A) Birth year and height B) Height and polygenic

height score (PHS). The black line indicates the fitted linear model for all individuals. Each color

represents a different quartile of Native American global ancestry. Polygenic height scores were

assessed utilizing UKBB summary statistics for 1,128 SNPs.

Table 1: Relationship of Native American global ancestry and birth year for Mexican

Americans stratified by recruitment region, US born vs non-US born status, sex and

educational attainment. For recruitment region, data stratification was limited to Chicago

and San Diego as sample size for the Bronx and Miami was limited: 124 and 25 individuals,

respectively. Education attainment was categorized as either less than a high school

diploma or equivalent degree (<HS), equal to a high school diploma or equivalent degree

(=HS), or post-secondary education (>HS).

Category	N	Mean	Median	R2	Effect	Std.Err	P
All	3622	0.489	0.468	0.027	0.0023	0.0002	3.58E-22
Chicago	1310	0.562	0.550	0.017	0.0016	0.0005	0.0006
San Diego	2163	0.428	0.422	0.012	0.0012	0.0002	4.29E-07
US born	634	0.427	0.418	0.063	0.0027	0.0004	1.77E-10
Not US born	2987	0.502	0.481	0.050	0.0032	0.0003	1.38E-30
Male	1500	0.494	0.475	0.038	0.0028	0.0004	3.83E-14
Female	2122	0.485	0.462	0.022	0.0019	0.0003	3.07E-10
<HS	1518	0.520	0.500	0.045	0.0026	0.0004	1.39E-12
=HS	960	0.501	0.479	0.022	0.0018	0.0005	0.0003
>HS	1140	0.436	0.422	0.045	0.0027	0.0004	6.53E-13

Supplementary Material

Supplementary Figure 1. Continental ancestral diversity of HCHS/SOL A) Ternary plot of global ancestry proportions colored by population for 10,268 HCHS/SOL individuals B) Uniform Manifold Approximation and Projection (UMAP) plot of HCHS/SOL and the reference panel (n=10,591) using 3 principal components, colored by HCHS/SOL population. C) UMAP plot of HCHS/SOL and the reference panel (n=10,591) using 3 principal components, colored by reference population.

Supplementary Figure 2. Concordance of ADMIXTURE and RFMix global ancestry estimates. A) Native American ancestry B) African ancestry and C) European ancestry.

Supplementary Figure 3. RFMix inferred Native American global ancestry proportions

plotted over time for HCHS/SOL Mexican Americans (n=3622).

Supplementary Figure 4: Distributions of Native American global ancestry means for

HCHS/SOL Mexican Americans (n=3622) generated by 1000 bootstrap resampling

iterations within each decade of binned birth years.

685
686

703 **Supplementary Figure 5. Replication in the Health and Retirement Study for 705 self-**

704 **identified Mexican Americans**. A) Ancestry over time B) Distribution of regression slopes after

705 1000 bootstrap resampling iterations C) Distribution of bootstrap regression p-values D) ECDF

706 of bootstrap regression p-values.

Supplementary Figure 6. Admixture mapping in HCHS/SOL Mexicans (n=3622) for Native

American ancestry and A) birth year and B) generation. Ancestry association testing was

performed at 211,151 markers using A) linear regression and B) logistic regression, both

including global Native American ancestry, sampling weight and center as covariates.

Supplementary Figure 7. Diversity of and within Native American ancestral

tracts. Diversity (π) of subcontinental Native American ancestry stratified by US born/not US

born status. π was calculated between pairs within each decade of birth years. 95% confidence

intervals are highlighted by the shaded regions for each group.

Supplementary Figure 8. Runs of homozygosity (ROH) in HCHS/SOL Mexican Americans.

A) ROH across all ancestries separated by ROH class B) ROH overlapping Native American

haplotypes separated by ROH class.

Supplementary Figure 9. Ancestry-related assortative mating in HCHS/SOL Mexican

Americans separated by decade. Each plot represents the correlation of parent's inferred

Native American ancestries using ANCESTOR by decade beginning with the 1930s (A) and

ending with the 1990s (G). Each point corresponds to one Mexican American couple and the

axes correspond to the inferred Native American ancestry of each partner.

  Supplementary Table 1: Association of global ancestries and birth year for all

  HCHS/SOL individuals. For each population, we tested for an association between global

  ancestry and birth year while accounting for the sampling design. NAM, AFR, and EUR

  refer to Native American, African, and European ancestry respectively.

Population	Ancestry	N	R2	Effect	Std.Err	P
Central American	NAM	1099	0.0139	0.0013	0.0004	0.0013
Central American	AFR	1099	0.0136	0.0002	0.0004	0.6561
Central American	EUR	1099	0.0138	-0.0015	0.0004	0.0002
Cuban	NAM	1536	0.0023	0.0002	0.0001	0.0938
Cuban	AFR	1536	0.0014	-0.0005	0.0004	0.1490
Cuban	EUR	1536	0.0005	0.0003	0.0004	0.3879
Dominican	NAM	954	0.0035	0.0002	0.0001	0.0663
Dominican	AFR	954	0.0030	-0.0007	0.0004	0.1287
Dominican	EUR	954	0.0022	0.0005	0.0004	0.2374
Mexican	NAM	3622	0.0268	0.0023	0.0002	0.0000
Mexican	AFR	3622	0.0008	0.0000	0.0000	0.4189
Mexican	EUR	3622	0.0285	-0.0023	0.0002	0.0000
Puerto Rican	NAM	1783	0.0014	0.0001	0.0001	0.1533
Puerto Rican	AFR	1783	0.0014	0.0003	0.0002	0.1743
Puerto Rican	EUR	1783	0.0027	-0.0005	0.0002	0.0355
South American	NAM	652	0.0110	0.0016	0.0007	0.0211
South American	AFR	652	0.0027	-0.0002	0.0004	0.5053
South American	EUR	652	0.0080	-0.0014	0.0006	0.0335

Supplementary Table 2 - Multiple regression table with traits that were significantly correlated with global Native American ancestry.

Trait	N	R2	Effect	Std.Err	P
Height	3615	0.588	-13.637	0.676	1.01E-85
Predicted FVC	3522	0.851	-695.376	36.867	1.13E-75
EGFR MDRD	3308	0.196	23.398	2.435	1.39E-21
EGFR CKD Epi	3308	0.441	14.752	1.581	1.90E-20
Waist to hip ratio	3617	0.195	0.043	0.007	6.96E-10
Glycosylated hemoglobin (HbA1c)	3609	0.081	9.184	1.590	8.30E-09
% Glycosylated hemoglobin	3609	0.080	0.837	0.146	9.88E-09
HDL cholesterol	3621	0.095	-6.740	1.372	9.39E-07
Total iron binding capacity	3620	0.066	23.540	5.467	1.71E-05
FEV1 FVC Ratio	3505	0.176	2.560	0.639	6.37E-05
% Lymphocytes	3442	0.021	3.768	1.023	2.35E-04
Hip girth	3617	0.077	-4.452	1.273	4.78E-04
% Neutrophils	3442	0.039	-3.649	1.178	1.96E-03
Monocyte count	3443	0.024	-0.048	0.019	1.27E-02
Neutrophil count	3442	0.043	-0.399	0.165	1.59E-02
LDL cholesterol	3529	0.047	-8.269	3.963	3.70E-02
Average diastolic blood pressure	3616	0.072	-2.067	1.136	6.88E-02
Total cholesterol	3622	0.065	-6.248	4.614	1.76E-01
White blood cell count	3442	0.032	-0.260	0.208	2.12E-01
Average systolic blood pressure	3619	0.211	1.590	1.712	3.53E-01
% Immature granulocytes	555	0.261	-0.090	0.110	4.14E-01
QRS duration	3596	0.168	0.062	1.268	9.61E-01

799 **Supplementary Table 3: Height over time: Height (cm) as a function of birth year**

800 **adjusting by sex, center, and sampling weight for 3614 Mexican Americans stratified by**

801 **the quartiles of global Native American ancestry.**

Group	N	R2	Effect	Std.Err	P
All	3614	0.54186	0.1200379	0.00905366	3.28E-39
NAM>0.58	929	0.5745454	0.158895	0.01783895	2.73E-18
0.46<=NAM<=0.58	955	0.5784636	0.1542762	0.01683919	3.07E-19
0.37<=NAM<0.46	842	0.5472275	0.1008054	0.01739819	9.73E-09
NAM<0.37	888	0.5369993	0.1161775	0.01815834	2.55E-10

805 **Supplementary Table 4: Predicted height vs. observed height.** Predicted height (cm) as

806 a function of observed height (cm) adjusting by sex, center, and sampling weight for 3614

807 Mexican Americans.

Group	N	R2	Effect	Std.Err	P
All	3614	0.0249	0.0045	0.0006	2.19E-12
NAM>0.58	929	0.0072	0.0022	0.0012	7.79E-02
0.46<=NAM<=0.58	955	0.0058	0.0011	0.0013	3.90E-01
0.37<=NAM<0.46	842	0.0144	0.0043	0.0015	3.22E-03
NAM<0.37	888	0.0164	0.0043	0.0013	7.91E-04

References:

1.   Nelson MR, Bryc K, King KS, Indap A, Boyko AR, Novembre J, et al. The Population

Reference Sample, POPRES: a resource for population, disease, and pharmacological genetics

research. Am J Hum Genet. 2008;83(3):347-58.

2.   Genomes Project C, Abecasis GR, Auton A, Brooks LD, DePristo MA, Durbin RM, et al.

An integrated map of genetic variation from 1,092 human genomes. Nature. 2012;491(7422):56-

65.

3.   Genomes Project C, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, et al. A

global reference for human genetic variation. Nature. 2015;526(7571):68-74.

4.   Gravel S, Zakharia F, Moreno-Estrada A, Byrnes JK, Muzzio M, Rodriguez-Flores JL, et

al. Reconstructing Native American migrations from whole-genome and whole-exome data.

PLoS Genet. 2013;9(12):e1004023.

5.   Homburger JR, Moreno-Estrada A, Gignoux CR, Nelson D, Sanchez E, Ortiz-Tello P, et

al. Genomic Insights into the Ancestry and Demographic History of South America. PLoS

Genet. 2015;11(12):e1005602.

6.   Moreno-Estrada A, Gignoux CR, Fernandez-Lopez JC, Zakharia F, Sikora M, Contreras

AV, et al. Human genetics. The genetics of Mexico recapitulates Native American substructure

and affects biomedical traits. Science. 2014;344(6189):1280-5.

7.   Moreno-Estrada A, Gravel S, Zakharia F, McCauley JL, Byrnes JK, Gignoux CR, et al.

Reconstructing the population genetic history of the Caribbean. PLoS Genet.

2013;9(11):e1003925.

8.   Reich D, Patterson N, Campbell D, Tandon A, Mazieres S, Ray N, et al. Reconstructing

Native American population history. Nature. 2012;488(7411):370-4.

9.   Bryc K, Durand EY, Macpherson JM, Reich D, Mountain JL. The genetic ancestry of

African Americans, Latinos, and European Americans across the United States. Am J Hum

Genet. 2015;96(1):37-53.

10.   Conomos MP, Laurie CA, Stilp AM, Gogarten SM, McHugh CP, Nelson SC, et al.

Genetic Diversity and Association Studies in US Hispanic/Latino Populations: Applications in the

Hispanic Community Health Study/Study of Latinos. Am J Hum Genet. 2016;98(1):165-84.

11.   Han E, Carbonetto P, Curtis RE, Wang Y, Granka JM, Byrnes J, et al. Clustering of

770,000 genomes reveals post-colonial population structure of North America. Nat Commun.

2017;8:14238.

12.   Baharian S, Barakatt M, Gignoux CR, Shringarpure S, Errington J, Blot WJ, et al. The

Great Migration and African-American Genomic Diversity. PLoS Genet. 2016;12(5):e1006059.

13.   Jordan IK, Rishishwar L, Conley AB. Native American admixture recapitulates

population-specific migration and settlement of the continental United States. PLoS Genet.

2019;15(9):e1008225.

14.   Agarwala V, Flannick J, Sunyaev S, Go TDC, Altshuler D. Evaluating empirical bounds

on complex disease genetic architecture. Nat Genet. 2013;45(12):1418-27.

15.   Eyre-Walker A. Evolution in health and medicine Sackler colloquium: Genetic

architecture of a complex trait and its implications for fitness and genome-wide association

studies. Proc Natl Acad Sci U S A. 2010;107 Suppl 1:1752-6.

16.   Maher MC, Uricchio LH, Torgerson DG, Hernandez RD. Population genetics of rare

variants and complex diseases. Hum Hered. 2012;74(3-4):118-28.

17.   Simons YB, Turchin MC, Pritchard JK, Sella G. The deleterious mutation load is

insensitive to recent population history. Nat Genet. 2014;46(3):220-4.

18.   Uricchio LH, Zaitlen NA, Ye CJ, Witte JS, Hernandez RD. Selection and explosive

growth alter genetic architecture and hamper the detection of causal rare variants. Genome

Res. 2016;26(7):863-73.

19.   Yang J, Bakshi A, Zhu Z, Hemani G, Vinkhuyzen AA, Lee SH, et al. Genetic variance

estimation with imputed variants finds negligible missing heritability for human height and body

mass index. Nat Genet. 2015;47(10):1114-20.

20.   Colby SLO, J.M. Projections of the Size and Compositon of the U.S. Population: 2014 to

2060. U.S. Census Bureau, Commerce USDo; 2015.

21.   Bustamante CD, Burchard EG, De la Vega FM. Genomics for the world. Nature.

2011;475(7355):163-5.

22.   Martin AR, Kanai M, Kamatani Y, Okada Y, Neale BM, Daly MJ. Clinical use of current

polygenic risk scores may exacerbate health disparities. Nat Genet. 2019;51(4):584-91.

23.   Popejoy AB, Fullerton SM. Genomics is failing on diversity. Nature. 2016;538(7624):161-

4.

24.   Mills MC, Rahal C. A scientometric review of genome-wide association studies.

Commun Biol. 2019;2:9.

25.   Sorlie PD, Aviles-Santa LM, Wassertheil-Smoller S, Kaplan RC, Daviglus ML, Giachello

AL, et al. Design and implementation of the Hispanic Community Health Study/Study of Latinos.

Ann Epidemiol. 2010;20(8):629-41.

26.   Collaboration NCDRF. A century of trends in adult human height. Elife. 2016;5.

27.   Yengo L, Sidorenko J, Kemper KE, Zheng Z, Wood AR, Weedon MN, et al. Meta-

analysis of genome-wide association studies for height and body mass index in approximately

700000 individuals of European ancestry. Hum Mol Genet. 2018;27(20):3641-9.

28.   Pasaniuc B, Price AL. Dissecting the genetics of complex traits using summary

association statistics. Nat Rev Genet. 2017;18(2):117-27.

29.   Sharp SA, Rich SS, Wood AR, Jones SE, Beaumont RN, Harrison JW, et al.

Development and Standardization of an Improved Type 1 Diabetes Genetic Risk Score for Use

in Newborn Screening and Incident Diagnosis. Diabetes Care. 2019;42(2):200-7.

30.   Schumacher FR, Al Olama AA, Berndt SI, Benlloch S, Ahmed M, Saunders EJ, et al.

Association analyses of more than 140,000 men identify 63 new prostate cancer susceptibility

loci. Nat Genet. 2018;50(7):928-36.

31. Maas P, Barrdahl M, Joshi AD, Auer PL, Gaudet MM, Milne RL, et al. Breast Cancer

Risk From Modifiable and Nonmodifiable Risk Factors Among White Women in the United

States. JAMA Oncol. 2016;2(10):1295-302.

32. Martin AR, Gignoux CR, Walters RK, Wojcik GL, Neale BM, Gravel S, et al. Human

Demographic History Impacts Genetic Risk Prediction across Diverse Populations. Am J Hum

Genet. 2017;100(4):635-49.

33. Lisker R, Ramirez E, Briceno RP, Granados J, Babinsky V. Gene frequencies and

admixture estimates in four Mexican urban centers. Hum Biol. 1990;62(6):791-801.

34. Martinez DZ-CIF. Is Mexico a Post-Racial Country? Inequality and Skin Tone across the

Americas. AmericasBarometer Insights Series by the Latin American Public Opinion Project

(LAPOP), wwwLapopSurveysorg [Internet]. 2017. Available from:

https://www.vanderbilt.edu/lapop/insights/ITB031en.pdf.

35. Fernández-Kelly P, Massey DS. Borders for Whom? The Role of NAFTA in Mexico-U.S.

Migration. The Annals of the American Academy of Political and Social Science. 2007;610:98-

118.

36. Verea M. Immigration Trends After 20 Years of NAFTA. 2014. 2014;9(2).

37. Contreras VR. THE ROLE OF DRUG-RELATED VIOLENCE AND EXTORTION IN

PROMOTING MEXICAN MIGRATION: Unexpected Consequences of a Drug War. Latin

American Research Review. 2014;49(3):199-217.

38. Obermeyer Z, Powers B, Vogeli C, Mullainathan S. Dissecting racial bias in an algorithm

used to manage the health of populations. Science. 2019;366(6464):447-53.

39. Thornton T, Tang H, Hoffmann TJ, Ochs-Balcom HM, Caan BJ, Risch N. Estimating

kinship in admixed populations. Am J Hum Genet. 2012;91(1):122-38.

40. Maples BK, Gravel S, Kenny EE, Bustamante CD. RFMix: a discriminative modeling

approach for rapid and robust local-ancestry inference. Am J Hum Genet. 2013;93(2):278-88.

41.   International HapMap C, Altshuler DM, Gibbs RA, Peltonen L, Altshuler DM, Gibbs RA,

et al. Integrating common and rare genetic variation in diverse human populations. Nature.

2010;467(7311):52-8.

42.   Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in

unrelated individuals. Genome Res. 2009;19(9):1655-64.

43.   Delaneau O, Zagury JF, Marchini J. Improved whole-chromosome phasing for disease

and population genetic studies. Nat Methods. 2013;10(1):5-6.

44.   Zou JY, Halperin E, Burchard E, Sankararaman S. Inferring parental genomic ancestries

using pooled semi-Markov processes. Bioinformatics. 2015;31(12):i190-6.

45.   Patterson N, Price AL, Reich D. Population structure and eigenanalysis. PLoS Genet.

2006;2(12):e190.

46.   PLUMMER M BN, COWLES K, VINES K. CODA: Convergence Diagnosis and Output

Analysis for MCMC. R News. 2012(6):7-11.

47.   Sobota RS, Shriner D, Kodaman N, Goodloe R, Zheng W, Gao YT, et al. Addressing

population-specific multiple testing burdens in genetic association studies. Ann Hum Genet.

2015;79(2):136-47.

48.   Szpiech ZA, Blant A, Pemberton TJ. GARLIC: Genomic Autozygosity Regions

Likelihood-based Inference and Classification. Bioinformatics. 2017;33(13):2059-62.

49.   Fisher GG, Ryan LH. Overview of the Health and Retirement Study and Introduction to

the Special Issue. Work Aging Retire. 2018;4(1):1-9.