1 Recent fluctuations in Mexican American genomes have altered the genetic architecture of
2 biomedical traits
3

Authors: Melissa L. Spear[1,2*], Alex Diaz-Papkovich[3,4], Elad Ziv[5,6,7,8], Joseph M. Yracheta[9,10]
Simon Gravel[4,11], Dara G. Torgerson[4,11], Ryan D. Hernandez[2,4,7,11,12,13*]

1. Biomedical Sciences Graduate Program, University of California, San Francisco, San

Francisco, CA, USA

2. Department of Bioengineering and Therapeutic Sciences, University of California, San

Francisco, San Francisco, CA, USA

3. Quantitative Life Sciences Program, McGill University, Montreal, QC, Canada

4. McGill University and Genome Quebec Innovation Center, McGill University, Montreal, QC,

Canada

5. Division of General Internal Medicine, University of California, San Francisco, San Francisco,

CA, USA

6. Department of Medicine, University of California, San Francisco, San Francisco, CA, USA

7. Institute of Human Genetics, University of California, San Francisco, San Francisco, CA, USA

8. Helen Diller Family Comprehensive Cancer Center, University of California, San Francisco,

San Francisco, CA, USA

9. Native BioData Consortium, Eagle Butte, SD, USA
20
21 10. Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD, USA
22
11. Department of Human Genetics, McGill University, Montreal, QC, Canada

12. Quantitative Biosciences Institute, University of California, San Francisco, San Francisco,

CA, USA

13. Institute for Human Genetics, University of California, San Francisco, CA, USA

*Corresponding authors: mlspear09@gmail.com and ryan.hernandez@me.com

Abstract

People in the Americas represent a diverse group of populations with varying degrees of

admixture among African, European, and Amerindigenous ancestries. In the United States, many

populations with non-European ancestry remain understudied, and thus little is known about the

genetic architecture of phenotypic variation in these populations. Using genome-wide genotype

data from the Hispanic Community Health Study/Study of Latinos, we find that Amerindigenous

ancestry has increased over time across Hispanic/Latino populations, particularly in Mexican

Americans where Amerindigenous ancestry increased by an average of ~20% over the 50-year

period spanning 1940s-1990s. We find similar patterns across American cities, and replicate our

observations in an independent sample of Mexican Americans. These dynamic ancestry patterns

are a result of a complex interaction of several population and cultural factors, including strong

ancestry-related assortative mating and subtle shifts in migration with differences in

subcontinental Amerindigenous ancestry over time. These factors have shaped patterns of

genetic variation, including an increase in runs of homozygosity in Amerindigenous ancestral

tracts, and also influenced the genetic architecture of complex traits within the Mexican American

population. We show for height, a trait correlated with ancestry, polygenic risk scores based on

summary statistics from a European-based genome-wide association study perform poorly in

Mexican Americans. Our findings reveal temporal changes in population structure within

Hispanics/Latinos that may influence biomedical traits, demonstrating a crucial need to improve

our understanding of the genetic diversity of admixed populations.

Introduction

The United States Census Bureau refers to the Hispanic/Latino ethnicity as a self-

identified category for individuals with ancestry deriving from Spain and the Spanish-speaking

countries of the Americas. As such, this broad ethnic group living in the United States is a

culturally, phenotypically, and genetically diverse continuum of populations. Individuals who

identify as Hispanic/Latino have varying proportions of Amerindigenous, African, and European

genetic ancestry, each with its own unique continental demographic history. Demographic forces

such as population bottlenecks and expansions, migration and adaptation to novel environments

resulted in observable differences in continental patterns of genetic variation (1-3). These differing

patterns were shaped by many historical events of migration which partially included the founding

of the Americas by Amerindigenous populations, the colonization by Europeans, and the African

slave trade (4-8), however additional complexities surrounding these events remain highly

understudied. These large-scale migrations and additional demographic events shaped the

genetic diversity of individuals currently living within the United States (9-13).

Demographic history has shaped the genetic architecture of modern human phenotypic

variation (14-19), and is critical to consider in the search for the genetic basis of complex diseases.

The demography of the United States has changed drastically over the 20th century, and by 2044

is predicted to become a 'minority-majority' country whereby no one racial/ethnic group comprises

more than 50% of the population (20). By 2060 Hispanics/Latinos are projected to make up the

largest of that share at 29% or 119 million individuals (20). However, to date, population-based

medical genomics research [and its subsequent benefits, including polygenic risk score (PRS)

profiling] have been disproportionately focused on individuals of European descent, with the

findings primarily benefiting European populations (21, 22). Despite the increases in sample

sizes, rates of discovery, and traits studied, Hispanic or Latin American participation in genome-

wide association studies (GWAS) has continued to hover around 1% (23, 24). This trend, along

with factors ranging from research abuse and community mistrust to community superstition and

apathy have led to a situation where these populations (and other non-European populations) are

particularly vulnerable to falling behind in receiving the benefits of the precision medicine

revolution (22, 23).

In this study we utilize the largest genetic study of Hispanics/Latinos in the U.S. to date --

the Hispanic Community Health Study/Study of Latinos (HCHS/SOL) (10) -- to understand how

patterns of genetic variation in Hispanic/Latino populations in the United States have changed

over the last century, and evaluate the impact such changes may be having on complex traits.

Results

Global ancestry proportions among HCHS/SOL Hispanic/Latino Populations

Using the subset of sites that overlapped with our African, European, and Amerindigenous

reference panels, we called 3-way global ancestry estimates for 10,268 unrelated HCHS/SOL

individuals (see methods). Figure 1A summarizes the global ancestry proportions shaded by

admixture estimates in a ternary plot, recapitulating the original HCHS/SOL analysis of continental

ancestry (10). However, while several population groups appear to have overlapping ancestry

proportions, this analysis masks more subtle structure in subcontinental ancestry. To investigate

subtle population structure across these self-identified population groups, we performed UMAP

on the top 3 principal components (see methods), and find substantial structure across self-

identified groups (Figure 1B and Supplementary Figure 1B). We find that Dominicans, who have

the highest average proportions of African ancestry, are in the middle, with Puerto Ricans and

Cubans, diverging in opposite directions (Supplementary Figure 1B) with clines of increasing

European ancestry proportions (Figure 1B). Further, while self-identified Mexican, Central, and

South American groups appear to have overlapping ancestry proportions in Figure 1A, UMAP

represents the Mexican Americans and Central/South American groups as large, separate wings

that diverge from self-identified Cubans and Dominicans, with both clusters diverging with clines

of increasing ancestry toward different Amerindigenous populations (Figure 1B and

Supplementary Figure 1B and 1C). While UMAP places each of the Amerindigenous populations

and the CEU population at the border of one or more HCHS/SOL clusters, UMAP isolates the YRI

samples into a distinct island suggesting that the use of this single African population may be

suboptimal for studying African ancestry in the populations sampled by HCHS/SOL

(Supplementary Figure 1C).

Dynamic Global Ancestry Proportions in Mexican Americans

For each of the HCHS/SOL populations, we evaluated differences in global ancestry estimates over time while accounting for the sampling method (referred to as "sampling weight", see methods) used for the design of the HCHS/SOL study (25). We found that in all populations, the effect size for Amerindigenous ancestry on birth year is positive, though only statistically significant after multiple testing in the Mexican American (β =0.0023; P=3.58E-22; Figure 1C) and Central American (β=0.0013; P=0.0013) cohorts (Supplementary Table 1). Due to the larger sample size, magnitude of the effect, and statistical significance, we shift our focus to Mexican Americans. In Mexican Americans, the increase in Amerindigenous global ancestry over time was consistent across multiple data stratifications including recruitment region, US born or not US born, educational attainment, and sex (Table 1), and was robust to alternative methods for estimating global ancestry proportions (e.g. based on the summation of RFMix local ancestry estimates; Supplementary Figures 2 and 3). We performed bootstrap resampling (n=1000) of global Amerindigenous ancestry for the Mexican Americans and observed a consistent increase in Amerindigenous ancestry with fitted LOESS smoothing (Figure 1D) and when individuals were binned by birth year decades (Supplementary Figure 4). On average, global Amerindigenous ancestry has increased ~20% over the last 50 years in Mexican Americans.

We replicated the increase in global Amerindigenous ancestry over time in a smaller, independent cohort of self-identified Mexican Americans (n=705) from the Health and Retirement Study (HRS) [34]. The HRS Mexican Americans in this study are older compared to the HCHS/SOL Mexican Americans (birth year distribution: 1915-1981; mean=1943,median:1942) and have lower levels of global Amerindigenous ancestry on average (mean=0.29), but we still observed an increase in global Amerindigenous ancestry over time (β=0.00082; P=0.02; SE=0.0003673; Supplementary Figure 5A). We performed 1000 bootstrap resampling iterations of the linear regression model (global Amerindigenous ancestry ~ birth year) fitted to the data.

From these resampling iterations, 98.2% of the tests had a slope > 0 (average β=0.00083) and

61.5% of the regression p-values were less than 0.05 as illustrated in Supplementary Figures 5B-

5D.

A previous study (12) identified ancestry biased migration in African Americans where

individuals with higher proportions of European ancestry migrated first out of the South during the

Great Migration followed by individuals with higher proportions African ancestry. We hypothesized

that earlier immigrants to the US had higher proportions of European ancestry followed by recent

immigrants having higher proportions of global Amerindigenous ancestry. In our non-US born

individuals (N=2987), we evaluated differences in ancestry estimates over time while accounting

for years in the US and sampling weight and identified a significant effect of years in the US (β=-

0.0009; P=0.0006; SE=0.0003). However, this did not change the effect of birth year on the

proportion of global Amerindigenous ancestry (β =0.0028; P<2e-16, SE=0.0003).

For US born individuals we assessed whether parental birth place could explain the

increases in global Amerindigenous ancestry. Of the 634 US born individuals, 385 had parents

both born outside of the US, 149 had one parent born outside of the US. 97 had both parents

born within the US. For the 385 individuals with both parents born abroad, we identified a

significant association between birth year and global Amerindigenous ancestry (β=0.004,

P=2.34e-10, SE=0.0006). For the remaining individuals we were unable to identify a significant

effect of birth year on global Amerindigenous ancestry, possibly due to a small sample size.

Individual loci are not driving global ancestry proportions

We used local ancestry estimates generated across the genome to perform admixture

mapping in HCHS/SOL Mexican Americans to determine if younger individuals harbored excess

Amerindigenous ancestry in certain regions of the genome. Although we tried two different models

(see methods), we did not find any loci to be significantly associated with birth year across the

genome (Supplementary Figure 6).

Little evidence for subcontinental population structure

It is possible that the increase in global Amerindigenous ancestry over time could be

biased by changes in the specific subcontinental Amerindigenous ancestries over time (though

such an effect is not visible in our UMAP analysis, Figure 1B). If it were the case, then we would

expect subtle signals of genetic divergence in Amerindigenous ancestry tracts over time. To

investigate this, we calculated F_{ST} between all pairs of birth-decades (see methods). Figure 2A

shows all pairwise comparisons among birth-decades, and demonstrates that while the estimates

of F_{ST} are negligible (with many estimates below 0), there is a subtle trend of increasing F_{ST} as

birth-decade differences increase (though individuals born in the 80s and 90s show a conflicting

pattern). We further investigated this pattern using genetic diversity, π, in Amerindigenous

ancestry tracts for each birth-decade (see methods). We hypothesized that if there were

increased migration from multiple Amerindigenous source populations (coupled with rapid

population growth in Mexican American communities), then genetic diversity should be increasing

over time. We found the opposite: Supplementary Figure 7 shows a subtle decrease in genetic

diversity (π) over time from the 1930s to the 1980s in non-US born Mexican Americans, and a

subtle decrease in US born Mexican Americans from the 70s to the 90s (while remaining roughly

constant from the 30s to the 70s).

Little evidence that Amerindigenous ancestry tract lengths have changed

We next sought to test whether differences at the local ancestry level could explain the

shift in global Amerindigenous ancestry over time in the Mexican Americans. We calculated the

length of each RFMix inferred local ancestry tract in each Mexican American individual, and tested

for differences in the distribution of tract lengths across birth-decades using a multiple linear

regression model (see methods). We found no significant associations between the decade bin

and the proportion of Amerindigenous ancestral tracts at various lengths (Figure 2B), even when

testing for violations of model assumptions (e.g. normalizing the tracts per bin by the number of

individuals, or excluding the 1930s and/or 1990s individuals due to the small sample size in each

bin).

Increased runs of homozygosity over time

Since genetic diversity has decreased over time in the Amerindigenous ancestry tracts of

Mexican Americans (despite rapid growth of the census population size), it is possible that this

population has also experienced increased haplotype homozygosity over time. We investigated

this possibility by exploring runs of homozygosity (ROH) in Amerindigenous ancestry tracts in

each of the 3622 Mexican Americans. We classified ROH into three categories: short, medium,

and long, based on the length distribution in the population. Generally, short ROH are tens of

kilobases in length and likely reflect the homozygosity of old haplotypes; medium ROH are

hundreds of kilobases in length and likely reflect background relatedness in the population; and

long ROH are hundreds of kilobases to several megabases in length and are likely the result of

recent parental relatedness. Figure 2C shows a fitted loess curve to the log of the total length of

ROH summed across each Mexican American's genome as a function of their birth year, broken

down by ROH size class (as well as the total of each size class that overlaps all ancestry tracts

(Supplementary Figure 8A) and Amerindigenous ancestry tracts (Supplementary Figure 8B)).

Overall, we find a significant positive correlation between birth year and the total summed ROH

across size classes (τ=0.0449, P=6.12e-5, Kendall's rank correlation), but this becomes more

significant when we restrict our analysis to ROH calls that overlap Amerindigenous ancestry tracts

(τ=0.0873, P=9.46e-15). When stratified by size class, the associations (all Kendall's rank

correlation) in ROH were primarily driven by the short (τ=0.0833, P=9.45e-14), and medium

(τ=0.0718, P=1.46e-10) size classes, and are again strongest when ROH overlap

Amerindigenous ancestry tracts (short τ=0.107, P<2.2e-16, medium τ=0.1003, P<2.2e-16). The

long ROH had a negative correlation with birth year, but was insignificant after multiple testing

(τ=-0.0291, P=0.01499; note that 1694 individuals did not have any long ROH calls in their

genome).

Strong ancestry-related assortative mating in HCHS/SOL Mexicans

Given that short and medium length ROH have increased over time, it appears that background

relatedness within Amerindigenous ancestry in Mexican Americans has increased over time (but

not an increase in recent parental relatedness). One way for this to occur is if individuals with

similar ancestry patterns tend to mate with one another more often than expected under a model

of random mating (i.e. assortative mating). To measure assortative mating, we estimated the

ancestral proportions of the biological parents of each HCHS/SOL Mexican American (see

methods). With individuals from all decades pooled together, we found the inferred biological

parental Amerindigenous ancestries to be significantly correlated (Figure 2D, r=0.708, 95%

CI:0.69-0.72, P<2.2e-16, Pearson correlation). When stratified by decade, the correlation in

inferred parental Amerindigenous global ancestry ranged from 0.65 to 0.74 (Supplementary

Figure 9), but were not statistically different from each other. This shows that there was a strong

parental ancestry correlation among Mexican Americans over different generations. This

signature of assortative mating is not due to recent parental relatedness, because there is no

trend in long ROH with birth year (and an overall low rate of long ROH among Mexican

Americans).

Genetic correlation of global Amerindigenous ancestry with biomedical traits

We have shown that genetic variation in the Mexican American population is dynamic, with

Amerindigenous ancestry increasing over a short period of time (combined with decreased

genetic diversity and increased short and medium length ROH within Amerindigenous ancestry

tracts). These features may have implications for the genetic architecture of complex traits within

Mexican Americans, a topic that is understudied and poorly understood. To further our

understanding of the genetic architecture of complex traits in Mexican Americans, we investigated

the relationship between Amerindigenous ancestry and various complex traits that may be

relevant to biomedical phenotypes. Specifically, we tested for a correlation between 66 complex

traits from the HCHS/SOL phenotypic dataset and global Amerindigenous ancestry (Kendall's τ).

As illustrated in Figure 3, 22 of these traits (33%) are significantly correlated after Bonferroni

correction (P<0.00076). We found that the effect of global Amerindigenous ancestry on many of

these phenotypes persisted when using multiple regression to account for age, sex, center, and

the sampling weight (Supplementary Table 2), highlighting the need for increased investigation

into the role of Amerindigenous genetic ancestry in admixed populations such as Mexican

Americans.

Assessing the genetic contribution of Amerindigenous ancestry to height

Among the traits we tested for a correlation with global Amerindigenous ancestry, height

had the strongest negative correlation, and our regression model indicated that height also had a

strong positive relationship with birth year (Figure 4A and Supplementary Table 3). Globally,

populations have grown taller over time due to a variety of non-genetic, environmental factors

(26). We find a similar trend in the HCHS/SOL Mexican Americans (Figure 4A). Indeed, when we

stratified individuals by quartiles of global Amerindigenous ancestry, we see that all quartiles have

increased in height by a similar amount over the period investigated (though individuals with lower

Amerindigenous ancestry were taller on average).

Height is one of the most highly studied complex traits, with GWAS sample sizes

numbering in the hundreds of thousands (27). Results for many of these studies have been made

readily available on public databases as summary association statistics that can be leveraged to

build genetic predictions through polygenic risk scores (PRS) (28). In Europeans, PRS have been

shown to have great predictive power for several traits, including breast cancer, prostate cancer,

and type 1 diabetes (22, 29-31). PRS are most effective in populations of European descent as

GWAS studies have been primarily performed in these populations (21-23) and are expected to

be biased when applied to other populations due to differences in the genetic architecture of traits

across diverse populations (32). Since Mexican Americans have some fraction of European

ancestry, we sought to determine whether PRS calculated utilizing GWAS summary statistics

from European populations could still provide useful insight.

To evaluate the effectiveness of PRS for height (i.e. the polygenic height score, or PHS,

see methods), we first tested whether there was an association between the observed height and

the predicted height estimates while controlling for sampling weight, sex, and recruitment center

(see methods). We identified a significant association between observed height and predicted

height for the population as a whole (β=0.0044881, P=2.19e-12; Figure 4B, Supplementary Table

4). However, when we stratified by quartiles of Amerindigenous global ancestry, the association

only remained for the individuals in the lower two quartiles of global Amerindigenous ancestry

proportions (AIA<0.37: β=0.004, P=0.0008 and 0.36<AIA<0.46: β=0.004, P=0.003,

Supplementary Table 4). The association between predicted height and observed height was no

longer significant for individuals in the upper two quartiles of global Amerindigenous ancestry

proportions (0.46<AIA<0.58: β=0.0011, P=0.39 and 0.58<AIA: β=0.0022, P=0.08,

Supplementary Table 4).

As we had found global Amerindigenous ancestry to be increasing over time, we

hypothesized that there would be a change in PHS over time as well. However, we find little

evidence supporting this hypothesis. While individuals born earlier than 1950 or in the 1950s have

a stronger correlation between their PHS and observed height (β=0.034 and 0.039; p=5.6e-4 and

2.7e-7 respectively) than individuals born in the 1960s, 1970s, or 1980s (β=0.016, 0.029, and

0.029; p=0.044, 0.0066, and 7.8e-5 respectively), there is no clear trend and we did not find a

significant effect of birth year on PHS (P=0.09) even when we stratified by the quartiles of global

Amerindigenous ancestry.

Discussion

The United States is a dynamic, rapidly changing population, and this will continue to occur

as the population size grows (20). Hispanics/Latinos are the largest and fastest growing minority

group, and are projected to comprise over 25% of the US population by 2060. They are a

genetically and phenotypically diverse population as a result of extensive admixture between

Amerindigenous populations and immigrants from multiple geographic locations around the world.

In this study, we identified additional population substructure complexities that may contribute to

phenotypic variation within Hispanics/Latinos.

Specifically, we demonstrated how the admixture dynamics of Mexican Americans have

changed over time, resulting in an increase of ~20% Amerindigenous ancestry on average over

the 50-year period studied. This change in ancestry is equivalent to a mean increase in

Amerindigenous ancestry of ~0.4% per year. While the effect sizes vary to some extent, we

replicate the underlying pattern across multiple data stratifications (two metropolitan cities, US

born and non-US born) and also replicate this feature in an independent cohort of Mexican

Americans. Further, we find that a similar trend holds across multiple self-identified

Hispanic/Latino populations in the US (and is statistically significant in Central Americans). This

effect does not appear to have a simple explanation: we do not see any statistically significant

increases at individual loci, we do not see more than a negligible degree of population

differentiation over time, and this increase cannot be entirely explained by very recent migration.

We do, however, find that as Amerindigenous ancestry has increased, genetic diversity within

Amerindigenous ancestry tracts across Mexican Americans has decreased over time, and is

associated with increased short and medium length ROH over time. This suggested that there

could be increased relatedness within Amerindigenous ancestries within Mexican Americans, and

we confirmed that there is a very high degree of ancestry-based assortative mating within the

Mexican American population.

What could be driving the increased Amerindigenous ancestry in Mexican Americans?

Population genetic theory suggests that while assortative mating could result in increased ROH

and decreased genetic diversity, ancestry-based assortative mating alone should not result in

mean changes in global ancestry proportions. Regardless of the underlying mechanisms driving

increased Amerindigenous ancestry in Mexican Americans, this additional source of temporal

substructure within this population has substantial consequences for phenotypic variation in

biomedical traits. We identify several biomedical traits that are correlated with Amerindigenous

ancestry, and show that in the case of height, there are both ancestry and temporal effects.

Further study is necessary to understand whether other biomedical traits are also changing over

time as the genomic ancestry proportions change in this population.

Interestingly, we identified another source of structure within HCHS/SOL, particularly in

the African ancestral component of Hispanics/Latinos. In our UMAP analysis, the YRI sample

form their own cluster as a reference population as compared to the Amerindigenous and

European reference populations which border the admixed samples with the highest proportion

of each ancestry, respectively (Supplementary Figure 1C). While most Latin Americans can trace

their African ancestry to Sub-Saharan Africa, previous studies have also identified hidden

Northern African ancestry in individuals from Southern Europe (33-35), who primarily colonised

the Americas. This may explain why the YRI sample is not at the boundary of the individuals with

the highest proportion of African ancestry in the HCHS/SOL sample as the African ancestral

component may be more complex. Our results suggest how careful consideration must be taken

into account when selecting reference populations to study the African ancestral component of

admixed individuals from the Americas.

In our study, we bring specific attention to the biases that continue to exist with using

European GWAS summary statistics to calculate polygenic risk scores in admixed populations

such as Mexican Americans that are comprised of European, Amerindigenous, and African

genetic ancestries. In particular, in the case of height, we found that the polygenic height score

(PHS) correlated with observed height only in the subset of individuals with the lowest levels of

Amerindigenous ancestry (i.e. the subset of individuals with highest European ancestry). As the

population dynamics of the US continue to change, it is imperative that we study diverse

populations, or we risk exacerbating the health disparities that currently exist. To date, population-

based medical genomics research (and its subsequent benefits) have been disproportionately

focused on populations of European ancestry. In order to improve the design and implementation

of medical genetics studies for the ethnically diverse U.S. population, we need detailed insights

into the population history of diverse U.S. populations. This includes characterizing the admixture

dynamics of Hispanic/Latino populations, as well as the evolutionary forces that shaped patterns

of genetic variation of the ancestral populations that contributed to modern day Hispanic/Latino

populations.

The genetic variation of the Hispanic community in the United States belies categorization

under a single label (10). The events that have shaped and continue to shape this genetic diversity

are complex, numerous, and nuanced, and the social history of such a diverse population is

intrinsic to any genetic study. Mexico's society was largely defined by an established social caste

system based on ancestry, which disappeared after Mexico's independence in 1821 (36). Even

so, social inequalities persist today with skin colour having a significant effect on wealth and

education (37). A multitude of factors within and outside Mexico — whether related to trade,

immigration policies, or armed conflicts — acted to influence who immigrated to the United States,

and the impact of each of these fluctuates over time (38-40). These changes shift the

demographics of immigration, which is inherently related to the genetic ancestry of the population.

Consequently, this shapes the genetic architecture of complex traits. Diverse populations

are at risk not only from underrepresentation in research, but because of poor understanding of

the temporal and spatial dynamics at play in genetic variation. The promise of equitable precision

medicine — one of the ultimate goals of medical genomics — cannot be kept without

understanding this interplay. Health disparities in the United States are fed by structural

inequalities. For example, studies that use modern Artificial Intelligence techniques have already

been shown to inflate existing disparities between Black Americans and White Americans (41).

Such biases, whether from algorithms, study designs, or misunderstandings of subtleties in

data, feed into the larger systemic pressures faced by minority populations in the United States.

While we have shown a dramatic shift in ancestry proportions in US Hispanic/Latinos, one

of the caveats of this study is that the HCHS/SOL cohort is not representative of all US

Hispanics/Latinos. HCHS/SOL participants were recruited at four primary centers: Bronx,

Chicago, Miami, and San Diego. There may be additional genetic diversity that has not been

captured by this dataset and trends exhibited in this dataset may not translate to Hispanic/Latino

populations living in other regions of the US (though the temporal increase in Amerindigenous

ancestry was replicated in an independent sample of Mexican Americans). Further, we have only

assembled a reference panel with limited numbers of individuals with various Amerindigenous,

European, and African ancestry. With better population genetic modeling and a deeper

understanding of the social and historical aspects of Hispanic/Latino populations, we will be able

to improve our understanding of the genetic and phenotypic diversity across these populations,

and subsequently improve our ability to understand genetic contributions to complex traits and

disease. These insights will lead to optimization of population sampling for the design of future

medical genetic studies, the identification of disease risk variants, and ultimately, precision

medicine for all.

Methods

Study dataset and initial quality control

The HCHS/SOL study is a community-based cohort study of self-identified Hispanic/Latino

individuals from four US metropolitan areas with the general goal of identifying risk and protective

factors for various medical conditions including cardiovascular disease, diabetes, pulmonary

disease, and sleep disorders (25). 12,434 participants with birth year estimates between 1934-

1993 who self-identified as being of Cuban, Dominican, Puerto Rican, Mexican, Central American,

or South American background consented to genetics studies and posting of their genetic and

phenotype data on the publicly available Database of Genotypes and Phenotypes (dbGaP)

through Study Accession phs000810.v1.p1. Samples were genotyped on an Illumina custom

array, SoL HCHS Custom 15041502 array (annotation B3, genome build 37), consisting of the

Illumina Omni 2.5M array and 148,353 custom single nucleotide polymorphisms (SNPs) (10).

Data posted to dbGaP had passed initial sample quality control filters, including removing samples

with differences in reported vs. genetic sex, call rates > 95%, and evidence for sample

contamination (e.g. heterozygosity and sample call rates). For initial SNP quality control, we

filtered out SNPs that were monomorphic, positional duplicates, or Illumina technical failures, as

well as SNPs that had cluster separation <= 0.3, call rate <=2%, >2 disconcordant calls in 291

duplicate samples, >3 Mendelian errors in parent-offspring pairs/trios, Hardy-Weinberg

Equilibrium combined P-value $<10^{-5}$, and sex differences in allele frequency ≥0.2. Our filtering

resulted in 1,763,935 genotyped SNPs with minor allele frequency (MAF) >0.01.

Additional sample quality control performed in the HCHS/SOL dataset included filtering

out samples with 1) large chromosomal anomalies, 2) substantial Asian ancestry as previously

identified in HCHS/SOL (*12*) and 3) individuals with up to third degree genetic relatedness in the

dataset as inferred by REAP (42). For genetic relatedness filtering, individuals from pairs were

kept to maximize representation of the birth year distribution, which resulted in 10,268 unrelated

remaining individuals.

From the original HCHS/SOL analysis, individuals were classified into genetic-analysis

groups, similar to self-identified background groups in that they share cultural and environmental

characteristics, but are also more genetically homogenous (10).

Birth year for all individuals was estimated by subtracting the difference between date of

first clinic visit for the baseline examination (25) and age. Year of arrival was estimated by

subtracting the difference between date of first clinic visit for the baseline examination and years

in the US.

Global, local, and parental ancestry inference

All ancestry analyses were restricted to the 211,152 autosomal SNP markers that overlapped

between the study and reference panel genotyping array. For the HCHS/SOL dataset, global

African, European, and Amerindigenous ancestries were inferred with ADMIXTURE, in an

unsupervised manner, with K=3. Amerindigenous ancestry refers to estimates of Indigenous

genetic ancestry from the Americas. For some analyses, HCHS/SOL individuals with greater than

95% of a single ancestry (e.g African, European, or Amerindigenous) were filtered out resulting

in 9,913 individuals: 1,099 Central American, 1,536 Cuban, 954 Dominican, 3,622 Mexican, 1,783

Puerto Rican, 652 South American and 267 "Other" individuals.

Ancestral tracts, known as 'local' ancestry, along the genome for all HCHS/SOL

individuals were inferred using RFMix (43) and a three population reference panel, comprised of

315 individuals: 104 HapMap phase 3 CEU (European) and 107 YRI (African) individuals (44)

and 112 Amerindigenous individuals from throughout Latin America (8). The reference panel was

limited to individuals with 99% continental ancestry as inferred by unsupervised ADMIXTURE

(45). Prior to local ancestry inference, HCHS/SOL individuals were merged with the reference

panels and then phased using SHAPEIT2 (46). For all HCHS/SOL Mexican American individuals,

parental genomic ancestry was inferred with ANCESTOR (47) using the local ancestry estimates

generated by RFMix.

Uniform Manifold Approximation and Projection (UMAP)

Principal components for HCHS/SOL and the reference panel were computed using smartPCA

(48). UMAP (version 0.3.8) was run using the Python script freely available at

https://github.com/diazale/gt-dimred with parameter specification set at 15 nearest neighbours

and a minimum distance between points of 0.5.

Admixture mapping

Local ancestry estimates for 211,151 SNPs across the genome were used to perform admixture

mapping in HCHS/SOL Mexican Americans to determine if younger individuals harbored excess

Amerindigenous ancestry in certain regions of the genome. Admixture mapping was performed

applying two different models: 1) a linear regression model with age as the dependent variable

adjusting for global Amerindigenous ancestry, sampling weight and center and 2) a logistic

regression model dividing the HCHS/SOL Mexican cohort in to an older vs younger generation

with 1965 set as the dividing point while also adjusting for global Amerindigenous ancestry,

sampling weight, and center. The threshold for genome-wide significance, 1.38×10^{-4} was

calculated using the empirical autoregression framework with the package *coda* in R to estimate

the total number of ancestral blocks (49, 50).

Tract Lengths

The multiple regression model: $\log(f) = \beta_0 + \beta_1\ T + \beta_2\ A + \beta_3\ TA + \varepsilon$, where f is a matrix

containing the proportion of lengths of all ancestral tracts across the genome for all 3622 Mexican

American individuals, T the tract length bin and A decade of birth year bin, was used to test for an

effect of birth decade on the proportion of Amerindigenous ancestral tract lengths. For

assessment between the fraction of ancestry tracts in an individual's genome and birth year, long

tract cutoffs were chosen based on tract separation between the birth year decades in Figure 2B.

Diversity Calculations

Subcontinental ancestry was assessed using the diversity measurements π and F_{ST}. π was

calculated as the average number of pairwise genetic differences among all pairs of overlapping

Amerindigenous ancestry tracts across individuals. F_{ST} was calculated as:

$$F_{ST} = (H_T - H_S)/ H_T$$

where H_T is the average heterozygosity when all individuals are pooled across decades and H_S is

the average heterozygosity within each decade of individuals.

Inference of Runs of Homozygosity

Runs of homozygosity (ROH) were called using the program GARLIC v1.1.4 (51) on 211,152 sites

for the Mexican American individuals. An analysis window size of 50 SNPs and an overlap fraction

of 0.25 were both chosen using GARLIC's rule of thumb parameter estimation. GARLIC chose a

LOD score cutoff of 0. Using a three-component Gaussian mixture, GARLIC determined class

A/B (short/medium) and class B/C (medium/long) size boundaries as 845,097 bp and 2,501,750

bp, respectively.

Imputation

481   Imputation for HCHS/SOL was performed locally using IMPUTE2 with the 1000 Genomes Project

482   Phase 3 haplotypes used as a reference panel. After filtering on an info score cutoff of 0.3, this

483   resulted in 33,041,084 SNPs.

Polygenic Risk Score Calculations

Polygenic risk scores for height were calculated using the publicly available UK Biobank

(UKBB) GWAS Round 2 Summary Statistics retrieved from http://www.nealelab.is/uk-biobank.

Briefly, for sample quality control, sample inclusion was limited to unrelated samples who passed

the sex chromosome aneuploidy filter. British ancestry was determined using the 1st 6 PCs;

individuals more than 7 standard deviations away from the 1st 6 PCs were excluded. Further

filtering included limiting to self -reported 'white-British' / 'Irish' / 'White' resulting in a QCed sample

count of 361,194 individuals https://github.com/Nealelab/UK_Biobank_GWAS#imputed-v3-

sample-qc. An imputation panel of ~90 million SNPs from HRC, UK10K and 1KG were used to

impute genotypes. 13.7 million autosomal and X-chromosome SNPs passed quality control

thresholds including Info score>0.8, MAF>0.0001, and HWE p-value>1e-10. For the phenotype,

a linear regression model in Hail was run for all individuals (both sexes) adjusting by the first 20

PCs + sex + age + age^2 + (sex*age) + (sex*age)^2. For height, there was complete phenotype

information for 360,388 individuals.

Risk scores were calculated by extracting the overlapping genome-wide significant hits

initially discovered in the UKBB GWASs of height and selecting SNPs with the lowest p-value in

each 1Mb window across the genome. For height this resulted in a dataset of 1,103 overlapping

SNPs that were present in our dataset of genotyped and imputed SNPs.

Health and Retirement Study (HRS)

For replication, we used genotype data from 705 self-identified Mexican-Americans from the

Health and Retirement Study (HRS) (52), genotyped on the Illumina Human Omni 2.5M platform.

HRS data was made available under IRB Study No. A11-E91-13B - The apportionment of genetic

diversity within the United States. Estimated global ancestry proportions for the Mexican American

population in the HRS were calculated as in Baharian et al. (12), which used an alternative

reference panel and alternative ancestry inference approach. Briefly, RFMix was used to infer

local ancestry estimates across the genome utilizing CHS, YRI, and CEU individuals from the

1000 Genomes Project as reference populations for Amerindigenous/Asian, African, and

European ancestries, respectively. Global ancestry estimates were calculated using the summed

RFMix calls.

Statistical Analyses and Plots

Statistical analyses and plot generation were performed within Rstudio using Version 1.1.463 and

R version 3.5.3. ternary and ggridges/ggplot2 packages were used to create the simplex and

ridgeline plots.

Acknowledgements

We thank many colleagues who commented on our preprint prior to submission, particularly Reed

Cartwright for suggestions on terminology. MLS was supported through the National Human

Genome Research Institute (NHGRI) of the National Institutes of Health (NIH) under Award

Number F31HG010104. ADP and SG were supported, in part, thanks to funding from the

Canada Research Chairs program and CIHR grant MOP-136855. RDH was supported, in part,

by NHGRI grant R01HG007644 and the Canadian Research Chairs program.

Declaration of Interests

The authors declare no competing interests.

Figure 1. Recent dynamics continually shape the continuum of continental ancestry Hispanic/Latino populations. A. Ternary plot of HCHS/SOL (n=10,268) colored by admixture proportions. B. Uniform Manifold Approximation and Projection (UMAP) plot depicting the genetic diversity of HCHS/SOL and the reference panel (n=10,591) using 3 principal components, colored by admixture proportions (see Supplemental Fig 1 for population labels). Within the legend, AFR, EUR, and AI refer to African, European, and Amerindigenous global ancestries, respectively. C. Global Amerindigenous ancestry proportions plotted by birth year for Mexican Americans (n=3,622). Fitted line is multiple regression of Amerindigenous ~ birth year + sampling weight. Bars represent 95% confidence intervals for individuals grouped by decade. D. Bootstrap resampling (n=1000 iterations) of Amerindigenous global ancestry for the Mexican American individuals with a fitted LOESS regression line for each iteration. Dashed lines represent the 95% confidence interval and the blue line represents the fitted regression line from Figure 1C.

Figure 2. Diversity of and within Amerindigenous ancestral tracts. A) F_{ST} estimates

calculated between each decade group. Bars represent the 95% CI. B) Proportion of total

Amerindigenous (AI) ancestral tracts in the HCHS/SOL Mexican American population by decade.

C) Loess regression of the log of the sum of total ROH and ROH overlapping Amerindigenous

(AI) ancestral tracts separated by ROH class. Total long ROH is not represented as an individual

group due to the high number of individuals missing long ROH (1694 for long ROH across

ancestries and 1987 for long AI ROH) but was included in the sum of "All ROH" and "All AI ROH".

D) Correlation of parent's inferred global Amerindigenous (AI) ancestries using ANCESTOR.

554
555
556

Figure 3. Correlation of 66 quantitative traits with global Amerindigenous ancestry.

Significance level was determined using Bonferroni correction adjusting by the number of

quantitative traits tested (0.05/66=0.00075).

585   **Figure 4: Height and global Amerindigenous ancestry in HCHS/SOL Mexican Americans.**

586   Each plot illustrates the relationship between A) Birth year and height B) Height and polygenic

587   height score (PHS). The black line indicates the fitted linear model for all individuals. Each color

588   represents a different quartile of Amerindigenous global ancestry. Polygenic height scores were

589   assessed utilizing UKBB summary statistics for 1,128 SNPs.

Table 1: Relationship of Amerindigenous global ancestry and birth year for Mexican

Americans stratified by recruitment region, US born vs non-US born status, sex and

educational attainment. For recruitment region, data stratification was limited to Chicago

and San Diego as sample size for the Bronx and Miami was limited: 124 and 25 individuals,

respectively. Education attainment was categorized as either less than a high school

diploma or equivalent degree (<HS), equal to a high school diploma or equivalent degree

(=HS), or post-secondary education (>HS).

Category	N	Mean	Median	R2	Effect	Std.Err	P
All	3622	0.489	0.468	0.027	0.0023	0.0002	3.58E-22
Chicago	1310	0.562	0.550	0.017	0.0016	0.0005	0.0006
San Diego	2163	0.428	0.422	0.012	0.0012	0.0002	4.29E-07
US born	634	0.427	0.418	0.063	0.0027	0.0004	1.77E-10
Not US born	2987	0.502	0.481	0.050	0.0032	0.0003	1.38E-30
Male	1500	0.494	0.475	0.038	0.0028	0.0004	3.83E-14
Female	2122	0.485	0.462	0.022	0.0019	0.0003	3.07E-10
<HS	1518	0.520	0.500	0.045	0.0026	0.0004	1.39E-12
=HS	960	0.501	0.479	0.022	0.0018	0.0005	0.0003
>HS	1140	0.436	0.422	0.045	0.0027	0.0004	6.53E-13

Supplementary Material

Supplementary Figure 1. Continental ancestral diversity of HCHS/SOL A) Ternary plot of

global ancestry proportions colored by population for 10,268 HCHS/SOL individuals B) Uniform

Manifold Approximation and Projection (UMAP) plot of HCHS/SOL and the reference panel

(n=10,591) using 3 principal components, colored by HCHS/SOL population. C) UMAP plot of

HCHS/SOL and the reference panel (n=10,591) using 3 principal components, colored by

reference population.

635
636

Supplementary Figure 2. Concordance of ADMIXTURE and RFMix global ancestry

estimates. A) Amerindigenous ancestry B) African ancestry and C) European ancestry.

648 **Supplementary Figure 3.** RFMix inferred Amerindigenous global ancestry proportions plotted

649 over time for HCHS/SOL Mexican Americans (n=3622).

668 **Supplementary Figure 4:** Distributions of Amerindigenous global ancestry means for

669 HCHS/SOL Mexican Americans (n=3622) generated by 1000 bootstrap resampling iterations

670 within each decade of binned birth years.

Supplementary Figure 5. Replication in the Health and Retirement Study for 705 self-identified Mexican Americans. A) Ancestry over time B) Distribution of regression slopes after 1000 bootstrap resampling iterations C) Distribution of bootstrap regression p-values D) ECDF of bootstrap regression p-values.

A

B

Supplementary Figure 6. Admixture mapping in HCHS/SOL Mexicans (n=3622) for

Amerindigenous ancestry and A) birth year and B) generation. Ancestry association testing

was performed at 211,151 markers using A) linear regression and B) logistic regression, both

including global Amerindigenous ancestry, sampling weight and center as covariates.

Supplementary Figure 7. Diversity of and within Amerindigenous ancestral tracts.

Diversity (π) of subcontinental Amerindigenous ancestry stratified by US born/not US born

status. π was calculated between pairs within each decade of birth years. 95% confidence

intervals are highlighted by the shaded regions for each group.

Supplementary Figure 8. Runs of homozygosity (ROH) in HCHS/SOL Mexican Americans.

A) ROH across all ancestries separated by ROH class B) ROH overlapping Amerindigenous

haplotypes separated by ROH class.

708 **Supplementary Figure 9. Ancestry-related assortative mating in HCHS/SOL Mexican**

709 **Americans separated by decade.** Each plot represents the correlation of parent's inferred

710 Amerindigenous (AI) ancestries using ANCESTOR by decade beginning with the 1930s (A) and

711 ending with the 1990s (G). Each point corresponds to one Mexican American couple and the

712 axes correspond to the inferred Amerindigenous (AI) ancestry of each partner.

720 **Supplementary Table 1: Association of global ancestries and birth year for all**

721 **HCHS/SOL individuals.** For each population, we tested for an association between global

722 ancestry and birth year while accounting for the sampling design. AI, AFR, and EUR refer to

723 Amerindigenous, African, and European ancestry respectively.

Population	Ancestry	N	R2	Effect	Std.Err	P
Central American	AI	1099	0.0139	0.0013	0.0004	0.0013
Central American	AFR	1099	0.0136	0.0002	0.0004	0.6561
Central American	EUR	1099	0.0138	-0.0015	0.0004	0.0002
Cuban	AI	1536	0.0023	0.0002	0.0001	0.0938
Cuban	AFR	1536	0.0014	-0.0005	0.0004	0.1490
Cuban	EUR	1536	0.0005	0.0003	0.0004	0.3879
Dominican	AI	954	0.0035	0.0002	0.0001	0.0663
Dominican	AFR	954	0.0030	-0.0007	0.0004	0.1287
Dominican	EUR	954	0.0022	0.0005	0.0004	0.2374
Mexican	AI	3622	0.0268	0.0023	0.0002	3.58E-22
Mexican	AFR	3622	0.0008	0.0000	0.0000	0.4189
Mexican	EUR	3622	0.0285	-0.0023	0.0002	0.0000
Puerto Rican	AI	1783	0.0014	0.0001	0.0001	0.1533
Puerto Rican	AFR	1783	0.0014	0.0003	0.0002	0.1743
Puerto Rican	EUR	1783	0.0027	-0.0005	0.0002	0.0355
South American	AI	652	0.0110	0.0016	0.0007	0.0211
South American	AFR	652	0.0027	-0.0002	0.0004	0.5053
South American	EUR	652	0.0080	-0.0014	0.0006	0.0335

734 **Supplementary Table 2 - Multiple regression table with traits that were significantly**

735 **correlated with global Amerindigenous ancestry.**

Trait	N	R2	Effect	Std.Err	P
Height	3615	0.588	-13.637	0.676	1.01E-85
Predicted FVC	3522	0.851	-695.376	36.867	1.13E-75
EGFR MDRD	3308	0.196	23.398	2.435	1.39E-21
EGFR CKD Epi	3308	0.441	14.752	1.581	1.90E-20
Waist to hip ratio	3617	0.195	0.043	0.007	6.96E-10
Glycosylated hemoglobin (HbA1c)	3609	0.081	9.184	1.590	8.30E-09
% Glycosylated hemoglobin	3609	0.080	0.837	0.146	9.88E-09
HDL cholesterol	3621	0.095	-6.740	1.372	9.39E-07
Total iron binding capacity	3620	0.066	23.540	5.467	1.71E-05
FEV1 FVC Ratio	3505	0.176	2.560	0.639	6.37E-05
% Lymphocytes	3442	0.021	3.768	1.023	2.35E-04
Hip girth	3617	0.077	-4.452	1.273	4.78E-04
% Neutrophils	3442	0.039	-3.649	1.178	1.96E-03
Monocyte count	3443	0.024	-0.048	0.019	1.27E-02
Neutrophil count	3442	0.043	-0.399	0.165	1.59E-02
LDL cholesterol	3529	0.047	-8.269	3.963	3.70E-02
Average diastolic blood pressure	3616	0.072	-2.067	1.136	6.88E-02
Total cholesterol	3622	0.065	-6.248	4.614	1.76E-01
White blood cell count	3442	0.032	-0.260	0.208	2.12E-01
Average systolic blood pressure	3619	0.211	1.590	1.712	3.53E-01
% Immature granulocytes	555	0.261	-0.090	0.110	4.14E-01
QRS duration	3596	0.168	0.062	1.268	9.61E-01

742 **Supplementary Table 3: Height over time: Height (cm) as a function of birth year**

743 **adjusting by sex, center, and sampling weight for 3614 Mexican Americans stratified by**

744 **the quartiles of global Amerindigenous ancestry (AIA).**

Group	N	R2	Effect	Std.Err	P
All	3614	0.54186	0.1200379	0.00905366	3.28E-39
AIA>0.58	929	0.5745454	0.158895	0.01783895	2.73E-18
0.46<=AIA<=0.58	955	0.5784636	0.1542762	0.01683919	3.07E-19
0.37<=AIA<0.46	842	0.5472275	0.1008054	0.01739819	9.73E-09
AIA<0.37	888	0.5369993	0.1161775	0.01815834	2.55E-10

748 **Supplementary Table 4: Predicted height vs. observed height.** Predicted height (cm) as

749 a function of observed height (cm) adjusting by sex, center, and sampling weight for 3614

750 Mexican Americans stratified by Amerindigenous ancestry (AIA).

Group	N	R2	Effect	Std.Err	P
All	3614	0.0249	0.0045	0.0006	2.19E-12
AIA>0.58	929	0.0072	0.0022	0.0012	7.79E-02
0.46<=AIA<=0.58	955	0.0058	0.0011	0.0013	3.90E-01
0.37<=AIA<0.46	842	0.0144	0.0043	0.0015	3.22E-03
AIA<0.37	888	0.0164	0.0043	0.0013	7.91E-04

References:

1.   Nelson MR, Bryc K, King KS, Indap A, Boyko AR, Novembre J, et al. The Population Reference Sample, POPRES: a resource for population, disease, and pharmacological genetics research. Am J Hum Genet. 2008;83(3):347-58.

2.   Genomes Project C, Abecasis GR, Auton A, Brooks LD, DePristo MA, Durbin RM, et al. An integrated map of genetic variation from 1,092 human genomes. Nature. 2012;491(7422):56-65.

3.   Genomes Project C, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, et al. A global reference for human genetic variation. Nature. 2015;526(7571):68-74.

4.   Gravel S, Zakharia F, Moreno-Estrada A, Byrnes JK, Muzzio M, Rodriguez-Flores JL, et al. Reconstructing Native American migrations from whole-genome and whole-exome data. PLoS Genet. 2013;9(12):e1004023.

5.   Homburger JR, Moreno-Estrada A, Gignoux CR, Nelson D, Sanchez E, Ortiz-Tello P, et al. Genomic Insights into the Ancestry and Demographic History of South America. PLoS Genet. 2015;11(12):e1005602.

6.   Moreno-Estrada A, Gignoux CR, Fernandez-Lopez JC, Zakharia F, Sikora M, Contreras AV, et al. Human genetics. The genetics of Mexico recapitulates Native American substructure and affects biomedical traits. Science. 2014;344(6189):1280-5.

7.   Moreno-Estrada A, Gravel S, Zakharia F, McCauley JL, Byrnes JK, Gignoux CR, et al. Reconstructing the population genetic history of the Caribbean. PLoS Genet. 2013;9(11):e1003925.

8.   Reich D, Patterson N, Campbell D, Tandon A, Mazieres S, Ray N, et al. Reconstructing Native American population history. Nature. 2012;488(7411):370-4.

9.   Bryc K, Durand EY, Macpherson JM, Reich D, Mountain JL. The genetic ancestry of African Americans, Latinos, and European Americans across the United States. Am J Hum Genet. 2015;96(1):37-53.

10.   Conomos MP, Laurie CA, Stilp AM, Gogarten SM, McHugh CP, Nelson SC, et al.

Genetic Diversity and Association Studies in US Hispanic/Latino Populations: Applications in the

Hispanic Community Health Study/Study of Latinos. Am J Hum Genet. 2016;98(1):165-84.

11.   Han E, Carbonetto P, Curtis RE, Wang Y, Granka JM, Byrnes J, et al. Clustering of

770,000 genomes reveals post-colonial population structure of North America. Nat Commun.

2017;8:14238.

12.   Baharian S, Barakatt M, Gignoux CR, Shringarpure S, Errington J, Blot WJ, et al. The

Great Migration and African-American Genomic Diversity. PLoS Genet. 2016;12(5):e1006059.

13.   Jordan IK, Rishishwar L, Conley AB. Native American admixture recapitulates

population-specific migration and settlement of the continental United States. PLoS Genet.

2019;15(9):e1008225.

14.   Agarwala V, Flannick J, Sunyaev S, Go TDC, Altshuler D. Evaluating empirical bounds

on complex disease genetic architecture. Nat Genet. 2013;45(12):1418-27.

15.   Eyre-Walker A. Evolution in health and medicine Sackler colloquium: Genetic

architecture of a complex trait and its implications for fitness and genome-wide association

studies. Proc Natl Acad Sci U S A. 2010;107 Suppl 1:1752-6.

16.   Maher MC, Uricchio LH, Torgerson DG, Hernandez RD. Population genetics of rare

variants and complex diseases. Hum Hered. 2012;74(3-4):118-28.

17.   Simons YB, Turchin MC, Pritchard JK, Sella G. The deleterious mutation load is

insensitive to recent population history. Nat Genet. 2014;46(3):220-4.

18.   Uricchio LH, Zaitlen NA, Ye CJ, Witte JS, Hernandez RD. Selection and explosive

growth alter genetic architecture and hamper the detection of causal rare variants. Genome

Res. 2016;26(7):863-73.

19.   Yang J, Bakshi A, Zhu Z, Hemani G, Vinkhuyzen AA, Lee SH, et al. Genetic variance

estimation with imputed variants finds negligible missing heritability for human height and body

mass index. Nat Genet. 2015;47(10):1114-20.

20.   Colby SLO, J.M. Projections of the Size and Compositon of the U.S. Population: 2014 to

2060. U.S. Census Bureau, Commerce USDo; 2015.

21.   Bustamante CD, Burchard EG, De la Vega FM. Genomics for the world. Nature.

2011;475(7355):163-5.

22.   Martin AR, Kanai M, Kamatani Y, Okada Y, Neale BM, Daly MJ. Clinical use of current

polygenic risk scores may exacerbate health disparities. Nat Genet. 2019;51(4):584-91.

23.   Popejoy AB, Fullerton SM. Genomics is failing on diversity. Nature. 2016;538(7624):161-

4.

24.   Mills MC, Rahal C. A scientometric review of genome-wide association studies.

Commun Biol. 2019;2:9.

25.   Sorlie PD, Aviles-Santa LM, Wassertheil-Smoller S, Kaplan RC, Daviglus ML, Giachello

AL, et al. Design and implementation of the Hispanic Community Health Study/Study of Latinos.

Ann Epidemiol. 2010;20(8):629-41.

26.   Collaboration NCDRF. A century of trends in adult human height. Elife. 2016;5.

27.   Yengo L, Sidorenko J, Kemper KE, Zheng Z, Wood AR, Weedon MN, et al. Meta-

analysis of genome-wide association studies for height and body mass index in approximately

700000 individuals of European ancestry. Hum Mol Genet. 2018;27(20):3641-9.

28.   Pasaniuc B, Price AL. Dissecting the genetics of complex traits using summary

association statistics. Nat Rev Genet. 2017;18(2):117-27.

29.   Sharp SA, Rich SS, Wood AR, Jones SE, Beaumont RN, Harrison JW, et al.

Development and Standardization of an Improved Type 1 Diabetes Genetic Risk Score for Use

in Newborn Screening and Incident Diagnosis. Diabetes Care. 2019;42(2):200-7.

30.   Schumacher FR, Al Olama AA, Berndt SI, Benlloch S, Ahmed M, Saunders EJ, et al.

Association analyses of more than 140,000 men identify 63 new prostate cancer susceptibility

loci. Nat Genet. 2018;50(7):928-36.

31.   Maas P, Barrdahl M, Joshi AD, Auer PL, Gaudet MM, Milne RL, et al. Breast Cancer

Risk From Modifiable and Nonmodifiable Risk Factors Among White Women in the United

States. JAMA Oncol. 2016;2(10):1295-302.

32.   Martin AR, Gignoux CR, Walters RK, Wojcik GL, Neale BM, Gravel S, et al. Human

Demographic History Impacts Genetic Risk Prediction across Diverse Populations. Am J Hum

Genet. 2017;100(4):635-49.

33.   Botigue LR, Henn BM, Gravel S, Maples BK, Gignoux CR, Corona E, et al. Gene flow

from North Africa contributes to differential human genetic diversity in southern Europe. Proc

Natl Acad Sci U S A. 2013;110(29):11791-6.

34.   Moorjani P, Patterson N, Hirschhorn JN, Keinan A, Hao L, Atzmon G, et al. The history

of African gene flow into Southern Europeans, Levantines, and Jews. PLoS Genet.

2011;7(4):e1001373.

35.   Chacon-Duque JC, Adhikari K, Fuentes-Guajardo M, Mendoza-Revilla J, Acuna-Alonzo

V, Barquera R, et al. Latin Americans show wide-spread Converso ancestry and imprint of local

Native ancestry on physical appearance. Nat Commun. 2018;9(1):5388.

36.   Lisker R, Ramirez E, Briceno RP, Granados J, Babinsky V. Gene frequencies and

admixture estimates in four Mexican urban centers. Hum Biol. 1990;62(6):791-801.

37.   Martinez DZ-CIF. Is Mexico a Post-Racial Country? Inequality and Skin Tone across the

Americas. AmericasBarometer Insights Series by the Latin American Public Opinion Project

(LAPOP), wwwLapopSurveysorg [Internet]. 2017. Available from:

https://www.vanderbilt.edu/lapop/insights/ITB031en.pdf.

38.   Fernández-Kelly P, Massey DS. Borders for Whom? The Role of NAFTA in Mexico-U.S.

Migration. The Annals of the American Academy of Political and Social Science. 2007;610:98-

118.

39.   Verea M. Immigration Trends After 20 Years of NAFTA. 2014. 2014;9(2).

40. Contreras VR. THE ROLE OF DRUG-RELATED VIOLENCE AND EXTORTION IN

PROMOTING MEXICAN MIGRATION: Unexpected Consequences of a Drug War. Latin

American Research Review. 2014;49(3):199-217.

41. Obermeyer Z, Powers B, Vogeli C, Mullainathan S. Dissecting racial bias in an algorithm

used to manage the health of populations. Science. 2019;366(6464):447-53.

42. Thornton T, Tang H, Hoffmann TJ, Ochs-Balcom HM, Caan BJ, Risch N. Estimating

kinship in admixed populations. Am J Hum Genet. 2012;91(1):122-38.

43. Maples BK, Gravel S, Kenny EE, Bustamante CD. RFMix: a discriminative modeling

approach for rapid and robust local-ancestry inference. Am J Hum Genet. 2013;93(2):278-88.

44. International HapMap C, Altshuler DM, Gibbs RA, Peltonen L, Altshuler DM, Gibbs RA,

et al. Integrating common and rare genetic variation in diverse human populations. Nature.

2010;467(7311):52-8.

45. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in

unrelated individuals. Genome Res. 2009;19(9):1655-64.

46. Delaneau O, Zagury JF, Marchini J. Improved whole-chromosome phasing for disease

and population genetic studies. Nat Methods. 2013;10(1):5-6.

47. Zou JY, Halperin E, Burchard E, Sankararaman S. Inferring parental genomic ancestries

using pooled semi-Markov processes. Bioinformatics. 2015;31(12):i190-6.

48. Patterson N, Price AL, Reich D. Population structure and eigenanalysis. PLoS Genet.

2006;2(12):e190.

49. PLUMMER M BN, COWLES K, VINES K. CODA: Convergence Diagnosis and Output

Analysis for MCMC. R News. 2012(6):7-11.

50. Sobota RS, Shriner D, Kodaman N, Goodloe R, Zheng W, Gao YT, et al. Addressing

population-specific multiple testing burdens in genetic association studies. Ann Hum Genet.

2015;79(2):136-47.

51.   Szpiech ZA, Blant A, Pemberton TJ. GARLIC: Genomic Autozygosity Regions

Likelihood-based Inference and Classification. Bioinformatics. 2017;33(13):2059-62.

52.   Fisher GG, Ryan LH. Overview of the Health and Retirement Study and Introduction to

the Special Issue. Work Aging Retire. 2018;4(1):1-9.