1   Classification: BIOLOGICAL SCIENCES: Biophysics and Computational

2   Biology

4   **The Nonequilibrium Mechanism of Noise Enhancer**

5   **synergizing with Activator in HIV Latency Reactivation**

7   Xiaolu Guo[1], Tao Tang[2], Minxuan Duan[3], Lei Zhang[4,5,*], Hao Ge[4,6,*]
8
[1]School of Mathematical Sciences, Peking University, Beijing, 100871, China
[2]School of Life Sciences, Peking University, Beijing, 100871, China
[3]Yuanpei College, Peking University, Beijing, 100871, China
[4]Beijing International Center for Mathematical Research, Peking University, Beijing,
100871, China
[5]Center for Quantitative Biology, Peking University, Beijing, 100871, China
[6]Biomedical Pioneering Innovation Center, Peking University, Beijing, 100871, China
16
17
*Correspondence: zhangl@math.pku.edu.cn (L.Z.), haoge@pku.edu.cn (H.G.)

Summary

Reactivating HIV latency and then simultaneously eliminating it by antiretroviral therapy

has become a leading strategy in curing HIV. Recently, single-cell screening experiments

have shown the drug synergy between two categories of biomolecules, Activators (AC)

and Noise Enhancers (NE): NE can amplify the reactivation of latent HIV induced by AC,

although NE itself cannot reactivate HIV latency. Based on an established LTR-two-state

effective model, we uncover two necessary conditions for this type of drug synergy: The

decreasing of the turning-on rate of LTR induced by NE is highly inhibited when

presented with AC; The timescale of LTR turning off without AC/NE presented should be

no slower than the timescale of Tat transactivation. Then we propose a detailed LTR-

four-state mechanistic model with both AC/NE regulation kinetics and Tat transactivation

circuit. We show that, in order to achieve drug synergy, the system of HIV gene state

transition must operate out of thermodynamic equilibrium, which is caused by energy

input. The direction of energy input determines whether the inhibition of NE upon the

reactivation rate of LTR-off states (unbinding of RNAP) can be successfully prevented in

the presence of AC. The drug synergy can also be significantly enhanced if the energy

input is appropriately distributed to more than one reaction. Our model reveals a generic

nonequilibrium mechanism underpinning the noise enhanced drug synergy, which may

apply to identify the same drug synergy on reactivating a diverse class of lentivirus

latency.

Keywords: HIV latency reactivation, drug synergy, nonequilibrium mechanism,

Tat transactivation, Markov jump process

Significance Statement

The "kick and kill" strategy has become a promising way to cure HIV by eliminating latent

HIV reservoirs, the main barrier to a clinical cure. Two categories of biomolecules,

Activators (AC) and Noise Enhancers (NE), have been found to have synergy on

reactivating latent HIV (kick strategy). We uncover the underlying non-equilibrium

mechanism of such drug synergy by developing mathematical models based on genetic

regulatory kinetics. We find that controlling the magnitude and direction of energy input

into genetic regulatory kinetics can prevent NE from reducing the turn-on rate of the

inactivated gene state in the presence of AC, which produces the synergy. This general

nonequilibrium mechanism can be useful for identifying other drug synergies on lentivirus

latency reactivation.

Introduction:

At the end of 2017, more than 36 million people were estimated to be infected with HIV (1). After HIV infects CD4+ cells, it can replicate or enter proviral latency (Figure 1A). Latent HIV reservoirs are the main obstacle to achieving a clinical cure (2). Reactivating latent HIV, quickly followed by antiretroviral therapy, has become a promising way to cure HIV-infected patients (3). Recently, synergistic combinations of noise enhancers and activators drugs were reported to beat other reactivation cocktails in reactivating HIV latency, yet induced less cytotoxicity (4). Two specific types of drugs used in (4): Activators (AC), a small biological molecule that increases the average expression level of HIV proteins, and Noise Enhancers (NE), a different type of molecule that increases the noise of HIV protein expression but does not affect the average expression level. While NE itself cannot reactivate latent HIV, it was shown to amplify AC-induced reactivation of HIV significantly (4). The synergy from adding NE with AC to latent HIV is shown in Figure 1B. However, the biomolecular mechanisms underlying the synergy between AC and NE have not been fully resolved.

The main ingredients of the HIV regulatory loop are the promoter long terminal repeats (LTR) and the Tat transactivation on LTR. LTR is the promoter of the HIV genome and has a larger expression noise than promoters of human genes (5, 6). Nucleosomes associated with the LTR often block the full transcription by RNA polymerase (RNAP), resulting in a

low basal expression rate. The rarely produced Tat protein complexes with

CDK9 and CyclinT1 to form the *positive transcriptional elongation factor b*

(pTEFb). pTEFb can bind to the transactivation response element (TAR) on

the initially transcribed part of the HIV mRNA and remodel downstream

nucleosomes. This remodeling assists the elongation of the mRNA, thus

forming positive feedback (7-9). In addition, a bimodal gene expression

("phenotypic bifurcation" (10)) pattern was found in the offspring of defective-

HIV infected cells with initially intermediate expression (10). However, it was

reported that the cooperativity coefficient (Hill coefficient) of Tat is only one

(11), which means that the mean-field deterministic dynamics of HIV gene

expression is monostable. This is distinct from the genetic toggle switch of the

lambda phage regulatory loop with stronger feedback and bi-stable

deterministic dynamics (12), or an oscillatory network with negative feedback

and a limit cycle (13). The deterministic dynamics of the HIV regulatory

network is insufficient to explain the observed bimodality, and a stochastic

description may be required. By combining the bimodality and noisiness of

HIV promoter gene expression, Weinberger et al. found that the bimodality

arises from a very slow rate of switching on LTR expression (10), resulting in

a much noisier dynamics than those observed for normal human promoters.

On the other hand, functions of certain Activators and Noise Enhancers are

partially known. For example, as Activators, TNF and prostratin can activate

the transcriptional factor NF-κB, and therefore antagonize HIV latency(14-16).

Some of these Noise Enhancers, such as ethinyl estradiol, can influence HIV

expression through another transcriptional factor SP1 or the structural state of

chromatin (17, 18). The molecular mechanisms of most noise enhancers are

still unclear, indicating the complicated regulation mechanism of HIV

dynamics. However, no matter how complex it is, what these activators or

noise enhancers regulated are just the rates between different gene/promoter

states described by TF binding or structural difference. Therefore, the

functions of Activators and Noise Enhancers can be analyzed by certain

minimal but general models.

Thermodynamic energy dissipation plays a crucial role in bioactivities and

bioreactions. General model considering the binding of multiple transcription

factors (TF) under thermodynamic equilibrium in prokaryotic cells and the

function that different pair TF interactions can achieve in gene expression of

cells was already studied extensively (19, 20). However, in studies of

eukaryotic transcriptional dynamics, a non-equilibrium mechanism is found

necessary(21), and many far-from-equilibrium models have been proposed

(22-25). In addition to biomolecule synthesis and cell motility, the regulatory

function of a living cell, such as adaptation and precise control of oscillations

was also found highly dissipative (26, 27). Hence, we are very curious about

whether certain energy input is necessary for this type of drug synergy.

In addition, in a self-positive-feedback gene regulatory network, the timescale

of DNA state transition, mRNA transcription, mRNA decay, protein translation,

and protein decay, will influence the cell fate landscape and phenotype

transitions (28-30). Post-integration HIV gene expression is one example

system of the TF regulatory mechanism of gene expression with self-positive-

feedback. Hence, we are also interested in how the timescales of gene-state

transition and protein dynamics influence the drug synergy.

In this essay, we first investigate an established LTR-two-state model and find

two necessary conditions for the synergy between AC and NE on reactivating

latently infected HIV: (i) AC inhibits NE's function of reducing the transition

rate k_{on} for the gene-state reactivation; (ii) the rate k_{off} for the promoter to turn

off is not slower than the rate for Tat transactivation when the promoter is

already turned on. We then propose an LTR-four-state model and prove that

the noise enhanced drug synergy is indeed a non-equilibrium phenomenon of

the regulation of HIV promoter LTR. We show that controlling the magnitude

and direction of the system energy input can deter NE from reducing the

turned-on rate of inactivated gene state in the presence of AC, which induces

the synergy between AC and NE on the LTR transcription. The drug synergy

can be significantly enhanced when we distribute the total energy input into

two specific different reactions. Moreover, we show that only when the

timescale of DNA unbinding RNAP (k_{off} or k_{unbindp}) is comparable to or faster

than the timescale of Tat transactivation at gene activated state, the synergy

of AC and NE on the transcription on genetic level can pass to the translation

on protein level in the Tat self-positive-feedback HIV system.

Materials and methods

To investigate the synergy between AC and NE, we first employ a well-

established LTR-two-state model with Tat transactivation from the previous

study (Figure 2A) (4, 31). In this model, LTR (the promoter of HIV) has two

states: on state and off state. LTR can transit to the transcriptionally inactive

state LTR-off at rate k_{off} and transit to the transcriptionally active state LTR-on

at rate k_{on}. Tat can transactivate LTR-on state promoting a higher

transcription rate k_{transact} than the basal transcription rate k_{m} due to Tat's

enhancing LTR transcriptional elongation (See Supplementary Section 1.1 for

more details). The values of parameters (k_{on}, k_{off}, k_{transact}) are the same as

those in (31), which is quantified by single-cell analysis (32-34).

As assumed in (4), adding only AC to the system increases k_{on}, while adding

only NE reduces k_{on} and k_{off} simultaneously with their ratio fixed. We use

parameter f_{inh} to quantify the degree of AC's inhibition upon the NE-induced

reduction of k_{on}, which is defined as

156
$$f_{\text{inh}} = \frac{\ln(k_{\text{on,AC,NE}}) - \ln(k_{\text{on,AC}})}{\ln(k_{\text{off,AC}}) - \ln(k_{\text{off,AC,NE}})} + 1$$

(See Supplementary Section 1.2-1.3 for more details). $f_{\text{inh}} = 0$ represents AC

does not inhibit NE's function of reducing k_{on} (Figure S1C, left panel), and

$f_{\text{inh}} > 0$ means that AC inhibits NE's function of reducing k_{on}. Particularly,

$f_{\text{inh}} = 1$ means that NE's function of reducing k_{on} is fully inhibited by AC

(Figure S1C, right panel).

We propose a more detailed LTR-four-state model with both transcript

activation and Tat transactivation (Figure 3A, Figure S2A). As the HIV

promoter, LTR controls HIV protein expression after the HIV genome

integrates into the DNA of human CD4+ cells (35, 36). In this model, there are

four different promoter states: LTR state is the free state; LTR* is the

activated state but without RNAP binding; LTR-P and LTR*-P are the

corresponding RNAP-bounded states. AC is assumed to promote LTR

transiting to the activated state LTR*, e.g., LTR bound with NF-κB, and LTR*

recruits RNA polymerase much easier than LTR itself, e.g., the NF-κB bound

to LTR acting as a Transcription Factor (TF) to recruit RNA polymerase

(RNAP) to LTR (37). Dar, Hosmane, Arkin, Siliciano and Weinberger (4) show

that the screened Noise Enhancers have no effect on post transcription.

Some NEs can increase transcription factors in cells, such as SP1 (17, 18).

Similar to (4), we assume that NE can slow down the switching rates between

LTR and LTR-P.

We use the Markov jumping process to model the transition among LTR

states with AC and/or NE added (Figure 3A). The four states can mutually

transit: in the absence of AC, RNAP binds to LTR at a relatively slow rate

$k_{\mathrm{bindp}}[P]$ and unbinds at a relatively fast rate k_{unbindp}; LTR transit to LTR*

state at an extremely slow rate k_{act} without AC, but at a much higher rate

$k_{\mathrm{act}}\gamma$ $(\gamma \gg 1)$ with AC presented; RNAP is recruited to LTR* at a higher rate

$\omega k_{\mathrm{bindp}}$, where ω is the attraction coefficient $(\omega > 1)$; when NE (or Noise

Suppressor, NS) is added, LTR will bind and unbind RNAP at slower (faster

for NS) rates ($k_{\mathrm{bindp}}e^{-\alpha}$, $k_{\mathrm{unbindp}}e^{-\alpha}$) with noise attenuation factor α. Note

that the setup of the model showed in Figure 3A is under the assumption of

the detailed balance. We will analyze both the models with or without this

assumption in the following sections.

We also combine the LTR-four-state model with the transcription/translation

module without Tat transactivation to characterize the procedure of screening

AC and NE (Figure 3B). In screening experiments of AC and NE, the LTR-

GFP vectors without Tat transactivation were used (4). For the LTR-GFP

vectors, with RNAP bond to LTR (i.e., LTR-P state or LTR*-P state), the

downstream DNA of LTR can be transcribed into mRNA at rate k_{m},

independent of the GFP copy number. Then the mRNA can be translated into

protein at rate k_{GFP}. Also, mRNA and GFP will degrade at rate d_{m} and d_{GFP},

respectively (Figure 3B).

In the experiments of testing the synergy between AC and NE on HIV latency

reactivation, the full-length HIV with Tat transactivation was used to infect host

cells instead of the LTR-GFP vector (4). For the full-length HIV vector, the Tat

protein expressed by the HIV genome enhances transcriptional elongation by

mediating RNAP hyperphosphorylation and thus transactivates the LTR

promoter (38-40).

Based on the biological evidence and previous models(31, 41), we construct

the Tat transactivation module (Figure 3C; see Supplementary Section 2.4 for

details). Only at LTR-on states (LTR-P and LTR*-P), HIV genome can be

transcribed into mRNA, and Tat transactivates LTR-on states in a Hill

equation form:

209
$$k_{\mathrm{m}} = k_{\mathrm{mbasal}} + k_{\mathrm{trs1}} \frac{\dfrac{Tat}{k_{\mathrm{trs2}}}}{1 + \dfrac{Tat}{k_{\mathrm{trs2}}}}$$

In addition, previous experiments also certified that Tat transactivation could

stabilize HIV activation state autonomous from the host cellular state (31, 34).

Based on this, we assume that Tat reduces the dissociating rate of RNAP

from LTR, $k_{unbindp}$, thereby stabilizing the HIV activation state (Figure 3C

inset) (See Supplementary Section 2.4 and 2.12 for details and parameter

value)

216
$$k_{\mathrm{unbindp}} = \frac{K_{\mathrm{threshold}}^3 + \delta Tat^3}{K_{\mathrm{threshold}}^3 + Tat^3} k_{\mathrm{unbindp0}}$$

Then in the following sections, we use the LTR-four-state model coupled with

Tat transactivation module to investigate how LTR expression and HIV fates

are regulated by NE and AC under the Tat transactivation. We also simulate

HIV expression trajectories in host T-cell from HIV latent initial state and

calculate the percentage of cells expressing HIV protein reached a threshold

(#Tat = 75) after a certain time (100 hours) as the reactivation ratio (Figure

S2C; see Supplementary Section 2.7 for more details).

Results

The LTR-two-state model with significant large k_{off} can exhibit drug

synergy between NE and AC

We simulate the LTR-two-state model showed in Figure 2A using Stochastic

Simulation Algorithm (SSA) (42), calculating the reactivation of HIV latently

infected T cells after adding NE and/or AC, with the above assumptions on

AC's and NE's functions.

Through simulation, we find that AC's inhibiting NE's function on k_{on} is

necessary for the synergy between AC and NE (Figure 2C-D). When AC has

no inhibiting effect on NE, corresponding to $f_{inh} = 0$, there is no synergy

between AC and NE in the whole reasonable parameter space (Figure 2C,

Figure 2B black arrow). Only when AC inhibits NE's function of reducing k_{on},

there can be synergy between AC and NE on reactivating latent HIV (Figure

2D, Figure 2B red arrow).

We also find out that another necessary condition for NE having synergy with

AC is k_{off} being greater than $10^{-2} \, \text{hour}^{-1}$, which is larger than the

parameters used in previous studies (31). However, other important results

shown in previous literature, the bimodal distribution of phenotype bifurcation

(10) and HIV latency establishment operating autonomously from the host

cellular state (31), are not sensitive to the increasing of k_{off} (Figure S1G-I).

When k_{off} is no greater than 10^{-2}, there is no synergy between AC and NE on

reactivating latent HIV (Figure 2E; Figure S1E). However, if k_{off} is greater

than 10^{-2}, then NE can have synergy with AC on reactivating latent HIV

(Figure 2F-G; Figure S1F). Furthermore, the synergy between AC and NE will

increase with the inhibiting effect quantity $f_{\text{inh}} > 0$ only when k_{off} is sufficiently

large, such as $k_{\text{off}} = 0.8 > 10^{-2}$ (red line in Figure 2G). There will be no

synergy between AC and NE with the inhibiting effect quantity $f_{\text{inh}} > 0$ if $k_{\text{off}} <$

10^{-2}(red line in Figure 2E). Actually, in a latent HIV system, k_{off} is more likely

to be larger than that previously used in literature. Direct sequencing of HIV

integration sites showed that latent HIV frequently integrated into

heterochromatin (38). The transcription is continuously turned off in

heterochromatin, a tightly packed form of DNA (43). Thus, in latent HIV, LTR

transcription-permissive state should be unstable, corresponding to larger k_{off},

as one characteristic of the HIV integrating into heterochromatin.

In summary, we find two necessary conditions of drug synergy from our

simulation results: (i) AC inhibits NE's function of reducing k_{on}; (ii) sufficiently

large k_{off} (rate of LTR turning off or unbinding RNAP). However, this LTR-two-

state model is oversimplified. We still do not know much about what causes

AC inhibiting NE's function of reducing k_{on} (the assumption made in (4)) and

why NE synergizes with AC only if k_{off} is large.

No drug synergy can be produced under the detailed balance condition

in the LTR-four-state model coupling with Tat transactivation

Under the detailed balance condition (Figure 3A), our LTR-four-state model

with the transcription/translation module without Tat transactivation (Figure

3B) illustrates that AC increases LTR mean expression level and that NE

increases LTR expression noise (Figure S4C-D), which is consistent with the

drug screening experimental results (Figure 1B) (4).

However, under the detailed balance condition, neither synergy between AC

and NE nor depression of Noise Suppressor (NS) on AC reactivating latent

HIV can be possible (Figure 4). This contradicts the experimental data

showing that NEs enhance AC's inducing latent HIV reactivation and that NSs

reduce AC's reactivating latent HIV (Figure 1B) (4). It is because under the

detailed balance condition, the probability P_{on} of RNA polymerase binding to

LTR (LTR-P and LTR*-P) (See Supplementary Section 2.1 and 2.5 for details)

only depends on the equilibrium constants of each reaction, and NE or NS

tunes the forward and backward rates simultaneously but keeps the

equilibrium constant unvaried. This conclusion is not dependent on concrete

models. It is a general physical and mathematical result. Hence it is not

possible to build another even more complicated Detailed-Balanced model to

overcome this obstacle.

The above theory can be illustrated by our LTR-four-state model (Figure 3A,

C, and Figure S2A, C) (See Supplementary Section 2.1,2.3 and 2.4 for

details). Under the detailed balance condition, P_{on} keeps the same when both

NE (or NS) and AC are added to the system ($\gamma \gg 1$, $\alpha = 1$ (or $\alpha = -1$))

compared to when only AC is added ($\gamma \gg 1$, $\alpha = 0$) (Figure 4A; see

Supplementary Section 2.5 for details). Thus, the Detailed-Balanced LTR-

four-state model predicts no synergy between NE and AC and predicts that

NS does not suppress the AC's function of increasing P_{on} (Figure 4A),

contradicting the experimentally observed synergy between AC and NE

(Figure 1B) (4).

We couple the Detailed-Balanced LTR-four-state model with Tat

transactivation, and there is still no synergy between Noise Enhancer and

Activator illustrated by the reactivation ratio of latent HIV (Figure 4B; see

Supplementary Section 2.7 for details). Note that the reactivation ratio of HIV

is calculated dynamically during a finite time starting from the latent state,

which is different from the steady-state probability P_{on}. However, they are

closely related to each other, since they both indicate the degree of

reactivation for latent HIV.

In addition, we calculate the mean duration time (MDT) of both the LTR-off

states (LTR and LTR*) and the LTR-on states (LTR-P and LTR*-P) (See

Supplementary Section 2.8 for details). The reciprocals of the MDTs

calculated from the LTR-four-state model can be regarded as the effective

transition rates in the reduced LTR-two-state model with only the LTR-off and

LTR-on state. Then we find out that AC can shorten the MDT at LTR-off

states (Figure S4E, Figure S4F), and that NE can lengthen the MDT at both

LTR-on states and LTR-off states with their ratio fixed (Figure 4C-D, S5G-H).

These results are consistent with the assumptions of the LTR-two-state model

in the previous section. However, in this Detailed-Balanced model, the

effective inhibiting effect quantity f_{inh} defined through the effective transition

rates, always vanishes (Figure 4E; see Supplementary Section 2.9 for the

exact definition of f_{inh}). This confirms the LTR-two-state model predictions,

that no synergy between AC and NE on reactivating latent HIV should be

observed when $f_{inh} = 0$.

Hence, for Noise Enhancers synergize with ACs, the regulation of HIV gene

expression must be a Non-Detailed-Balanced process with energy dissipation.

The direction of the cycle flux caused by energy input in the Non-

Detailed-Balanced LTR-four-state model determines the synergy

Inside a living cell, continuous energy consumption is necessary for executing

different vital functions. We already know that systems with the drug synergy

must be energy dissipative, but how energy input, i.e., breaking the detailed

balance, influences the drug synergy remains poorly understood.

We mainly investigate how the cycle flux direction and the energy input

distribution, as the features of the non-equilibrium system, effect the synergy.

Breaking the detailed balance is equivalent to having non-vanishing cycle

fluxes. In our LTR-four-state model, the cycle fluxes can be either counter-

clockwise or clockwise. Energy input can be distributed on one or more

reactions. Here, we first consider the case when energy input is only through

one single reaction (Figure 5A; see Supplementary Section 2.2 and Table S2-

1 for details). In the real biological system, the energy input can be realized

through ATP hydrolysis or other reversible covalent modification (44).

We prove that the Non-Detailed-Balanced LTR-four-state model can produce

the drug synergy between NE and AC on P_{on}, if and only if the direction of

cycle flux is clockwise. Mathematical analysis (See Supplementary Section

2.10 for details) and numerical simulations illustrate the same phenomenon.

The model with counter-clockwise cycle flux predicts no synergy between NE

and AC on P_{on} or HIV latency reactivation, and no reduction of P_{on} or latent

HIV reactivation is observed when NS is added with AC (Figure 5B). On the

other hand, with clockwise cycle flux, the model predicts 100% of NE can

synergize with AC on P_{on}, and about 75% of NE synergize with AC on latent

HIV reactivation (Figure 5C). 50% (4 out of 8) of the ways to break the

detailed balance through a single reaction to produce a clockwise cyclic

probability flux predicts significant synergy ($> 5\%$) on P_{on} between NE and AC

(Figure 5C up panel), and that NS reduces P_{on} with AC added. 25% (2 out of

8, increasing transition rate from LTR*-P to LTR-P or reducing transition rate

from LTR-P to LTR*-P) of the ways predicts significant synergy ($> 5\%$) on

reactivation of latent HIV between NE and AC (Figure 5C down panel), and

that NS reduces AC-induced HIV latency reactivation. In the experiments,

around 80% of the NSs reduce AC-induced HIV latency reactivation, while

more than 64% of NEs have synergistic effects with AC on reactivating latent

HIV (4). Hence the results of our model are consistent with the experimental

fact that the majority of NEs amplify AC reactivating latent HIV, while the

majority of NSs suppresses reactivation of latent HIV with AC added. Thus,

the Non-Detailed-Balanced LTR-four-state model reveals a general

mechanism of the synergy between NEs and ACs on the reactivation of latent

HIV, instead of a particular mechanism of a specific NE.

We also show that in the above cases producing significant drug synergy

between AC and NE, the clockwise cyclic probability flux always promotes

LTR turn on mainly through the LTR*-to-LTR*-P pathway strengthened by AC

and turn off through the LTR-P-to-LTR pathway weakened directly by NE

(Figure 5D-E). It explains why NE can further amply the HIV latency

reactivation induced by AC, as long as the energy input provides clockwise

cyclic probability flux.

In addition, for the equilibrium system, the distribution of the dwell time at

LTR-off states is predicted to be monotonically decreasing and convex (45)

(Figure S10, solid black lines). The monotonicity or convexity can be

maintained for the non-equilibrium system with a low magnitude of energy

input (small disturbance from the equilibrium system) (Figure S10, dashed red

line). However, as the magnitude of energy dissipation increases, the

nonmonotonicity or concavity of the distribution of dwell time could appear,

similar to the phenomenon of phase transition (Figure S10D, H, dotted red

line, and solid red line).

375 **The LTR-four-state model with distributed energy input may achieve**

376 **much stronger synergy than that with energy input only on a single**

377 **reaction**

378 One possible strategy, through which strong synergy can be achieved, is to

379 drive the LTR promoter to turn on mostly through the LTR-to-LTR*-to-LTR*-P

380 pathway, whose rate can be significantly increased by AC, and to turn off

381 mostly through the LTR*P-to-LTR-P-to-LTR pathway (Figure 5D-E), whose

382 rate can be distinctly decreased by NE. This way, the promoter is more likely

383 to transit to the state LTR-P rather than the state LTR* once it is at the state

384 LTR*-P. Here, we build an EITST (Energy Input on the Two Specific

385 Transition rates) LTR-four-state model, in which part of the energy input

386 reduces the transition rate from LTR*-P to LTR* (β_2) and the other part

387 increases the transition rate from LTR*-P to LTR-P (β_1) (Figure 6A; see

388 Supplementary Section 2.2 for details), with the total energy fixed ($\beta_1 + \beta_2 =$

389 β).

390 We find that there is an optimal energy input distribution ($\beta_1 = 1.8$, $\beta_2 = 8.2$)

391 for the system to perform the strongest synergy between AC and NE on P_{on}

392 (Figure 6B). The drug synergy on HIV latency reactivation dependent on the

393 energy input distribution is qualitatively quite similar. (Figure 6C). Overall, the

394 certain distributed energy input with $0 < \beta_1 < 10$ may achieve stronger

395 synergy than that on only a single reaction ($\beta_1 = 0$ or $\beta_1 = 10$). Without loss of

generality, we set $\beta_1 = 5$, $\beta_2 = 5$ for the EITST LTR-four-state model, and all

the following simulation results are based on this value.

In such a Non-Detailed-Balanced model, simulation results of adding AC or

NE alone with GFP present are consistent with the drug screening

experimental data, that AC increases LTR mean expression level and NE

increases LTR expression noise (Figure 6D-E, Figure 1B).

The synergy between Noise Enhancer and Activator on both P_{on} (Figure 6G)

and HIV latency reactivation (Figure 6H) are observed and much stronger

than the scenario where energy input is only on a single reaction. Noise

Enhancer can increase the HIV latency reactivation from approximately 7%

when Activator is already added to 13% when both are added (Figure 6H).

These numbers are quite similar to the best cases observed in experiments

with prostratin as AC (experimental data from Figure 3A in (4)). And Noise

suppressor reduces the AC-induced HIV latency reactivation from 7% to less

than 1% (Figure 6H). This synergy between AC and NE on both P_{on} and HIV

latency reactivation is found to be positively correlated with the magnitude $|\beta|$

of the energy input, but will reach maximum (for P_{on}) and saturation (for HIV

latency reactivation) when $|\beta|$ is getting sufficiently large ($\beta > 5$) (Figure S6A-

B). In addition, the synergy is found to be positively correlated with the noise

of NEs (Figure S6C), this is also consistent with the experimental data (Figure

3B in (4)).

We also calculate the mean duration time (MDT) of the LTR-off and LTR-on

states. Different from the Detailed-Balanced situation, NEs can lengthen the

MDT at the LTR-on states more significantly than lengthen the MDT at the

LTR-off states (Figure 6F, Figure S7C-F). Further, the effective inhibiting

parameter $f_{\text{inh}} \approx 1 > 0$ (See Supplementary Section 2.9 for the exact

definition of this effective parameter) means that AC does inhibit NE's function

of reducing the transition rate from the LTR-off states to the LTR-on states

(Figure 6F). These simulation results verified the conclusion we made in the

LTR-two-state model, that NE can synergize with AC and NE on reactivating

latent HIV only when $f_{\text{inh}} > 0$. Now we know that AC's inhibiting NE's function

of reducing effective k_{on} is achieved by the clockwise cycle flux driven by the

energy input.

However, same as in the LTR-two-state model, Noise Enhancer can amplify

Activator's reactivating latent HIV only if k_{unbindp} (equivalent to k_{off} in the

LTR-two-state model) is greater than 10^{-2} (Figure S8A). To explain this

necessary condition, we analyze the timescale of Tat transactivation

dynamics and the timescale of LTR transitions. We find that it takes about

$\tau_0 \approx 20$ hours on average for Tat transactivation and then LTR can maintain

the activated state for a long time (Figure S8B-C). Therefore, if k_{unbindp} is very

small compared to the time scale of $1/\tau_0$, the duration time of LTR-on state

without NE presented will be long enough for Tat transactivation with a large

possibility, so further reducing k_{unbindp} by NE will have little effect (Figure

S8B). Only when k_{unbindp} is not small compared to $1/\tau_0$, the duration time of

LTR-on state without NE presented is typically not long enough for Tat

transactivation. In this case, noise enhancer lengthening the duration time at

the LTR-on state will provide Tat more time to reactivate latent HIV (Figure

S8B), resulting in the drug synergy with Activator.

Finally, to verify the model applicability, we use the same EITST model to

explain other important previous experimental observations (Figure S3A-C)

including Tat-transactivation-controlled HIV latency establishment operating

autonomously from the host cellular state (31) and the bimodal distribution of

phenotype bifurcation (10) (See supplementary Section 2.13 for parameter

values). Also, in our EITST model, nonmonotonicity and concavity of the

distribution of the dwell time are also observed with large magnitude of energy

dissipation (Figure S10F).

Discussion

Retroviruses are mild at beginning infection. Most replicable retroviruses do

not cause cytopathic effects, and infected cells hardly produce a defensive

response to such viral replication. Animals rarely experience acute symptoms

due to infection with retroviruses, but viremia occurs; the host immune

responses reduce the production of the virus. The virus cannot be eradicated

even though can be suppressed by the host's immune response. Therefore,

low levels of viremia usually accompany the life of the host (46). As a typical

example of the retrovirus, the long-lived latent HIV-1 are the main obstacles to

a clinical cure (2). Recently, noise enhanced synergistic combinations of

drugs (Activators and Noise Enhancers) were reported more effective than

other reactivation cocktails on reactivating HIV latency (4). We here propose

an LTR-four-state model with Tat transactivation (the cooperativity is only

one) to reveal the mechanism of this synergy, produced by the combination of

Activator and Noise Enhancer. By analysis and simulation of the model, we

find that the drug synergy on HIV latency reactivation depends the distribution

of energy input and the direction of cycle flux of the system. Such a proposed

non-equilibrium mechanism should be useful for identifying new drug synergy

or regulating the existing drug synergy in HIV and a diverse class of

retroviruses infection therapies, most of which have latency states preventing

from a complete cure.

Design principle for specific biological functions have been extensively

explored, such as reliable cell decisions (47), adaptation (48), robust and

tunable biological oscillation (49), and dual function of adaptation and noise

attenuation (50). Some of these functions, such as precise biochemical

oscillations and accurate adaptation, were found crucially depending on the

energy dissipation (26, 27). Here, we show that the drug synergy between NE

and AC on reactivating HIV latency also essentially depends on the directional

dissipated chemical energy on the HIV LTR-state transition. This non-

equilibrium property could also be used as a potential target for lentivirus

latency reversal synergetic therapeutic interventions. The optimization

principle of energy input distribution for the highest drug synergy might also

apply to the network designing.

Our LTR-four-state model is a minimal model in which the effects of AC and

NE are distinguished. This LTR-four-state model can be expanded into a

more detailed LTR-six-state model where the Tat positive feedback is

modeled through Tat binding to LTR and forming two new states, LTR-P-Tat

and LTR*-P-Tat as modeled in the previous study (31). In the LTR-six-state

model, the same synergy can be predicted (Figure S9, Supplementary

Section 2.4). Hence, our uncovered non-equilibrium mechanism of drug

synergy is not dependent on the specific Tat positive feedback mechanism.

The synergy that we studied here does not require cooperative binding

between the two categories of drugs (AC and NE or NS), which is essentially

different from the drug synergy based on classical equilibrium binding

mechanism. It might be the reason why the tested drug synergies in *vitro*

often fail in *vivo.* The nonequilibrium model proposed here provides a new

perspective to understand the drug synergy mechanism.

Acknowledgments

We thank Profs. Yuhai Tu and Tom Chou for helpful discussions. The work

was supported by National Natural Science Foundation of China

(11861130351, 11622101, 11622102,11971037).

Reference

1. UNAIDS (2018) UNAIDS Data 2018.
2. Richman DD, *et al.* (2009) The Challenge of Finding a Cure for HIV Infection. *Science* 323(5919):1304-1307.
3. Deeks SG (2012) HIV Shock and kill. *Nature* 487(7408):439-440.
4. Dar RD, Hosmane NN, Arkin MR, Siliciano RF, & Weinberger LS (2014) Screening for noise in gene expression identifies drug synergies. *Science* 344(6190):1392-1396.
5. Pai A & Weinberger LS (2017) Fate-Regulating Circuits in Viruses: From Discovery to New Therapy Targets. *Annu Rev Virol* 4(1):469-490.
6. Dar RD, *et al.* (2012) Transcriptional burst frequency and burst size are equally modulated across the human genome. *Proc Natl Acad Sci U S A* 109(43):17454-17459.
7. Kao SY, Calman AF, Luciw PA, & Peterlin BM (1987) Anti-termination of transcription within the long terminal repeat of HIV-1 by tat gene product. *Nature* 330(6147):489-493.
8. Kaehlcke K, *et al.* (2003) Acetylation of Tat defines a cyclinT1-independent step in HIV transactivation. *Mol Cell* 12(1):167-176.
9. Karn J (2000) Tat, a novel regulator of HIV transcription and latency. in *HIV Sequence Compendium 2000*.
10. Weinberger LS, Burnett JC, Toettcher JE, Arkin AP, & Schaffer DV (2005) Stochastic gene expression in a lentiviral positive-feedback loop: HIV-1 Tat fluctuations drive phenotypic diversity. *Cell* 122(2):169-182.
11. Weinberger LS & Shenk T (2007) An HIV feedback resistor: auto-regulatory circuit deactivator and noise buffer. *PLoS Biol* 5(1):e9.
12. Gardner TS, Cantor CR, & Collins JJ (2000) Construction of a genetic toggle switch in Escherichia coli. *Nature* 403(6767):339-342.
13. Elowitz MB & Leibler S (2000) A synthetic oscillatory network of transcriptional regulators. *Nature* 403(6767):335-338.
14. Williams SA, *et al.* (2004) Prostratin antagonizes HIV latency by activating NF-kappa B. *J Biol Chem* 279(40):42008-42017.
15. Duh EJ, Maury WJ, Folks TM, Fauci AS, & Rabson AB (1989) Tumor Necrosis Factor a Activates Human Immunodeficiency Virus Type-1 through Induction of Nuclear Factor Binding to the Nf-Kappa-B Sites in the Long Terminal Repeat. *P Natl Acad Sci USA* 86(15):5974-5978.
16. Osborn L, Kunkel S, & Nabel GJ (1989) Tumor Necrosis Factor-Alpha and Interleukin-1 Stimulate the Human Immunodeficiency Virus Enhancer by Activation of the Nuclear Factor Kappa-B. *P Natl Acad Sci USA* 86(7):2336-2340.
17. Asin SN, Heimberg AM, Eszterhas SK, Rollenhagen C, & Howell AL (2008) Estradiol and progesterone regulate HIV type 1 replication in peripheral blood cells. *Aids Res Hum Retrov* 24(5):701-716.

18.   Katagiri D, Hayashi H, Victoriano AFB, Okamoto T, & Onozaki K (2006)
Estrogen stimulates transcription of human immunodeficiency virus type 1
(HIV-1). *Int Immunopharmacol* 6(2):170-181.
19.   Buchler NE, Gerland U, & Hwa T (2003) On schemes of combinatorial
transcription logic. *P Natl Acad Sci USA* 100(9):5136-5141.
20.   Bintu L*, et al.* (2005) Transcriptional regulation by the numbers: models. *Curr*
Opin Genet Dev 15(2):116-124.
21.   Coulon A, Chow CC, Singer RH, & Larson DR (2013) Eukaryotic transcriptional
dynamics: from single molecules to cell populations. *Nat Rev Genet*
14(8):572-584.
22.   Scholes C, DePace AH, & Sanchez A (2017) Combinatorial Gene Regulation
through Kinetic Control of the Transcription Cycle. *Cell Syst* 4(1):97-+.
23.   Estrada J, Wong F, DePace A, & Gunawardena J (2016) Information
Integration and Energy Expenditure in Gene Regulation. *Cell* 166(1):234-244.
24.   Ahsendorf T, Wong F, Eils R, & Gunawardena J (2014) A framework for
modelling gene regulation which accommodates non-equilibrium
mechanisms. *Bmc Biol* 12.
25.   Li CX, Cesbron F, Oehler M, Brunner M, & Hofer T (2018) Frequency
Modulation of Transcriptional Bursting Enables Sensitive and Rapid Gene
Regulation. *Cell Syst* 6(4):409-+.
26.   Cao Y, Wang H, Ouyang Q, & Tu Y (2015) The free energy cost of accurate
biochemical oscillations. *Nat Phys* 11(9):772-778.
27.   Lan G, Sartori P, Neumann S, Sourjik V, & Tu YH (2012) The energy-speed-
accuracy trade-off in sensory adaptation. *Nat Phys* 8(5):422-428.
28.   Potoyan DA & Wolynes PG (2017) Stochastic dynamics of genetic
broadcasting networks. *Phys Rev E* 96(5).
29.   Jia C, Qian H, Chen M, & Zhang MQ (2018) Relaxation rates of gene
expression kinetics reveal the feedback signs of autoregulatory gene
networks. *J Chem Phys* 148(9).
30.   Ge H, Qian H, & Xie XS (2015) Stochastic Phenotype Transition of a Single Cell
in an Intermediate Region of Gene State Switching. *Phys Rev Lett* 114(7).
31.   Razooky BS, Pai A, Aull K, Rouzine IM, & Weinberger LS (2015) A Hardwired
HIV Latency Program. *Cell* 160(5):990-1001.
32.   Dar RD*, et al.* (2012) Transcriptional burst frequency and burst size are
equally modulated across the human genome. *P Natl Acad Sci USA*
109(43):17454-17459.
33.   Singh A, Razooky B, Cox CD, Simpson ML, & Weinberger LS (2010)
Transcriptional Bursting from the HIV-1 Promoter Is a Significant Source of
Stochastic Noise in HIV-1 Gene Expression. *Biophys J* 98(8):L32-L34.
34.   Weinberger LS, Dar RD, & Simpson ML (2008) Transient-mediated fate
determination in a transcriptional circuit of HIV. *Nat Genet* 40(4):466-470.
35.   Klaver B & Berkhout B (1994) Comparison of 5' and 3' Long Terminal Repeat
Promoter Function in Human-Immunodeficiency-Virus. *Journal of Virology*
68(6):3830-3840.

36. Cullen BR, Lomedico PT, & Ju G (1984) Transcriptional Interference in Avian
Retroviruses - Implications for the Promoter Insertion Model of
Leukemogenesis. *Nature* 307(5948):241-245.
37. Barboric M, Nissen RM, Kanazawa S, Jabrane-Ferrat N, & Peterlin BM (2001)
NF-kappa B binds P-TEFb to stimulate transcriptional elongation by RNA
polymerase II. *Mol Cell* 8(2):327-337.
38. Jordan A, Bisgrove D, & Verdin E (2003) HIV reproducibly establishes a latent
infection after acute infection of T cells in vitro. *Embo J* 22(8):1868-1877.
39. Lassen K, Han YF, Zhou Y, Siliciano J, & Siliciano RF (2004) The multifactorial
nature of HIV-1 latency. *Trends Mol Med* 10(11):525-531.
40. Han YF, Wind-Rotolo M, Yang HC, Siliciano JD, & Siliciano RF (2007)
Experimental approaches to the study of HIV-1 latency. *Nat Rev Microbiol*
5(2):95-106.
41. Weinberger LS, Burnett JC, Toettcher JE, Arkin AP, & Schaffer DV (2005)
Stochastic gene expression in a lentiviral positive-feedback loop: HIV-1 Tat
fluctuations drive phenotypic diversity. *Cell* 122(2):169-182.
42. Gillespie DT (1977) Exact Stochastic Simulation of Coupled Chemical-
Reactions. *J Phys Chem-Us* 81(25):2340-2361.
43. Volpe TA*, et al.* (2002) Regulation of heterochromatic silencing and histone
H3 lysine-9 methylation by RNAi. *Science* 297(5588):1833-1837.
44. Berg JM, Tymoczko JL, & Stryer L (2002) Covalent Modification Is a Means of
Regulating Enzyme Activity. *Biochemistry*, (W H Freeman, New York), 5th Ed.
45. Tu Y (2008) The nonequilibrium mechanism for ultrasensitivity in a biological
switch: Sensing by Maxwell's demons. *P Natl Acad Sci USA* 105(33):11737-
11741.
46. Geng Y (2012) *Retrovirus-induced Diseases*. *Molecular Biology of Lentivirus*
and Foamy virus, eds Xia L & Zhang L (Science Press, Beijing, China), p 23.
47. Brandman O, Ferrett JE, Li R, & Meyer T (2005) Interlinked fast and slow
positive feedback loops drive reliable cell decisions. *Science* 310(5747):496-
498.
48. Ma WZ, Trusina A, El-Samad H, Lim WA, & Tang C (2009) Defining Network
Topologies that Can Achieve Biochemical Adaptation. *Cell* 138(4):760-773.
49. Tsai TYC*, et al.* (2008) Robust, tunable biological oscillations from interlinked
positive and negative feedback loops. *Science* 321(5885):126-129.
50. Qiao LX, Zhao W, Tang C, Nie Q, & Zhang L (2019) Network Topologies That
Can Achieve Dual Function of Adaptation and Noise Attenuation. *Cell Syst*
9(3):271-+.
51. Dar RD, Hosmane NN, Arkin MR, Siliciano RF, & Weinberger LS (2014)
Screening for noise in gene expression identifies drug synergies. *Science*
344(6190):1392-1396.

 FIGURE LEGENDS

Figure 1. **HIV-infected cell fates and biological function of biomolecules**

reactivating latent HIV.

(A) Schematic of different fates of cells infected by HIV: HIV active replication,

HIV proviral latency, and HIV latency reactivation (adapted from Figure 1A of

(51)).

(B) Diagram of experimentally screening Activator (AC) and Noise Enhancer

(NE) (up) and experimental synergy test on the reactivation of latent HIV after

adding AC or/and NE (down). In experiments, ACs and NEs were selected by

to detecting the mean and noise of LTR expression using cells infected by the

LTR-GFP vector. The synergy between ACs and NEs on HIV reactivation was

tested using cells infected by full-length HIV with Tat transactivation.

Untreated (grey bar) represents a control group. Adding Activator (green bar)

increases LTR expression compared to the control group. Adding Noise

Enhancer (magenta bar) increases LTR noise compared to the control group.

Adding AC and NE simultaneously (red bar) has a strong synergy on HIV

reactivation (increases Reactivation of latent HIV infected Cells). Adding AC

and Noise Suppressors (NS, blue bar) has a depressing effect on HIV latency

reactivation compared to adding AC only. (51)

Figure 2. **Two necessary conditions for drug synergy**

(A) Modified from Figure 3A of (31). The LTR-two-state model with Tat

feedback to explain the effects of NE and AC molecules on HIV. LTR has two

states, on and off, which convert to each other at the rate of k_{on} and k_{off}; the

LTR-on state transcribes HIV mRNA at a rate k_{m}, mRNA degrades at d_{m} rate

or translates to protein at a rate of k_{Tat}, and Tat degrades at rate d_{Tat}. NE

decreases k_{on} and k_{off} with their ratio fixed; AC molecules will increase k_{on};

AC inhibits NE's function on k_{on} when added together as assumed in (51). Tat

transactivate LTR through enhancing the transcriptional rate $k_{trasact}$.

(B) The heat map of Reactivation across different values of LTR turning off

rate k_{off} and LTR turning on rate k_{on}. The green arrow corresponds to adding

AC. Red arrow corresponds to adding NE only decreasing k_{off} without

changing k_{on}, i.e., AC inhibits NE's function on k_{on}. Black arrow corresponds

to adding NE decreasing k_{off} and k_{on} with their ratio fixed, i.e., AC does not

inhibit NE decreasing k_{on}. (See Table S1-1 for parameter values)

(C-D) The heat map of Synergy on Reactivation without AC's inhibition on NE

($f_{inh} = 0$) and with AC's complete inhibition on NE ($f_{inh} = 1$), respectively,

across different values of k_{off} and k_{on} with AC added. The synergy is the

reactivation with NE and AC added subtracting the reactivation with only AC

added. (See Table S1-1 for parameter values)

(E-G) The plots of synergy on reactivation changing with different f_{inh} values

for small k_{off}, intermediate k_{off}, and large k_{off}, respectively. Red lines indicate

adding NE and AC simultaneously; green lines indicate adding AC only; blue

lines indicate adding NS and AC simultaneously. (See Table S1-1 and Table

S1-2 for parameter values)

Figure 3. **The LTR-four-state model under the detailed balance condition**

and transcription/translation modules without/with Tat transactivation

(A) Schematic of the LTR-four-state model with the Tat-feedback circuit. The

LTR promoter is modeled to have four states: a transcriptional silence state

(LTR state) extremely slowly binding RNAP polymerase or activation

transcription factors such as NF-κB, an activated state (LTR*) such as LTR

with NF-κB bond, a transcription-permissive state (LTR-P), a transcription-

permissive state with NF-κB bond (LTR*-P). k_{act} is the rate for LTR binding

NF-κB. k_{unact} is the rate for LTR unbinding NF-κB. γ models AC increases the

rate for LTR binding NF-κB. $\gamma = 1$ for untreated HIV infected cells, and $\gamma \gg 1$

for adding AC. ω is the attraction coefficient between NF-κB and RNAP, $\omega =$

10 ($\omega > 1$ means NF-κB attracts RNAP); k_{bindp} is the rate at which RNAP

binds to LTR; k_{unbindp} is the rate at which RNAP unbinds from LTR; α is the

noise attenuation factor ($\alpha > 0$ corresponds to Noise Enhancer, and $\alpha < 0$

corresponds to Noise Suppressor). The parameter setting here is under the

detailed balance condition. The case of breaking the detailed balance is in

Figure 5 and Table S2-1.

(B) Schematic of the LTR-four-state model coupled with the transcription and

translation module without feedback. LTR-on states (red, including LTR-P

state and LTR*-P state) transcribes mRNA at rate k_{m}; mRNA decays at rate

d_{m} or can be translated at rate k_{p} into GFP; GFP decays at rate d_{p}.

(C) Schematic of the LTR-four-state model coupled with the transcription and

translation module with the Tat-transactivation circuit. LTR-on states (red,

including LTR-P state and LTR*-P state) transcribes mRNA at rate k_{m}; mRNA

decays at rate d_{m} or can be translated at rate k_{Tat} into Tat; Tat decays at rate

d_{Tat}; Tat has positive feedback on k_{m}; Tat stabilizes the state of LTR-on state

through negative feedback on k_{unbindp}.

Figure 4. No synergy is predicted under the detailed balance condition

(A-B) Probability of LTR-on states (LTR-P state and LTR*-P state), P_{on}, and

Reactivation ratio of Latent HIV, respectively, in the Detailed-Balanced LTR-

four-state model. Y-axis is the P_{on} and the reactivation ratio value,

respectively, and X-axis is the categories of different combinations of ACs and

NEs/NSs. Untreated (grey bars) corresponds to $\gamma = 1$, $\alpha = 0$; adding only AC

(green bars) corresponds to $\gamma \gg 1$, $\alpha = 0$; adding only NE (magenta bars)

corresponds to $\gamma = 1$, $\alpha = 1$; adding NE and AC (red bars) corresponds to

$\gamma \gg 1$, $\alpha = 1$; adding NS and AC (blue bars) corresponds to $\gamma \gg 1$, $\alpha = -1$.

(B) We use Stochastic Simulation Algorithm (SSA) to calculate the

reactivation ratio of the LTR-four-state model coupled with Tat feedback.

Reactivation Ratio is the ratio of reactivated HIV trajectory number by time

100h to all trajectory number, starting from latency state (LTR=1, all other

species=0, simulated 5000 cells).

(C-D) Mean Duration Time at LTR-off states and LTR-on states, respectively,

under the detailed balance condition.

(E) f_{inh}, the degree of AC's inhibition upon the reduction of k_{on} induced by NE

under the detailed balance condition. We first calculate the reciprocal of the

Mean Duration Time as the transition rate between LTR-on and LTR-off

states, λ_{on} and λ_{off}, respectively. Then we calculate using the formula: $f_{\text{inh}} =$

$\frac{\ln\left(k_{\text{on}}([N],[A])\right)-\ln\left(k_{\text{on}}(0,[A])\right)}{\ln\left(k_{\text{off}}(0,[A])\right)-\ln\left(k_{\text{off}}([N],[A])\right)} + 1$. (C-E) The red lines correspond to adding AC

and NE ($\alpha > 0$); the cyan lines correspond to adding AC only ($\alpha = 0$); the blue

lines correspond to adding AC and NS ($\alpha < 0$). (See Table S2-3 and Table

S2-4 for parameter values.)

Figure 5. The LTR-four-state model with energy input on a single transition

produces synergy between AC and NE if and only if the system has

clockwise cyclic probability flux

(A) Schematic of the Non-Detailed-Balanced LTR-four-state model. Energy

input can influence any single transition rate of the Detailed-Balanced LTR-

four-state model with the corresponding rate multiplying by e^{β} or $e^{-\beta}$. Such an

energy input will cause clockwise (c.w.) cyclic probability flux or counter-

clockwise (c.c.w.) cyclic probability flux.

(B-C) Probability of LTR-on states, P_{on} (up panels), and Reactivation ratio of

Latent HIV (down panels) calculated from the Non-Detailed-Balanced models

with energy input on different single transitions causing a c.c.w. cyclic

probability flux (B) or c.w. cyclic probability flux (C). R stands for LTR state; R*

stands for LTR* state; P stands for LTR-P state; R*P stands for LTR*-P state.

Each group of x-axis represents the Non-Detailed-Balanced model with the

corresponding transition rate multiplying by e^β ($\beta > 0$ for orange groups, $\beta <$

0 for blue groups). (See Table S2-1 for model details. See Table S2-3 and

Table S2-4 for parameter values.)

(D-E) The distributions of fluxes for LTR turning on (left panels) and turning off

(right panels).

(See Table S2-1 and Supplementary Section 2.5-7 for model details. See

Table S2-3 and Table S2-4 for parameter values.)

Figure 6. The LTR-four-state model with distributed energy input exhibit

strong synergy between AC and NE

(A) Schematic of the EITST model with distributed energy input ($\beta = \beta_1 +$

$\beta_2 = 10$). The first part of the energy β_1 increases the LTR*-P-to-LTR-P

transition rate by multiplying e^{β_1}, the other part of the energy β_2 reduces the

LTR*-P-to-LTR* transition rate by multiplying $e^{-\beta_2}$.

(B-C) The synergy on P_{on} (B) and HIV latency reactivation (C) vary with

energy distribution ($\beta_1 + \beta_2 = \beta = 10$). (C) Each point is average of 250

simulation experimental data points with 10000 cells simulated for each

simulation experiment. Error bars show the Standard Deviation.

(D-J) $\beta_1 = \beta_2 = 5$.

(D-E) The mean and noise of GFP expression calculated from the EITST

model without positive feedback.

(F) f_{inh}, the degree of AC's inhibition upon the reduction of k_{on} induced by NE

of the EITST model.

(G-H) Probability of LTR-on states (LTR-P state and LTR*-P state), P_{on}, and

Reactivation ratio of Latent HIV, calculated from the EITST model. (See Table

S2-1 for model details. See Table S2-3, S2-4, and S2-5 for parameter values.)

A   **Detailed Balance**

$k_{act}\,\gamma$

LTR $\longrightarrow$ LTR*

k_{unact}

$k_{unbindp}\,e^{-\alpha}$ $k_{bindp}\,e^{-\alpha}$ $k_{unbindp}$ ωk_{bindp}

$\omega k_{act}\,\gamma$

LTR-P $\longrightarrow$ LTR*-P

k_{unact}

$\gamma \gg 1$: AC; $\alpha > 0$: NE; $\alpha < 0$: NS

B

LTR OFF

LTR ON $\longrightarrow$ mRNA $\xrightarrow{k_p}$ GFP
$\qquad\quad k_m \qquad\;\; \downarrow d_m \qquad \downarrow d_p$
$\qquad\qquad\qquad$ Decay $\qquad$ Decay

C

$k_{unbindp}$ vs Tat

LTR OFF

LTR ON $\longrightarrow$ mRNA $\xrightarrow{k_{Tat}}$ Tat
$\quad k_m \qquad\;\; \downarrow d_m \qquad\qquad \downarrow d_{Tat}$
$\qquad\qquad$ Decay $\qquad\qquad$ Decay

A **Energy Input :** e^β

$\beta \neq 0$: **Non-Detailed-Balance**

- untreated
- AC
- NE
- AC+NE
- AC+NS

B Energy on a single transtion with c.c.w. probability flux

C Energy on a single transtion with c.w. probability flux

D Energy on the transtion from LTR*-P to LTR-P with $\beta=10$

E Energy on the transtion from LTR*-P to LTR* with $\beta=-10$