Integrating Proteomes for Lung Tissue and Lavage Reveals Pathways that Link Responses

in Allergen-Challenged Mice

Thomas H Mahood[1,2,3,7,*], Christopher D Pascoe[1,2,3,7,*], Tobias K Karakach[4], Aruni Jha[1,2,3,7],

Sujata Basu[1,2,3,7], Peyman Ezzati[5], Victor Spicer[5], Neeloffer Mookherjee[2,3,5,6,7], Andrew J

Halayko[1,2,3,7]

[1]*Department of Physiology & Pathophysiology, University of Manitoba, Winnipeg, MB, Canada*

[2]*DEVOTION Network, Winnipeg, Manitoba, Canada*

[3]*Biology of Breathing Group, Children's Hospital Research Institute of Manitoba, Winnipeg,*

MB, Canada

[4]*Bioinformatics Core Laboratory, Children's Hospital Research Institute of Manitoba,*

Winnipeg, MB, Canada

[5]*Manitoba Centre for Proteomics and Systems Biology, Department of Internal Medicine,*

University of Manitoba, Winnipeg, MB, Canada

[6]*Department of Immunology, University of Manitoba, Winnipeg, MB, Canada*

[7] *Canadian Respiratory Research Network, http://respiratoryresearchnetwork.ca/, Ottawa,*

Ontario, Canada

* These authors contributed equally to this work

Corresponding Author: Dr. Andrew J Halayko, Department of Physiology and Pathophysiology,

University of Manitoba, 606-John Buhler Research Centre, 715 McDermott Ave, Winnipeg, MB

Canada R3E 3P4 E-mail- andrew.halayko@umanitoba.ca

Running Title: Lung Tissue and Lavage Proteome

Abstract

To capture interplay between biological pathways we analyzed the proteome from matched lung tissue and bronchoalveolar lavage fluid (BALF) of individual allergen-naïve and house dust mite (HDM)-challenged BALB/c mice, a model of allergic asthma. Unbiased label-free LC-MS/MS analysis quantified 2,675 proteins from tissue and BALF of allergen-naïve and HDM-exposed mice. In comparing the four datasets we found significantly greater diversity in proteins between lung tissue and BALF than in the changes induced by HDM challenge. The biological pathways enriched after allergen exposure were compartment-dependent. Lung tissue featured innate immune responses and oxidative stress, while BALF most strongly revealed changes in metabolism. We combined lung tissue and BALF proteomes, which principally highlighted oxidation reduction (redox) pathways, a finding influenced chiefly by the lung tissue dataset. Integrating lung and BALF proteomes also uncovered new proteins and biological pathways that may mediate lung tissue and BALF interactions after allergen challenge, for example, B Cell Receptor signaling. We demonstrate that enhanced insight is fostered when different biological compartments from the lung are investigated in parallel. Integration of proteomes from lung tissue and BALF compartments reveals new information about protein networks in the response to environmental challenge and interaction between intracellular and extracellular process.

Keywords: bronchoalveolar lavage fluid / asthma / bioinformatics / allergen / proteomics

Introduction

The biological underpinnings of chronic inflammation in asthma involve a complex molecular interplay between structural cells, recruited inflammatory cells, and the extracellular mediators that they release. Understanding this interplay is important to understand pathobiology and support pre-clinical research to develop new therapies. Mice exposed to allergen are a mainstay for asthma research, and a common approach is to challenge animals with inhaled house dust mite (HDM). HDM is an aeroallergen that is clinically relevant, and is a complex stimulus with multiple immunogens and stressors, including fungal spores, bacterial endotoxins, lipid-binding proteins and proteases [1,2]. Repeated HDM challenge induces pathophysiological symptoms that are hallmarks of human asthma [3]. However, the scope and complexity of the interplay between lung cells, autocrine and paracrine pathways, and the integrated signaling networks associated with the pathobiology that determines the efficacy of pre-clinical therapeutics, has not been refined.

To understand complex disease mechanisms, a number of omics technologies have been employed, including proteomics. To the best of our knowledge, most studies examine the proteome of individual biological compartments, including airway spaces - collected as bronchoalveolar lavage fluid (BALF) or sputum - and lung tissue from asthmatic patients and murine models of the disease [4-6]. These studies have been important for endotyping patients and animal models, identifying biomarkers of disease, and providing direction for new therapeutic strategies. However, analysis of these biological compartments in isolation can only partially identify the molecular networks and biomarkers that are critical for disease expression in a complex system like the lung. Nonetheless, understanding the molecular systems that determine the interaction between secreted proteins in the lung and the pathways that are regulated in the resident cells of the lung has not been fully established in conventional proteomics studies.

To address this knowledge gap, we used unbiased proteomics and unsupervised

exploratory data analysis to establish a molecular signature in matched lung tissue and BALF

from individual mice. We then combined these to create integrated proteomes that allow unique

signals from each sample type to be identified, and the nature of the interactions between

biological compartments to be deciphered in describing the response to HDM challenge. We

found that the proteomes of lung tissue and of BALF are distinct and that inhaled HDM

challenge induces unique alterations in each compartment. Moreover, by integrating proteomes

we uncovered novel signaling networks that were not evident from individual proteome dataset

of lung tissue or BALF.

Results

Lung function & differential cell count analysis

HDM challenge of adult female BALB/c mice resulted in pathophysiological features

that are consistent with human asthma. This included significant increases in airway resistance,

tissue elastance and tissue resistance in methacholine challenge (Supplemental Figure S1A).

Differential counting of BALF immune cells confirmed that HDM challenge triggered

significant accumulation of eosinophils and neutrophils that is also consistent with prior studies [3]

and mimics human asthma (Supplemental Figure S1B).

Lung tissue and BALF proteome disparities

In total, our analysis of lung tissue and BALF samples in all mice, yielded 2,675 uniquely

identifiable proteins (Protein IDs). On a per mouse basis, we obtained $1,594 \pm 154$ protein IDs

(mean $\pm$ SD) from lung tissue and 641 ± 207 protein IDs from BALF samples (Supplemental

Figure S2A). To confirm that secreted proteins were enriched in the BALF samples, we

characterized protein IDs using the Uniprot keyword, "secreted", which annotated >75% of the

proteins in the BALF. In contrast, ~only 25% of the protein ID's were characterized as

"secreted" in the tissue lysate. To confirm that membrane-associated proteins are enriched in

lung tissue samples, we classified proteins using the Uniprot annotated keyword

"transmembrane". This identified 69% of proteins in lung tissue, and only 23% of BALF

proteins.

To investigate if the proteome of lung tissue and BALF from allergen-naïve and HDM-

exposed mice are distinct, we performed Principal Component Analysis (PCA) on the top 500

most variable proteins across all samples. This cut-off was chosen arbitrarily to capture the high

intensity (and therefore highly variable) signals from our dataset. Our comparison matrix of all

samples from allergen-naïve and HDM-challenged mice using the top two principal components

(PC) accounted for ~80% of variability between samples (Figure 1A). Lung tissue and BALF

samples are separated along the 1st PC (x-axis), which accounts for approximately 59% of the

data variance. Lung tissue and BALF samples from naïve and HDM-challenged mice were also

distinct, with the 2nd PC accounting for about 20% of data variance, with HDM effects seemingly

much greater in BALF than in lung tissue (Figure 1A, expanded panel).

We next examined the distribution of proteins across all samples. A clustering heatmap

(k-means) shows that BALF and tissue clustered separately, as did naïve and HDM-challenged

for tissue and BALF (Figure 1B). We also observed that the differences in lung tissue and BALF

samples in allergen-naïve and HDM-challenged mice were associated with the unique protein

IDs in individual datasets. Approximately 72.7% of all protein IDs were unique to either lung

tissue or BALF samples.

We next examined the distribution of Uniprot IDs in lung tissue and BALF, and the

effects of HDM-challenge (Figure 1C). Venn diagrams show that HDM exposure increased the

absolute number of proteins in lung tissue and BALF by 114% and 173%, respectively. The

fraction of protein IDs common to BALF and lung tissue was 18% in allergen-naïve mice, and

this increased to 28% after HDM exposure. HDM exposure reduced the fraction of unique lung

tissue Uniprot IDs from 75% to 63%. The proportion of proteins that were unique to BALF was

8-10%; this was not changed by HDM-challenge.

We next examined patterns of protein abundance of the specific subset that were common

to BALF and lung tissue from allergen-naïve and HDM-challenged mice (Figure 1D). The

dominating feature of the heat map is that it discerns large clusters of proteins that are relatively

more prominent in BALF or lung tissue, independent of the effects of HDM challenge. Thus, for

these proteins that are common to lung and BALF, k-means clustering is consistent with PC

analysis of all proteins (Figure 1A) which showed that data variability was chiefly the result of

differences between lung tissue and BALF. Though effects of HDM challenge within lung tissue

or BALF are evident in this common protein subset, our analysis highlights the disparities

between lung and BALF that likely affect the biological manifestation of HDM challenge.

Independent validation of proteins selected from the proteomic analysis

To validate our proteomic data, immunoblotting was performed on the same samples for

three proteins (Figure 2). ARG1 (Arginase-1), CLCA1 (Calcium-activated chloride channel

regulator 1) and FDPS (Farnesyl pyrophosphate synthase) were selected based upon their

enrichment by HDM exposure and the availability of reliable immunoblotting grade antibodies.

Immunoblotting confirmed our proteomic data for ARG1 and CLCA1, as they were undetectable

in allergen-naïve samples from lung tissue and BALF, but prominent bands were evident for

samples from allergen-challenged mice (Figures 2A and 2B). Immunoblotting analysis

confirmed presence of FDPS in all samples, with a significant enrichment of 270% (p.adj $\leq$ 0.05)

in the lung tissue after HDM-challenge (Figures 2A and 2B). This is consistent with proteomic

data for FDPS, which was detected in only one BALF and lung tissue sample from an allergen-

naïve mouse, while it was present and increased in all samples from HDM-challenged mice.

Combining Tissue-BALF proteomes to uncover whole lung responses to HDM challenge

To enable investigation of how HDM-challenge affects protein interactions within the

whole lung, we combined all protein IDs from BALF and tissue proteomes (n=2,675). To

validate and reduce variability due to missing values, we filtered the dataset to include only those

proteins that were detected in all biological replicates for each sample type and experimental

condition. This strict criterion reduced the data set to 1,237 proteins for network and pathways

analyses. The combined dataset of all tissue and BALF proteins included allergen-naïve (n=558)

and HDM-challenged (n=1,064) proteins both containing 385 proteins in common. Using this

full dataset we performed PCA to examine the effects of HDM-challenge, and this showed that

only 8.1% of the variation in protein abundance correlated with differences between naïve and

HDM-treated mice (Y-axis, Figure 3A), whereas 83.8% of variation could be attributed to

differences in protein abundance between lung tissue and BALF (X-axis, Figure 3A).

Venn diagram mapping of protein IDs from the combined total lung tissue and BALF

dataset showed that 69% of proteins were unique to naïve (173) or HDM-exposed samples (679),

and 31% (385) were common to the treatment groups (Figure 3B). Differential expression

analysis of the 385 common proteins using LIMMA showed that two proteins were depleted, and

eight proteins were enriched after HDM challenge (Table 1 and Figure 3B). To identify which

biological processes were represented by these changes, we performed Gene Ontology analysis

on a dataset that included the group of eight proteins *enriched by HDM* and the group of 679

proteins that were unique to samples from HDM-challenged mice (see Figure 3B). We identified

36 significantly enriched pathways, with the top 5 biological processes being: "oxidation-

reduction process", "translational initiation", "protein transport", "spliceosomal snRNP

assembly" and "GDP-mannose metabolic process" (Figure 3C). To complement this analysis, we

performed a protein interaction analysis using NetworkAnalyst. The top three nodes, based on

the number of connections with other proteins, included: SMARCA4 (Transcription activator

BRG1), PIK3R1 (Phosphatidylinositol 3-kinase regulatory subunit alpha) and NFKB1 (Nuclear

factor NF-kappa-B p105 subunit) (Figure 3D). This set of networks was most significantly

associated with "Interleukin-3, 5 and GM-CSF signaling" (p.adj = 1.40×10^{-8}), forming a

backbone for the whole lung response to HDM-challenge.

Biological significance of proteome changes induced specifically in lung tissue by allergen

challenge

To assess lung tissue-specific effects of HDM-challenge we analyzed 1,787 proteins that

were unique to all lung tissue samples (Figure 4A). A large fraction of these proteins (n=1,304,

75%) was common to allergen-naïve and HDM-challenged mice, and differential abundance

analysis uncovered 43 that were significantly changed after HDM exposure (28 depleted, 17

enriched) (Figure 4B). The proteins most significantly *depleted* by HDM challenge included

SERPINA3K (Serine protease inhibitor A3K), CHAD (Chondroadherin) and ADSS

(Adenylosuccinate synthetase isozyme 2). We combined the 28 significantly depleted proteins

with 138 proteins that were unique to allergen-naïve lung tissue (therefore absent after HDM

challenge) (Figure 4A). From this set of proteins that are *uniquely depleted* in the lung by HDM

challenge, Gene Ontology analysis revealed that "negative regulation of endopeptidase activity"

was the most significantly enriched biological process (Figure 4C).

Of the 17 proteins that were *enriched* (>log$_2$ Fold change) by HDM challenge and that

were common to allergen-naïve lung tissue samples, CHIL3 (Chitinase-like protein 3), EPX

(Eosinophil peroxidase) and LGALS3 (Galectin-3) increased most significantly. We combined

the 17 significantly enriched proteins with 345 proteins (Figure 4A) that were unique to HDM-

challenged lung tissue samples. Using this set of 362 proteins *uniquely enriched* in lung tissue

from HDM-challenged mice, Gene Ontology analysis revealed that "oxidation-reduction"

regulation was the most significantly enriched process (Figure 4D).

To extend understanding of the differential abundance of the 362 proteins that we

identified as being uniquely altered in allergen exposed lung tissue, we used NetworkAnalyst to

develop interactome maps that reveal interaction nodes and signaling networks for biological

responses (Figure 5A). The top three interaction nodes for lung tissue were HDAC1 (Histone

deacetylase 1), MAPK1 (Mitogen-activated protein kinase 1) and PLCG2 (1-

phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-2), and these nodes generated

an interactome involving multiple pathways. Using the Reactome database linked within

NetworkAnalyst, the top 5 pathways were: antigen mediated B-cell receptor activation and

secondary messenger generation; hemostasis; signaling by interleukins; proteins associated with

G0 and Early G1; and, activation of circadian expression through BMAL1:CLOCK/NPAS2.

Biological significance of proteome changes induced specifically in BALF by allergen

challenge

To assess effects of HDM challenge on the BALF-specific proteome, we analyzed 706

proteins that were unique to all BALF samples (Figure 4E). Approximately half (n=372; 53%) of

these protein IDs were common to BALF from allergen-naïve and HDM-challenged mice.

Further analysis for differential abundance in these common proteins did not reveal any proteins

that were significantly enriched or depleted by allergen challenge (Figure 4F). Therefore, we

performed Reactome pathway analysis (linked within NetworkAnalyst) using the 23 proteins that

had only been identified in allergen-naïve BALF samples (Figure 4E). This discriminated 10

biological processes that are uniquely *depleted* by HDM challenge in BALF, with "rRNA

pseudouridine synthesis" being the most significantly affected (Figure 4G). To determine

pathways in BALF that were *enriched* by HDM challenge, we also performed Gene Otology

analysis in InnateDB using in the 311 proteins that were unique to BALF from HDM-challenged

mice (Figure 4E). This revealed a top 10 biological processes *uniquely enriched* by challenge to

inhaled allergen in BALF , and this included "proteolysis involved in cellular protein catabolic

process(es)" as the most significantly enriched (Figure 4H).

To extend understanding of the 311 proteins that are uniquely altered in the allergen

exposed BALF, we developed interactome maps to identify primary interaction nodes and

signaling networks (Figure 5B). Using the Reactome database in NetworkAnalyst, the top three

protein-protein interaction (PPI) nodes were SKP1A (S-phase kinase-associated protein 1A),

HIST1H4B (H4 clustered histone 2), and PSMB6 (Proteasome subunit beta type-6). This PPI

network included diverse pathways, with the top 5 pathways being the SCF(Skp2)-mediated

degradation of p27/p21, Cyclin E associated events during G1/S transition, Cyclin A:Cdk2-

associated events at S phase entry, activation of NF-κB in B cells, and SCF-beta-TrCP mediated

degradation of Emi1.

Predicting interactions between lung tissue and BALF

To broaden the scope of our study, we interrogated protein interactions across biological

compartments using the lung tissue and BALF samples collected from each mouse. To uncover

points of integration of biological activities in lung tissue and the surrounding extracellular space

in response to allergen challenge, we integrated the 362 proteins that were uniquely enriched by

HDM challenge in lung tissue (see Figures 4A, 4B and 5A) with the 311 proteins that were

uniquely enriched in BALF (see Figures 4E, 4F and 5B). Importantly, as this integration directly

compares the interactions between the lung tissue and BALF, we excluded the 49 proteins that

are shared between the two datasets, resulting in 311 and 262 proteins found exclusively in the

allergen-exposed lung tissue and BALF datasets respectively. We next identified protein

interaction nodes in the combined dataset using NetworkAnalyst, enhancing the rigour of our

analyses by filtering the results to only include protein nodes with five or more first-order

interactions. From the 37 proteins that met this criterion (12 from lung tissue, 25 from BALF),

we selected the top 5 proteins from both the lung tissue and BALF datasets that become enriched

in the combined dataset. We found that the integration of BALF proteins with lung tissue-

specific proteins resulted in a significant enrichment, $142.1 \pm 19.1\%$ (range of 122 - 185%), of

first order interactions for nodes that emerged from lung-derived seed proteins (the proteins that

start the growth of a network) (Table 2). In contrast, combining lung tissue proteins with those

from BALF did not affect the number of interactions for seed protein nodes that originated from

BALF (Table 3).

To uncover points of crosstalk between lung tissue and BALF proteins we built a

combined interaction network using NetworkAnalyst (Figure 6). Once we removed the proteins

that are common between the two datasets, we calculated the number of interactions each protein

has from within lung tissue (n=311) and BALF (n=262) datasets. We then extracted the top five

proteins from the lung tissue that increased as a result of combining with the BALF dataset

(Table 2). These included HDAC1, CBX1 (Chromobox protein homolog 1), EVL (Ena/VASP-

like protein), ABI1 (Abl interactor 1) and COPS2 (COP9 signalosome complex subunit 2). Since

combining these datasets did not enrich the number of interactions from each protein node

derived from BALF, for interactome mapping we included the top five seed proteins that we

maintained interactions when combined with lung tissue proteins. These included, SPTBN1

(Spectrin beta chain, non-erythrocytic 1), RPS3 (40S ribosomal protein S3), HNRNPAB

(Heterogeneous nuclear ribonucleoprotein A/B), PSMA5 (Proteasome subunit alpha type-5), and

PSMA2 (Proteasome subunit alpha type-2) (Table 3).

267   A unique aspect of our analytical approach is that it uncovered six predicted integration

proteins between key pathway nodes in lung tissue and BALF. As shown in Figure 6, these

included NANOG (Homeobox protein NANOG), POU5F1 (POU domain, class 5, transcription

factor 1), TFCP2I1(Transcription factor CP2-like protein 1), SMAD4 (Mothers against

decapentaplegic homolog 4), TAL1 (T-cell acute lymphocytic leukemia protein 1 homolog) and

YWHAZ (14-3-3 protein zeta/delta), and each protein had two to four interactions spanning lung

tissue and BALF. Among lung-derived seed proteins, HDAC1 interacted with seven predicted

integrating proteins, and thus associated with all of the BALF-origin protein nodes. Neither

COPS2 nor ABI1 signalling nodes from lung tissue were associated with any of the predicted

integration proteins from the BALF dataset, although indirect linkage of ABI1 via EVL and

CBX1 is apparent. Among BALF-derived seed nodes, SPTBN1 and HNRNPAB exhibited six

and two links, respectively, to predicted integration proteins, thus creating interactions to

HDAC1 and EVL in lung tissue. As well, PSMA2 and PSMA5 from BALF projected three

interactions to lung tissue nodes (HDAC1 and EVL). Similarly, SPTBN1 in BALF predicted

direct linkages to SMAD4, TAL1 and YWHAZ along with indirect linkages between EVL,

HDAC1 and CBX1 (Figure 6A). Together, these results reveal that combining proteins in a

manner that allows identification of the specific biological compartment from which they were

derived enables the critical discrimination of pathways that link the allergic response in lung

tissue with that in the surrounding extracellular space.

The impact of combining data for proteins that are uniquely enriched by HDM challenge

in matched BALF and lung tissue samples is also evident in the divergence of predicted

biological processes that emerges, compared to those based on lung tissue or BALF datasets

alone. Figures 6B and 6C list the top 10-biological processes predicted using the Reactome

database in NetworkAnalyst for lung tissue and BALF, using the top five seed proteins derived

from each, but as influenced by the combined dataset. Prominent lung tissue processes are

associated with immune system regulation (innate and adaptive), including T-cell signalling and

hemostasis, as well as tissue remodelling (ROBO receptors), and cell division. BALF seed

proteins are linked to processes associated with cell stasis (apoptosis, protein turnover, and DNA

damage control pathways, including ornithine decarboxylase), as well as antigen presentation.

Figure 6D represents the predicted biological processes using all ten seed protein nodes from

BALF and lung tissue combined. The list of biological processes reflects a strong dominance by

lung tissue-linked processes, in particular immune responses, and the regulation of mitosis and

cell division. Of note, two areas that are not evident in lung tissue and BALF alone, but are

known components of allergic asthma pathobiology were revealed by the integration of datasets,

including B-cell receptor signaling and inflammation, featuring NF-κB activation, pathways

induced by Wnt, and degradation of β-catenin.

Discussion

Murine models of allergic airways inflammation employ inhaled aeroallergen such as

HDM to support preclinical and discovery research. Asthma pathobiology involves interaction

between recruited inflammatory cells and lung structural cells. In this study we used label-free

proteomics and multivariate bioinformatics to describe and compare the molecular interactome

in lung tissue and BALF from HDM challenged mice. We used matched samples from individual

mice for this process, and in so doing have been able to discriminate responses in tissue and

extracellular spaces. Using *in silico* re-integration of datasets, we predict the interactions

between the lung tissue and the BALF. We demonstrate that the proteomes of lung tissue and

BALF are remarkably different, and this is further enhanced by disparate effects resulting after

HDM challenge, highlighting inherent biological diversity in these samples. Examining protein-

protein interactions between lung tissue and BALF reveals points for crosstalk between lung

tissue and extracellular proteins after HDM exposure. Our study reveals the scope and limits of

biological insight that can be obtained from lung tissue or BALF sample proteins alone, and

offers potential to interrogate network interactions between the lung tissue and extracellular

airway space during allergen challenge.

Confirming a B cell signaling immune signature in BALF and lung tissue

In the current study, we confirmed that HDM challenge led to predominant eosinophil

and neutrophil infiltration of BALF, and we found B cell receptor signaling to be a key

component of, and crosstalk between, lung tissue and BALF in allergen exposed mice (Figures

5A and 6D). The immune signature evident in lung tissue and BALF after HDM challenge is

predominantly of a Th2-polarizied phenotype, in part dependent on endotoxin levels in HDM [7, 8].

Murine B cells are antigen presenting cells for HDM antigens, and B cell depletion profoundly

reduces allergic responses in the lung due to their interactions with resident lung cells and

inflammatory cell that infiltrate the lung and airways [9]. Importantly, memory B-cell populations

remain in the lung tissue and epithelial cell layers well after HDM sensitization, forming a

localized population that can quickly become activated by further allergen challenge [10]. This

pattern of a localized memory B cell population contributes to inflammatory responses that

associate with airway remodeling in asthmatic patients [11]. As a biological validation of our

proteomic investigation, our chief findings reveal a B-cell signaling signature after repeated

allergen challenge. Moreover, our network analysis of integrated lung tissue and BALF

(collected post-lavage) proteomes reveals multiple levels for crosstalk between the lung tissue

and the extracellular microenvironment. In addition to the immunoblot validation of proteins in

lung tissue and BALF that we provide, these observations provide evidence of the reliability of

our proteome measurement and analysis protocols, and further support an important role for B

cells in coordination of biological responses in lung tissue and BALF.

Protein nodes that link lung tissue and BALF

In lung tissue samples we detected markedly increased HDAC1 (Histone deacetylase 1)

after allergen exposure, and it was a seed protein for a major lung tissue interaction node.

Moreover, via all six proteins we predicted to integrate lung tissue and BALF responses (Figure

6), HDAC1 appears to interact with all of the BALF protein nodes we identified in HDM-

challenge mice. This suggests a fundamental role for HDAC 1 in allergic pathophysiology,

which is consistent with its primary role, in conjunction with histone acetyltransferase (HAT), to

modulate chromatin accessibility for transcription. Our finding is consistent with evidence from

ovalbumin sensitized and challenged mice, in which HDAC1 abundance and activity is increased

[12]. Skewed HDAC and HAT activity has been documented in both asthmatic and COPD patients

[13, 14], and HDAC1 activity negatively correlates with disease severity and IL-8 expression [14].

This link with inflammation is due, in part, to transcriptional control of NF-κB signaling and its

transcriptional activation of pro-inflammatory genes and antimicrobial peptides [15–17]. Together

our study reveals the complexity of HDAC regulation and suggests that modification of

transcriptional events leads to physiological and inflammatory changes that span beyond a single

biological compartment.

358   CBX1 abundance was increased in lung tissue after allergen exposure, but exhibited

limited interactions with BALF proteins. CBX1 (Chromobox protein homolog 1), a

heterochromatin binding protein associated with transcriptional regulation of cell proliferation,

also interacts with nuclear lamin (LMNB1) and EMSY (BRCA2-interacting transcriptional

repressor) [18, 19]. EMSY, is an interferon-α/β gene regulator [20], and a genome-wide association

study recently identified EMSY (*C11orf30*-LRRC32) to be significantly associated with

moderate-to-severe asthma [21]. Similar to CBX1, ABI1 increased significantly in lung tissue after

HDM challenge, but network analysis did not reveal interactions with protein nodes in BALF.

ABI1 associates with N-WASP to regulate filamentous actin dynamics, as well as acetylcholine

mediated contraction of airway smooth muscle [22]. ABI1 is regulated by the tyrosine kinase ABL

[22], which is increased in lungs of asthmatic donors and allergen challenged in mice [23]. Moreover,

conditional ABL knockout mice are refractory to allergen challenge-associated increased airway

resistance, airway smooth muscle mass, and tracheal contractility [23]. Collectively, our finding of

lung tissue-specific induction of CBX1 and ABI1 is consistent with the presence of inherent

biological pathways that are primary drivers of fibro-proliferative changes associated with

airway remodeling [3], and airway hyper-responsiveness.

374   In BALF samples from allergen-challenged mice, SPTBN1 (Spectrin β chain, non-

erythrocytic 1; also known as ELF) was increased in abundance, and our network analyses

revealed linkage to HDAC1 in tissue via SMAD4 and TAL1. Spectrin is associated with

cytoskeletal maintenance with ankyrin and anion exchanger 1 (AE1) to stabilize cortical actin

links to the cell membrane [24]. Spectrin repeat and Ph domains of SPTBN1 bind cell membrane

phospholipids, thus is positioned to respond to, and effect, interactions between cells in complex

biological systems [25, 26]. SPTBN1 also has a critical role in TGF-β induced gene transcription, as

SMAD3/4 become mis-localized and dysfunctional in SPTBN1 knockout mouse [27]. This is

critical as TGF-β induced SMAD3/4 signaling can induce human airway smooth muscle cell

shortening and hyperresponsiveness, and is a primary extracellular driver of asthma pathobiology

[28]. Based on its key role in mediating cytoskeletal structure and TGF-β signaling, our finding that

SPTBN1 is induced in BALF likely reflects its role in cell-cell interactions, including paracrine

signaling systems.

In BALF from allergen exposed mice we detected increased PSMA2 and PSMA5

(proteasome subunit α -2 and -5). These proteins were distinct in that interactions with lung

tissue protein nodes were limited to single links with HDAC1 and EVL, suggesting their primary

roles lie within the BALF compartment. PSMA2 and PSMA5 maintain structure and function of

the 20S proteasomal complex, and has increased activity in BALF from acute respiratory distress

patients [29, 30]. Intratracheal instillation of a proteasome inhibitor reduces NF-κB signaling and

eosinophil number in the BALF from OVA exposed rats [31]. A definitive role for proteasomal

complexes in BALF after allergen challenge is unclear, but they have been implicated in diverse

processes such as the formation exogenous peptide antigens for immune cells, degradation of

oxidatively damaged proteins, and activation of pre-cursor proteins [32]. We identified PSMA2 and

PSMA5 as a significant node in BALF that is relatively disconnected from lung tissue protein

nodes, suggesting their principal role lies in regulation of inflammation, involving inflammatory

cells, and perhaps their interaction with structural cells of the lung.

Proteins that mediate lung tissue-BALF crosstalk in allergen exposed mice

We identified remarkable differences in the protein profile between lung tissue and

BALF, even more so than the proteome changes that develop in each sample type after allergen

challenge. This highlights need to consider a role for protein interactions between lung tissue and

BALF after allergen challenge in order to fully interrogate lung system responses to allergic

insult. To meet this need, mandated that we use a unique design in which we collected lung

tissue and BALF samples from each mouse, performed proteomic analysis in each sample, and

then re-combined datasets to predict how the most significant protein nodes in lung tissue and

BALF interact. Protein interaction network analysis identified 6 proteins, NANOG, POU5F1,

TFCP2I1, SMAD4, TAL1 and YWHAZ, that are present in both lung tissue and BALF datasets,

and that could be pivotal mediators of coordinated responses involving both biological

compartments.

NANOG (Homeobox protein NANOG) and POU5F1 (POU domain, class 5, transcription

factor 1; also known as OCT4) are transcription factors that can function in concert to regulate

several genes, including EPHA, FGF2, SMAD1, and SKIL, that are of relevance to lung

diseases. These targets regulate cell matrix production and cell motility (EPHA) [33], induce

airway smooth muscle hyperplasia (FGF2) [34], and provide transcriptional control of TGF-β,

either directly by receptor activated SMAD1 [35], and indirectly, by the inhibitory protein, SKIL

(also called SnoN) [36]. SKIL, which is part of a family of proteins including SKI and Sno that

repress transcription of TGF-β response genes, by interacting with SMAD3 and SMAD4.

Interestingly, as noted above, network analysis identified SMAD4, which binds TGF-β induced

R-Smads to form transcriptional activator complexes, and TAL1, a hematopoietic transcription

factor, as regulators of coordinated responses in BALF and lung tissue. SKI, SKIL and other

TGF-β signaling inhibitors regulate cell responses dependent on SMAD4 and TAL1 [37, 38]. TGF-β

is a central driver of asthma pathobiology, and its elevated levels in lung tissue and BALF is

associated with diverse effects. Highlighting this, in mice, though neutralizing antibodies for

TGF-β have little effect on HDM challenge-induced lung dysfunction and airway smooth muscle

thickening, they do alter the nature and degree of airway inflammation [39]. Overall, our analysis

reveals that coordination of responses in BALF and lung tissue compartments involves

considerable transcriptional regulation, in particular related to the broad biological effects of

TGF-β on tissue repair and remodeling, inflammation, immunity and hematopoietic cell

maturation and activation.

Our inter-compartment network analysis also points to the airway epithelium and a key

zone for lung tissue-BALF crosstalk, which is consistent with the fact that epithelial barrier

function disruption and mucus hypersecretion are hallmark features of the response to repeated

allergen inhalation. We identified the transcription factor, TFCP2I1 (Transcription factor CP2-

like protein 1; also known as Crtr-1) as a novel co-ordinating protein node. Notably, it has a

diverse roles in epithelial cell maturation and modulating β-catenin signaling [40, 41]. To our

knowledge, a role for TFCP2I1 in allergic lung pathobiology has not been reported, but in

kidneys and salivary glands it is crucial regulator of epithelial cell maturation [40]. There is

evidence from lung cancer studies that TFCP2I1 directly interacts with YWHAZ, and protects β-

catenin from degradation [42]. YWHAZ, also called 14-3-3ζ, is another protein we predict to be a

coordinator of lung tissue-BALF biological responses. It is a binding partner with numerous

cytoplasmic and nuclear proteins and modulates signaling associated with apoptosis and cell

cycle progression [43, 44]. Furthermore, 14-3-3 proteins can bind the 5' untranslated region of

human surfactant protein A2 mRNA, which may be liked to surfactant deficiency in neonatal

respiratory distress syndrome [45]. A role for YWHAZ in β-catenin signaling is potentially

important in asthma pathobiology, as β-catenin is associated regulation of TGF-β regulated cell-

cell adhesion and epithelial to mesenchyme transition [41], and Wnt/β-catenin signaling is a

significant determinant of airway remodeling in asthma [46]. Our finding that TFCP2I1 and

YWHAZ may be associated with coincident allergic pathobiology in lung tissue and BALF

suggests that the initiation and progression of epithelial barrier disruption and tissue remodeling

results from an integrated network of pathways that are associated, in part, with TGF-β and

Wnt/β-catenin.

Limitations

Interpretation of our study is limited by a number of factors. Though we carefully

controlled collection methods to reduce variability, such as minimizing lung tissue damage

caused by collecting BALF, we cannot discount that BALF samples included a small fraction of

cellular proteins. We also only collected samples at one time point (48 hours after final allergen

challenge), a strategic choice, as this is when lung dysfunction is greatest. Thus, our work

provides only a snapshot of the proteome response, and future studies should adopt a design to

assess temporal patterns of the response to HDM challenge. We also only used a single allergen,

HDM, but this was a strategic selection as HDM a common human aeroallergen, and has a

complex composition that includes multiple allergens, proteases and toxins, therefore offers the

advantage of inducing inflammation of a broad profile [7, 8].

Summary

We characterized the proteome of lung tissue and BALF from HDM-challenged mice that

mimic allergic asthma pathophysiology. Using matched samples from individual animals, our

work reveals that lung tissue and BALF proteomes are diverse, and that integrating both datasets

reveals additional novel biological processes and protein interaction nodes, in particular

highlighting transcriptional regulation as a key integrating parameter for coincident pathobiology

in lung tissue and extracellular spaces of the lung. This work provides a resource and approach

for identifying new proteins and pathways, and a basis to interrogate interactions between sample

compartments to identify mechanisms for airways pathophysiology and, perhaps, new targets for

developing therapeutic approaches.

Methods

Animal experiments

(a) Murine HDM allergen challenge

All animal experiments were planned and performed following the approved protocols and guidelines of the animal ethics board at the University of Manitoba. Female, 8-week old BALB/c mice (6-8 weeks, $n = 3$) were intranasally challenged with HDM (25 µg per mouse, 35 µL saline) five times a week for two weeks (Supplemental Figure S3). HDM extract (Greer Labs; Lenoir, NC) was prepared in sterile phosphate buffered saline (pH 7.4; Life Technologies; Waltham, MA). The HDM extract we used contained 36,000 endotoxin units (EU; 7877 EU/mg of protein or 196.9 EU/dose) and 4.9% Der p 1 protein per vial.

(b) Lung function, inflammatory differential cell counts and sample collection

Lung function was performed 48 h after the last HDM challenge. Mice were anesthetized with sodium pentobarbital (90 mg/kg) given intraperitoneally, and tracheotomized with a 20-gauge polyethylene catheter. The catheter was connected to a flexiVent small animal ventilator (Scireq; Montréal, Canada) and mechanically ventilated with a tidal volume of 10 mL/kg body weight, 150 times/min. Forced oscillation technique and positive end expiratory pressure of 3 cm·H_2O was used for the entire study. A nebulized methacholine (MCh) challenge (0 to 50 mg/mL) was administered to assess concentration dependent response of the respiratory mechanics. Values for each parameter (newtonian resistance, Rn; peripheral tissue damping, G; tissue elastance, H; total resistance, R) were calculated as the peak of all 12 perturbation cycles performed after each MCh challenge. Statistical analysis was conducted using a two-way nested ANOVA with Tukey's multiple comparison and FDR correction (performed in R).

Post lung function measurement, lungs were lavaged with 1.0 mL of saline two times, for a total of 2 mL containing 0.1% ethylenediaminetetraacetic acid (EDTA; Sigma-Aldrich; St.

Louis, MO). BALF was centrifuged to collect the immune cell pellet (1,000 xg, 10 min, 4 ℃)

and the supernatant was collected and aliquoted prior to flash freezing in liquid nitrogen and

storage at -80 ℃. Immune cell pellet was resuspended in saline and the total immune cell count

was estimated using a hemocytometer. For differential counts, cells were stained with a modified

Wright-Giemsa stain (HEMA 3 STAT PACK, Fisher Scientific; Waltham, MA). Cell

distribution was analyzed by manually identifying and counting eosinophils, neutrophils,

macrophages and lymphocytes in six randomly chosen fields of view examined under a light

microscope at 200 x magnification. Post-BAL lung tissue from the left lung and half the right

lung were portioned (~35 mg/each), flash frozen in liquid nitrogen and stored at -80 ℃.

Preparation of lung tissue

A randomly selected portioned lung tissue was thawed and weighed before wash the lung

tissue of residual blood contamination. Lung tissue was submerged in a 15 mL centrifuge tube

containing 15 mL PBS (-CaCl$_2$, -MgCl$_2$, pH 7.4; Invitrogen, Waltham, MA) with

protease/phosphatase inhibitors and mixed in an end-over-end mixer (4 ℃, 30 min). Inhibitors

including Phenylmethylsulfonyl Fluoride (PMSF, 100 mM stock), Phosphatase Inhibitor

Cocktail 2 (Sigma-Aldrich; St. Louis, MO), and Protease Inhibitor (Sigma-Aldrich) each at

1:100 dilution. The tissue surface was cut with scissors to increase effectiveness of the lung

tissue wash.

Tissues were homogenized using a TissueRuptor (Qiagen; Venlo, Netherlands) in

siliconized centrifuge tubes (Thomas Scientific; Swedesboro, NJ) containing ice cold lysis

buffer. Lysis buffer composition: 150 mM NaCl (Fisher Scientific), 50 mM Tris-HCL (pH 7.5;

Fisher Scientific, Waltham, MA), 5% glycerol (Sigma-Aldrich), 1% sodium deoxycholate

(Sigma-Aldrich), 1% benzonase (25 U/µL; Merck,; Kenilworth, NJ), 1% sodium dodecyl sulfate

(SDS; Fisher Scientific), Protease Inhibitor Cocktail 2 (1:100 dilution, Sigma-Aldrich), 1 mM

PMSF (Sigma-Aldrich), Phosphatase Inhibitor Cocktail 2 (1:100 dilution, Sigma-Aldrich), 2 mM

$MgCl_2$ (Fisher Scientific) built up with molecular grade water (Invitrogen). Cell free supernatant

(21,000 xg, 10 min, 4 ºC, no break) was incubated at room temperature for 30 min to permit

benzonase activity before storage at -80ºC. All chemicals used for tissue preparation were of

molecular/electrophoresis grade.

Assessment of protein extraction proficiency

From each randomly chosen portion of frozen lung tissue, our protein extraction process

yielded an average of 1.146 mg of total protein (µBCA protein assay; Pierce; Waltham, MA).

Our rationale for randomizing our selection for a portion of frozen lung tissue resides in the

heterogenous nature of allergen induced airway inflammation throughout the lung - there is no

region of the lung where the extent of allergen exposure can induce uniform changes across

animals. BALF yielded an average of 600 µg of total protein (100 µg per 250 µL aliquot) per

mouse (µBCA, Pierce). A coomassie blue stained (GelCode, Invitrogen) gradient LDS-PAGE

(NuPAGE; Invitrogen) was imaged (ChemiDock, Bio-Rad) to qualitatively assess total protein

molecular weight diversity of both lung tissue (Supplemental Figure S4A) and BALF

(Supplemental Figure S4B) protein lysates. No high abundance protein depletion methods were

employed to reduce potential elimination bias as shown by the dark band at ~66.5 kDa

approximating the molecular weight of albumin (Supplemental Figure S4A,B).

Replicate analysis comparing both technical and biological replicates used mouse BALF

HDM sample in parallel protein Filter Assisted Sample Preparation (FASP) procedures. BALF

was divided into equal parts (100 µg total protein each) and processed individually and in

parallel through our FASP protocol. Our results show that technical variation (Supplemental

Figure S4E) is lower than our biological variation (Supplemental Figure S4F) and therefore, we

negated the use of technical replicates for our proteomic analysis.

Filter assisted sample preparation (FASP) of lung tissue and BALF

We modified a previously used FASP protocol for use in lung tissue [47]. Tissue

homogenate (300 µg total protein) was supplemented with DTT (final concentration 100 mM)

before boiling for 5 min, cooled and supernatant (21,000 xg, 10 min, 4 ºC, no break) collected.

Tissue homogenates were built with urea buffer (8 M urea, 100 mM Tris) to 800 µL before

loading onto 30 kDa molecular weight cut off filters (Amicron Ultra 0.5; Milipore; Burlington,

MA). Once samples were loaded (10,000 xg, 10 min, room temperature), columns were washed

twice with urea buffer (450 µL; 10,000 xg, 10 min, room temperature). Columns were alkylated

(400 µL, 50 mM iodoacidimide in urea buffer; Sigma-Aldrich) for 45 min (protected from light,

room temperature) before being stopped by the addition of DTT (20 mM) followed by

centrifugation (13,000 xg, 10 min, room temperature). Columns were washed twice (450 µL urea

buffer) before a drying the column (14,000 xg, 15 min, room temperature). Trypsin (Trypsin

Gold; Promega; Madison, WI) was suspended in digestion buffer (50 mM Tris, 2 mM $CaCl_2$,

LCMS water) and was added to the column at a protein:trypsin ratio of 50:1. Samples were

incubated under shaking conditions at 37 ºC for 16 h before being halted by the addition of TFA

(1% final concentration) and placing the samples at 4 ºC. Columns were incubated (400 µL, 50%

methanol, 5 min) prior to being eluted (10,000 xg, 10 min, room temperature). A final wash of

the column was also added to the eluted sample (300 µL 15% acetonitrile). Frozen samples (-80

ºC) were lyophilized via speed vac and stored at -80 ºC.

Mouse BALF samples were processed in a similar manner to that of mouse lung tissue

with some minor changes to the protocol. We resuspended 100 µg total protein from thawed

BALF supernatant in urea buffer (450 µL; containing 100 mM DTT final concentration) under

mixing conditions for 1 h (room temperature) before loading onto columns.

Peptide desalting by reverse phase 1D-HPLC

Warmed samples were re-suspended in 800 µL TFA (0.5%, LCMS grade water) and

centrifuged (21,000 xg, 10 min, 4 ºC) to check for undissolved peptides. BALF were loaded onto

a C18 column (Luna 10 µM C18(2), 100 Å, 50 x 4.6 mm; Phenomenex, Torrance, CA) while

tissue lysate samples were loaded onto a separate C18 column (Phenomenex) offline. Column

efficiency and elution conditions were tested using a specialized 6 peptide solution prior to

starting the samples [48]. Samples were fractionated across six 1.5 mL tubes and were collected at

a flow rate of 500 µL/min with an additional 30 s before and after the eluted peptide spectra was

detected. Manual loading was used with no gradient. Samples were frozen at -80 ºC. An Agilent

1110 HPLC System using ChemStation for Control and Data Analysis (Santa Clara, CA) was

used to analyze the chromatograms from Reverse Phase – HPLC (high pressure liquid

chromatography).

Reconstitution of sample & LC-MS/MS run

Samples were desiccated by speed vac (as previously mentioned) and reconstituted in 50

µL formic acid (0.1%). Peptide concentration was determined by UV spectrophotometry

(Nanodrop Spectrophotometer 2000, Thermofisher) at 280 nm. Peptides (2 µg) were diluted in

formic acid (0.1%) and injected into an online LC-MS/MS workflow (500 nL/min) using a 3 h

gradient run using a Sciex TripleTOF 5600 instrument (Sciex; Framingham, MA). Raw spectra

files were converted into Mascot Generic File format (MGF) for protein identification using the

tools bundled by the manufacturer. All chemicals used were mass spec grade.

Data Processing

The MGF files were processed by X!Tandem [49] against single-missed-cleavage tryptic

peptides from the *Mus musclus* Uniprot database (16,704 proteins). The following X!Tandem

search parameters were used: 20 ppm and 50 ppm mass tolerance for parent and fragment ions,

respectively; constant modification of Cys with iodoacetamide; default set post-translational

modifications: oxidation of Met, Trp; N-terminal cyclization at Qln, Cys; N-terminal acetylation,

phosphorylation (Ser, Thr, Tyr), deamidation (Asn and Gln); an expectation value cut-off of $\log_e$

< -1 for both proteins and peptides. Each MS run yielded a list of protein expression values in a

$\log_2$ scale, quantified based on their member peptide MS2 fragment intensity sums. Quantified

proteins were detected if at least two non-redundant (unique) peptides with identification scores

$\log_e$ < -1.5 each were identified.

Data Acquisition: replication, randomization and blocking

In order to homogenize the variance in our studies and avoid confounding technical

variability with the biological question, we randomized the order of all our experimental sample

work-up. In each treatment group, biological variability was captured with three replicate

animals. Moreover, in the case of LC/MS experiments, the sample order was randomized in

order to ameliorate effects related to instrumental drift and other variations. The hierarchical

order for the magnitude of variability in many omics experiments is well documented and it is

this principle that guided our data acquisition strategy [50].

Immunoblotting Validation of Proteomics

Protein targets were selected based upon proteomic enrichment after HDM exposure

(hypergeometric model with FC >= 2, p.adj <= 0.05, Benjamini-Hochberg multiple comparison)

and by antibody availability. Immunoblotting was performed as previously mentioned [51]. Briefly,

SDS-PAGE was performed on the same proteins samples used for proteomic analysis. Once 20

µg (total protein) of each sample was transferred onto nitrocellulose membranes, blots were

stained with Ponseau S total protein stain (Sigma) prior to blocking at room temperature (5%

skim milk, TBST 0.2%, 2 h, RT) and immunoblotted overnight at 4 ℃ (1% milk, TBST 0.2%).

After washing (3 x 15 min, TBST 0.2%), secondary antibody was added (Goat anti-rabbit HRP,

1% milk, TBST 0.2%, 2 h, RT, 1:5000; Sigma). Once washed, blots were flooded with ECL

(Amersham; GE Healthcare, Chicago, IL) and bands detected using chemiluminescent film

(Hyperfilm ECL; GE Healthcare). Semiquantitative densitometric analysis was performed using

scanned film and AlphaEase FC software (Alpha Innotech, San Leandro, CA). Antibodies used:

FDPS (PA5-28228, 1:1,000; Thermofisher), ARG1 (ab91279, 1:1,000; Abcam, Cambridge,

United Kingdom), CLCA1 (ab180851, 1:2,500; Abcam). To measure average protein content,

pooled samples (Naïve Pool and HDM Pool) were prepared by combining equal protein content

across biological replicates to equal 20 µg.

Bioinformatics & Statistical Analysis

Our raw data dataset (n=2,675) across treatment groups: allergen-naïve (n=1,996), and

HDM exposed (n=2,502). To limit the number of comparisons we employed a strict filtering

criteria; we selected only proteins that are detected in both the compartments, i.e., we kept only

those proteins that are identified in all replicates of at least one treatment (HDM or naive).

Together this dataset is referred to the Integrated Tissue-BALF dataset (ITB, n=1,237).

All the data analyses were conducted in the R, version 3.6.1 (*Action of the Toes*), environment

using in-house written code employing packages: limma, vsn, ggplots and pheatmap. The

stepwise workflow for the analyses was as follows.

(a) Data pre-processing and filtering and normalization

The BALF proteomic measurement of one HDM exposed mouse (labeled #379) was

found to be missing over 80% of Uniprot IDs compared to other allergen exposed BALF

samples. This sample was also found to be a technical outlier on examination of within group

clusters visualized by Principal Component Analysis (PCA). This sample was therefore excluded

from all downstream analysis leaving a total of 13 samples to comprise our full dataset (n=13)

from which our results are observed (Figure S2B,C). To retain a robust biological signature for

the remaining samples, we instituted a data filtering criterion in which we retained proteins

whose signal was identified in at least 2 out of 3 technical replicates (or at least 3 out of 4 for

HDM BALF) of at least one condition. This then allowed us to impute the rest of the missing

values using the k-nearest neighbour (knn) imputation method. To proportionally compare the

protein abundances from lung tissue and BALF samples, we performed background correction

and normalization by variance stabilizing normalization (vsn) [52]. Using this approach, the

variance of the dataset remains nearly constant over the whole dynamic range of our proteomic

samples.

(b) Exploratory Data Analysis and Differential Analysis (LIMMA)

In order to visualize the (dis)similarities between our proteomics measurements of the

samples under different conditions, we used principal component analysis (PCA) as a dimension

reduction technique. The variance of all the sample measurements for each protein was

calculated, and the top 500 proteins with the greatest variance was selected. We examined the

relationship among samples by projecting the original measurements on the two PCs that account

for over 80% of the data variance, and plotted these projections as shown in Figure 1A and

Figure 3A.

Moreover, in order to identify significantly enriched or depleted proteins we performed

differential abundance analysis on the vsn-normalized data using the Linear Models for

Microarray Data (LIMMA) approach [53]. We used a statistical threshold of log2 fold changes ≥ 1

and reported FDR adjusted p-values using Benjamini-Hochberg multiple comparison correction.

Threshold for statistical significance was p.adj ≤ 0.05. The cutoff criterion for all LIMMA results

are uniform across this study.

(c) Data visualization

To identify the distribution of individual protein IDs across groups, proportional area

venn diagrams were constructed using the eulerr R package (v6.1.0).

Heatmaps were constructed to identify the distribution of protein IDs across conditions.

These heatmaps were constructed using either normalized expression values from LIMMA

(Figure 1D and 2B) or using a presence/absence classification to identify unique protein IDs

(Figure 1B). In addition, grouping analysis (k-means clustering) was constructed for each

heatmap identifying groupings across both protein ID and sample condition. Figures were

generated using the pheatmap package in R (v1.0.12). Histograms, bargraphs, boxplots, line

plots, and linear regressions were constructed using DataGraph (DataGraph v4.5, Visual Data

Tools, Inc., Chapel Hill, NC, USA, https://www.visualdatatools.com/).

(d) Functional Analyses

Using the Gene Ontology database in InnateDB we first performed over-representation

analysis using the default parameters: hypergeometric model with Benjamini-Hochberg multiple

comparison correction [54]. To retain non-redundant biological processes we used Revigo with a

similarity search pattern of 0.5 [55]. Statistical significance of biological pathway enrichment was

set at p.adj ≤ 0.05. Moreover, we performed further functional analysis by conducting a protein-

protein interaction networks analysis (NetworkAnalyst) and Biological Pathway Assessment

(Reactome). Protein-protein interaction network analysis was performed by inputting UniprotIDs

into NetworkAnalyst (v3.0, accessed April 2020) and analyzed using the InnateDB protein-

interaction network database [54, 56]. The top 1^{st} order network (by number of connections) was

selected for further analysis. Within each network, the most interconnected protein nodes (from

the input Uniprot ID list) that are unique to either BALF or tissue, were selected and subset from

the larger networks to examine direct connections between the selected nodes. Biological

pathway analysis was then performed using the Reactome database module inside

NetworkAnalyst (hypergeometric model with p.adj ≤ 0.05 by Benjamini-Hochberg multiple

comparison). The protein UBC not detected in any of our analyses and was removed from all

protein-protein interaction analyses.

(e) Dataset preparation for predicting lung tissue/BALF interactions

We first integrated the 362 proteins that were uniquely enriched by HDM challenge in

lung tissue with the 311 proteins that were uniquely enriched after allergen challenge in BALF.

After excluding the 49 proteins that are shared between the two datasets, we input the lung tissue

(n=311), BALF (n=262), and combined (n=573) datasets into NetworkAnalyst. To focus on the

nodes that are the most interconnected between the BALF and lung tissue datasets we filtered the

combined dataset to include proteins with five or more 1^{st} order interactions [57]. From the 37

proteins that met this criterion (12 from lung tissue, 25 from BALF), we selected the top 5

proteins from the BALF and lung tissue dataset that become enriched (% enrichment) in the

combined dataset. Selecting the top 5 proteins from each dataset was essential to focus network

complexity on the top contributors to protein-interactions.

Data Storage

We have made all proteomic data freely available in the Mass Spectrometry Interactive

Virtual Environment (MassIVE) database housed at UCSD under the id MSV000086003. Files

are available at reviewers can grab the files at: ftp://MSV000086003@massive.ucsd.edu

(password: a)

Acknowledgements

Work for this study was supported in part by the CIHR-Canadian Respiratory Research Network (CRRN), Research Manitoba, the DEVOTION Network, and the Biology of Breathing Group, Children's Hospital Research Institute of Manitoba. THM was supported by studentships from Research Manitoba, CRRN, AllerGen NCE Inc. and Asthma Canada. CDP was supported by a CIHR Banting Fellowship and a Research Manitoba Fellowship. AJ was supported by studentships from Research Manitoba and CRRN. AJH is supported through the Canada Research Chairs Program.

Author Contributions

THM performed all proteomics experiments, primary and secondary data analysis, prepared all figures and tables and wrote the manuscript draft. CDP provide supervision, guidance and assisted in the completed biostatistics and bioinformatics, design of figures, and editing and writing of the manuscript. TKK provided secondary data quality control, statistical and bioinformatic support as well as support for design and editing of both figures and the final manuscript. AJ and SB were involved with experimental design and completion of animal studies. PE guided sample preparation and performed mass spectrometry. VS contributed to experimental design, primary proteomic data analysis, proteomic data quality control and primary statistics. NM contributed to experimental design and direction, and edited the manuscript. AJH conceived and led design of the study, including scope of biostatistics and bioinformatics, and contributed to writing and editing of the final manuscript.

Conflict of Interest

The authors declare no competing or financial interests.

References

References

(1) Calderón, M. A.; Linneberg, A.; Kleine-Tebbe, J.; De Blay, F.; Hernandez Fernandez de

Rojas, D.; Virchow, J. C.; Demoly, P. Respiratory allergy caused by house dust mites: What do

we really know. *J Allergy Clin Immunol*. **2015**, *136*, 38-48.

(2) Choopong, J.; Reamtong, O.; Sookrung, N.; Seesuay, W.; Indrawattana, N.; Sakolvaree, Y.;

Chaicumpa, W.; Tungtrongchitr, A. Proteome, Allergenome, and Novel Allergens of House Dust

Mite, Dermatophagoides farinae. *J Proteome Res*. **2016**, *15*, 422-430.

(3) Jha, A.; Ryu, M. H.; Oo, O.; Bews, H. J.; Carlson, J. C.; Schwartz, J.; Basu, S.; Wong, C. S.;

Halayko, A. J. Prophylactic benefits of systemically delivered simvastatin treatment in a house

dust mite challenged murine model of allergic asthma. *Br J Pharmacol*. **2018**, *175*, 1004-1016.

(4) Wu, J.; Kobayashi, M.; Sousa, E. A.; Liu, W.; Cai, J.; Goldman, S. J.; Dorner, A. J.; Projan,

S. J.; Kavuru, M. S.; Qiu, Y.; Thomassen, M. J. Differential proteomic analysis of

bronchoalveolar lavage fluid in asthmatics following segmental antigen challenge. *Mol Cell*

Proteomics. **2005**, *4*, 1251-1264.

(5) O'Neil, S. E.; Sitkauskiene, B.; Babusyte, A.; Krisiukeniene, A.; Stravinskaite-Bieksiene, K.;

Sakalauskas, R.; Sihlbom, C.; Ekerljung, L.; Carlsohn, E.; Lötvall, J. Network analysis of

quantitative proteomics on asthmatic bronchi: effects of inhaled glucocorticoid treatment. *Respir*

Res. **2011**, *12*, 124.

(6) Burg, D.; Schofield, J. P. R.; Brandsma, J.; Staykova, D.; Folisi, C.; Bansal, A.; Nicholas, B.;

Xian, Y.; Rowe, A.; Corfield, J.; Wilson, S.; Ward, J.; Lutter, R.; Fleming, L.; Shaw, D. E.;

Bakke, P. S.; Caruso, M.; Dahlen, S. E.; Fowler, S. J.; Hashimoto, S.; Horváth, I.; Howarth, P.;

Krug, N.; Montuschi, P.; Sanak, M.; Sandström, T.; Singer, F.; Sun, K.; Pandis, I.; Auffray, C.;

Sousa, A. R.; Adcock, I. M.; Chung, K. F.; Sterk, P. J.; Djukanović, R.; Skipp, P. J.; The, U.-B.

S. G. Large-Scale Label-Free Quantitative Mapping of the Sputum Proteome. *J Proteome Res*.

2018, *17*, 2072-2091.

(7) Pascoe, C. D.; Jha, A.; Basu, S.; Mahood, T.; Lee, A.; Hinshaw, S.; Falsafi, R.; Hancock, R.

E. W.; Mookherjee, N.; Halayko, A. J. The importance of reporting house dust mite endotoxin

abundance: impact on the lung transcriptome. *Am J Physiol Lung Cell Mol Physiol*. **2020**, *318*,

L1229-L1236.

(8) Piyadasa, H.; Altieri, A.; Basu, S.; Schwartz, J.; Halayko, A. J.; Mookherjee, N. Biosignature

for airway inflammation in a house dust mite-challenged murine model of allergic asthma. *Biol*

Open. **2016**, *5*, 112-121.

(9) Wypych, T. P.; Marzi, R.; Wu, G. F.; Lanzavecchia, A.; Sallusto, F. Role of B cells in T_H cell

responses in a mouse model of asthma. *J Allergy Clin Immunol*. **2018**, *141*, 1395-1410.

(10) Turner, D. L.; Goldklang, M.; Cvetkovski, F.; Paik, D.; Trischler, J.; Barahona, J.; Cao, M.;

Dave, R.; Tanna, N.; D'Armiento, J. M.; Farber, D. L. Biased Generation and In Situ Activation

of Lung Tissue-Resident Memory CD4 T Cells in the Pathogenesis of Allergic Asthma. *J*

Immunol. **2018**, *200*, 1561-1569.

(11) Svenningsen, S.; Kirby, M.; Starr, D.; Coxson, H. O.; Paterson, N. A.; McCormack, D. G.;

Parraga, G. What are ventilation defects in asthma. *Thorax*. **2014**, *69*, 63-71.

(12) Su, X. M.; Ren, Y.; Li, M. L.; Zhao, X.; Kong, L. F.; Kang, J. Performance evaluation of

histone deacetylases in lungs of mice exposed to ovalbumin aerosols. *J Physiol Pharmacol*.

2018, *69*, 265-273.

(13) Ito, K.; Caramori, G.; Lim, S.; Oates, T.; Chung, K. F.; Barnes, P. J.; Adcock, I. M.

Expression and activity of histone deacetylases in human asthmatic airways. *Am J Respir Crit*

Care Med. **2002**, *166*, 392-396.

(14) Ito, K.; Ito, M.; Elliott, W. M.; Cosio, B.; Caramori, G.; Kon, O. M.; Barczyk, A.; Hayashi,

S.; Adcock, I. M.; Hogg, J. C.; Barnes, P. J. Decreased histone deacetylase activity in chronic

obstructive pulmonary disease. *N Engl J Med*. **2005**, *352*, 1967-1976.

(15) Poynter, M. E.; Irvin, C. G.; Janssen-Heininger, Y. M. Rapid activation of nuclear factor-

kappaB in airway epithelium in a murine model of allergic airway inflammation. *Am J Pathol*.

2002, *160*, 1325-1334.

(16) Kallsen, K.; Andresen, E.; Heine, H. Histone deacetylase (HDAC) 1 controls the expression

of beta defensin 1 in human lung epithelial cells. *PLoS One*. **2012**, *7*, e50000.

(17) Redhu, N. S.; Saleh, A.; Lee, H. C.; Halayko, A. J.; Ziegler, S. F.; Gounni, A. S. IgE

induces transcriptional regulation of thymic stromal lymphopoietin in human airway smooth

muscle cells. *J Allergy Clin Immunol*. **2011**, *128*, 892-896.e2.

(18) Huang, Y.; Myers, M. P.; Xu, R. M. Crystal structure of the HP1-EMSY complex reveals an

unusual mode of HP1 binding. *Structure*. **2006**, *14*, 703-712.

(19) Lu, J.; Gilbert, D. M. Proliferation-dependent and cell cycle regulated transcription of

mouse pericentric heterochromatin. *J Cell Biol*. **2007**, *179*, 411-421.

(20) Ezell, S. A.; Polytarchou, C.; Hatziapostolou, M.; Guo, A.; Sanidas, I.; Bihani, T.; Comb,

M. J.; Sourvinos, G.; Tsichlis, P. N. The protein kinase Akt1 regulates the interferon response

through phosphorylation of the transcriptional repressor EMSY. *Proc Natl Acad Sci U S A*. **2012**,

109, E613-21.

(21) Chen, J.; Chen, Q.; Wu, C.; Jin, Y. Genetic variants of the C11orf30-LRRC32 region are

associated with childhood asthma in the Chinese population. *Allergol Immunopathol (Madr)*.

2019, *48*, 390-394.

(22) Wang, T.; Cleary, R. A.; Wang, R.; Tang, D. D. Role of the adapter protein Abi1 in actin-

associated signaling and smooth muscle contraction. *J Biol Chem*. **2013**, *288*, 20713-20722.

(23) Cleary, R. A.; Wang, R.; Wang, T.; Tang, D. D. Role of Abl in airway hyperresponsiveness

and airway remodeling. *Respir Res*. **2013**, *14*, 105.

(24) Nestor, M. W.; Cai, X.; Stone, M. R.; Bloch, R. J.; Thompson, S. M. The actin binding

domain of βI-spectrin regulates the morphological and functional dynamics of dendritic spines.

PLoS One. **2011**, *6*, e16197.

(25) Ray, S.; Chakrabarti, A. Membrane interaction of erythroid spectrin: surface-density-

dependent high-affinity binding to phosphatidylethanolamine. *Mol Membr Biol*. **2004**, *21*, 93-

100.

(26) Diakowski, W.; Sikorski, A. F. Interaction of brain spectrin (fodrin) with phospholipids.

Biochemistry. **1995**, *34*, 13252-13258.

(27) Tang, Y.; Katuri, V.; Dillner, A.; Mishra, B.; Deng, C. X.; Mishra, L. Disruption of

transforming growth factor-beta signaling in ELF beta-spectrin-deficient mice. *Science*. **2003**,

299, 574-577.

(28) Ojiaku, C. A.; Cao, G.; Zhu, W.; Yoo, E. J.; Shumyatcher, M.; Himes, B. E.; An, S. S.;

Panettieri, R. A. TGF-β1 Evokes Human Airway Smooth Muscle Cell Shortening and

Hyperresponsiveness via Smad3. *Am J Respir Cell Mol Biol*. **2018**, *58*, 575-584.

(29) Keller, I. E.; Vosyka, O.; Takenaka, S.; Kloß, A.; Dahlmann, B.; Willems, L. I.; Verdoes,

M.; Overkleeft, H. S.; Marcos, E.; Adnot, S.; Hauck, S. M.; Ruppert, C.; Günther, A.; Herold, S.;

Ohno, S.; Adler, H.; Eickelberg, O.; Meiners, S. Regulation of immunoproteasome function in

the lung. *Sci Rep*. **2015**, *5*, 10230.

(30) Sixt, S. U.; Adamzik, M.; Spyrka, D.; Saul, B.; Hakenbeck, J.; Wohlschlaeger, J.; Costabel,

U.; Kloss, A.; Giesebrecht, J.; Dahlmann, B.; Peters, J. Alveolar extracellular 20S proteasome in

patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med*. **2009**, *179*, 1098-

1106.

(31) Elliott, P. J.; Pien, C. S.; McCormack, T. A.; Chapman, I. D.; Adams, J. Proteasome

inhibition: A novel mechanism to combat asthma. *J Allergy Clin Immunol*. **1999**, *104*, 294-300.

(32) Sixt, S. U.; Peters, J. Extracellular alveolar proteasome: possible role in lung injury and

repair. *Proc Am Thorac Soc*. **2010**, *7*, 91-96.

(33) Yamazaki, T.; Masuda, J.; Omori, T.; Usui, R.; Akiyama, H.; Maru, Y. EphA1 interacts

with integrin-linked kinase and regulates cell morphology and motility. *J Cell Sci*. **2009**, *122*,

243-255.

(34) Jeon, S. G.; Lee, C. G.; Oh, M. H.; Chun, E. Y.; Gho, Y. S.; Cho, S. H.; Kim, J. H.; Min, K.

U.; Kim, Y. Y.; Kim, Y. K.; Elias, J. A. Recombinant basic fibroblast growth factor inhibits the

airway hyperresponsiveness, mucus production, and lung inflammation induced by an allergen

challenge. *J Allergy Clin Immunol*. **2007**, *119*, 831-837.

(35) Liu, F.; Hata, A.; Baker, J. C.; Doody, J.; Cárcamo, J.; Harland, R. M.; Massagué, J. A

human Mad protein acting as a BMP-regulated transcriptional activator. *Nature*. **1996**, *381*, 620-

623.

(36) Nyman, T.; Trésaugues, L.; Welin, M.; Lehtiö, L.; Flodin, S.; Persson, C.; Johansson, I.;

Hammarström, M.; Nordlund, P. The crystal structure of the Dachshund domain of human SnoN

reveals flexibility in the putative protein interaction surface. *PLoS One*. **2010**, *5*, e12907.

(37) He, J.; Tegen, S. B.; Krawitz, A. R.; Martin, G. S.; Luo, K. The transforming activity of Ski

and SnoN is dependent on their ability to repress the activity of Smad proteins. *J Biol Chem*.

2003, *278*, 30540-30547.

(38) Meng, X.; Lu, P.; Bai, H.; Xiao, P.; Fan, Q. Transcriptional regulatory networks in human

lung adenocarcinoma. *Mol Med Rep*. **2012**, *6*, 961-966.

(39) Fattouh, R.; Midence, N. G.; Arias, K.; Johnson, J. R.; Walker, T. D.; Goncharova, S.;

Souza, K. P.; Gregory, R. C.; Lonning, S.; Gauldie, J.; Jordana, M. Transforming growth factor-

beta regulates house dust mite-induced allergic airway inflammation but not airway remodeling.

Am J Respir Crit Care Med. **2008**, *177*, 593-603.

(40) Yamaguchi, Y.; Yonemura, S.; Takada, S. Grainyhead-related transcription factor is

required for duct maturation in the salivary gland and the kidney of the mouse. *Development*.

2006, *133*, 4737-4748.

(41) Giangreco, A.; Lu, L.; Vickers, C.; Teixeira, V. H.; Groot, K. R.; Butler, C. R.; Ilieva, E. V.;

George, P. J.; Nicholson, A. G.; Sage, E. K.; Watt, F. M.; Janes, S. M. β-Catenin determines

upper airway progenitor cell fate and preinvasive squamous lung cancer progression by

modulating epithelial-mesenchymal transition. *J Pathol*. **2012**, *226*, 575-587.

(42) Chen, C. H.; Chuang, S. M.; Yang, M. F.; Liao, J. W.; Yu, S. L.; Chen, J. J. A novel

function of YWHAZ/β-catenin axis in promoting epithelial-mesenchymal transition and lung

cancer metastasis. *Mol Cancer Res*. **2012**, *10*, 1319-1331.

(43) Weerasekara, V. K.; Panek, D. J.; Broadbent, D. G.; Mortenson, J. B.; Mathis, A. D.; Logan,

G. N.; Prince, J. T.; Thomson, D. M.; Thompson, J. W.; Andersen, J. L. Metabolic-stress-

induced rearrangement of the 14-3-3ζ interactome promotes autophagy via a ULK1- and AMPK-

regulated 14-3-3ζ interaction with phosphorylated Atg9. *Mol Cell Biol*. **2014**, *34*, 4379-4388.

(44) Lee, J. J.; Lee, J. S.; Cui, M. N.; Yun, H. H.; Kim, H. Y.; Lee, S. H.; Lee, J. H. BIS targeting

induces cellular senescence through the regulation of 14-3-3 zeta/STAT3/SKP2/p27 in

glioblastoma cells. *Cell Death Dis*. **2014**, *5*, e1537.

(45) Noutsios, G. T.; Ghattas, P.; Bennett, S.; Floros, J. 14-3-3 isoforms bind directly exon B of

the 5'-UTR of human surfactant protein A2 mRNA. *Am J Physiol Lung Cell Mol Physiol*. **2015**,

309, L147-57.

(46) Kumawat, K.; Koopmans, T.; Gosens, R. β-catenin as a regulator and therapeutic target for

asthmatic airway remodeling. *Expert Opin Ther Targets*. **2014**, *18*, 1023-1034.

(47) Wisniewski, J. R.; Zougman, A.; Nagaraj, N.; Mann, M. Universal sample preparation

method for proteome analysis. *Nat Methods*. **2009**, *6*, 359-362.

(48) Krokhin, O. V.; Spicer, V. Peptide retention standards and hydrophobicity indexes in

reversed-phase high-performance liquid chromatography of peptides. *Anal Chem*. **2009**, *81*,

9522-9530.

(49) Craig, R.; Beavis, R. C. TANDEM: matching proteins with tandem mass spectra.

Bioinformatics. **2004**, *20*, 1466-1467.

(50) Karakach, T. K.; Wentzell, P. D.; Walter, J. A. Characterization of the measurement error

structure in 1D 1H NMR data for metabolomics studies. *Anal Chim Acta*. **2009**, *636*, 163-174.

(51) Tran, T.; McNeill, K. D.; Gerthoffer, W. T.; Unruh, H.; Halayko, A. J. Endogenous laminin

is required for human airway smooth muscle cell maturation. *Respir Res*. **2006**, *7*, 117.

(52) Välikangas, T.; Suomi, T.; Elo, L. L. A systematic evaluation of normalization methods in

quantitative label-free proteomics. *Brief Bioinform*. **2018**, *19*, 1-11.

(53) Ritchie, M. E.; Phipson, B.; Wu, D.; Hu, Y.; Law, C. W.; Shi, W.; Smyth, G. K. limma

powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic*

Acids Res. **2015**, *43*, e47.

(54) Breuer, K.; Foroushani, A. K.; Laird, M. R.; Chen, C.; Sribnaia, A.; Lo, R.; Winsor, G. L.;

Hancock, R. E.; Brinkman, F. S.; Lynn, D. J. InnateDB: systems biology of innate immunity and

beyond--recent updates and continuing curation. *Nucleic Acids Res*. **2013**, *41*, D1228-33.

(55) Supek, F.; Bošnjak, M.; Škunca, N.; Šmuc, T. REVIGO summarizes and visualizes long

lists of gene ontology terms. *PLoS One*. **2011**, *6*, e21800.

(56) Zhou, G.; Soufan, O.; Ewald, J.; Hancock, R. E. W.; Basu, N.; Xia, J. NetworkAnalyst 3.0:

a visual analytics platform for comprehensive gene expression profiling and meta-analysis.

Nucleic Acids Res. **2019**, *47*, W234-W241.

(57) Vallabhajosyula, R. R.; Chakravarti, D.; Lutfeali, S.; Ray, A.; Raval, A. Identifying hubs in

protein interaction networks. *PLoS One*. **2009**, *4*, e5344.

Tables

928 **Table 1:** Significantly enriched or depleted proteins after HDM challenge in the combined

929 tissue-BALF dataset. Statistical threshold ($log_2FC \geq 1$, p.adj ≤ 0.05). Abbreviations: House Dust

930 Mite (HDM), log_2 fold change (LogFC)

Enriched/Depleted in HDM	Uniprot ID	Name	Description	Log_2FC	p.adj
Enriched	O35744	CHIL3	Chitinase-like protein 3	6.89	0.0018
Enriched	P97425	EAR2	Eosinophil cationic protein 2	2.69	0.0125
Enriched	Q9D7Z6	CLCA1	Calcium-activated chloride channel regulator 1	2.68	0.0133
Enriched	Q9CPY7	LAP3	Cytosol aminopeptidase	2.44	0.00181
Enriched	P24452	CAPG	Macrophage-capping protein	2.39	0.0172
Enriched	Q61233	LCP1	Plastin-2	2.08	0.00692
Enriched	Q00612	G6PDX	Glucose-6-phosphate 1-dehydrogenase X	1.81	0.0102
Enriched	O88342	WDR1	WD repeat-containing protein 1	1.26	0.014
Depleted	Q9DCT8	CRIP2	Cysteine-rich protein 2	2.33	0.00181
Depleted	P10649	GSTM1	Glutathione S-transferase Mu 1	1.59	0.00679

934   **Table 2:** Network complexity increases when combining the lung tissue and BALF datasets that

935   are uniquely altered after HDM exposure. The number of protein interactions (both 1st and 2nd

936   order) between the lung tissue and BALF datasets are shown (far right column).

Dataset Uniquely Altered by HDM	Uniprot ID	Name	Description	# of Interactions		Protein Node Enrichment in the Combined (%)	Number of 1st and 2nd Order Tissue-BALF Interactions
				Tissue	Combined		
Tissue	P83917	CBX1	Chromobox protein homolog 1	4	7	175.00%	3
Tissue	P61202	COPS2	COP9 signalosome complex subunit 2	3	5	166.67%	0
Tissue	Q8CBW3	ABI1	Abl interactor 1	5	8	160.00%	0
Tissue	O09106	HDAC1	Histone deacetylase 1	23	34	147.83%	7
Tissue	P70429	EVL	Ena/VASP-like protein	5	7	140.00%	3

940   **Table 3:** Network complexity does not change when combining the BALF and lung tissue

941   datasets that are uniquely altered after HDM exposure. The number of protein interactions (both

942   1st and 2nd order) between the BALF and lung tissue datasets are shown (far right column).

Dataset Uniquely Altered by HDM	Uniprot ID	Name	Description	# of Interactions		Protein Node Enrichment in the Combined (%)	Number of 1st and 2nd Order BALF-Tissue Interactions
				BALF	Combined		
BALF	Q62261	SPTBN1	Spectrin beta chain, non-erythrocytic 1	10	10	100.00%	7
BALF	P62908	RPS3	40S ribosomal protein S3	7	7	100.00%	1
BALF	Q99020	HNRNPAB	Heterogeneous nuclear ribonucleoprotein A/B	6	6	100.00%	2
BALF	Q9Z2U1	PSMA5	Proteasome subunit alpha type-5	5	5	100.00%	2
BALF	Q9R1P3	PSMB2	Proteasome subunit beta type-2	5	5	100.00%	3

Figure Legends

Figure 1: Characterization of the distinct proteomic signatures of the lung tissue and BALF. **A)** PCA of the top 500 most variable proteins identifies that BALF and tissue datasets have greater variation (59.4%) than the datasets comparing allergen-naïve and HDM challenged samples (22.4%). Note that the variability between the highly divergent tissue and BALF datasets masks the differences shown between allergen-naïve and HDM-exposed lung tissue as described in the figure inset (PCA of the top 500 most variable proteins). **B)** The differences between BALF and tissue samples is driven by missing proteins in BALF compartment as shown by K-means clustering heatmap of detectable proteins. **C)** The distribution of Uniprot ID's across lung tissue and BALF in both naïve and HDM exposed mice. **D)** K-means clustering heatmap showing protein abundance differences between lung tissue and BALF from allergen naive and HDM mice. K-means clustering identified 6 clusters (numbered on the left hand side). Red – high abundance, Blue – low abundance. Abbreviations used: House Dust Mite (HDM), Bronchial Alveolar Lavage Fluid (BALF), Principal Component Analysis (PCA).

Figure 2: Western blot validation of lung tissue and BALF proteomics. **A-B)** images of scanned chemiluminescent film for ARG1, CLC1, FDPS. Protein bands were normalized to total protein loading (Ponseau S) for quantification. Abbreviations used: Bronchial Alveolar Lavage Fluid (BALF), * (Welch's unpaired t-test p.adj ≤ 0.05), ND (Not Detected). Protein Names: ARG1 (Arginase-1), CLCA1 (Calcium-activated chloride channel regulator 1), FDPS (Farnesyl pyrophosphate synthase).

Figure 3: Combining tissue and BALF proteomes identified global processes in the lung of HDM challenged animals. **A)** PCA analysis highlights tissue sample and allergen driven differences in combined proteomes. **B)** Venn diagram showing proteins common to naïve and

HDM proteomes and heatmap identifying differentially abundant proteins common to both

proteomes. Red – high abundance, Green – low abundance. **C)** Gene ontology biological

processes enriched in the HDM proteome (p.adj $\leq$ 0.05). Vertical line indicates significance

threshold. **D)** Protein-protein interaction network of the top 3 most interconnected protein-

protein interactions nodes in the HDM proteome. Numbers beside each protein indicate number

of direct protein-protein interactions (1[st] order). Abbreviations used: Bronchial Alveolar Lavage

Fluid (BALF). Protein Names: SMARCA4 (Transcription activator BRG1), PIK3R1

(Phosphatidylinositol 3-kinase regulatory subunit alpha), NFKB1 (Nuclear factor NF-kappa-B

p105 subunit).

Figure 4: Individually analyzed HDM influenced lung tissue or BALF proteomes do not

recapitulate the biology of the Integrated-Tissue-BALF proteome. **A)** Venn diagram of the

distribution of protein IDs between allergen-naïve and HDM exposed lung tissue datasets. **B)**

Volcano plot showing significantly altered proteins that were common between naïve and HDM

tissue proteomes. Red points are significant (p.adj <0.05) and have a $\log_2 FC > 1$. Green points

have a $\log_2 FC > 1$ but are not significant. **C-D)** Gene ontology analysis for enriched biological

processes on differentially abundant and unique to HDM proteins in tissue. **E)** Venn diagram of

the distribution of protein IDs between allergen-naïve and HDM exposed BALF datasets. **F)**

Volcano plot showing significantly altered proteins that were common between naïve and HDM

BALF proteomes. No significantly altered proteins were detected. **G-H)** Gene ontology analysis

for enriched biological processes on unique to HDM proteins in BALF. For brevity, only the Top

10 non-redundant biological processes are shown. Vertical black line indicates significance (p =

0.05).

Figure 5: Most connected proteins in the HDM influenced lung tissue and BALF have distinct

biological signatures. **A)** Protein-protein interaction network of the top 3 most connected

proteins in the HDM influenced lung tissue proteome. **B)** Protein-protein interaction network of

the top 3 most connected proteins in the HDM influenced BALF proteome. Biological pathway

assessment from each network (Reactome) is shown the right hand side. Protein nodes and lines

are coloured to identify direct protein-protein interactions. Other parameters such as node size

and line distance is used for illustrative purposes only. Numbers beside each protein indicate

number of direct protein-protein interactions (1^{st} order). Abbreviations used: House Dust Mite

(HDM), Bronchial Alveolar Lavage Fluid (BALF). Protein Names: HDAC1 (Histone

deacetylase 1), MAPK1 (Mitogen-activated protein kinase 1), PLCG2 (1-phosphatidylinositol

4,5-bisphosphate phosphodiesterase gamma-2), SKP1A (S-phase kinase-associated protein 1A),

HIST1H4B (H4 clustered histone 2), PSMB6 (Proteasome subunit beta type-6).

1007

Figure 6: Integrating proteomes not only enrich protein-protein interactions but also connects

otherwise separate proteomes together. **A)** Protein-protein interaction network of the top 5 most

connected proteins in the HDM influenced lung tissue and BALF proteomes. Lung tissue nodes

(light red) are shown on the left. BALF nodes (light blue) are shown on the right. Protein nodes

that are directly involved (1^{st} order) in lung tissue-BALF crosstalk are shown in the center

shaded region (purple) and are labelled in grey. The Combined dataset is representative of all

protein nodes within green box. **B-D)** Biological processes derived from the Reactome dataset

are colour co-ordinated to their respective networks and each includes the 6 proteins in common

(purple). Circles represent individual proteins while protein-protein interactions are shown as

lines. Line colour that matches seed nodes indicate direct protein interactions with seed nodes.

Circles with split colours indicate shared 1^{st} order interactions between multiple seed nodes.

Black lines indicate 2^{nd} order (indirect) inter-dataset lung tissue-BALF crosstalk protein-

interactions. Broken lines indicate intra-dataset protein node interactions (2^{nd} order). Number of

direct protein-protein interactions (1st order) for each seed node is shown in parentheses. Other

parameters such as node size, node distribution and line distance are used for illustrative

purposes only. Abbreviations used: Bronchial Alveolar Lavage Fluid (BALF). Protein Names:

HDAC1 (Histone deacetylase 1), CBX1 (Chromobox protein homolog 1), EVL (Ena/VASP-like

protein), ABI1 (Abl interactor 1), COPS2 (COP9 signalosome complex subunit 2), SPTBN1

(Spectrin beta chain, non-erythrocytic 1), RPS3 (40S ribosomal protein S3), HNRNPAB

(Heterogeneous nuclear ribonucleoprotein A/B), PSMA5 (Proteasome subunit alpha type-5),

PSMA2 (Proteasome subunit alpha type-2), NANOG (Homeobox protein NANOG), POU5F1

(POU domain, class 5, transcription factor 1), TFCP2I1 (Transcription factor CP2-like protein 1),

SMAD4 (Mothers against decapentaplegic homolog 4), TAL1 (T-cell acute lymphocytic

leukemia protein 1 homolog) and YWHAZ (14-3-3 protein zeta/delta).

1032

Supporting Information

Supplemental 1: HDM exposure induces altered lung function and increased inflammatory cell counts. **A)** Lung mechanics were measured 48 h after last HDM treatment using a flexivent small animal ventilator. Increasing doses of methacholine (3-50 mg/mL) intranasally administered to measure changes in airway resistance, tissue resistance and tissue elastance. **B)** Differential immune cell counts from Bronchial Alveolar Lavage Fluid (BALF) of naïve and HDM exposed mice. Each value is representative of the mean and SEM of three biological replicates. Statistical significance was determined by a two-way nested ANOVA with Tukey's post-hoc test for lung function and unpaired t-test with welch's correction for cell counts. FDR corrected p-values are reported. *(p.adj ≤ 0.05), **(p.adj ≤ 0.01), ****(p.adj ≤ 0.0001).

Supplemental 2: Summary of MS/MS analysis from X!Tandem informatic pipeline. **A)** Summary of raw MS/MS data. An EV cutoff of -1.5 and a minimum of two peptides were needed for each peptide and protein ID. **B)** Distribution of unique protein IDs as distributed over the 13 replicates. **C)** Number of protein IDs on a per sample basis. Abbreviations used: Mouse (In House Mouse ID), number of spectra (SPEC), number of peptides (PEPS), number of non-redundant peptides (NR-PEPS), number of proteins (PROTS), number of quantified proteins (QPROT), number of quantified peptides (QPEPS), Mean and Standard Deviation of Log_2 MS/MS Intensity (MEAN & SD), Bronchial Alveolar Lavage Fluid (BALF), House Dust Mite (HDM). (*) Denotes the biological replicate that was removed due to technical reasons.

Supplemental 3: Experimental Workflow. Six mice were split into two groups and exposed to either HDM (House Dust Mite) or PBS (naïve) for two weeks. On day 14, mice were anesthetized and lung function data was collected using a small animal ventilator. Bronchial Alveolar Lavage Fluid (BALF) was collected, spun down to collect immune cells and flash

frozen in liquid nitrogen. Post-BAL lung tissue was portioned into 5 equal segments and flash

frozen in liquid nitrogen. Lung tissue was washed in PBS to remove excess blood and

subsequently homogenized. Samples were processed using FASP, trypsinized, desalted by 1D

HPLC and quantified using an online LC-MS/MS proteomic system. Protein IDs were identified

using our X!Tandem informatic pipeline and subsequent bioinformatic analysis was conducted

on the data. Protein IDs were then filtered retaining $\geq 2/3$ or $\geq 3/4$ biological replicates per

condition prior to variance stabilizing normalization. Protein IDs shared across conditions used

normalized protein abundance values as input for differential protein abundance analysis using

LIMMA. Protein IDs unique to a condition and those protein IDs which met LIMMA filtering

criteria were kept for protein functional analysis using biological pathway enrichment tools

(Gene Ontology, Reactome) and protein-protein interaction analysis (NetworkAnalyst).

Abbreviations used: Linear Models for Microarray Data (LIMMA), 1D HPLC (1 Dimension

High Pressure Liquid Chromatography), PBS (Phosphate Buffered Saline), LC-MS/MS (Liquid

Chromatography – Dual Mass Spectrometry), FASP (Filter Assisted Sample Preparation).

1072

Supplemental 4: Proteomic quality control and reproducibility tests indicate proteomic

variability resides in biological and not technical replicates. **A,B)** Representative SDS-PAGE

gels stained for total protein loading with coomassie blue of tissue and BALF protein

homogenates (20 µg loading for tissue, 7 µg for BALF). Each lane represents a different mouse.

C,D) Histograms of detected proteins (mean Log_2 protein abundance) for the tissue **(C)** and

BALF **(D)** datasets are all above the lower limit of detection (vertical black dotted line, 11.67).

Technical vs biological replicate assessment of Log_2 MS/MS intensities from HDM exposed

mouse BALF. **E,F)** Biological replicate assessment of Log_2 MS/MS intensities from mouse

BALF. Red line indicates linear goodness of fit (**D,F**).

1082

FIGURE 1

A

PCA plot - top 500 variable proteins

B

Missing values pattern
- Missing value
- ■ Valid value

C

D

log2 Centered intensity

FIGURE 2

FIGURE 3

Figure 4

Tissue (Naïve vs HDM)

A

Tissue (1787)

Naïve Tissue (1442)

HDM Tissue (1649)

138 | 1304 | 345

B

C Depleted in Tissue HDM

D Enriched in Tissue HDM

negative regulation of endopeptidase activity

complement activation alternative pathway

bioluminescence

regulation of intestinal cholesterol absorption

mannose metabolic process

positive regulation of cholesterol esterification

cellular aldehyde metabolic process

phosphatidylcholine metabolic process

acute-phase response

carbohydrate metabolic process

$-\log_{10}(p.adj)$

oxidation-reduction process

innate immune response

metabolic process

glutathione metabolic process

labyrinthine layer blood vessel development

chitin catabolic process

intracellular signal transduction

post-embryonic development

plasma membrane organization

ferric iron import into cell

$-\log_{10}(p.adj)$

Figure 4

BALF (Naïve vs HDM)

E

F

G

Depleted in BALF HDM

H

Enriched in BALF HDM

Figure 5

A Tissue

Antigen Activates B Cell Receptor Leading
to Generation of Second Messengers

Hemostasis

Signaling by Interleukins

G0 and Early G1

BMAL1:CLOCK/NPAS2
Activates Circadian Expression

TRAF6 Mediated Induction
of proinflammatory cytokines

MyD88:Mal cascade initiated
on plasma membrane

Toll Like Receptor TLR1:TLR2 Cascade

MyD88 cascade initiated
on plasma membrane

Toll Like Receptor 10 (TLR10) Cascade

$-\log_{10}(\text{p.adj})$

B BALF

SCF(Skp2)-mediated
degradation of p27/p21

Cyclin E associated
events during G1/S transition

Cyclin A:Cdk2-associated
events at S phase entry

Activation of NF-kappaB in B Cells

SCF-beta-TrCP mediated
degradation of Emi1

APC/C-mediated
degradation of cell cycle proteins

Regulation of mitotic cell cycle

Antigen processing:
Ubiquitination & Proteasome degradation

Regulation of APC/C activators
between G1/S and early anaphase

Degradation of beta-catenin by
the destruction complex

$-\log_{10}(\text{p.adj})$

Figure 6

A

Tissue

BALF

CBX1(7)

HDAC1(34)

EVL(7)

ABI1(8)

COPS2(5)

NANOG
POU5F1
TFCP2l1
SMAD4
TAL1
YWHAZ

RPS3(7)

HNRNPAB(6)

PSMA2(5)

PSMA5(5)

SPTBN1(10)

B   Tissue

Immune System

Signaling by Robo receptor

Cyclin D associated
events in G1

G1 Phase

Factors involved in megakaryocyte
development and platelet production

Adaptive Immune System

Hemostasis

Fcgamma receptor (FCGR)
dependent phagocytosis

TCR signaling

Innate Immune System

$-\log_{10}(p.adj)$

C   BALF

Apoptosis

Cross-presentation of soluble
exogenous antigens (endosomes)

Regulation of mRNA Stability
by Proteins that Bind AU-rich Elements

Regulation of ornithine
decarboxylase (ODC)

Regulation of activated PAK-2p34
by proteasome mediated degradation

CDK-mediated phosphorylation
and removal of Cdc6

p53-Independent DNA
Damage Response

p53-Independent G1/S
DNA damage checkpoint

Ubiquitin Mediated Degradation
of Phosphorylated Cdc25A

Ubiquitin-dependent degradation
of Cyclin D1

$-\log_{10}(p.adj)$

D   Combined

Immune System

Signaling by the
B Cell Receptor (BCR)

Mitotic G1-G1/S phases

Adaptive Immune System

Activation of NF-kappaB
in B Cells

Degradation of beta-catenin
by the destruction complex

Signaling by Wnt

Ubiquitin-dependent
degradation of Cyclin D1

Cell Cycle, Mitotic

S Phase

$-\log_{10}(p.adj)$