TITLE PAGE

Title: Evolutionary and ecological processes influencing chemical defense variation in an

aposematic and mimetic *Heliconius* butterfly

Short title: Chemical defense variation in a butterfly

Authors: Anniina L. K. Mattila[1,6*]; Chris D. Jiggins[2]; Øystein H. Opedal[1,3]; Gabriela Montejo-

Kovacevich[2]; Érika de Castro[2]; William O. McMillan[4]; Caroline Bacquet[5]; Marjo

Saastamoinen[1,6]

Author affiliations:

13
1. Research Centre for Ecological Change, Organismal and Evolutionary Biology Research

Programme, University of Helsinki, Helsinki, Finland

2. Department of Zoology, University of Cambridge, Cambridge, UK

3. Department of Biology, Lund University, Lund, Sweden

4. Smithsonian Tropical Research Institute, Gamboa, Panama

5. Universidad Regional Amazónica de Ikiam, Tena, Ecuador

6. Helsinki Life Science Institute, University of Helsinki, Helsinki, Finland

Corresponding author: Anniina L. K. Mattila: anniina.mattila@helsinki.fi, *current affiliation:

Finnish Museum of Natural History (LUOMUS), University of Helsinki, Helsinki, Finland

ABSTRACT

Chemical defences against predators underlie the evolution of aposematic coloration and mimicry, which are

classic examples of adaptive evolution. Surprisingly little is known about the roles of ecological and

evolutionary processes maintaining defence variation, and how they may feedback to shape the evolutionary

dynamics of species. Cyanogenic *Heliconius* butterflies exhibit diverse warning color patterns and mimicry,

thus providing a useful framework for investigating these questions. We studied intraspecific variation in *de*

novo biosynthesized cyanogenic toxicity and its potential ecological and evolutionary sources in wild

populations of *Heliconius erato* along environmental gradients, in common-garden broods and with feeding

treatments. Our results demonstrate substantial intraspecific variation, including detectable variation among

broods reared in a common garden. The latter estimate suggests considerable evolutionary potential in this

trait, although predicting the response to selection is likely complicated due to the observed skewed distribution

of toxicity values and the signatures of maternal contributions to the inheritance of toxicity. Larval diet

contributed little to toxicity variation. Furthermore, toxicity profiles were similar along steep rainfall and

altitudinal gradients, providing little evidence for these factors explaining variation in biosynthesized toxicity

in natural populations. In contrast, there were striking differences in the chemical profiles of *H. erato* from

geographically distant populations, implying potential local adaptation in the acquisition mechanisms and

levels of defensive compounds. The results highlight the extensive variation and potential for adaptive

evolution in defense traits for aposematic and mimetic species, which may contribute to the high diversity

often found in these systems.

KEYWORDS: chemical defenses, aposematism, mimicry, evolvability, *Heliconius*

MAIN TEXT

INTRODUCTION

Chemical defenses are a common means for animals to gain protection against predators. Defensive chemicals

can be acquired from the larval or adult diet (sequestered) and/or biosynthesized *de novo* by the organism

(Bowers, 1992; de Castro et al., 2019; Opitz & Müller, 2009). Chemical defenses are often coupled with bright

color patterns, which local predators learn to associate with toxicity or unpalatability of the prey, a

phenomenon known as aposematism (Ruxton et al., 2018). This can also lead to mimicry among co-occurring

prey species, one of the best studied examples of Darwin's theory of evolution by natural selection and the

subject of one of evolution's oldest mathematical models (Müller, 1879). Because palatability can vary along

a continuum (Arias et al., 2016; Prudic et al., 2019; Speed, 1999; Speed et al., 2012; Turner, 1984), mimicry

can take several forms; Batesian, where a palatable mimic exploits an unpalatable model, Müllerian, in which

unpalatable species copy one another for mutual benefit, or Quasi-Batesian, where a less well protected species

acts in a parasitic manner, diluting the protection of the better defended species (Mallet, 1999; Speed, 1993,

1999; Turner, 1987).

These different mimicry relationships all imply predictable differences in toxicity among

mimetic species and individuals (Speed, 1999; Turner, 1987), and yield different predictions about the origins

of diversity in warning coloration and mimicry, which is often greater than predicted by classic aposematic

theory (Briolat et al., 2019; Joron & Mallet, 1998; Speed, 1999). For example, alternation between different

mimicry types could influence patterns of convergent vs. advergent selection (where selective pressures cause

phenotypic convergence of one species on another, but not vice versa) (Mallet, 1999; Mallet & Joron, 1999;

Speed, 1999). However, there are many open questions related to the roles of ecological and evolutionary

processes shaping chemical defense variation in aposematic and mimetic species (Mallet & Joron, 1999; Speed

et al., 2012; Ruxton et al., 2018; Briolat et al., 2019). One outstanding current question is "what drives variation

in defensive toxins within and among prey populations?"

Extensive variation in defensive toxins within species and across mimicry rings has been

demonstrated repeatedly (Speed et al., 2012). Indeed, unpalatability levels of mimic species can range from

equal to very uneven (Arias et al., 2016; Bowers, 1992; Ritland & Brower, 1991). Potential sources of this

variation include host-plant chemistry, which often varies with the environment and is influenced by

resource availability (Massad et al., 2011). Host-plant chemistry can be a major source of intraspecific

variation particularly in sequestered defensive compounds (Bowers, 1992; Opitz & Müller, 2009), as seen in

many butterfly species (Camara, 1997; Gardner & Stermitz, 1988; Hay-Roe & Nation, 2007). Detoxification

of host plant defensive chemicals can also negatively affect an individual's resource budget, leading to

variation in investment in defenses (Lindstedt et al., 2010; Reudler et al., 2015). Similarly, species that have

evolved the ability to biosynthesize defensive compounds often vary extensively in their defensive

chemicals. In this case, the origin of both inter- and intraspecific variation is less clear (Ruxton et al., 2018;

Burdfield-Steel et al., 2019; Zvereva & Kozlov, 2016). Sources of variation may include biotic factors, such

as the energy and resources available for toxin synthesis, interactions with life-history traits including age

and reproductive stage, and general condition as assayed by e.g. immunological status (Bowers, 1992;

Smilanich et al., 2009; Zvereva & Kozlov, 2016). This variation could simply be maintained by drift if

toxicity profiles are selectively neutral, for example if acquiring and maintaining defenses does not incur

costs for the individual, and variation in defense level does not affect predator behavior (Speed et al., 2012;

Briolat et al., 2019).

88   In those species that sequester and those that synthesize defensive compounds, there are also

several potential adaptive explanations for the persistence of variation in nature (Speed et al., 2012). The

selection pressures acting on chemical defense traits are expected to be complex, as chemically defended

herbivores are often involved in close coevolutionary arms-races with their host plants, predators and

parasitoids (Turner, 1984; Ruxton et al., 2018). The costs of producing and maintaining toxicity may also

result in frequency- or density-dependent selection for resource optimization (Speed et al., 2012). Such costs

could lead to energetic trade-offs (Bowers, 1992; Fordyce & Nice, 2008; Lindstedt et al., 2020) and

automimicry (the occurrence of palatable "cheaters" in a chemically defended population; Brower et al.,

1970; Speed et al., 2012). Evolutionary explanations for chemical defense variation are only plausible if such

variation is genetically determined, and sufficient genetic variation is available for selection to act on.

Therefore, knowledge about the heritability and evolvability of chemical defense traits is fundamental in

evaluating the roles of evolutionary *vs.* ecological processes in the origins of the extensive variation observed

in these traits. Overall, empirical data unraveling the patterns and origins of chemical defense variation are

needed to help build a more complete understanding of the evolution and diversification of aposematic and

mimetic systems.

Here, we investigate aspects of chemical defense variation in neotropical *Heliconius*

butterflies. With their incredible diversity in wing coloration, they are a well-known example of

aposematism and mimicry (Jiggins, 2017; Merrill et al., 2015), and provide a powerful study system in

which to assess the role of ecology and evolution in chemical defense variation within species and

populations. *Heliconius* larvae feed exclusively on passion vines (*Passiflora*), with which they have radiated

and speciated through a coevolutionary arms race (Benson et al., 1975; de Castro et al., 2018). Almost 30

different cyanogenic glucoside structures have been reported in the *Passiflora* genus (de Castro et al., 2019)

and only the most simple of these compounds can be sequestered by *Heliconius* larvae (de Castro et al.,

2019; Engler et al., 2000). *Heliconius* larvae and adult butterflies can additionally biosynthesize the

cyanogens linamarin and lotaustralin *de novo* from aliphatic amino acids (Nahrstedt & Davis, 1983).

Although biosynthesis is not uncommon in chemically defended animals (Bowers, 1992; Ruxton et al.,

2018), it is uncommon among chemically defended Lepidoptera (Nishida, 2002). There is substantial

variation in the amounts and types of cyanogens among different *Heliconius* species (Arias et al., 2016;

Cardoso & Gilbert, 2013; de Castro et al., 2019; Engler-Chaouat & Gilbert, 2007; Sculfort et al., 2020).

Some of this is explained by the level of diet specialization, which is a major determinant of the relative roles

of cyanogen sequestration from plants and *de novo* biosynthesis (de Castro et al., 2019; Engler-Chaouat &

Gilbert, 2007). The level of sequestered defenses is also known to be affected by variation in the chemical

content of the host plants (Engler-Chaouat & Gilbert, 2007; Hay-Roe & Nation, 2007), whereas the sources

of variation in biosynthesis are less well understood.

We studied ecological and evolutionary aspects of intraspecific variation in biosynthesized

cyanogenic toxicity in *Heliconius erato* using a combination of extensive field collections and common-

garden rearing experiments. *Heliconius erato* provides an interesting study subject in terms of variation in

chemical defenses, warning coloration patterns and mimicry relationships, as it has a diverse array of color

forms associated with local color pattern mimicry rings, and is found in ecologically diverse habitats across a

large geographic range. Our field collections exploit two strong environmental gradients: a rainfall gradient

with a five-fold difference in rainfall, and a steep altitudinal gradient in the Andes. We use these data to

tackle three questions: 1) What is the extent of natural variation in toxicity within and among populations of

H. erato? 2) Can some of this variation be explained by environmental gradients or highly divergent habitat

types across a wider geographic scale? 3) How much of the variation in biosynthesized chemical defenses is

heritable and evolvable, and how much is governed by environmental (particularly dietary) variance? By

examining our results in the context of the well-studied mimicry system of *Heliconius* butterflies, we discuss

the potential wider implications of our findings for aposematic and mimetic systems.

METHODS

Study sites and field collection of butterflies and host plants

Heliconius erato demophoon and its typical host plant *P. biflora* were collected in Panama along a steep

rainfall gradient (the Smithsonian Tropical Research Institute Physical Monitoring Program;

https://biogeodb.stri.si.edu/physical_monitoring/) from three populations (Wet, Intermediate and Dry study

sites; Fig 1A, Supplementary File 4), and *H. erato lativitta* from High and Low altitude sites on the Eastern

slope of the Andes in Ecuador (Fig. 1A, Supplementary File 4). Butterflies were caught using a hand net and

transported live to the laboratory. All individuals caught from the Panamanian study areas ($n = 92$, of which

34, 28 and 30 individuals were caught from Dry, Intermediate and Wet sites, respectively) were sexed,

weighed, and preserved in 1 ml 100% MeOH (body excluding wings and one half of the thorax) on the day

of capture (males) or after being allowed to oviposit in an insectary (females, see Supplementary File 4). At

the Ecuadorian sites, fertilized females were brought to a common garden environment, allowed to oviposit

and raised separately according to the altitude where they were collected. The thorax of 21 F1 individuals

from each altitude (total $n=42$) were sampled in 1 ml 100% MeOH.

150   *Passiflora biflora* leaf samples (one to four leaves pooled per plant individual, see

Supplementary File 4) were collected from the Panama Dry ($n = 20$), Intermediate ($n = 11$) and Wet ($n = 20$)

study areas (Fig. 1A), and preserved fresh in 1ml 100% MeOH. In addition, 50 cm cuttings, all from

different plant individuals, were sampled from the Dry ($n = 12$) and Wet ($n = 12$) study areas for greenhouse

cultivation.

Host plant greenhouse cultivation and treatments

Passiflora biflora were cultivated in greenhouses (75% relative humidity; 8-20 hrs: 25°C, light; 20-8 hrs:

20°C, dark) in a soil mixture (50% compost, 20% coir, 15% perlite, 15% sand or gravel) (see Supplementary

File 4 for details on cultivation methods and watering treatments). Standard (std)-treated *P. biflora* were used

as oviposition plants for the *H. erato* parental generation and larval diet for the F1 families not included in

feeding treatments (see below). The quality of host plants in *H. erato* feeding treatments was manipulated

with watering treatments, motivated by the importance of water availability for many cyanogenic plants

(Gleadow & Woodrow, 2002; Hayden & Parker, 2002; Liang, 2003; Vandegeer et al., 2013). The *P. biflora*

cuttings collected from the Dry and Wet study areas in Panama were cultivated in a full-factorial

combination of origins and water treatments, resulting in four host plant types. Plants in the dry treatment

were kept at 75% relative humidity and watered two to three times weekly with about 1/3 of the water

volume compared to the wet treatment (constant water availability, relative humidity up to 100%), such that

the surface soil was allowed to dry and the leaves slightly droop between watering events. The dry treatment

thus aimed to mimic conditions of drought stress.

Butterfly rearing, breeding design and feeding treatments

Laboratory *H. erato* populations descendent of the wild Dry population were established in greenhouse

conditions (see Supplementary File 4 for details on rearing methods). Larvae were reared on std-type *P.*

biflora. All butterflies were individually marked, and were allowed to fly, mate, and oviposit freely in 2×2×2

m mesh cages with potted std-type *P. biflora* and a standardized diet of sugar solution supplemented with

amino acids provided *ad libitum*. The butterfly cages were observed throughout the day for mating pairs, and

mated females were moved to individual 2×2×2 m mesh cages and allowed to oviposit on std-type *P. biflora*.

Eggs of 14 females (eight mated with a known father, average no. of full-sibs/family = 8, see Supplementary

File 2) were collected, and hatched larvae were reared in full-sib groups fed on std-type *P. biflora* until

pupation.

181   The effect of larval dietary cyanogen content and quality on adult butterfly biosynthesized

toxicity was studied with feeding treatments. We expected larval dietary quality to affect butterfly toxicity

due to potential variation in the acquisition of nitrogenous resources for cyanogen biosynthesis (Cardoso and

Gilbert 2013), possible detoxification costs (Lindstedt et al., 2010; Reudler et al., 2015) and the general

importance of larval diet for fitness traits in holometabolous insects (e.g. Boggs & Freeman, 2005). The

offspring of six mated pairs (both parental identities known, average no. of full-sibs/family = 29,

Supplementary File 2) were included in the feeding treatments, where eggs from each mother were divided

equally into four groups and transferred onto each of the four *P. biflora* treatment types. Individuals of the

F1 generation were collected within three days following emergence (unmated and unfed), weighed, and

preserved in 100% MeOH (body excluding wings and one-half of thorax).

Butterfly toxicity analyses with [1]H-NMR

The concentrations of the two cyanogenic compounds biosynthesized by *Heliconius* larvae and adults,

linamarin and lotaustralin, were analyzed from the butterfly samples using nuclear magnetic resonance ([1]H-

NMR) with Bruker Avance III HD NMR spectrometer (Bruker BioSpin, Germany) (see Supplementary File

4 for details on sample extraction and [1]H-NMR methods). The data were processed and analyzed using

Bruker TopSpin software. The peaks of cyanogen compounds were identified based on comparison of

sample [1]H spectra with the [1]H spectra of authentic reference samples of linamarin and lotaustralin. Note that

we analyzed only the biosynthesized cyanogenic compounds (not sequestered cyanogenic compounds nor

other chemicals potentially influencing unpalatability in *Heliconius*; see Jiggins, 2017), and *Heliconius*

"cyanogenic toxicity" or "toxicity" refer here to biosynthesized cyanogenic toxicity. Our study species has

been considered generalist or oligophagous on *Passiflora*, largely acquiring its cyanogenic toxicity through

de novo biosynthesis of cyanogens (Engler-Chaouat & Gilbert, 2007; Merrill et al., 2013), and the cyanogen

profile of its main host plant *P. biflora*, also used in our experimental setup, consists of complex compounds

which *Heliconius* larvae are unable to sequester (de Castro et al., 2019; Engler-Chaouat & Gilbert, 2007).

206   All statistical analyses of the study were undertaken using R 3.4.4 (R Core Team, 2018). The

207   values of cyanogen concentrations in the wild-collected samples were normalized with a square-root-

208   transformation before applying ANOVA to test for population differences. Distributions of cyanogen traits

209   were also explored by inspection of truncated weighted density distributions (accounting for the skewness of

210   data towards near-zero values) using the "sm" R package (Bowman & Azzalini, 2014). For the analyses

211   using common-garden data, the data were normally distributed and untransformed values were used.

Host plant toxicity and quality analyses

The cyanogenic content of the *Passiflora biflora* samples was analyzed using liquid chromatography-mass spectrometry (LC-MS/MS) with Agilent 1100 Series LC (Agilent Technologies, Germany) hyphenated to a Bruker HCT-Ultra ion trap mass spectrometer (Bruker Daltonics, Germany) following the procedure of de Castro et al. (2019)(see Supplementary File 4 for details on sample extraction, spectrometry and analyses). Mass spectral data were analyzed with the native data analysis software (Compass DataAnalyses, Bruker Daltonics). The cyanogen compounds of *P. biflora* were detected and quantified following de Castro et al. (2019).

Plant quality traits assumed to be proxies of nutritional value, including nitrogen/carbon ratio (a proxy of protein/carbohydrate-ratio), were measured from the greenhouse-cultivated *P. biflora* treatment types (average $n = 63$) with Dualex® Scientific+ leaf clip meter (Cerovic et al. 2012; Supplementary File 4). We tested for differences in cyanogen levels and plant quality traits between the treatment groups in an ANOVA including plant origin, treatment and their interaction.

Estimation of variance components, heritability, evolvability and maternal effects in *Heliconius* biosynthesized cyanogenic toxicity

The concentration of the two tested biosynthesized cyanogens, linamarin and lotaustralin were strongly correlated with each other ($r = 0.897$). Cyanogen concentration was not correlated with adult body mass in either sex ($F_{1,193} = 0.52$, $P = 0.470$ for total cyanogen concentration; mass data available for $n = 196$ individuals). Adult mass also did not differ between the feeding treatments ($F_{4,186} = 0.88$, $P = 0.480$; model including feeding treatment, sex and their interaction), and was not included as a variable in further analyses.

Variance components of *Heliconius erato* biosynthesized cyanogenic toxicity were analyzed based on the two-generation full-sib/half-sib breeding design (Lynch & Walsh, 1998b) by fitting linear mixed models applying the "animal model" framework (Kruuk, 2004; Wilson et al., 2010), in which the full pedigree (relatedness matrix) is incorporated as an individual-level random effect. The models were fitted using ASReml-R Version 4 (Butler et al., 2017), an R package that fits linear mixed models using Residual Maximum Likelihood (REML) estimation. The two-generation pedigree consisted of 20 broods and their parents, and included mother-offspring, father-offspring, full-sibling and paternal half-sibling relationships

(n = 322 individuals, Supplementary File 2). Of the 20 broods, six were included in the feeding treatments

receiving four different *P. biflora* types as larval diet (Supplementary File 2). Individuals of the parental

generation were assumed to be unrelated. With the materials available, we maximized our power to detect

genetic variation by generating as many family groups as possible. Because the number of families was

limited, we also analyzed a large number of offspring per family, which allowed precise estimates of family

means. Because our breeding design resulted mostly in full-sib families, we have limited power to

distinguish additive-genetic from other genetic variance components (dominance, epistasis, maternal/paternal

effects). We thus consider our genetic-variance estimate as a 'broad-sense' genetic variance, although the

estimate approaches the narrow-sense estimate more than a simple full-sib design due to the parent-offspring

and half-sib relations in the breeding design (Lynch & Walsh, 1998b).

Animal models were fitted separately for the three biosynthesized cyanogenic toxicity traits

(concentration of linamarin, lotaustralin and total cyanogens). The primary models (referred to as "Animal

Model1") included the individual breeding value as a random effect, and the fixed effects of sex and feeding

treatment (variance explained by fixed effects computed using the method of Villemereuil et al. (2018); see

also Supplementary File 4). "Animal Model2" included additionally the random effect of maternal identity.

We computed broad-sense heritabilities (H^2) as the proportion of the total phenotypic variance (V_P)

explained by the (broad-sense) genetic variance component (V_G). When calculating heritabilities we included

the variance due to sex in the total phenotypic variance, but we did not include the variance due to the

treatment because this component of the variance is unlikely to represent natural variation (Villemereuil et

al., 2018). Heritabilities are useful measures of the resemblance between offspring and parents, but are poor

standardized measures of evolutionary potential (Hansen et al., 2011). As a measure of evolutionary

potential, we therefore computed mean-scaled evolvabilities (e_μ) as $e_\mu = V_G/\mu^2$, where μ is the trait mean (of

common garden brood data trait values), which gives the expected percent change in the trait mean per

generation under a unit strength of selection (Hansen & Houle, 2008; Hansen et al., 2011).

Given that our design included mostly full-sib groups, we obtained an additional estimate of

the component of toxicity variance explained by brood by estimating the intra-class correlation coefficient

(ICC), i.e. repeatability, with a linear mixed model approach. For this, we implemented the "rptR" package

(Stoffel et al., 2017), with brood identity as the grouping factor, a Gaussian distribution and 1000 parametric

bootstraps to quantify uncertainty.

To further explore the role of maternal contributions to the inheritance of cyanogenic toxicity,

we estimated maternal-effect coefficients following Falconer´s dilution model, which assumes that offspring

phenotype is a direct, linear function of maternal phenotype, beyond the contribution of maternally

transmitted genes (Lande & Price, 1989; Lynch & Walsh, 1998a; Walsh & Lynch, 2018b). Thus, the

interpretation of this 'maternal effect' is related to specific measured traits (here, cyanogenic toxicity), and is

different from the typical formulation of 'maternal environmental effects' (influence of the maternal

environment on offspring traits). We estimated maternal effect coefficients (m) as the difference in slope

between a mother-offspring (mo) and a father-offspring (fo) regression, i.e. $m = \beta_{mo} - \beta_{fo}$ (Lande & Price,

1989; Lynch & Walsh, 1998a). We fitted the parent-offspring regressions as linear mixed models with the

lme4 R package (Bates et al., 2014), including family as a random factor and maternal or paternal toxicity

phenotype as explanatory variables of offspring toxicity phenotype.

RESULTS

First, we investigated the ecological origins of variation in defensive traits by studying toxicity of butterflies

along steep gradients in rainfall and altitude (Fig. 1A). Toxicity levels were similar between populations

within the gradients, and variation in butterfly toxicity was not detectably associated with toxicity variation

of the major host plant species along the rainfall gradient (Fig. 1B). A comparison of geographically distant

natural populations of *H. erato* from Panama and Ecuador showed that populations of the same species can

have substantially different levels of biosynthesized toxicity (Fig. 1B). Furthermore, within natural *H. erato*

populations sampled along the rainfall gradient in Panama, the distribution of biosynthesized cyanogenic

toxicity was consistently skewed, so that a large proportion of individuals had a noticeably low level of

toxicity compared with a smaller proportion of highly toxic individuals within all sampled populations (Fig.

1C). Finally, we partitioned the variation in toxicity into genetic and environmental components, with

particular focus on the variance originating from diet. Common-garden pedigree data comprising 20 broods

and more than 300 *Heliconius erato* individuals revealed a moderate but detectable component of variance

explained by brood identity (around 12% of the total variance; Fig. 1D). However, further analyses

Figure 1. Natural variation and evolutionary potential of biosynthesized cyanogenic toxicity in *Heliconius erato*. **A)** Top panel: Locations of the natural study populations in Central and South America (Panama and Ecuador, respectively), with a more detailed map of study areas along the rainfall gradient in Panama. Bottom panel: Experimental setups for common garden populations of *H. erato* and host plant *Passiflora biflora*. **B)** Variation among natural populations in total- and total biosynthesized cyanogenic (CNglc) toxin concentrations in *P. biflora* and *H. erato*, respectively. Small-case letters indicate statistically significant differences between populations (Tukey HSD *P*<0.05). **C)** Zero-truncated kernel density distributions of total biosynthesized cyanogenic toxicity in natural populations of *H. erato* in Panama. **D)** Top panel: Variance components of *H. erato* biosynthesized cyanogenic toxicity (with S.E.) based on pedigree data (see panel A and Table 2). Bottom panel: Repeatability estimates with brood identity as the grouping factor and 1000 parametric bootstraps to quantify uncertainty.

suggested that the among-brood variance may not primarily represent additive genetic variance but rather be

driven by maternal environmental or genetic influences on offspring toxicity phenotype. In contrast,

nutritional quality indicators and cyanogenic toxicity of the host plant *Passiflora biflora*, which varied in

response to water availability treatments, had no effect on the biosynthesized toxicity level of the adult

butterflies that had fed on these plants as larvae. Together, these results demonstrate high levels of

intraspecific variation that is not detectably driven by the ecological factors studied here. They also indicate

complex modes of inheritance in this chemical defense trait of an aposematic and mimetic *Heliconius*

butterfly.

Wide natural variation in toxicity within populations and between regions, but not along

environmental gradients

The distribution of biosynthesized cyanogen levels, as estimated from the concentrations of biosynthesized

compounds linamarin and lotaustralin, was similar across natural populations of *H. erato* sampled along an

environmental rainfall gradient in Panama (Fig. 1C). A large proportion of individuals had low toxicity

values (e.g. 40% of individuals had total cyanogen concentrations falling within the lower 25% of toxicity

values), while a small proportion of individuals had a considerably higher toxicity level (e.g. 5% of

individuals had total cyanogen concentrations within the highest 25% of toxicity values). Total cyanogen

concentration and concentration of linamarin did not differ among the three Panamanian populations (Fig.

1B, Table 1, Supplementary file 1). The concentration of lotaustralin differed among populations ($F_{2,\,89} =$

7.97, $P < 0.001$), with the Intermediate population having higher concentrations than the two populations at

the opposite extremes along the rainfall gradient (Supplementary file 1).

		H. erato total cyanogens %						*P. biflora* total cyanogens (µg/mg)				
Country	**Population**	Mean	SD	Dry	Int	Wet	Low	High	Mean	SD	Dry	Int	Wet
Panama	*Dry*	1.064	0.709						11.850	10.073			
Panama	*Intermediate*	1.271	0.631	0.432					21.544	13.586	**0.038**		
Panama	*Wet*	0.907	0.521	0.905	0.098				18.005	7.953	0.146	0.627	
Ecuador	*Lowland*	0.011	0.028	**<0.001**	**<0.001**	**<0.001**							
Ecuador	*Highland*	0.026	0.033	**<0.001**	**<0.001**	**<0.001**	0.936						

Table 1. Total cyanogen concentrations (mean, standard deviation (SD) and *P*-values of Tukey HSD pairwise comparisons) in natural populations of *Heliconius erato* and *Passiflora biflora* sampled along environmental gradients in Panama and Ecuador. Statistically significant pairwise comparisons are marked in bold. See Supplementary file 1 for full results table with different cyanogen compounds shown separately.

The total cyanogen levels of the host plant *Passiflora biflora* differed among the natural

populations along the rainfall gradient in Panama ($F_{2,\,48} = 3.65$, $P = 0.034$; Fig. 1B, Table 1). The Dry

population had substantially lower cyanogen levels compared with both the Intermediate and Wet

populations (Fig 1B, Table 1). In contrast, levels of cyanogen did not vary substantially across populations of

H. erato (Fig. 1B, Table 1, Supplementary file 1).

The populations along the altitudinal gradient in Ecuador did not differ in biosynthesized

cyanogenic toxicity from each other, but differed markedly from their Panamanian conspecifics (Fig. 1B,

Table 1). Most Ecuadorian samples lacked both linamarin and lotaustralin (Supplementary file 1). Linamarin

was detected in only 29% of samples, most of which originated from the high-altitude site (75% of those

with any signal), and the average biosynthesized cyanogen level in these samples was very low (0.063%;

Fig. 1B, Table 1, Supplementary file 1), which is below the minimum level observed in the Panamanian

populations.

Biosynthesized toxicity does not vary with diet but varies among broods

The level of biosynthesized cyanogenic toxicity varied two-fold among the 20 common-garden reared broods (range of brood averages = 0.48% - 0.88% of dry mass, Fig. 2A). Phenotypic variance components of cyanogenic toxicity estimated with an animal model including the full pedigree as a random effect and fixed effects of sex and feeding treatment (Model1) revealed a statistically significant genetic variance component (V_G) for total biosynthesized cyanogen concentration level (Fig. 1D, Table 2, Supplementary File 1), corresponding to a broad-sense heritability (H^2) of 0.115 (H^2(linamarin) = 0.139, H^2(lotaustralin) = 0.067; Table 2, Supplementary File 1). We confirmed the presence of detectable among-brood variance in a simpler brood-level analysis, in which brood identity explained 12.5% of the variation in biosynthesized cyanogen concentration across common-garden reared broods (repeatability $R = 0.125$, $SE = 0.059$, $CI = [0.015, 0.245]$, $P = 6.02e-05$). Among-brood variance could originate from several sources including 1) additive genetic variance, 2) maternal environmental or genetic effects, and 3) shared environment, and below we explore these options further.

First, we consider the estimated broad-sense genetic variance component and estimate evolutionary potential of toxicity, assuming the genetic component consists of largely additive genetic variance. The broad-sense evolvability of total cyanogen concentration was e_μ=1.55%, $SE \pm 0.89$ (e_μ(linamarin) = 1.89%, e_μ(lotaustralin)= 1.04%; Table 2, Supplementary File 1), meaning that the trait mean is expected to change by 1.55% per generation per unit strength of selection (Hansen & Houle, 2008; Hansen et al., 2011).

Model type	Cyanogen trait	Trait mean (μ)	No. Obs.	No. broods	V_G	$P\,V_G$	H^2 (V_G/V_P)	e_μ (V_G/μ^2); %	V_{mat}/V_P	V_{trm}/V_P	V_{sex}/V_P	V_R/V_P	V_P
1	Total CNglc %	0.772	322	20	0.0093	0.037	0.115	1.555		0.011	0.011	0.874	0.080
1	Linamarin %	0.577	322	20	0.0063	0.020	0.139	1.887		0.001	0.013	0.848	0.045
1	Lotaustralin %	0.195	322	20	0.0004	0.130	0.067	1.039		0.008	0.004	0.929	0.006
2	Total CNglc %	0.772	322	20	1.0e-7	1.000	1.4e-6	1.7e-5	0.102	0.043	0.002	0.895	0.073

Column groups: Genetic variance component, heritability and evolvability (V_G, $P\,V_G$, H^2, e_μ); Other variance components (V/V_P) (V_{mat}/V_P, V_{trm}/V_P, V_{sex}/V_P, V_R/V_P); Total phenotypic variance (V_P).

Table 2. Summary of variance components of *Heliconius* biosynthesized cyanogenic (CNglc) toxicity estimated with REML animal models. V_{mat}, V_{trm}, V_{sex} and V_R are the variance components for maternal effects, host plant feeding treatment, sex, and residual variance, respectively. See Supplementary File 1 for full model results.

Our baseline estimates of repeatability, as well as broad-sense heritability and evolvability (Model 1) thus showed that there is moderate but detectable variation among broods in biosynthesized

Figure 2. A) Variation in total biosynthesized cyanogen (CNglc) concentration (% dry mass) among full-sibling broods of *Heliconius erato* by increasing maternal CNglc % (one pair of full-sib broods are paternal half-siblings, in grey) **B)** Total biosynthesized CNglcs concentration (% dry mass) in newly-emerged adult individuals fed with different *Passiflora biflora* host plant treatment types (see Fig. 1A) as larvae. **C)** Parent-offspring regressions of total biosynthesized CNglcs concentration.

toxicity. We then estimated maternal and paternal contributions to this variation. When including maternal

identity as a second random effect in the animal model (Model2), this maternal variance component

explained most of the variance explained by the genetic variance component in Model1, whereas V_G became

negligible, illustrating that the two random terms explain essentially the same variance component (i.e. the

among-brood variance; Table 2, Supplementary File 1). Parent-offspring regressions revealed that offspring

cyanogen concentration was positively associated with the cyanogen concentration of the mother (slope =

0.181 ± 0.071, $t = 2.539$, $P = 0.012$ for total cyanogen concentration; Fig. 2C), while the association between

fathers and offspring showed a negative, but statistically non-significant trend (slope = -0.122 ± 0.083, $t = -$

1.469, $P = 0.143$ for total cyanogen concentration). In turn, the midparent-offspring regression was nearly

flat (slope = 0.046 ± 0.107, $t = 0.435$, $P = 0.664$ for total cyanogen concentration; Fig. 2C). These regression

slopes translate into a maternal-effect coefficient for total cyanogen concentration of $m = 0.261$, $CI = [-$

$0.099, 0.614]$ (m(linamarin)= 0.288, CI = [-0.094, 0.687]; m (lotaustralin)= 0.193, $CI = [-0.135, 0.520]$),

which implies the existence of positive maternal effects in biosynthesized cyanogenic toxicity. However,

uncertainty in these maternal effects remains notable, as illustrated by the wide confidence intervals.

Finally, we investigated other sources, particularly environmental origins of variance. Larvae

reared on different host plant types (Fig. 1A) did not differ in biosynthesized cyanogenic toxicity as adults

(Fig. 2B), and the host plant type explained a negligible proportion of variance in butterfly toxicity level

($V_{trm}/V_P = 0.011$; Fig. 1D, Table 2, Supplementary File 1). This is despite the strong effect of the watering

treatments on the cyanogenic toxicity and quality indicators of the host plants (e.g. nitrogen-carbon balance;

see Supplementary File 3), and also indicates that among-brood variance is unlikely to be explained by

common environment effects related to dietary quality. Sex was also a negligible source of variation in

toxicity ($V_{sex}/V_P = 0.011$; Fig. 1D, Table 2, Supplementary File 1). Despite the common-garden conditions

(outside of dietary treatments) that all individuals experienced throughout their life cycle, the residual

(unexplained) variance component of cyanogen concentration remained large ($V_{res}/V_P = 0.874$ for total

cyanogens, Fig. 1D, Table 2, Supplementary File 1). This large amount of unexplained variance is also

demonstrated by a difference in toxicity level between the parental and offspring generation ($F_{1, 320} = 27.42$,

$P < 0.001$, effect size (Hedges' g) = 0.962 for total cyanogens), despite the two generations sharing common

garden conditions.

DISCUSSION

We have documented extensive variation in chemical defenses, both within and across, natural populations

of *H. erato*. Wide variation in defensive toxins has often been found in chemically defended prey, yet the

sources of this variation remains largely unclear (Speed et al., 2012). Our data shows that intraspecific

variation in *de novo* synthesized chemicals was not strongly associated with environmental conditions and

levels of biosynthesized toxicity were similar in butterflies collected across steep environmental gradients.

Similarly, our experimental work consistently showed that variation in host plant biochemical profiles and

nutritional quality do not contribute to variation in *H. erato* biosynthesized toxicity. Instead, the variation has

a moderate but detectable broad-sense genetic component (12.5% of variation explained by brood), which

could, given that it was largely composed of additive genetic effects, imply substantial evolutionary potential

in biosynthesized chemical defenses of *H. erato*. Our broad-sense heritability estimate of biosynthesized

cyanogenic toxicity (0.115 ± 0.086) is close to the average heritability value of physiological traits ($0.12 \pm$

0.05), based on a meta-analysis of a wide range of traits and taxa (Hansen et al., 2011). This estimate differs

markedly, though, from that obtained in a recent study on toxicity in warningly colored wood tiger moths

(*Arctia plantaginis*), in which no evidence for a genetic component to variation in the secretion amounts of

de novo synthesized chemical defenses could be detected (Burdfield-Steel et al., 2018; Lindstedt et al.,

2020). A few other studies have estimated genetic components of insect chemical defense variation, with

some detecting a genetic component (Eggenberger & Rowell-Rahier, 1992; Holloway et al., 1993; Yezerski

et al., 2004) and others not (Müller et al., 2003). Our study adds on these previous work by investigating

chemical defense variation across environments and mimicry rings.

400   Our estimated value for mean-scaled evolvability of cyanogenic toxicity, which gives a better

standardized measure of evolutionary potential (Hansen & Houle, 2008; Hansen et al., 2011) is high ($1.55 \pm$

0.89%), especially when compared to the average value of physiological traits ($0.49 \pm 0.14\%$), as well as the

median evolvability of all studied traits across taxa ($0.26 \pm 0.03\%$; see Hansen et al., 2011). The evolvability

of $e_\mu = 1.55\%$ means that the mean cyanogenic toxicity can be expected to change by 1.55% per unit strength

of selection per generation. This high estimate of evolvability would imply that the trait has the potential to

evolve rapidly, especially in conditions of strong selection. The effect would be magnified given the short

generation time (usually one to two months) of these tropical butterflies. For example, assuming *Heliconius*

has six generations per year, and increased cyanogenic toxicity is selected for with selection as strong as

selection on fitness as a trait ($\beta_\mu = 100\%$, see Hereford et al., 2004), the trait mean would have the potential to

increase by 9.7% in just one year.

411   However, our data from experimental crosses indicate that the inheritance of biosynthesized

cyanogenic toxicity has a more complex basis, beyond pure additive genetic effects. Particularly, there is

evidence for a central role for maternal effects. This is supported by our observations that 1) the maternal

effect component fitted in our animal model seemed to explain the same component of variance as the broad-

sense genetic component, as well as that 2) midparent-offspring and father-offspring regressions were non-

significant while mother and offspring toxicity values were positively correlated. Given that mothers were

raised in common garden conditions, and environmental variance is thus expected to be low, these effects

likely have a genetic or epigenetic basis. At this point, it is unclear if the maternal effects that we document

here are the result of maternally transmitted genes, epigenetic effects, direct transfer of toxins from mother to

offspring in eggs (Nahrstedt & Davis, 1983), or other types of maternal influence on the offspring phenotype

(Lande & Price, 1989; Lynch & Walsh, 1998a; Walsh & Lynch, 2018b). Females are the heterogametic sex

in Lepidoptera, but gene content of the highly heterochromatic W chromosome is poorly explored.

Heliconius eggs have very high concentrations of cyanogenic toxins (de Castro et al., 2020; Nahrstedt &

Davis, 1983) and the relationship between the provisioning of compounds in eggs by mothers and the level

of these compounds in adults could be explored in future studies. We did not detect any paternal effects,

even though *Heliconius* males are known to transfer a nuptial gift containing biosynthesized cyanogenic

compounds to females at mating (Cardoso & Gilbert, 2007).

Independent of what is driving the maternal effects, the occurrence of a maternal effect of the

type estimated here, defined simply as an effect of the mothers phenotype on the offspring's phenotype (as in

Falconer´s dilution model; Lande & Price, 1989; Lynch & Walsh, 1998a), could considerably strengthen the

response of the trait to selection. Although there is much uncertainty in how maternal effects will affect the

response to selection (Walsh & Lynch, 2018a), the predicted cumulative selection response with maternal

effects under Falconer´s model for a trait with a maternal effect coefficient of $m = 0.25$, which is comparable

to the $m = 0.261$ estimated here for cyanogenic toxicity, would over time exceed that of a trait with $m = 0$ by

as much as 50% (assuming constant heritability) (Walsh & Lynch, 2018a). This effect, however, is expected

to dilute over generations. The maternal coefficient should also be interpreted with caution because of

remaining uncertainty in our estimate. Predicting the response of toxicity to selection in the wild could also

be complicated by the observed skewed distribution of toxicity, as skewed breeding values could generate an

asymmetrical selection response. Furthermore, we cannot rule out that some of the observed among-brood

variance is explained by common environment effects other than maternal effects and the measured effect of

diet, e.g. differences in group size during larval development. However, the demonstrated negligible plastic

responses to diet, common-garden rearing, and the temporal spread of egg-laying of each brood (eggs laid

singly over a time period of up to two weeks per brood) are expected to reduce the likelihood of such

common environment effects among broods.

Implications of evolutionary potential in chemical defenses for aposematic and mimetic systems.

Studies of mimicry often focus on color pattern, i.e. the warning signal (Benson, 1972; Jiggins, 2017; Kapan,

2001; Mallet & Barton, 1989; Merrill et al., 2015). However, it is important to note that contrasting selective

agents could be acting on the defense trait associated with the warning signal, both traits equally required for

aposematism and mimicry. The selection pressures acting on chemical defense traits can be expected to be

complex and dynamic (Briolat et al., 2019; Jones et al., 2016; Speed et al., 2012). The role of predators as the

major selective agent is clearly a key component of possible adaptive evolutionary processes herein (Briolat

et al., 2019; Ruxton et al., 2018). Even high evolutionary potential will only lead to evolutionary changes if

the defense trait variation influences predation rates (Speed et al., 2012). In general, empirical data on the

impact of variation in defenses and its signaling on predators is lacking, but there is evidence for variation

among predators in traits including perceptual abilities, individual motivation and tolerance to chemical

defenses (Briolat et al., 2019; Halpin et al., 2017). Furthermore, prey toxicity level does not necessarily

correlate with actual unpalatability, the ultimate cue for predator choices (Winters et al., 2018). Predator-

mediated selection is therefore expected to be dynamic and difficult to estimate. In addition, bottom-up

selection related to co-evolutionary relationships with host plants, as well as selection on energy optimization

may also play a role (Turner, 1984; Ruxton et al., 2018).

Species may only respond adaptively to such selection pressures given heritable genetic

variation in these traits. Evolutionary potential in chemical defenses could allow species to respond to the

expected dynamic top-down and bottom-up selection, resulting in diverse patterns of adaptive evolution in

chemical defense traits. This could have important consequences for the evolutionary trajectories of

aposematic and mimetic species, and ultimately play a role in the evolution of warning signal diversity. This

is because evolutionary changes in chemical defense levels could alter mimetic relationships among groups

of species across geographic space, including shifts between Müllerian, Batesian and quasi-Batesian

relationships, which could in turn influence advergent vs. convergent selection and contribute to the

observed diversity in mimicry systems (Briolat et al., 2019; Mallet & Joron, 1999; Speed, 1999). With

regards to *Heliconius,* the combination of a locally dynamic selective environment, flexible genetic

mechanisms for generating warning signals (Van Belleghem et al., 2017; Wallbank et al., 2016), and

evolutionary potential in toxicity could explain the rapid origin of inter- and intraspecific variation. In this

light, the ability of *Heliconius* to biosynthesize defense compounds could allow for increased genetic control

of chemical defenses compared to most other chemically defended Lepidoptera, which rely solely on host

plant compound sequestration (Bowers, 1992; Nishida, 2002; Zvereva & Kozlov, 2016). Because *Heliconius*

can sequester cyanogenic compounds of only some *Passiflora* species (de Castro et al., 2019; Engler et al.,

2000), selection would be targeted toward *de novo* synthesized defenses particularly in environments lacking

such host species. Furthermore, previous investigations both confirmed that the cyanogenic glucoside

compounds in *Heliconius* deter vertebrate predators (Cardoso, 2020) and that predator avoidance learning

rate increases when comparing low to moderate levels of cyanogenic compound concentrations (Chouteau et

al., 2019). Thus, the level of cyanogen defenses is likely to be a direct target of selection by predators in

some circumstances.

By altering the nature of mimetic relationships or selection for increased defense levels on a

new warning signal form, evolvability of defense traits could drive the evolution and spread of new color

patterns and the reproductive isolation that may follow, potentially even leading to the formation of new

species (Jiggins et al., 2001; Mallet et al., 1998). A phylogenetic signal in the cyanogenic glucoside chemical

profile in the Heliconiini tribe supports this idea (de Castro et al., 2019; Sculfort et al., 2020) and

underscores the potential for adaptive evolution in chemical defense traits to influence evolutionary

trajectories in mimicry systems. However, more empirical data is called for in future studies to estimate

levels of evolutionary potential and patterns of selective responses in defensive traits in aposematic and

mimetic species.

Evolutionary processes shaping chemical defense variation: the role of automimicry

A further complication in chemical defense evolution is the interplay between individual and public good

(Jones et al., 2016). In this respect, our data demonstrating a broad-sense genetic component in the defense

trait coupled with patterns of natural variation of cyanogen levels in *Heliconius erato* populations are

consistent with balancing selection related to automimicry, in which automimic "cheaters" exploit the

protection given by their warning coloration with investing less in chemical defenses themselves.

Automimicry may occur if producing and maintaining chemical defenses is costly (Brower et al., 1970;

Speed et al., 2012), leading to selection reducing individual-level toxicity (Bowers, 1992; Ruxton et al.,

2018; Zvereva & Kozlov, 2016), and if there are also individual benefits of investing in chemical defenses

(e.g. Ruxton & Speed, 2006; Svennungsen & Holen, 2007). Previous studies indicate that *de novo* synthesis

of defenses may generally be more costly than the sequestration of plant compounds (Zvereva & Kozlov,

2016), which may make automimicry more likely in species with biosynthesized defenses. The occurrence of

automimicry could lead to a decrease in population average unpalatability, which could theoretically dilute

the protection conveyed by the warning signal (Gibson, 1984; Jones et al., 2016; Speed et al., 2012), as in

Batesian and quasi-Batesian mimicry, and influence the evolutionary trajectories of the species sharing the

warning signal. Thus, empirical data on within-population defense variation and automimicry can give

important insights into the diversity of mimetic systems (Jones et al., 2016).

511   Our results show that all three butterfly populations along the studied rainfall gradient

exhibited similar skewed distributions of biosynthesized cyanogens. The majority of individuals in these

populations had relatively low, although not zero, levels of toxicity (24% of individuals had less than 0.5%

concentration of biosynthesized cyanogenic compounds from an overall range of 0.1 - 2.9%), whereas a

smaller proportion had substantially higher levels, and the pattern was surprisingly similar in all populations.

Such a pattern could be explained by age distributions, as previous studies suggest that *Heliconius*

accumulate cyanogenic toxins with age (de Castro et al., 2020; Nahrstedt & Davis, 1985), or by individual

differences in the balance between sequestration and biosynthesis of cyanogenic compounds (de Castro et

al., 2019; Engler-Chaouat & Gilbert, 2007; de Castro et al., 2020). However, as many individuals in the

study populations had near zero levels of biosynthesized toxins, the pattern could also be indicative of

automimicry. In *Heliconius*, the costs of chemical defense remain poorly known, but some cases of

automimicry have previously been reported (Arias et al., 2016). Here, toxicity and mass were not associated,

which could imply that potential costs of cyanogen biosynthesis are not related to or mediated by a trade-off

with body size. In terms of benefits of chemical protection, requirements for automimicry are likely to be

met, as empirical data on *Heliconius* imply direct benefits (surviving predator attacks) for chemically

defended individuals (Boyden, 1976; Pinheiro, 1996; Pinheiro & Campos, 2019). Furthermore, an increase in

protection level is found especially when comparing low to moderate cyanogenic toxicity levels (Chouteau et

al., 2019), applicable to the observed cyanogen levels of *H. erato* (Arias et al., 2016; de Castro et al., 2019).

However, future studies on natural defense variation and automimicry in *Heliconius* should measure the

entire range of cyanogens (both sequestered and biosynthesized) and other chemical defense compounds

potentially interacting to produce the unpalatability level perceived by relevant predators (Jiggins, 2017;

Winters et al., 2018). In more general terms, future research in this framework should investigate the costs of

defensive compound biosynthesis (see also Zvereva & Kozlov, 2016) and its role in the evolution of

automimicry, as well as automimicry-originating defense variation as a potential source introducing diversity

into mimetic systems.

The roles of phenotypic plasticity and ecological sources of defense variation in wild populations

It has been suggested that the often wide variation in defense toxins could be explained by the non-

adaptiveness of toxicity profiles, in which case the variation would stem from chance effects and plastic

responses to the environment (Speed et al., 2012). Empirical data on chemical defense variation in wild

populations e.g. along environmental gradients could help to evaluate ecological vs. evolutionary origins of

chemical defense variation, and the expected consequences for diversity in aposematic and mimetic systems

(Briolat et al., 2019). Although the large amount of unexplained variance in our quantitative genetics model

may suggest the presence of unknown environmental sources of variation in biosynthesized toxicity, we

found very little evidence for a role of known ecological factors in explaining variation in toxicity profiles

along our studied environmental gradients, despite the steepness of these gradients. For example, the

chemical profiles were very similar in the Panamanian Wet and Dry populations, which experience a five-

fold difference in rainfall. Furthermore, despite water availability inducing substantial differences in the

cyanogen content and quality indicators of *P. biflora*, both our field and experimental data indicate that

variation within the host plant species does not contribute to variation in *H. erato* biosynthesized toxicity.

This is in contrast to our expectations of dietary quality as a major source of plastic chemical defense

variation.

However, there were striking differences in the chemical profiles between the two distinct

warning color forms of *H. erato* that we sampled from geographically distant populations in Panama and

Ecuador. Compared to their Panamanian conspecifics sampled along the gradient of rainfall at sea level, the

Ecuadorian populations inhabit a very different and geographically distant biotope along the eastern slope of

the Andes (Montejo-Kovacevich et al., 2020). These environments differ in larval host plant and

nectar/pollen availability, the community of predators and of co-mimics, which are far more diverse in

Ecuador. Lower levels of cyanogen biosynthesis in Ecuadorian populations of *H. erato* may suggest that

these populations have specialized on a *Passiflora* host which provides more cyanogens for sequestration (de

Castro et al., 2019; Engler-Chaouat & Gilbert, 2007; de Castro et al., 2020). Alternatively, the more diverse

mimetic ring in Ecuador could also provide greater protection by more toxic model species, relaxing

selection on defenses in co-mimics (Mallet, 1999; Speed, 1993, 1999; Turner, 1987).

564   In this light, our data may provide an example of how ecological variation could lead to

differential selective environments between conspecific populations, thereby leading to local adaptation of

the acquisition mechanisms and levels of defensive compounds. Such adaptive radiations would be

facilitated by evolutionary potential in defense traits. More data on different aposematic and mimetic systems

will be needed to confirm whether ecological variation may have a greater influence on chemical defense

variation through introducing spatially and temporally varying selection pressures, rather than by introducing

"ecological noise" through plastic responses in defense traits.

CONCLUSION

The role of ecological and evolutionary processes in the origins and maintenance of the wide variation in

chemical defenses, and the consequences for the evolution of aposematism and mimicry holds many

unanswered questions. Our analysis of biosynthesized cyanogenic toxicity in *Heliconius erato,* studied along

natural environmental gradients and in a common garden, shows that this important chemical defense trait

varies substantially within and among populations. We found no evidence for a role of dietary variation

(within host species) or steep environmental gradients in explaining toxicity profiles, despite a large

component of unexplained variance in common-garden reared broods. Instead, the results suggest that

adaptive processes originating from e.g. larval host species availability, mimicry environment, and

individual-level energy optimization may play important roles in the variation of cyanogenic defenses. This

is in line with our data indicating substantial broad-sense evolvability in cyanogenic toxicity, although more

data will be needed to confirm the origins of the observed among-brood variance and shed light on how the

indicated maternal effects will affect the overall selection response. Our study suggests that adaptive

evolution could be an important force driving variation in defensive traits. Such adaptive processes could

play a part in explaining the incredible diversity in aposematic and mimetic systems.

ACKNOWLEDGEMENTS

Personnel and researchers at the Smithsonian Tropical Research Institute (STRI Panama) and Ikiam at

Universidad Regional Amazónica (Ecuador) are acknowledged for help with permits and the field collection

of the study species. We are especially grateful to Marleny Rivera (STRI) for reliable and friendly assistance

in field sampling. Samples were processed at the Molecular Ecology Laboratory at the University of Helsinki

(UH), and analysed at the NMR facility at the Institute of Biotechnology (UH), which is supported by

Biocenter Finland and Helsinki Institute of Life Science (HiLIFE). Søren Bak and David Pattison

(Department for Plant and Environmental Sciences, University of Copenhagen) are acknowledged for

collaboration on the LC-MS analyses of *Passiflora* samples. Juha-Matti Pitkänen, Tuomas Niemi-Aro and

Annukka Ruokolainen are thanked for technical assistance, and Guillaume Minard, Anne Duplouy, Richard

Wallbank, Richard Merrill, Oscar Paneso, Krzysztof Kozak and Sophia Gripenberg for advice on study

design and lab and field work. Mika Zagrobelny and Marcio Cardoso are thanked for advice on cyanogenic

assays and Mika Zagrobelny also for providing reference samples of the CNglcs compounds. Research and

collecting permits were granted by the Ministerio de Ambiente, Republica de Panama (Permit SE/AP-21-16)

and the Ministerio del Ambiente, Ecuador (under the Contrato Marco MAE-DNB-CM-2017-0058). Financial

support was provided by the Academy of Finland (Grant no. 286814 to A.M.), ERC (SpeciationGenetics

339873 to J.C.), NERC Doctoral Training Partnership (NE/L002507/1 to G.M.K.), and Spanish Agency for

International Development Cooperation (AECID, grant number 2018SPE0000400194 to C.B.).

AUTHOR CONTRIBUTIONS

A.M., M.S. and C.J. conceived the study. A.M., M.S. and G.M.K. conducted fieldwork. A.M. conducted

experimental work. A.M. and E.C. conducted laboratory work. A.M. and Ø.O. collated the dataset and

conducted the analyses. W.M. and C.B. provided facilities and materials. A.M., C.J. and W.M. wrote the

manuscript. All authors read and approved the final manuscript.

DATA ACCESSIBILITY

All data generated and analysed during this study will be deposited in a digital data repository (Dryad data

repository) at acceptance for publication.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

iii. REFERENCES

Arias, M., Meichanetzoglou, A., Elias, M., Rosser, N., de-Silva, D. L., Nay, B., & Llaurens, V. (2016). Variation
in cyanogenic compounds concentration within a Heliconius butterfly community: Does mimicry
explain everything? *BMC Evolutionary Biology*, *16*(1), 272. https://doi.org/10.1186/s12862-016-
0843-5

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). *Fitting Linear Mixed-Effects Models using lme4*.
https://arxiv.org/abs/1406.5823v1

Benson, W. W. (1972). Natural Selection for Miillerian Mimicry in Heliconius erato in Costa Rica. *Science*,
176(4037), 936–939. https://doi.org/10.1126/science.176.4037.936

Benson, W. W., Brown, K. S., & Gilbert, L. E. (1975). Coevolution of Plants and Herbivores: Passion Flower
Butterflies. *Evolution*, *29*(4), 659–680. https://doi.org/10.1111/j.1558-5646.1975.tb00861.x

Boggs, C. L., & Freeman, K. D. (2005). Larval food limitation in butterflies: Effects on adult resource
allocation and fitness. *Oecologia*, *144*(3), 353–361. https://doi.org/10.1007/s00442-005-0076-6

Bowers, M. D. (1992). The evolution of unpalatibility and the cost of chemical defense in insects. In *Insect
Chemical Ecology: An Evolutionary Approach* (pp. 216–244). Chapman Hall.

Bowman, A. W., & Azzalini, A. (2014). *R package "sm": Nonparametric smoothing methods (version 2.2-5.5)*.
http://www.stats.gla.ac.uk/~adrian/sm

Boyden, T. C. (1976). Butterfly Palatability and Mimicry: Experiments with Ameiva Lizards. *Evolution*, *30*(1),
73–81. JSTOR. https://doi.org/10.2307/2407673

Briolat, E. S., Burdfield-Steel, E. R., Paul, S. C., Rönkä, K. H., Seymoure, B. M., Stankowich, T., & Stuckert, A.
M. M. (2019). Diversity in warning coloration: Selective paradox or the norm? *Biological Reviews*,
94(2), 388–414. https://doi.org/10.1111/brv.12460

Brower, L. P., Pough, F. H., & Meck, H. R. (1970). Theoretical Investigations of Automimicry, I. Single Trial
Learning. *Proceedings of the National Academy of Sciences*, *66*(4), 1059–1066.
https://doi.org/10.1073/pnas.66.4.1059

Burdfield-Steel, E., Brain, M., Rojas, B., & Mappes, J. (2019). The price of safety: Food deprivation in early
life influences the efficacy of chemical defence in an aposematic moth. *Oikos*, *128*(2), 245–253.
https://doi.org/10.1111/oik.05420

Burdfield-Steel, E., Pakkanen, H., Rojas, B., Galarza, J. A., & Mappes, J. (2018). De novo Synthesis of
Chemical Defenses in an Aposematic Moth. *Journal of Insect Science*, *18*(2).
https://doi.org/10.1093/jisesa/iey020

Butler, D. G., Cullis, B. R., Gilmour, A. R., Gogel, B. G., & Thompson, R. (2017). *ASReml-R Reference Manual
Version 4*. VSN International Ltd, Hemel Hempstead, HP1 1ES, UK.

Camara, M. D. (1997). Predator Responses to Sequestered Plant Toxins in Buckeye Caterpillars: Are
Tritrophic Interactions Locally Variable? *Journal of Chemical Ecology*, *23*(9), 2093–2106.
https://doi.org/10.1023/B:JOEC.0000006431.34359.c2
Cardoso, M. Z., & Gilbert, L. E. (2013). Pollen feeding, resource allocation and the evolution of chemical
defence in passion vine butterflies. *Journal of Evolutionary Biology*, *26*(6), 1254–1260.
https://doi.org/10.1111/jeb.12119
Cardoso, Márcio Zikán. (2020). The effect of cyanogenic glucosides and their breakdown products on
predation by domestic chicks. *Chemoecology*, *30*(3), 131–138. https://doi.org/10.1007/s00049-020-
00304-6
Cardoso, Márcio Zikán, & Gilbert, L. E. (2007). A male gift to its partner? Cyanogenic glycosides in the
spermatophore of longwing butterflies (Heliconius). *Naturwissenschaften*, *94*(1), 39–42.
https://doi.org/10.1007/s00114-006-0154-6
Cerovic, Z. G., Masdoumier, G., Ghozlen, N. B., & Latouche, G. (2012). A new optical leaf-clip meter for
simultaneous non-destructive assessment of leaf chlorophyll and epidermal flavonoids. *Physiologia
Plantarum*, *146*(3), 251–260. https://doi.org/10.1111/j.1399-3054.2012.01639.x
Chouteau, M., Dezeure, J., Sherratt, T. N., Llaurens, V., & Joron, M. (2019). Similar predator aversion for
natural prey with diverse toxicity levels. *Animal Behaviour*, *153*, 49–59.
https://doi.org/10.1016/j.anbehav.2019.04.017
de Castro, É. C. P., Demirtas, R., Orteu, A., Olsen, C. E., Motawie, M. S., Zikan Cardoso, M., Zagrobelny, M.,
& Bak, S. (2020). The dynamics of cyanide defences in the life cycle of an aposematic butterfly:
Biosynthesis versus sequestration. *Insect Biochemistry and Molecular Biology*, *116*, 103259.
https://doi.org/10.1016/j.ibmb.2019.103259
de Castro, É. C. P., Zagrobelny, M., Cardoso, M. Z., & Bak, S. (2018). The arms race between heliconiine
butterflies and Passiflora plants – new insights on an ancient subject. *Biological Reviews*, *93*(1),
555–573. https://doi.org/10.1111/brv.12357
de Castro, É. C. P., Zagrobelny, M., Zurano, J. P., Cardoso, M. Z., Feyereisen, R., & Bak, S. (2019).
Sequestration and biosynthesis of cyanogenic glucosides in passion vine butterflies and
consequences for the diversification of their host plants. *Ecology and Evolution*, *9*(9), 5079–5093.
https://doi.org/10.1002/ece3.5062
Eggenberger, F., & Rowell-Rahier, M. (1992). Genetic component of variation in chemical defense ofOreina
gloriosa (Coleoptera: Chrysomelidae). *Journal of Chemical Ecology*, *18*(8), 1375–1387.
https://doi.org/10.1007/BF00994363
Engler, H. S., Spencer, K. C., & Gilbert, L. E. (2000). Preventing cyanide release from leaves. *Nature*,
406(6792), 144. https://doi.org/10.1038/35018159
Engler-Chaouat, H. S., & Gilbert, L. E. (2007). De novo Synthesis vs. Sequestration: Negatively Correlated
Metabolic Traits and the Evolution of Host Plant Specialization in Cyanogenic Butterflies. *Journal of
Chemical Ecology*, *33*(1), 25–42. https://doi.org/10.1007/s10886-006-9207-8
Fordyce, J. A., & Nice, C. C. (2008). Antagonistic, Stage-Specific Selection on Defensive Chemical
Sequestration in a Toxic Butterfly. *Evolution*, *62*(7), 1610–1617. https://doi.org/10.1111/j.1558-
5646.2008.00388.x
Gardner, D. R., & Stermitz, F. R. (1988). Host plant utilization and iridoid glycoside sequestration
byEuphydryas anicia (Lepidoptera: Nymphalidae). *Journal of Chemical Ecology*, *14*(12), 2147–2168.
https://doi.org/10.1007/BF01014022
Gibson, D. O. (1984). How is Automimicry Maintained? In *The Biology of Butterflies* (pp. 163–165).
Academic Press.
Gleadow, R. M., & Woodrow, I. E. (2002). Defense chemistry of cyanogenic Eucalyptus cladocalyx seedlings
is affected by water supply. *Tree Physiology*, *22*(13), 939–945.
https://doi.org/10.1093/treephys/22.13.939
Halpin, C. G., Skelhorn, J., Rowe, C., Ruxton, G. D., & Higginson, A. D. (2017). The Impact of Detoxification
Costs and Predation Risk on Foraging: Implications for Mimicry Dynamics. *PLoS ONE*, *12*(1).
https://doi.org/10.1371/journal.pone.0169043

Hansen, T. F., & Houle, D. (2008). Measuring and comparing evolvability and constraint in multivariate characters. *Journal of Evolutionary Biology*, *21*(5), 1201–1219. https://doi.org/10.1111/j.1420-9101.2008.01573.x

Hansen, Thomas F., Pélabon, C., & Houle, D. (2011). Heritability is not Evolvability. *Evolutionary Biology*, *38*(3), 258. https://doi.org/10.1007/s11692-011-9127-6

Hayden*, K. J., & Parker, I. M. (2002). Plasticity in cyanogenesis of Trifolium repens L.: Inducibility, fitness costs and variable expression. *Evolutionary Ecology Research*, *4*(2), 155–168.

Hay-Roe, M. M., & Nation, J. (2007). Spectrum of Cyanide Toxicity and Allocation in Heliconius erato and Passiflora Host Plants. *Journal of Chemical Ecology*, *33*(2), 319–329. https://doi.org/10.1007/s10886-006-9234-5

Hereford, J., Hansen, T. F., & Houle, D. (2004). Comparing strengths of directional selection: How strong is strong? *Evolution; International Journal of Organic Evolution*, *58*(10), 2133–2143. https://doi.org/10.1111/j.0014-3820.2004.tb01592.x

Holloway, G. J., de Jong, P. W., & Ottenheim, M. (1993). The Genetics and Cost of Chemical Defense in the Two-Spot Ladybird (Adalia bipunctata L.). *Evolution*, *47*(4), 1229–1239. JSTOR. https://doi.org/10.2307/2409988

Jiggins, C. D. (2017). *The Ecology and Evolution of Heliconius Butterflies* (1 edition). Oxford University Press.

Jiggins, C. D., Naisbit, R. E., Coe, R. L., & Mallet, J. (2001). Reproductive isolation caused by colour pattern mimicry. *Nature*, *411*(6835), 302–305. https://doi.org/10.1038/35077075

Jones, R. S., Speed, M. P., & Mappes, J. (2016). Parameterising a public good: How experiments on predation can be used to predict cheat frequencies. *Evolutionary Ecology*, *30*(5), 825–840. https://doi.org/10.1007/s10682-016-9851-6

Joron, M., & Mallet, J. L. (1998). Diversity in mimicry: Paradox or paradigm? *Trends in Ecology & Evolution*, *13*(11), 461–466. https://doi.org/10.1016/S0169-5347(98)01483-9

Kapan, D. D. (2001). Three-butterfly system provides a field test of müllerian mimicry. *Nature*, *409*(6818), 338–340. https://doi.org/10.1038/35053066

Kruuk, L. E. B. (2004). Estimating genetic parameters in natural populations using the 'animal model.' *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, *359*(1446), 873–890. https://doi.org/10.1098/rstb.2003.1437

Lande, R., & Price, T. (1989). Genetic correlations and maternal effect coefficients obtained from offspring-parent regression. *Genetics*, *122*(4), 915–922.

Liang, W.-S. (2003). Drought stress increases both cyanogenesis and β-cyanoalanine synthase activity in tobacco. *Plant Science*, *165*(5), 1109–1115. https://doi.org/10.1016/S0168-9452(03)00306-6

Lindstedt, C., Suisto, K., Burdfield-Steel, E., Winters, A. E., & Mappes, J. (2020). Defense against predators incurs high reproductive costs for the aposematic moth Arctia plantaginis. *Behavioral Ecology*. https://doi.org/10.1093/beheco/araa033

Lindstedt, C., Talsma, J. H. R., Ihalainen, E., Lindström, L., & Mappes, J. (2010). Diet Quality Affects Warning Coloration Indirectly: Excretion Costs in a Generalist Herbivore. *Evolution*, *64*(1), 68–78. https://doi.org/10.1111/j.1558-5646.2009.00796.x

Lynch, M., & Walsh, B. (1998a). Maternal Effects. In *Genetics and Analysis of Quantitative Traits* (pp. 687–714). Oxford University Press.

Lynch, M., & Walsh, B. (1998b). Resemblance Between Relatives. In *Genetics and Analysis of Quantitative Traits*. Oxford University Press.

Mallet, J. (1999). Causes and Consequences of a Lack of Coevolution in Müllerian mimicry. *Evolutionary Ecology*, *13*(7), 777–806. https://doi.org/10.1023/A:1011060330515

Mallet, J., & Barton, N. H. (1989). Strong Natural Selection in a Warning-Color Hybrid Zone. *Evolution*, *43*(2), 421–431. https://doi.org/10.1111/j.1558-5646.1989.tb04237.x

Mallet, J., & Joron, M. (1999). Evolution of Diversity in Warning Color and Mimicry: Polymorphisms, Shifting Balance, and Speciation. *Annual Review of Ecology and Systematics*, *30*(1), 201–233. https://doi.org/10.1146/annurev.ecolsys.30.1.201

Mallet, J., McMillan, W. O., & Jiggins, C. D. (1998). Mimicry and warning color at the boundary between microevolution and macroevolution. In *Endless Forms: Species and Speciation*. Oxford University Press.

757   Massad, T. J., Fincher, R. M., Smilanich, A. M., & Dyer, L. (2011). A quantitative evaluation of major plant
758   defense hypotheses, nature versus nurture, and chemistry versus ants. *Arthropod-Plant*
759   *Interactions*, *5*(2), 125–139. https://doi.org/10.1007/s11829-011-9121-z

760   Merrill, R. M., Dasmahapatra, K. K., Davey, J. W., Dell'Aglio, D. D., Hanly, J. J., Huber, B., Jiggins, C. D., Joron,
761   M., Kozak, K. M., Llaurens, V., Martin, S. H., Montgomery, S. H., Morris, J., Nadeau, N. J.,
762   Pinharanda, A. L., Rosser, N., Thompson, M. J., Vanjari, S., Wallbank, R. W. R., & Yu, Q. (2015). The
763   diversification of Heliconius butterflies: What have we learned in 150 years? *Journal of Evolutionary*
764   *Biology*, *28*(8), 1417–1438. https://doi.org/10.1111/jeb.12672

765   Merrill, R. M., Naisbit, R. E., Mallet, J., & Jiggins, C. D. (2013). Ecological and genetic factors influencing the
766   transition between host-use strategies in sympatric Heliconius butterflies. *Journal of Evolutionary*
767   *Biology*, *26*(9), 1959–1967. https://doi.org/10.1111/jeb.12194

768   Montejo-Kovacevich, G., Martin, S. H., Meier, J. I., Bacquet, C. N., Monllor, M., Jiggins, C. D., & Nadeau, N. J.
769   (2020). Microclimate buffering and thermal tolerance across elevations in a tropical butterfly.
770   *Journal of Experimental Biology*, *223*(8). https://doi.org/10.1242/jeb.220426

771   Müller, C., Zwaan, B. J., de Vos, H., & Brakefield, P. M. (2003). Chemical defence in a sawfly: Genetic
772   components of variation in relevant life-history traits. *Heredity*, *90*(6), 468–475.
773   https://doi.org/10.1038/sj.hdy.6800265

774   Nahrstedt, A., & Davis, R. H. (1983). Occurrence, variation and biosynthesis of the cyanogenic glucosides
775   linamarin and lotaustralin in species of the Heliconiini (Insecta: Lepidoptera). *Comparative*
776   *Biochemistry and Physiology Part B: Comparative Biochemistry*, *75*(1), 65–73.
777   https://doi.org/10.1016/0305-0491(83)90041-X

778   Nahrstedt, A., & Davis, R. H. (1985). Biosynthesis and quantitative relationships of the cyanogenic
779   glucosides, linamarin and lotaustralin, in genera of the Heliconiini (Insecta: Lepidoptera).
780   *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry*, *82*(4), 745–749.
781   https://doi.org/10.1016/0305-0491(85)90519-X

782   Nishida, R. (2002). Sequestration of Defensive Substances from Plants by Lepidoptera. *Annual Review of*
783   *Entomology*, *47*(1), 57–92. https://doi.org/10.1146/annurev.ento.47.091201.145121

784   Opitz, S. E. W., & Müller, C. (2009). Plant chemistry and insect sequestration. *Chemoecology*, *19*(3), 117.
785   https://doi.org/10.1007/s00049-009-0018-6

786   Pinheiro, C. E. G. (1996). Palatablility and escaping ability in Neotropical butterflies: Tests with wild
787   kingbirds (Tyrannus melancholicus, Tyrannidae). *Biological Journal of the Linnean Society*, *59*(4),
788   351–365. https://doi.org/10.1111/j.1095-8312.1996.tb01471.x

789   Pinheiro, C. E. G., & Campos, V. C. (2019). The responses of wild jacamars (Galbula ruficauda, Galbulidae) to
790   aposematic, aposematic and cryptic, and cryptic butterflies in central Brazil. *Ecological Entomology*,
791   *44*(4), 441–450. https://doi.org/10.1111/een.12723

792   Prudic, K. L., Timmermann, B. N., Papaj, D. R., Ritland, D. B., & Oliver, J. C. (2019). Mimicry in viceroy
793   butterflies is dependent on abundance of the model queen butterfly. *Communications Biology*,
794   *2*(1), 68. https://doi.org/10.1038/s42003-019-0303-z

795   R Core Team. (2018). *R: A language and environment for statistical computing. R Foundation for Statistical*
796   *Computing*. R Foundation for Statistical Computing, Vienna,   Austria. https://www.R-project.org/

797   Reudler, J. H., Lindstedt, C., Pakkanen, H., Lehtinen, I., & Mappes, J. (2015). Costs and benefits of plant
798   allelochemicals in herbivore diet in a multi enemy world. *Oecologia*, *179*(4), 1147–1158.
799   https://doi.org/10.1007/s00442-015-3425-0

800   Ritland, D. B., & Brower, L. P. (1991). The viceroy butterfly is not a batesian mimic. *Nature*, *350*(6318), 497.
801   https://doi.org/10.1038/350497a0

802   Ruxton, G. D., & Speed, M. P. (2006). How can automimicry persist when predators can preferentially
803   consume undefended mimics? *Proceedings of the Royal Society B: Biological Sciences*, *273*(1584),
804   373–378. https://doi.org/10.1098/rspb.2005.3238

805   Ruxton, G. D., William Allen L., Sherratt, T. N., & Speed, M. P. (2018). *Avoiding Attack: The Evolutionary*
806   *Ecology of Crypsis, Aposematism, and Mimicry* (Second Edition). Oxford University Press.

807   Sculfort, O., Castro, E. C. P. de, Kozak, K. M., Bak, S., Elias, M., Nay, B., & Llaurens, V. (2020). Variation of
808   chemical compounds in wild Heliconiini reveals ecological factors involved in the evolution of

chemical defenses in mimetic butterflies. *Ecology and Evolution*.
https://doi.org/10.1002/ece3.6044
Smilanich, A. M., Dyer, L. A., Chambers, J. Q., & Bowers, M. D. (2009). Immunological cost of chemical
defence and the evolution of herbivore diet breadth. *Ecology Letters*, *12*(7), 612–621.
https://doi.org/10.1111/j.1461-0248.2009.01309.x
Speed, M. P. (1993). Muellerian mimicry and the psychology of predation. *Animal Behaviour*, *45*(3), 571–
580. https://doi.org/10.1006/anbe.1993.1067
Speed, M. P. (1999). Batesian, quasi-Batesian or Müllerian mimicry? Theory and data in mimicry Research.
Evolutionary Ecology, *13*(7), 755–776. https://doi.org/10.1023/A:1010871106763
Speed, M. P., Ruxton, G. D., Mappes, J., & Sherratt, T. N. (2012). Why are defensive toxins so variable? An
evolutionary perspective. *Biological Reviews*, *87*(4), 874–884. https://doi.org/10.1111/j.1469-
185X.2012.00228.x
Stoffel, M. A., Nakagawa, S., & Schielzeth, H. (2017). rptR: Repeatability estimation and variance
decomposition by generalized linear mixed-effects models. *Methods in Ecology and Evolution*,
8(11), 1639–1644. https://doi.org/10.1111/2041-210X.12797
Svennungsen, T. O., & Holen, Ø. H. (2007). The evolutionary stability of automimicry. *Proceedings of the
Royal Society B: Biological Sciences*, *274*(1621), 2055–2063.
https://doi.org/10.1098/rspb.2007.0456
Turner, J. R. G. (1984). Mimicry: The palatability spectrum and its consequences. In *The Biology of
Butterflies* (pp. 141–161). Academic Press.
Turner, J. R. G. (1987). The evolutionary dynamics of batesian and muellerian mimicry: Similarities and
differences. *Ecological Entomology*, *12*(1), 81–95. https://doi.org/10.1111/j.1365-
2311.1987.tb00987.x
Van Belleghem, S. M., Rastas, P., Papanicolaou, A., Martin, S. H., Arias, C. F., Supple, M. A., Hanly, J. J.,
Mallet, J., Lewis, J. J., Hines, H. M., Ruiz, M., Salazar, C., Linares, M., Moreira, G. R. P., Jiggins, C. D.,
Counterman, B. A., McMillan, W. O., & Papa, R. (2017). Complex modular architecture around a
simple toolkit of wing pattern genes. *Nature Ecology & Evolution*, *1*(3), 1–12.
https://doi.org/10.1038/s41559-016-0052
Vandegeer, R., Miller, R. E., Bain, M., Gleadow, R. M., & Cavagnaro, T. R. (2013). Drought adversely affects
tuber development and nutritional quality of the staple crop cassava (Manihot esculenta Crantz).
Functional Plant Biology, *40*(2), 195–200. https://doi.org/10.1071/FP12179
Villemereuil, P. de, Morrissey, M. B., Nakagawa, S., & Schielzeth, H. (2018). Fixed-effect variance and the
estimation of repeatabilities and heritabilities: Issues and solutions. *Journal of Evolutionary Biology*,
31(4), 621–632. https://doi.org/10.1111/jeb.13232
Wallbank, R. W. R., Baxter, S. W., Pardo-Diaz, C., Hanly, J. J., Martin, S. H., Mallet, J., Dasmahapatra, K. K.,
Salazar, C., Joron, M., Nadeau, N., McMillan, W. O., & Jiggins, C. D. (2016). Evolutionary Novelty in a
Butterfly Wing Pattern through Enhancer Shuffling. *PLOS Biology*, *14*(1), e1002353.
https://doi.org/10.1371/journal.pbio.1002353
Walsh, B., & Lynch, M. (2018a). Selection in the presence of heritable maternal effects. In *Evolution and
Selection of Quantitative Traits* (pp. 537–547).
Walsh, B., & Lynch, M. (2018b). *Evolution and Selection of Quantitative Traits*. Oxford University Press.
Wilson, A. J., Réale, D., Clements, M. N., Morrissey, M. M., Postma, E., Walling, C. A., Kruuk, L. E. B., &
Nussey, D. H. (2010). An ecologist's guide to the animal model. *Journal of Animal Ecology*, *79*(1),
13–26. https://doi.org/10.1111/j.1365-2656.2009.01639.x
Winters, A. E., Wilson, N. G., van den Berg, C. P., How, M. J., Endler, J. A., Marshall, N. J., White, A. M.,
Garson, M. J., & Cheney, K. L. (2018). Toxicity and taste: Unequal chemical defences in a mimicry
ring. *Proceedings of the Royal Society B: Biological Sciences*, *285*(1880), 20180457.
https://doi.org/10.1098/rspb.2018.0457
Yezerski, A., Gilmor, T. P., & Stevens, L. (2004). Genetic Analysis of Benzoquinone Production in Tribolium
confusum. *Journal of Chemical Ecology*, *30*(5), 1035–1044.
https://doi.org/10.1023/B:JOEC.0000028465.37658.ae

Zvereva, E. L., & Kozlov, M. V. (2016). The costs and effectiveness of chemical defenses in herbivorous
insects: A meta-analysis. *Ecological Monographs*, *86*(1), 107–124. https://doi.org/10.1890/15-
0911.1
863
864

SUPPLEMENTARY FILES

Supplementary File 1. A. Full results table of between-population variation of cyanogenic glucoside

compounds in natural populations of *Heliconius erato* and *Passiflora biflora*. **B.** Full results table of REML

animal models.

Supplementary File 2. Summary table of *Heliconius erato* broods.

Supplementary File 3. Supplementary results: Cyanogenic toxicity and overall quality of host plant *Passiflora*

biflora watering treatment groups.

Supplementary File 4. Supplementary Methods. Detailed information about methods used in the study.