TITLE

Distinct components of photoperiodic light are differentially encoded by the mammalian

circadian clock

AUTHORS

Michael C. Tackenberg[1], Jacob J. Hughey[2,3], and Douglas G. McMahon[1,2*]

[1]Vanderbilt Brain Institute, Vanderbilt University, Nashville, TN, USA

[2]Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA

[3]Department of Biomedical Informatics, Vanderbilt University School of Medicine, Nashville, TN,

USA

Correspondence: douglas.g.mcmahon@vanderbilt.edu

RUNNING TITLE

Components of photoperiodic light

---

[*] Lead Contact

ABSTRACT

Seasonal light cycles influence multiple physiological functions and are mediated

through photoperiodic encoding by the circadian system. Despite our knowledge of the strong

connection between seasonal light input and downstream circadian changes, less is known

about the specific components of seasonal light cycles that are encoded and induce persistent

changes in the circadian system. Using combinations of three T cycles (23, 24, 26 hr.) and two

photoperiods per T cycle (Long and Short, with duty cycles scaled to each T cycle), we

investigate after-effects of entrainment to these six light cycles on locomotor behavior duration

(α), period (τ), and entrained phase angle (ψ) *in vivo,* and SCN phase distribution ($\sigma_\square$), τ, and ψ

ex vivo in order to refine our understanding of critical light components for influencing particular

circadian properties. We find that photoperiod and T cycle length both drive determination of *in*

vivo ψ but differentially influence after-effects in α and τ, with photoperiod driving changes in α

and photoperiod length and T cycle length combining to influence τ. Using skeleton

photoperiods, we demonstrate that *in vivo* ψ is determined by both parametric and non-

parametric components, while changes in α are driven non-parametrically. Within the *ex vivo*

SCN, we find that ψ and $\sigma_\square$ of the PER2::LUCIFERASE rhythm follow closely with their likely

behavioral counterparts (ψ and α of the locomotor activity rhythm), while also confirming

previous reports of τ after-effects of gene expression rhythms showing negative correlations

with behavioral τ after-effects in response to T cycles. We demonstrate that within-SCN $\sigma_\square$

changes, thought to underly α changes *in vivo*, are induced primarily non-parametrically. Taken

together, our results demonstrate distinct components of seasonal light input differentially

influence ψ, α, and τ, and suggest the possibility of separate mechanisms driving the persistent

changes in circadian behaviors mediated by seasonal light.

KEYWORDS

INTRODUCTION

The duration of daylight, or photoperiod, represents a predictable and dynamic signal of seasonal change throughout the year across much of the planet. The predictability of this signal is harnessed by organisms to initiate or avoid a variety of biological functions at specific times of year, including reproductive behaviors (Elliott and Goldman, 1981), resource conservation (Bartness et al., 1989; Bartness and Wade, 1984), and migration (Gwinner, 1990). In humans, a variety of disorders and non-communicable diseases have seasonal correlations (Basnet et al., 2016; Oh et al., 2010; Wehr, 2001), with perhaps the most striking seasonal influence being on depression-anxiety behaviors (Koorengevel et al., 2003). Despite clear associations between photoperiodic exposure and physiological function and dysfunction, the roles of specific components of seasonal light cycles that induce those changes are incompletely understood.

One factor critical to our current understanding of the mechanism of photoperiodic induction is the persistence of seasonal changes. Physiological hallmarks of exposure to particular light schedules can be observed, for varying intervals, beyond direct exposure to that light schedule (Pittendrigh and Daan, 1976a). These persistent changes, referred to as after-effects, include characteristic changes to core circadian locomotor behavior patterns in the form of activity duration (alpha, α), free-running period (tau, τ), and phase angle of entrainment (psi, ψ). In nocturnal animals, prior exposure to long photoperiods results in a compressed α, a shortened τ, and an advanced ψ relative to short photoperiod counterparts (Pittendrigh and Daan, 1976a).

The existence of after-effects suggests that seasonal light cycle information is encoded within the brain and can persist beyond the interval of direct exposure. The master circadian pacemaker of the brain, the suprachiasmatic nucleus (SCN), likely plays a major role in this encoding, particularly with regards to after-effects in circadian behavior (for review, see

Tackenberg and McMahon, 2018). In addition, recent work has identified a distinct non-SCN

target for photoperiodic influence, the peri-habenular area, influencing seasonal response in

mood (Fernandez et al., 2018). Like circadian locomotor behavior patterning, the SCN

undergoes substantial changes in organization following exposure to different photoperiods,

constituting network-level after-effects. These responses include changes in phase distribution

($\sigma_\square$) of its constituent neurons as well as in τ and ψ of SCN gene expression rhythms. Prior

exposure to long photoperiod has been shown to increase $\sigma_\square$ of *Period1::Luciferase* (Inagaki et

al., 2007), PERIOD2::LUCIFERASE (Buijink et al., 2016), and electrical (VanderLeest et al.,

2007) rhythms compared to short photoperiod counterparts. As in locomotor behavior after-

effects, some reports have shown that prior exposure to long photoperiod shortens the SCN τ

compared to short photoperiod counterparts (16:8 L:D, Ciarleglio et al., 2011; 20:4 L:D, Evans

et al., 2013) while others have found non-significant decreases in SCN τ length following

exposure to long photoperiod, particularly in the anterior SCN (Buijink et al., 2016; Mickman et

al., 2008).

Induction of persistent circadian after-effects is not limited to changes in photoperiod.

Manipulation of total day-night cycle (T cycle) length also induces after-effects in circadian

locomotor behavior, with entrainment to short T cycles (T < 24 hours) resulting in contracted α

(Azzi et al., 2014) and shortened τ upon release into constant conditions (Azzi et al., 2014;

Schwartz et al., 2011). However, unlike photoperiodic after-effects, in which the changes in the τ

of the explanted SCN itself match those on behavior, there is a negative correlation between

behavioral and SCN explant τ following entrainment to T cycles. Exposure to short T cycles

results in a long SCN τ and exposure to long T cycles results in a short SCN τ (Aton et al.,

2004; Azzi et al., 2017; Molyneux et al., 2008).

The findings described above reveal a potential pattern of circadian after-effects, with

long photoperiods and short T cycles producing one set of behavioral responses (short α, short

τ), and short photoperiods and long T cycles producing another (long α, long τ). Because recent

work has revealed a role for DNA methylation in establishing τ after-effects of extended T cycle

entrainment, the potential association between τ and α after-effects provides a promising

biochemical mechanism for both changes (Azzi et al., 2014). The linkage between the two types

of after-effects, however, has not yet been sufficiently interrogated. The discordance between

the SCN after-effects of T cycle and photoperiod offers a hint that the relationship may not be so

simple. Here, we seek to further examine the alignment of T cycle and photoperiodic after-

effects by measuring *in vivo* (α, τ, and ψ of locomotor behavior rhythms) and *ex vivo* ($\sigma_\square$, τ, and

ψ of SCN gene expression rhythms) responses following exposure to six combinations of

photoperiod length and T cycle. We find that after-effects in each of these rhythm characteristics

respond to the six different light cycles in distinct patterns, suggesting that they are each

induced by different aspects of the input light cycle. Using skeleton photoperiods, we

demonstrate that parametricity differs between after-effects in α, τ and ψ, further indicating

encoding of distinct components of light input by each of these characteristics. Examining the

rhythms of explanted SCN *ex vivo* following entrainment to these light schedules, we find that

the pattern of SCN $\sigma_\square$ reflects that of α *in vivo*, providing additional correlative evidence that

SCN $\sigma_\square$ may be a determining factor in setting α.

106   **MATERIALS AND METHODS**

Animals

Animal experiments were conducted in accordance with Vanderbilt University Institutional

Animal Care and Use Committee regulations. All animals used for behavioral experiments were

heterozygous for the *Per2::Luciferase* allele (*Per2::Luciferase$^{+/-}$*) or were *Per2* wild-type

(*Per2::Luciferase$^{-/-}$*). All animals used for SCN slice culture experiments were heterozygous for

the *Per2::Luciferase* allele (*Per2::Luciferase$^{+/-}$*).

Activity Monitoring and Housing

Animals were singly-housed in light-tight boxes with activity monitoring by wheel revolutions

detected by ClockLab acquisition software (Actimetrics, Inc.). Complete photoperiods used were

15:8 and 16:7 (T23 Long, ~67% duty cycle), 8:15 and 7:16 (T23 Short, ~33% duty cycle), 16:8

(T24 Long, ~67% cycle), 8:16 (T24 Short, ~33% cycle), 17:9 (T26 Long, ~67% cycle), and 9:17

(T26 Short, ~33 cycle).

Animals were transferred to the specified light schedule after at least 3 weeks of age, and

housed there for at least 28 days before transfer to DD or brain extraction for slicing. Six T26

Short animals included in the datasets (three used in behavioral experiments, three used in slice

experiments) experienced a multi-hour light failure during the 28 days of entrainment, but

received 14 days of the correct light cycle following the failure.

Skeleton photoperiods were set up by housing animals in 12:12 LD complete photoperiods for

approximately 5 days before transitioning to a 12:12 skeleton (1:10:1:12 L:D:L:D). After several

cycles on this skeleton, the onset pulse was gradually advanced (0.5 hour – 1 hour per day) or

delayed (1 hour per day) until a skeleton long (1:14:1:8 L:D:L:D) or skeleton short (1:6:1:16

L:D:L:D) photoperiod, respectively, was established. Once the final skeleton was established,

animals were given 28 days of exposure to that skeleton until transfer to DD or slicing.

Actogram Analysis

Locomotor behavior was analyzed using ClockLab Analysis software and R. Period was

determined using the Chi Square periodogram over the first 7 days of DD. α was measured

using automated onset/offset detection for each cycle of the actogram. Full actograms can be

found in Supplemental Data 1.

Automated α Measurement

Automated α measurement was performed in R. The activity (bin size 6 minutes) is smoothed

by a Savitzky-Golay filter with a span of 25 and degree of 3. A threshold is set relative to the

maximum smoothed activity (0.01 * the maximum for each day). Points at which the smoothed

activity crosses this threshold are identified as potential onsets and offsets. Segments of activity

less than 2 hours are considered inactive. The longest inactive segment is identified, with the

start of that segment recorded as the offset and the end of that segment recorded as the onset.

α is calculated as activity offset – onset, and ψ as the timing of the activity onset in the first cycle

in DD relative to the projected time of lights off from the previous light dark cycle (projected

Zeitgeber Time 12 [ZT12]). Identified onsets and offsets for each cycle can be found in

Supplemental Data 2. R scripts used for analysis can be found at

10.6084/m9.figshare.11513892.

Slicing and Slice Cultures

Within 4 hours of lights off, animals were sacrificed by cervical dislocation and the brain

removed. SCN slices of 200-µm were made, and the SCN further isolated by scalpel cut under a

dissecting microscope. Slices were transferred to a 6-well plate with cell culture membrane

insert (Millipore, PICMORG50) and 1.2-mL of slice culture media. Slice chambers were sealed

with a coverslip and vacuum grease and placed into an incubated light-tight inverted microscope

(Zeiss Axioskop). Luminescence was detected using an intensified CCD (Stanford Photonics)

controlled by µ-Manager recorded for 2' every 10' for at least 1 week.

Ex-Vivo **Analyses**

Luminescence recordings were analyzed using Fiji and R. OME-TIFF files from μ-Manager were

opened in Fiji and smoothed using a two-frame minimization (frame rate reduction from 6/hour

to 3/hour). A 102x102 grid of 10x10-pixel (5.175 x 5.175 μm) ROIs was overlaid on the images

and brightness measured for each of these 10,404 ROIs measured for every frame. These grid

measurements were then exported to R for further analysis. The first 8 hours of recording were

excluded from peak-finding to prevent slicing artifacts from interfering with measurements. First,

the ROIs comprising the SCN were identified by using pixels within the top 50% of brightness.

The timing of each peak of luminescence was then detected for each ROI and used to calculate

the initial phase. Distribution measurements were made by median absolute deviation across all

SCN ROIs. Phase maps of each SCN used can be found in Supplemental Data 3. R scripts

used for analysis can be found at 10.6084/m9.figshare.11513892.

Statistical Analysis

For measurements of α, τ, $\sigma_\square$, and ψ, the effects of photoperiod and T cycle (or completeness

and onset/offset interval) were analyzed by two-way ANOVA. For each comparison, the *p* value

and percent of total variation of each factor, as well as their interaction, are reported. Where

indicated in the text, Sidak's multiple comparisons test was used to further examine within- and

between-factor effects. For correlation analysis shown in Figure 3, Pearson's r test was used.

RESULTS

Distinct response patterns of ψ, and after-effects on α and τ

To examine the persistent encoding of photoperiods and T cycles *in vivo*, as represented

by entrained phase angle (ψ) and after-effects in activity duration (α) and free-running period (τ),

we exposed mice to six different light schedules consisting of combinations of short or long

photoperiod and short, standard, and long T cycles (T23 Long, T23 Short, T24 Long, T24 Short,

T26 Long, T26 Short). Following 14-28 days of exposure to these light schedules (see

Methods), animals were transferred into constant darkness (DD) for 7 days. Based on previous

studies examining the after-effects of T cycles (Azzi et al., 2014, 2017; Pittendrigh and Daan,

1976a; Schwartz et al., 2011) and photoperiod (Buijink et al., 2016; Ciarleglio et al., 2009;

Evans et al., 2013; Pittendrigh and Daan, 1976a), we hypothesized that the response of τ and α

after-effects to these six light schedules would produce one of three general patterns (Figure 1).

First, if after-effects in these rhythm characteristics were strictly dependent on the duty

cycle of the light interval, then the groups would segregate into two levels, corresponding to long

or short photoperiods, regardless of T cycle (Figure 1A). A primary influence of the photoperiod

could indicate a parametric effect of the total light duration per cycle and/or a non-parametric

effect from the phase shifts caused by light onset and light offset. Second, if plasticity is strictly

dependent on net daily phase advances or delays driven by entrainment to the T cycles, then

the measurements would segregate into three groups based on T cycle (Figure 1B). This

pattern of results would most readily suggest that after-effects were induced by non-parametric

effects of the daily entraining phase shifts induced by the T cycles. Third, significant main

effects of both T cycle and photoperiod would result in after-effects with each of the 6 groups

having a characteristic level (Figure 1C), while interactions between T cycles and photoperiod

would produce many possible variations of the pattern in Figure 1C.

Indeed, ψ of locomotor rhythms relative to projected ZT12 of the previous light:dark cycle

(see Methods) was strongly influenced by both T cycle length and photoperiod length, with a

significant interaction between these two main effects (Figure 2B). As T cycle length increases,

ψ becomes more advanced. Within each T cycle, the short photoperiod version produces a

more delayed ψ than its long photoperiod counterpart, but the difference between the two

photoperiods decreases as T cycle length increases. This pattern resembled the example in

Figure 1C, with the interaction providing an additional layer of complexity. In contrast to the

multifaceted inputs to ψ, we found that photoperiod was the sole significant influence on α, with

persistently reduced α in DD following long photoperiods compared to short photoperiods

across all three T cycles (Figure 2C). This pattern most closely resembled the example in Figure

1A, and suggested that a parametric effect of the duration of light per cycle and/or a non-

parametric effect of the phase shifts caused at light onset and offset induced the after-effect in

α. After-effects on locomotor τ were significantly influenced by both T cycle length and

photoperiod (Figure 2D), but unlike ψ, there was no detectable interaction between the main

effects. Within each T cycle, the long photoperiod group had consistently shorter τ compared to

its partner short photoperiod. Across T cycles, however, there was a direct relationship with τ,

with τ increasing along with T cycle length regardless of photoperiod. This pattern most closely

resembles the example in Figure 1C.

To further examine the relationship between the α and τ after-effects, we plotted the two

measurements against one another (Figure 3). The two factors correlated significantly in two

specific cases: T23 Long (Figure 3, top left) and T26 Short (Figure 3, bottom right). These two

photoperiod/T cycle combinations represent the instances where in previous reports T cycle and

photoperiod independently produce similar *in vivo* after-effects on α and τ, with long photoperiod

or short T decreasing α and τ, and short photoperiod or long T increasing α and τ. Interestingly,

there was a lack of correlation of these measures in T cycle/photoperiod combinations where

the predicted after-effects of the cycle τ and photoperiod would be in nominal conflict (Long

photoperiod and Long T, and Short photoperiod and Short T, Azzi et al., 2014; Pittendrigh and

Daan, 1976a).

ψ and α after-effects differ in their parametricity

The after-effects described in Figure 2 for all parameters showed significant main effects

of photoperiod, revealing that either the interval between light transitions or the light duration is

inducing persistent effects on ψ, α, and τ. To determine whether the critical aspect of the light

cycle is the timing of the light/dark and dark/light transitions or the duration of light within each

cycle, we used skeleton photoperiods consisting of two 1-hour pulses of light, separated by

intervals of darkness, that mimic the timing of the light/dark transitions on full photoperiods of

16:8 and 8:16 (Figure 4A). Skeleton photoperiods have been used to assess the mechanism of

entrainment through seasonal changes in ψ (Pittendrigh and Minis, 1964), as well as to

determine the proximal effect on locomotor behavior duration (Pittendrigh and Daan, 1976a,

1976b). In order to determine whether skeleton photoperiods can induce plasticity in α, τ, and ψ,

we assessed after-effects in locomotor behavioral rhythms in DD following 28 days of exposure

to skeleton long or short photoperiods in comparison to similar entrainment to full 16:8 and 8:16

photoperiods.

ψ exhibited main effects of both the duration of light in complete photoperiods

(Completeness, Figure 4B), and the timing of light transitions in skeleton photoperiods (Interval,

Figure 4B). This result demonstrates that the ψ induced by photoperiodic entrainment has a

significant non-parametric component. In addition to the main effect of the interval between light

transitions on skeleton photoperiods, there is also a significant effect of the light duration of

complete photoperiods, indicating that there is a parametric influence on the ψ of locomotor

behavior rhythms. In contrast, we found a significant main effect only of the onset-offset interval

of skeleton photoperiods on α after-effects, suggesting that this effect is primarily non-

parametric (Figure 4C). We found that the skeleton photoperiods had no significant effect on τ

(Figure 4D), however previous investigators have reported τ after-effects of skeleton

photoperiods (Pittendrigh and Daan, 1976a). Thus we found that there are significant parametric

and non-parametric effects on ψ, non-parametric effects of onset-offset interval on α, and

relatively weak non-significant after-effects on τ of skeleton photoperiods.

***Ex vivo* SCN after-effects**

Using *ex vivo* imaging of PER2::LUCIFERASE (PER2::LUC) rhythms following

entrainment to the same six light schedules described above, we investigated the effects of

photoperiod and T cycles on plasticity of the SCN pacemaker. We housed heterozygous

PER2::LUC animals in identical conditions to our complete photoperiod *in vivo* experiments,

including running wheels, and collected SCN slices from the animals after 14-28 days of

exposure to each of the light schedules described above. Slices were collected within 4 hours of

lights off, maintained in organotypic culture, and PER2::LUC luminescence recorded *ex vivo* for

5-7 days (see Methods). Slice recordings were analyzed by detecting peak PER2::LUC

expression in each of a 102 x 102 grid of sub-cellular (~5 x 5 µm) regions of interest (ROIs)

spread across the entire SCN image. These PER2::LUC peak times were then used to calculate

and map $\sigma_{\Box}$ (Figure 5A), while the average PER2::LUC luminescence profile over time was

used to calculate SCN τ.

We found that the ψ of SCN PER2::LUC rhythms (relative to projected ZT12 of the

previous light cycle) was significantly influenced by both T cycle and photoperiod, with a

significant interaction between the two factors (Figure 5B), similar to the ψ of locomotor rhythms.

As T cycle length increases, SCN ψ also becomes more advanced, and within each T cycle the

long photoperiod is more advanced than the short. Again, like the *in vivo* ψ, the *ex vivo* ψ

featured a narrowing difference between long and short as T cycle length increases. Similarly,

mirroring *in vivo* α, we found that the photoperiod modulated the degree of synchrony of

PER2::LUC rhythms within explanted SCN (σ_{ϕ}, Figure 5C), with increased phase dispersion in

long photoperiods compared to short. However, we also found a significant main effect of T

cycle length on the SCN σ_{ϕ} after-effect. This effect was driven by increased σ_{ϕ} in both T26

groups compared to both T24 groups (post-hoc multiple comparisons test between T cycle

lengths, T24 vs. T26 adjusted *p* = 0.0329). There is no single consistent linear effect of T cycle

on the *ex vivo* σ_{ϕ} after-effect. As such, the pattern most resembled that of the example in

Supplemental Figure 2A, but with the T26 groups elevated.

SCN τ after-effects following T cycle entrainment have been a source of significant

interest in part because of the apparent disconnect between after-effects of T cycle (where *ex*

vivo SCN and behavioral measurements negatively correlate) and of photoperiod (where *ex vivo*

SCN and behavioral measurements agree). In our combined T cycle/photoperiod paradigm we

found a significant negative relationship between T cycle length and SCN τ length. Entrainment

to short T cycles *in vivo* resulted in lengthened SCN τ *ex vivo*, and *vice versa* as previously

reported (Aton et al., 2004; Azzi et al., 2017; Molyneux et al., 2008). Visually, there is an

intriguing reversal in trend direction of possible photoperiod effects between T24 and the two

non-24 T cycles that could indicate an interaction. In T24 cycles, the previously reported

relationship between long and short photoperiod (shortened, behavior-like τ after-effect for long

photoperiod) is present, while in T23 and T26 cycles, the relationship between long and short is

inverted. This relationship results in the longest and shortest τ aligning with T23 Long and T26

Short, the same two groups highlighted above (Figure 3) as overlap groups where the effects of

T cycle and photoperiod are in line with one another. However, no significant photoperiod

effects or interactions were revealed by the ANOVA, or by multiple post-hoc tests for differences

within the T cycle groups, so these visual trends require further testing.

***Ex vivo* SCN after-effects have similar parametricity to behavioral after-effects**

To investigate the parametricity of SCN after-effects *ex vivo*, we collected slices from

animals entrained to skeleton long and short photoperiods (Figure 6A). Similar to *in vivo* ψ of the

behavioral rhythm to skeleton photoperiods, there was a significant effect of skeleton onset-

offset interval on the ψ of the PER2::LUC rhythm (Figure 6B). Unlike *in vivo*, the PER2::LUC ψ

did not have a significant effect of photoperiod completeness, although there was a trend

towards a decrease in the effect size on ψ of skeleton photoperiods compared to the complete

photoperiods. The $\sigma_\square$ after-effect was significantly influenced by the onset-offset interval (Figure

6C), similar to *in vivo*. The $\sigma_\square$ after-effect *ex vivo* was also significantly influenced by

photoperiod completeness. However, the effect size of the skeleton photoperiods on SCN $\sigma_\square$

was larger compared to the complete photoperiods. Though the interaction that this implies was

not statistically significant, it does suggest that skeleton interval is a principal driver of $\sigma_\square$ after-

effects similar to *in vivo* α (Figure 4C). As with behavioral τ after-effects, we found no significant

effect of onset-offset interval or photoperiod completeness on the *ex vivo* τ after-effect, again

suggesting that the effects of photoperiod on τ after-effects are modest compared to T cycle

entrainment.

DISCUSSION

Measurements in constant conditions reveal properties of the SCN and downstream

circadian outputs free of light input, and as such the study of after-effects is one means to

understand the plasticity of the SCN in response to seasonal light conditions. Understanding

after-effects is also useful *per se*, as the encoding of light information within the SCN allows

seasonal circadian properties to remain stable across ephemeral changes in the environment.

Organisms in the wild experience fluctuations in light exposure during the annual photoperiodic

cycle because of weather, nesting, hibernation, and other factors. Stable but flexible encoding of

seasonal light inputs within the circadian system would allow for consistency across these light

exposure fluctuations.

To improve our understanding of how after-effects are induced, we examined the relative

influences of T cycle length, photoperiod, and onset-offset interval on *in vivo* (α, τ) and *ex vivo*

($\sigma_\square$, τ) after-effects, as well as on ψ (both *in vivo* and *ex vivo*). Using eight light cycle input

groups, including six T cycle/photoperiod combinations and two skeleton photoperiods, we

tested which specific components of light cycle input (light duration, onset-offset interval,

repeated phase shifts) are encoded by the SCN in a persistent manner by measuring after-

effects in circadian behavior and in SCN rhythms assayed *ex vivo*.

Comparison of *in vivo* and *ex vivo* responses

We found that ψ is strongly influenced by T cycle and photoperiod length, both *in vivo*

(Figure 2B) and *ex vivo* (Figure 5B), with a significant interaction between T cycles and

photoperiod in setting subsequent ψ in both cases. Using skeleton photoperiods to assess the

influence of non-parametric (light onset-offset interval) and parametric (light interval

completeness) input, we found both input types significantly influence ψ *in vivo* (Figure 4B) while

observing significant influences of non-parametric input and a trend toward an influence of

parametric input *ex vivo* (Figure 6B). Determination of ψ has long been considered to be non-

parametric, with the exception of the continuity of light on full light cycles disambiguating onset

from offset on near symmetrical light cycles (e.g. near T24 12:12, Pittendrigh and Minis, 1964),

however our results suggest a more complex picture akin to the circadian surface entrainment

model of Roenneberg for *Neurospora* (Rémi et al., 2010; Roenneberg et al., 2010).

After-effects in α were strongly influenced by photoperiod length (Figure 2C) in a non-

parametric manner (Figure 4C). This result was mirrored by the significant main effect of

photoperiod on *ex vivo* SCN σ$_\square$ (Figure 5C), which was also primarily non-parametric (Figure

6C). One notable difference between *in vivo* α and *ex vivo* SCN σ$_\square$, however, was the general

increase in SCN σ$_\square$ measured in both T26 photoperiods. The increase in the T26 Short group

may be related to the contracted α observed in LD during entrainment to T26 Long and Short

(Figure 2A, bottom row): Though the free-running α expands on the first day of DD, there is

apparent negative masking occurring in the early active phase of these mice during LD.

Because slices were made during the last LD cycle rather than on the first day of DD, the σ$_\square$

could reflect that negatively-masked pattern and imply that σ$_\square$ is correlated with overt "net" α,

rather than the underlying assumed endogenous active phase length.

We found that behavioral τ after-effects are significantly influenced by both T cycle and

photoperiod length, but that the T cycle influence is of greater statistical significance and effect

size (Figure 2D), suggesting that the T cycle input to τ after-effects is more robust. Our results

with skeleton photoperiods *in vivo* support this notion, as we did not observe significant τ after-

effects with skeleton photoperiods (Figure 4D). Previous studies have described τ after-effects

of skeleton photoperiods on behavioral τ, but with modest effect sizes (Pittendrigh and Daan,

1976a). While locomotor τ after-effects reflected the length of the entraining T cycle, we

observed negative correlations between the τ after-effects of *ex vivo* SCN and the length of the

entraining T cycle, consistent with previous findings (Aton et al., 2004; Azzi et al., 2017;

Molyneux et al., 2008). We did not detect any statistically significant modulation of the T cycle

effects on τ by varying photoperiod, but visually, there are trend reversals in the expected

direction of photoperiodic effects on τ in the non-24 hour T cycle groups that suggest the

possibility of interactions that we did not detect statistically (T24 Long vs. T23 Long, p = 0.1513;

T24 Short vs. T26 Short, p = 0.0584; Sidak's Multiple Comparisons, see further discussion

below).

Differences between T cycle- and photoperiod-induced responses

Because α and τ after-effects have been observed together (shorter α, shorter τ; longer

α, longer τ), previous studies have attempted to explain both changes with a single underlying

mechanism (Beersma et al., 2017; Gu et al., 2016). We observed different patterns of

responses for after-effects in α and τ in photoperiod/T cycle combinations (Figure 2A, C-D) as

well as in skeleton photoperiods (Figure 4A, C-D), and correlation of changes in α and τ were

specific to certain T and photoperiod combinations (Figure 3). When measuring the

correspondence of τ and α after-effects *in vivo*, correlations were only observed in cases where

T cycle and photoperiod effects are expected to align (Figure 3). These results suggest that

while the two after-effects may frequently be observed together, their distinct patterning in

response to particular T cycles and photoperiods could be driven by disparate but conditionally

convergent mechanisms.

Changes in SCN σ remain a likely explanation for alterations in downstream locomotor

behavior duration, as an extended population-level high-firing phase of SCN neurons (for which

the PER2::LUC σ is a proxy here) would logically underlie the regulation of timing activity or

inactivity (Ciarleglio et al., 2009, 2011; Inagaki et al., 2007; VanderLeest et al., 2007). Changes

in τ, however, shown here to be induced primarily through T cycle entrainment and likely the

requisite repeated phase shifts thereof, may be regulated in a different manner. Mechanisms

suggested recently by other groups (Azzi et al., 2014, 2017) surrounding the epigenetic

changes that may occur after extended entrainment to T cycles may help explain alterations in τ

that we observed.

Regional phase differences and their influence on after-effects

Regional phase differences within the SCN network have been observed in response to

T cycles and photoperiod, and have been proposed to drive τ after-effects (Azzi et al., 2017;

Buijink et al., 2016; Evans et al., 2013; Myung et al., 2015). In SCN explants from mice

entrained to long photoperiods, the ventral or ventromedial region SCN phase leads in some

studies (Buijink et al., 2016; Evans et al., 2013), but the reverse regional phase relationship was

observed in another (Myung et al., 2015). SCN τ shortening after entrainment to long

photoperiods was observed to correlate with ventral phase lead (Evans et al., 2013), with

ventral phase lag (Myung et al., 2015), or not be present (Buijink et al., 2016). Similarly, Azzi *et

al*. (2017) found that following entrainment to T cycles, there was an anticorrelation between

SCN τ after-effect and the previous T cycle τ, as expected from previous results (Aton et al.,

2004; Molyneux et al., 2008). The same study also found that shortened τ correlated with

ventral phase lag, while lengthened τ correlated with ventral phase lead – the opposite pattern

previously found for photoperiodic τ after-effects by the same group (Evans et al., 2013). Thus,

there is no clear consensus in the literature as to whether SCN regional phase differences

mediate τ after-effects.

We also observed the anticorrelation of *ex vivo* SCN τ after T cycle entrainment *in vivo*,

as well as a variety of regional $\sigma_\square$ across our experimental conditions that included examples of

ventral core phase lead in response to short T cycles (see T23 Long in Figure 6) similar to Azzi

et al. (2017), and examples of more ventromedial phase clustering similar to that of Buijink *et al*.

(2016) in response to long T cycles (see T26 Short in Figure 6). Differences in regional patterns

(ventral core vs. ventromedial) may be explained by slice position on the rostral caudal axis of

the SCN. Azzi et al. (2017) show that taking sections more caudal in the SCN changes the

regional phase pattern from ventral core grouping to a ventromedial grouping similar to Buijink

(2016). We speculate that we may not have controlled well enough for slice depth, resulting in

our observation of variations of both patterns in our data.

Conceptually, the anticorrelation of SCN *ex vivo* τ with *in vivo* τ after-effects of T cycles

presents a challenge. One interpretation of these results is that the observed *ex vivo* τ reflects a

deafferentation-induced reorganization of the SCN network, indicating extra-SCN input is key to

T cycle after-effects *in vivo* (Aton et al., 2004; Molyneux et al., 2008). Support for this view

comes from our previous results showing an anticorrelation between *in vivo* behavioral and *ex*

vivo SCN τ effects of genetic serotonergic de-afferentation of the SCN (Ciarleglio et al., 2014).

In addition, part of the confound regarding previous *ex vivo* SCN T-cycle after-effects appears to

be a bias in the reporting of SCN regional rhythms by the standard PER2::LUC reporter that

more highly weights τ of the dorsal region (Azzi et al., 2017). In any case, the lack of

correspondence between *in vivo* and *ex vivo* T-cycle τ after-effects in the SCN makes it difficult

to attribute *ex vivo* findings of regional patterns or mechanisms to the *in vivo* case, although the

system can be used to study SCN *ex vivo* reorganization and its mechanisms (Azzi et al., 2017).

SCN $\sigma_{\square}$ and its relationship with α

The degree of SCN neuron $\sigma_{\square}$ is a characteristic attribute of photoperiodic encoding, as

it represents a change in state of the SCN induced by the duration of the light interval of the

daily cycle and is maintained after transfer to constant conditions (Buijink et al., 2016; Ciarleglio

et al., 2011; Inagaki et al., 2007; VanderLeest et al., 2007). Our results suggest that the timing

of light/dark transitions is a key factor in influencing SCN $\sigma_{\square}$ (Figure 6C) and *in vivo* α (Figure

4C), although we also detected an effect of light duration on SCN σ_ϕ (Figure 6C). The spread in

the timing of neuronal rhythms allows for the daily duration of high firing activity of the SCN to

be broadened or contracted without required alteration to the waveforms of individual SCN

neurons, although this aspect is modulated by developmental inputs (Ciarleglio et al., 2011). As

such, the degree of σ_ϕ within the SCN is a potential target for artificial SCN manipulation –

driving neurons to adopt the σ_ϕ characteristic of a photoperiod, such as by pharmacologically,

optogenetically, or chemogenetically extending the high firing phase, would be expected to

induce the behavioral α characteristic of that particular photoperiod, and perhaps other

photoperiod-dependent responses, such as reproductive state and affective behaviors.

Because current methods for *in vivo* imaging are limiting, the measure of α after-effects

may be a useful proxy for inferring the waveform of the SCN. The integrity of this SCN

waveform-α connection depends on the correlation between the two. Our findings that α and *ex*

vivo SCN neuron σ_ϕ associate across multiple lighting conditions regardless of net daily phase

shift do not address causality, but strengthen the association between the two measurements.

As the broadened electrical waveform of the SCN is associated with the compressed behavioral

α seen in long photoperiods (and the contracted waveform with extended α in short

photoperiods), the spread in phase of SCN neurons and the corresponding widening of overall

firing that occurs as a result should account for the observed changes in α (Houben et al.,

2009).

Changes to SCN network synchrony may have additional ramifications in the form of

altered responses to light input, potentially explaining previously observed larger phase shifts

after exposure to short days, and smaller phase shifts after exposure to long days (Refinetti,

2002). The role of SCN neuron σ_ϕ in mediating those altered responses and altered τ has been

explored computationally (Gu et al., 2016) and through altered GABA signaling (Farajnia et al.,

2014). In their analysis of after-effects on τ, Pittendrigh and Daan (1976b) suggested that this

change in circadian property may provide functional significance in terms of priming future

responses to light, but it remains to be seen whether any of these after-effects constitutes an

adaptive advantage or if they are simply a side effect of altered network synchrony.

460   **CONCLUSION**

461   Taken together, our results show that distinct components of light cycles can have

distinct effects on the circadian system, with (a) the timing of light/dark transitions in

photoperiods driving α plasticity; (b) combined photoperiod and T cycle length driving plasticity

in τ and ψ of both *in vivo* behaviors and *ex vivo* SCN gene expression; and (c) the light/dark

transition interval and long T cycle lengths affecting *ex vivo* SCN $\sigma_{\square}$. These results open the

door to selective manipulation of circadian plasticity in α or τ for further experimentation and for

potential translation to treat photoperiodic syndromes. The results shown here also demonstrate

the utility of deconvolving overlapping types of input when studying complex processes within

the SCN. Our results indicate that α and τ after-effects are possibly driven by distinct, though

perhaps related, mechanisms, and such information can be incorporated into experimental and

computational approaches to test the system in the future. With new techniques currently being

established in our field to measure SCN neuronal firing rate, calcium signaling, and activity

levels with high cellular specificity and extended timeframe, there will be an improved ability to

characterize the specific effect of photoperiodic influences on these circadian components.

476   **ACKNOWLEDGMENTS**

477   The authors would like to thank Maria Luísa Jabbur, Carl Johnson, Terry Page, and Chloe

478   Snider for their helpful comments and discussions. This work was supported by NIH R01

479 GM117650 to D.G.M., NIH F31 NS096813 to M.C.T., and NSF GRFP 0909667 to M.C.T.

481   **AUTHOR CONTRIBUTIONS**

484   **DECLARATION OF INTERESTS**

485 The authors have no competing interests to report.

487   **REFERENCES**

Aton SJ, Block GD, Tei H, Yamazaki S and Herzog ED (2004) Plasticity of circadian behavior

and the suprachiasmatic nucleus following exposure to non-24-hour light cycles. *Journal of*

biological rhythms 19(3): 198–207.

Azzi A, Dallmann R, Casserly A, Rehrauer H, Patrignani A, Maier B, Kramer A and Brown SA

(2014) Circadian behavior is light-reprogrammed by plastic DNA methylation. *Nature*

neuroscience 17(3): 377–82.

Azzi A, Evans JA, Leise T, Myung J, Takumi T, Davidson AJ and Brown SA (2017) Network

Dynamics Mediate Circadian Clock Plasticity. *Neuron* 93(2): 441–450.

Bartness TJ and Wade GN (1984) Photoperiodic control of body weight and energy metabolism

in Syrian hamsters (Mesocricetus auratus): role of pineal gland, melatonin, gonads, and

diet. *Endocrinology* 114(2): 492–8.

Bartness TJ, Elliott JA and Goldman BD (1989) Control of torpor and body weight patterns by a

seasonal timer in Siberian hamsters. *The American journal of physiology* 257(1 Pt 2):

R142-9.

Basnet S, Merikanto I, Lahti T, Männistö S, Laatikainen T, Vartiainen E and Partonen T (2016)
Seasonal variations in mood and behavior associate with common chronic diseases and
symptoms in a population-based study. *Psychiatry research* 238: 181–188.

Beersma DGM, Gargar KA and Daan S (2017) Plasticity in the Period of the Circadian
Pacemaker Induced by Phase Dispersion of Its Constituent Cellular Clocks. *Journal of*
Biological Rhythms 32(3): 237–245.

Buijink MR, Almog A, Wit CB, Roethler O, Olde Engberink AHO, Meijer JH, Garlaschelli D,
Rohling JHT and Michel S (2016) Evidence for Weakened Intercellular Coupling in the
Mammalian Circadian Clock under Long Photoperiod. Yamazaki S (ed.) *PLOS ONE*
11(12): e0168954.

Ciarleglio CM, Gamble KL, Axley JC, Strauss BR, Cohen JY, Colwell CS and McMahon DG
(2009) Population Encoding by Circadian Clock Neurons Organizes Circadian Behavior.
Journal of Neuroscience 29(6): 1670–1676.

Ciarleglio CM, Axley JC, Strauss BR, Gamble KL and McMahon DG (2011) Perinatal
photoperiod imprints the circadian clock. *Nature Neuroscience* 14(1). Nature Publishing
Group: 25–27.

Ciarleglio CM, Resuehr HES, Axley JC, Deneris ES and McMahon DG (2014) Pet-1 Deficiency
Alters the Circadian Clock and Its Temporal Organization of Behavior. Mistlberger RE (ed.)
PLoS ONE 9(5): e97412.

Elliott JA and Goldman BD (1981) Seasonal Reproduction. In: Adler NT (ed.)
Neuroendocrinology of Reproduction. Boston, MA: Springer US, pp. 377–423.

Evans JA, Leise TL, Castanon-Cervantes O and Davidson AJ (2013) Dynamic interactions

mediated by nonredundant signaling mechanisms couple circadian clock neurons. *Neuron*

80(4): 973–983.

Farajnia S, van Westering TLE, Meijer JH and Michel S (2014) Seasonal induction of

GABAergic excitation in the central mammalian clock. *Proceedings of the National*

Academy of Sciences 111(26): 9627–9632.

Fernandez DC, Fogerson PM, Lazzerini Ospri L, Thomsen MB, Layne RM, Severin D, Zhan J,

Singer JH, Kirkwood A, Zhao H, Berson DM and Hattar S (2018) Light Affects Mood and

Learning through Distinct Retina-Brain Pathways. *Cell* 175(1): 71-84.e18.

Gu C, Rohling JHT, Liang X and Yang H (2016) Impact of dispersed coupling strength on the

free running periods of circadian rhythms. *Physical Review E* 93(3): 032414.

Gwinner E (1990) Circannual Rhythms in Bird Migration: Control of Temporal Patterns and

Interactions with Photoperiod. In: Gwinner E (ed.) *Bird Migration*. Berlin, Heidelberg:

Springer Berlin Heidelberg, pp. 257–268.

Houben T, Deboer T, van Oosterhout F and Meijer JH (2009) Correlation with Behavioral

Activity and Rest Implies Circadian Regulation by SCN Neuronal Activity Levels. *Journal of*

Biological Rhythms 24(6): 477–487.

Inagaki N, Honma S, Ono D, Tanahashi Y and Honma K (2007) Separate oscillating cell groups

in mouse suprachiasmatic nucleus couple photoperiodically to the onset and end of daily

activity. *Proc Natl Acad Sci USA* 104(18): 7664–7669.

Koorengevel KM, Beersma DG., den Boer J a and van den Hoofdakker RH (2003) Mood

regulation in seasonal affective disorder patients and healthy controls studied in forced

desynchrony. *Psychiatry Research* 117(1): 57–74.

Mickman CT, Stubblefield JS, Harrington ME and Nelson DE (2008) Photoperiod alters phase

difference between activity onset in vivo and mPer2::luc peak in vitro. *American Journal of*

Physiology-Regulatory, Integrative and Comparative Physiology 295(5): R1688–R1694.

Molyneux PC, Dahlgren MK and Harrington ME (2008) Circadian entrainment aftereffects in

suprachiasmatic nuclei and peripheral tissues in vitro. *Brain Research* 1228: 127–134.

Myung J, Hong S, DeWoskin D, De Schutter E, Forger DB and Takumi T (2015) GABA-

mediated repulsive coupling between circadian clock neurons in the SCN encodes

seasonal time. *Proceedings of the National Academy of Sciences* 112(29): E3920–E3929.

Oh E-Y, Ansell C, Nawaz H, Yang C-H, Wood PA and Hrushesky WJM (2010) Global breast

cancer seasonality. *Breast Cancer Research and Treatment* 123(1): 233–243.

Pittendrigh CS and Daan S (1976a) A functional analysis of circadian pacemakers in nocturnal

rodents (I). *Journal of Comparative Physiology A* 106(3). Springer-Verlag: 223–252.

Pittendrigh CS and Daan S (1976b) A functional analysis of circadian pacemakers in nocturnal

rodents (IV). *Journal of Comparative Physiology A* 106(3). Springer-Verlag: 291–331.

Pittendrigh CS and Minis DH (1964) The Entrainment of Circadian Oscillations by Light and

Their Role as Photoperiodic Clocks. *The American Naturalist* 98(902): 261–294.

Refinetti R (2002) Compression and expansion of circadian rhythm in mice under long and short

photoperiods. *Integrative Physiological & Behavioral Science* 37(2): 114–127.

Rémi J, Merrow M and Roenneberg T (2010) A Circadian Surface of Entrainment: Varying T, τ,

and Photoperiod in Neurospora crassa. *Journal of Biological Rhythms* 25(5): 318–328.

Roenneberg T, Rémi J and Merrow M (2010) Modeling a Circadian Surface. *Journal of*

Biological Rhythms 25(5): 340–349.

Schwartz WJ, Tavakoli-Nezhad M, Lambert CM, Weaver DR and de la Iglesia HO (2011)

Distinct patterns of Period gene expression in the suprachiasmatic nucleus underlie

circadian clock photoentrainment by advances or delays. *Proceedings of the National*

Academy of Sciences 108(41): 17219–17224.

Tackenberg MC and McMahon DG (2018) Photoperiodic Programming of the SCN and Its Role

in Photoperiodic Output. *Neural Plasticity* 2018: 1–9.

VanderLeest HT, Houben T, Michel S, Deboer T, Albus H, Vansteensel MJ, Block GD and

Meijer JH (2007) Seasonal Encoding by the Circadian Pacemaker of the SCN. *Current*

Biology 17(5): 468–473.

Wehr TA (2001) Photoperiodism in Humans and Other Primates: Evidence and Implications.

Journal of Biological Rhythms 16(4): 348–364.

                                    Figures

Figure 1. Schematic of possible patterns of measurements across T23 L/S, T24 L/S, and T26
L/S. **A**, a T-cycle dependent response where the measurement is graded depending on the T
cycle and photoperiod within the T-cycle has negligible effects. **B**, a photoperiod-dependent
response where light duty cycle dictates the measurement regardless of T cycle. **C**, a
combination of photoperiod and T cycle influences, with each of the 6 combinations having a
characteristic measurement level. Note that in all examples, the polarity of the measurement
change is arbitrary (e.g., T cycle response in A can increase or decrease with T cycle,
photoperiod response in B can have either Long or Short measure higher).These patterns are
illustrations of individual and combined main effects; possible interaction effects are not shown.

Figure 2. Behavioral after-effects of six photoperiod/T cycle combinations. **A**, representative actograms from long (left column) and short (right column) photoperiods combined with T23 (top row), T24 (middle row), and T26 (bottom row). **B**, ψ following exposure to the six light schedules. Values are relative to projected ZT12 (positive values advanced, negative values delayed). Interaction $p = 0.0014$ (7.242%), T Cycle Length $p < 0.0001$ (51.30%), Photoperiod Legth $p < 0.0001$ (20.97%). **C**, α after-effect across all six schedules as measured by automated α detection (see Methods). Interaction $p = 0.4256$ (2.081%), T Cycle Length $p = 0.1648$ (4.468%), Photoperiod Length $p < 0.0001$ (21.67%). **D**, period after-effect across the six schedules as measured by chi square periodogram. Interaction $p = 0.7587$ (0.2835%), T Cycle Length $p < 0.0001$ (56.17%), Photoperiod Length $p = 0.0087$ (3.778%). Two-way ANOVA results represent p value and percent variation ($p < 0.0001$ ****, $p < 0.001$ ***, $p < 0.01$ **, $p < 0.05$ *). T23 Long, n = 10; T23 Short, n = 12; T24 Long, n = 12; T24 Short, n = 12; T26 Long, n = 6; T26 Short, n = 14.

Figure 3. Within-individual correlations between τ and α after-effects following the given T-
cycle/photoperiod combination. T23 Long 0.8238 (p = 0.0034, **), T24 Long -0.06835 (p = 0.8328, ns),
T26 Long 0.004966 (p = 0.9926, ns), T23 Short -0.3147 (p = 0.4477, ns), T24 Short 0.2671 (p = 0.4013,
ns), T26 Short 0.5818 (p = 0.0291, *); Values reported are Pearson r and p value (p < 0.0001 ****, p <
0.001 ***, p < 0.01 **, p < 0.05 *).

Figure 4. Behavioral after-effects following exposure to skeleton long and short photoperiods compared
to T24 complete long and short photoperiods. **A**, representative actograms from long (top) and short
(bottom) skeleton photoperiods. **B**, ψ following exposure to skeleton long and short photoperiods. Values
are relative to projected ZT12 (positive values advanced, negative values delayed). Interaction $p = 0.3161$
(1.33%), Completeness $p = 0.0003$ (21.24%), Onset-Offset Interval $p < 0.0001$ (29.32%). **C**, α after-effect
of skeleton long and short photoperiods. Interaction $p = 0.1861$ (4.332%), Completeness $p = 0.9858$
(0.0008%), Onset-Offset Interval $p = 0.0376$ (11.15%). **D**, τ after-effect of skeleton long and short
photoperiods. Interaction $p = 0.7835$ (0.2063%), Completeness $p = 0.7006$ (0.4044%), Onset-Offset
Interval $p = 0.0765$ (8.988%). Long and Short complete photoperiod values in B-D are replotted from
Figure 2. Two-way ANOVA results represent p value and percent variation ($p < 0.0001$ ****, $p < 0.001$ ***,
$p < 0.01$ **, $p < 0.05$ *). Complete Long, n = 12; Complete Short, n = 12; Skeleton Long, n =7, Skeleton
Short, n = 6.

Figure 5. PER2::LUC after-effects following exposure to six photoperiod/T cycle combinations. **A**,
representative relative phase maps from long (left column) and short (right column) photoperiod
combined with T23 (top row), T24 (middle row), or T26 (bottom row). **B**, ψ of the PER2::LUC rhythm
following entrainment to the specified T cycle/photoperiod combination. Values represent the time of the
first PER2::LUC peak relative to the projected lights off for the first cultured cycle (projected ZT 12).
Interaction $p = 0.0052$ (8.185%), T Cycle Length $p < 0.0001$ (49.47%), Photoperiod Length $p < 0.0001$
(24.94%). **C**, σ_ϕ (median absolute deviation of PER2::LUC peak times) across SCN slices from animals
entrained to each of the specified light schedules. Interaction $p = 0.7168$ (1.433%), T Cycle Length $p =$
0.0353 (15.89%), Photoperiod Length $p = 0.0114$ (15.42%). **D**, τ of the PER2::LUC rhythm over 5 cycles
ex vivo. Interaction $p = 0.4104$ (4.109%), T Cycle Length $p = 0.0111$ (23.40%), Photoperiod Length $p =$
0.4151 (1.529%). Two-way ANOVA results represent p value and percent variation ($p < 0.0001$ ****, $p <$
0.001 ***, $p < 0.01$ **, $p < 0.05$ *). T23 Long, n = 6; T23 Short, n = 4; T24 Long, n = 6; T24 Short, n = 8;
T26 Long, n = 7; T26 Short, n = 6.

Figure 6. PER2::LUC after-effects following entrainment to skeleton long and short photoperiods compared to T24 complete long and short photoperiods. **A**, representative relative phase maps from skeleton long (top) and short (bottom) photoperiod. **B**, ψ of the PER2::LUC rhythm following entrainment to skeleton long and short photoperiods. Values represent the time of the first PER2::LUC peak relative to the projected lights off for the first cultured cycle (projected ZT 12). Interaction $p = 0.7514$ (0.1326%), Completeness $p = 0.1660$ (2.651%), Onset-Offset Interval $p < 0.0001$ (70.79%). **C**, σ_ϕ (median absolute deviation of PER2::LUC peak times) across SCN slices from animals entrained to skeleton long and short photoperiod. Interaction $p = 0.0654$ (7.695%), Completeness $p = 0.0318$ (10.77%), Onset-Offset Interval $p = 0.0003$ (37.26%). **D**, τ of the PER2::LUC rhythm over 5 cycles *ex vivo*. Interaction $p = 0.4867$ (2.277%), Completeness $p = 0.5608$ (1.587%), Onset-Offset Interval $p = 0.6959$ (0.7131%). Long and Short complete photoperiod values in B-D are replotted from Figure 5. Two-way ANOVA results represent p value and percent variation ($p < 0.0001$ ****, $p < 0.001$ ***, $p < 0.01$ **, $p < 0.05$ *). Complete Long, n = 6; Complete Short, n = 8; Skeleton Long, n = 6, Skeleton Short, n = 5.