TITLE

Distinct components of photoperiodic light are differentially encoded by the mammalian

circadian clock

AUTHORS

Michael C. Tackenberg[1], Jacob J. Hughey[2,3], and Douglas G. McMahon[1,2*]

[1]Vanderbilt Brain Institute, Vanderbilt University, Nashville, TN, USA

[2]Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA

[3]Department of Biomedical Informatics, Vanderbilt University School of Medicine, Nashville, TN,

USA

Correspondence: douglas.g.mcmahon@vanderbilt.edu

RUNNING TITLE

Components of photoperiodic light

---

[*] Lead Contact

ABSTRACT

Seasonal light cycles influence multiple physiological functions and are mediated through

photoperiodic encoding by the circadian system. Despite our knowledge of the strong connection

between seasonal light input and downstream circadian changes, less is known about the specific

components of seasonal light cycles that are encoded and induce persistent changes in the

circadian system. Using combinations of three T cycles (23, 24, 26 hr.) and two photoperiods per

T cycle (Long and Short, with duty cycles scaled to each T cycle), we investigate after-effects of

entrainment to these six light cycles. We measure locomotor behavior duration (α), period (τ), and

entrained phase angle (ψ) *in vivo,* and SCN phase distribution (σ_ϕ), τ, and ψ *ex vivo* in order to

refine our understanding of critical light components for influencing particular circadian properties.

We find that photoperiod and T cycle length both drive determination of *in vivo* ψ but differentially

influence after-effects in α and τ, with photoperiod driving changes in α and photoperiod length

and T cycle length combining to influence τ. Using skeleton photoperiods, we demonstrate that *in*

vivo ψ is determined by both parametric and non-parametric components, while changes in α are

driven non-parametrically. Within the *ex vivo* SCN, we find that ψ and σ_ϕ of the

PER2::LUCIFERASE rhythm follow closely with their likely behavioral counterparts (ψ and α of

the locomotor activity rhythm), while also confirming previous reports of τ after-effects of gene

expression rhythms showing negative correlations with behavioral τ after-effects in response to T

cycles. We demonstrate that within-SCN σ_ϕ changes, thought to underly α changes *in vivo*, are

induced primarily non-parametrically. Taken together, our results demonstrate distinct

components of seasonal light input differentially influence ψ, α, and τ, and suggest the possibility

of separate mechanisms driving the persistent changes in circadian behaviors mediated by

seasonal light.

KEYWORDS

INTRODUCTION

The duration of daylight, or photoperiod, represents a predictable and dynamic signal of seasonal change throughout the year across much of the planet. The predictability of this signal is harnessed by organisms to initiate or avoid a variety of biological functions at specific times of year, including reproductive behaviors (Elliott and Goldman, 1981), resource conservation (Bartness et al., 1989; Bartness and Wade, 1984), and migration (Gwinner, 1990). In humans, a variety of disorders and non-communicable diseases have seasonal correlations (Basnet et al., 2016; Oh et al., 2010; Wehr, 2001), with perhaps the most striking seasonal influence being on depression-anxiety behaviors (Koorengevel et al., 2003). Despite clear associations between photoperiodic exposure and physiological function and dysfunction, the roles of specific components of seasonal light cycles that induce those changes are incompletely understood.

One factor critical to our current understanding of the mechanism of photoperiodic induction is the persistence of seasonal changes. Physiological hallmarks of exposure to particular light schedules can be observed, for varying intervals, beyond direct exposure to that light schedule (Pittendrigh and Daan, 1976a). These persistent changes, referred to as after-effects, include characteristic changes to core circadian locomotor behavior patterns in the form of activity duration (alpha, α), free-running period (tau, τ), and phase angle of entrainment (psi, ψ). In nocturnal animals, prior exposure to long photoperiods results in after-effects of compressed α, a shortened τ, and an advanced ψ relative to short photoperiod counterparts as measured upon release into DD (Pittendrigh and Daan, 1976a).

The existence of after-effects suggests that seasonal light cycle information is encoded within the brain and can persist beyond the interval of direct exposure. The master circadian pacemaker of the brain, the suprachiasmatic nucleus (SCN), likely plays a major role in this encoding, particularly with regards to after-effects in circadian behavior (for review, see

Tackenberg and McMahon, 2018). In addition, recent work has identified a distinct non-SCN

target for photoperiodic influence, the peri-habenular area, influencing seasonal response in

mood (Fernandez et al., 2018). Like circadian locomotor behavior patterning, the SCN undergoes

substantial changes in organization following exposure to different photoperiods, constituting

network-level after-effects. These responses include changes in phase distribution (σ_ϕ) of its

constituent neurons as well as in τ and ψ of SCN gene expression rhythms. Prior exposure to

long photoperiod has been shown to increase σ_ϕ of *Period1::Luciferase* (Inagaki et al., 2007),

PERIOD2::LUCIFERASE (Buijink et al., 2016), and electrical (VanderLeest et al., 2007) rhythms

compared to short photoperiod counterparts. As in locomotor behavior after-effects, some reports

have shown that prior exposure to long photoperiod shortens the SCN τ compared to short

photoperiod counterparts (16:8 L:D, Ciarleglio et al., 2011; 20:4 L:D, Evans et al., 2013) while

others have found non-significant decreases in SCN τ length following exposure to long

photoperiod, particularly in the anterior SCN (Buijink et al., 2016; Mickman et al., 2008).

Induction of persistent circadian after-effects is not limited to changes in photoperiod.

Manipulation of total day-night cycle (T cycle) length also induces after-effects in circadian

locomotor behavior, with entrainment to short T cycles (T < 24 hours) resulting in contracted α

(Azzi et al., 2014) and shortened τ upon release into constant conditions (Azzi et al., 2014;

Schwartz et al., 2011). However, unlike photoperiodic after-effects, in which the changes in the τ

of the explanted SCN itself match those on behavior, there is a negative correlation between

behavioral and SCN explant τ following entrainment to T cycles. Exposure to short T cycles results

in a long SCN τ and exposure to long T cycles results in a short SCN τ (Aton et al., 2004; Azzi et

al., 2017; Molyneux et al., 2008).

The findings described above reveal a potential pattern of circadian after-effects, with long

photoperiods and short T cycles producing one set of behavioral responses (short α, short τ), and

short photoperiods and long T cycles producing another (long α, long τ). Because recent work

has revealed a role for DNA methylation in establishing τ after-effects of extended T cycle

entrainment, the potential association between τ and α after-effects provides a promising

biochemical mechanism for both changes (Azzi et al., 2014). The linkage between the two types

of after-effects, however, has not yet been sufficiently interrogated. The discordance between the

SCN after-effects of T cycle and photoperiod offers a hint that the relationship may not be so

simple. Here, we seek to further examine the alignment of T cycle and photoperiodic after-effects

by measuring *in vivo* (α, τ, and ψ of locomotor behavior rhythms) and *ex vivo* (σ_ϕ, τ, and ψ of

SCN gene expression rhythms) responses following exposure to six combinations of photoperiod

length and T cycle. We find that after-effects in each of these rhythm characteristics respond to

the six different light cycles in distinct patterns, suggesting that they are each induced by different

aspects of the input light cycle. Using skeleton photoperiods, we demonstrate that parametric vs.

non-parametric responses differ between after-effects in α, τ and ψ, further indicating encoding

of distinct components of light input by each of these characteristics. Examining the rhythms of

explanted SCN *ex vivo* following entrainment to these light schedules, we find that the pattern of

SCN σ_ϕ reflects that of α *in vivo*, providing additional correlative evidence that SCN σ_ϕ may be a

determining factor in setting α.

MATERIALS AND METHODS

Animals

Animal experiments were conducted in accordance with Vanderbilt University Institutional Animal

Care and Use Committee regulations. All animals used for behavioral experiments were

heterozygous for the *Per2::Luciferase* allele (*Per2::Luciferase*[+/-]) or were *Per2* wild-type

(*Per2::Luciferase*[-/-]). All animals used for SCN slice culture experiments were heterozygous for

the *Per2::Luciferase* allele (*Per2::Luciferase*[+/-]).

Activity Monitoring and Housing

Animals were singly-housed in light-tight boxes with activity monitoring by wheel revolutions

detected by ClockLab acquisition software (Actimetrics, Inc.). Complete photoperiods used were

15:8 and 16:7 (T23 Long, ~67% duty cycle), 8:15 and 7:16 (T23 Short, ~33% duty cycle), 16:8

(T24 Long, ~67% cycle), 8:16 (T24 Short, ~33% cycle), 17:9 (T26 Long, ~67% cycle), and 9:17

(T26 Short, ~33 cycle). Animal numbers by group were as follows: 15:8 (*in* vivo n = 5, *ex vivo* n =

4), 16:7 (5, 2), 8:15 (6, 4), 7:16 (2, 0), 16:8 (12, 6), 8:16 (12, 8), 17:9 (6, 7), 9:17 (14,6). The two

photoperiod duty cycles used in T23 (15:8/8:15 vs. 16:7/7:16) gave similar results and were

combined into T23 Long and T23 Short groups for analysis.

Animals were transferred to the specified light schedule after at least 3 weeks of age, and housed

there for at least 28 days before transfer to DD or brain extraction for slicing. Six T26 Short animals

included in the datasets (three used in behavioral experiments, three used in slice experiments)

experienced a multi-hour light failure during the 28 days of entrainment, but received 14 days of

the correct light cycle following the failure.

Skeleton photoperiods were set up by housing animals in 12:12 LD complete photoperiods for

approximately 5 days before transitioning to a 12:12 skeleton (1:10:1:12 L:D:L:D). After several

cycles on this skeleton, the onset pulse was gradually advanced (0.5 hour – 1 hour per day) or

delayed (1 hour per day) until a skeleton long (1:14:1:8 L:D:L:D) or skeleton short (1:6:1:16

L:D:L:D) photoperiod, respectively, was established. Once the final skeleton was established,

animals were given 28 days of exposure to that skeleton until transfer to DD or slicing. Animal

numbers by group were as follows: skeleton short (*in vivo* n = 6, *ex vivo* n = 5), skeleton long (7,

6).

Actogram Analysis

Locomotor behavior was analyzed using ClockLab Analysis software and R. Period was determined using the Chi Square periodogram over the first 7 days of DD. α was measured using automated onset/offset detection for each cycle of the actogram. Full actograms can be found in Supplemental Data 1.

Automated α Measurement

Automated α measurement was performed in R. The activity (bin size 6 minutes) is smoothed by a Savitzky-Golay filter with a span of 25 and degree of 3. A threshold is set relative to the maximum smoothed activity (0.01 * the maximum for each day). Points at which the smoothed activity crosses this threshold are identified as potential onsets and offsets. Segments of activity less than 2 hours are considered inactive. The longest inactive segment is identified, with the start of that segment recorded as the offset and the end of that segment recorded as the onset. α is calculated as activity offset – onset, and ψ as the timing of the activity onset in the first cycle in DD relative to the projected time of lights off from the previous light dark cycle (projected Zeitgeber Time 12 [ZT12]). Identified onsets and offsets for each cycle can be found in Supplemental Data 2. R scripts used for analysis can be found at 10.6084/m9.figshare.11513892.

Slicing and Slice Cultures

Within 4 hours of lights off, animals were sacrificed by cervical dislocation and the brain removed. SCN slices of 200-μm were made, and the SCN further isolated by scalpel cut under a dissecting microscope. Slices were transferred to a 6-well plate with cell culture membrane insert (Millipore, PICMORG50) and 1.2-mL of slice culture media. Slice chambers were sealed with a coverslip and vacuum grease and placed into an incubated light-tight inverted microscope (Zeiss Axioskop). Luminescence was detected using an intensified CCD (Stanford Photonics) controlled by μ-Manager recorded for 2' every 10' for at least 1 week.

Ex-Vivo Analyses

Luminescence recordings were analyzed using Fiji and R. OME-TIFF files from μ-Manager were

opened in Fiji and smoothed using a two-frame minimization (frame rate reduction from 6/hour to

3/hour). A 102x102 grid of 10x10-pixel (5.175 x 5.175 μm) ROIs was overlaid on the images and

brightness measured for each of these 10,404 ROIs measured for every frame. These grid

measurements were then exported to R for further analysis. The first 8 hours of recording were

excluded from peak-finding to prevent slicing artifacts from interfering with measurements. First,

the ROIs comprising the SCN were identified by using pixels within the top 50% of brightness.

The timing of each peak of luminescence was then detected for each ROI and used to calculate

the initial phase. Distribution measurements were made by median absolute deviation across all

SCN ROIs. Phase maps of each SCN used can be found in Supplemental Data 3. R scripts used

for analysis can be found at 10.6084/m9.figshare.11513892. Cycle-by-cycle interpeak-calculated

period and phase distribution are shown in Supplementary Figure 1 and 2, respectively.

Statistical Analysis

For measurements of α, τ, σ_ϕ, and ψ, the effects of photoperiod and T cycle (or completeness and

onset/offset interval) were analyzed by two-way ANOVA. For each comparison, the *p* value and

percent of total variation of each factor, as well as their interaction, are reported. Where indicated

in the text, Sidak's multiple comparisons test was used to further examine within- and between-

factor effects. For correlation analysis shown in Figure 3, Pearson's r test was used.

                                                                RESULTS

Distinct response patterns of ψ, and after-effects on α and τ

To examine the persistent encoding of photoperiods and T cycles *in vivo*, as represented

by entrained phase angle (ψ) and after-effects in activity duration (α) and free-running period (τ),

we exposed mice to six different light schedules consisting of combinations of short or long

photoperiod and short, standard, and long T cycles (T23 Long, T23 Short, T24 Long, T24 Short,

T26 Long, T26 Short). Following 14-28 days of exposure to these light schedules (see Methods),

animals were transferred into constant darkness (DD) for 7 days. Based on previous studies

examining the after-effects of T cycles (Azzi et al., 2014, 2017; Pittendrigh and Daan, 1976a;

Schwartz et al., 2011) and photoperiod (Buijink et al., 2016; Ciarleglio et al., 2009; Evans et al.,

2013; Pittendrigh and Daan, 1976a), we hypothesized that the response of τ and α after-effects

to these six light schedules would produce one of three general patterns (Figure 1).

First, if after-effects in these rhythm characteristics were strictly dependent on the duty

cycle of the light interval, then the groups would segregate into two levels, corresponding to long

or short photoperiods, regardless of T cycle (Figure 1A). A primary influence of the photoperiod

could indicate a parametric effect of the total light duration per cycle and/or a non-parametric

effect from the phase shifts caused by light onset and light offset. Second, if plasticity is strictly

dependent on net daily phase advances or delays driven by entrainment to the T cycles, then the

measurements would segregate into three groups based on T cycle (Figure 1B). This pattern of

results would most readily suggest that after-effects were induced by non-parametric effects of

the daily entraining phase shifts induced by the T cycles. Third, significant main effects of both T

cycle and photoperiod would result in after-effects with each of the 6 groups having a

characteristic level (Figure 1C), while interactions between T cycles and photoperiod would

produce many possible variations of the pattern in Figure 1C.

Indeed, ψ of locomotor rhythms relative to projected ZT12 of the previous light:dark cycle

(see Methods) was strongly influenced by both T cycle length and photoperiod length, with a

significant interaction between these two main effects (Figure 2B). As T cycle length increases, ψ

becomes more advanced. Within each T cycle, the short photoperiod version produces a more

delayed ψ than its long photoperiod counterpart, but the difference between the two photoperiods

decreases as T cycle length increases. This pattern resembled the example in Figure 1C, with

the interaction providing an additional layer of complexity. In contrast to the multifaceted inputs to

ψ, we found that photoperiod was the sole significant influence on α, with persistently reduced α

in DD following long photoperiods compared to short photoperiods across all three T cycles

(Figure 2C). This pattern most closely resembled the example in Figure 1A, and suggested that

a parametric effect of the duration of light per cycle and/or a non-parametric effect of the phase

shifts caused at light onset and offset induced the after-effect in α. After-effects on locomotor τ

were significantly influenced by both T cycle length and photoperiod (Figure 2D), but unlike ψ,

there was no detectable interaction between the main effects. Within each T cycle, the long

photoperiod group had consistently shorter τ compared to its partner short photoperiod. Across T

cycles, however, there was a direct relationship with τ, with τ increasing along with T cycle length

regardless of photoperiod. This pattern most closely resembles the example in Figure 1C.

To further examine the relationship between the α and τ after-effects, we plotted the two

measurements against one another (Figure 3). The two factors correlated significantly in two

specific cases: T23 Long (Figure 3, top left) and T26 Short (Figure 3, bottom right). These two

photoperiod/T cycle combinations represent the instances where in previous reports T cycle and

photoperiod independently produce similar *in vivo* after-effects on α and τ, with long photoperiod

or short T decreasing α and τ, and short photoperiod or long T increasing α and τ. Interestingly,

there was a lack of correlation of these measures in T cycle/photoperiod combinations where the

predicted after-effects of the cycle τ and photoperiod would be in nominal conflict (Long

photoperiod and Long T, and Short photoperiod and Short T, Azzi et al., 2014; Pittendrigh and

Daan, 1976a).

ψ and α after-effects differ in their parametric vs. non-parametric responses

The after-effects described in Figure 2 for all parameters showed significant main effects

of photoperiod, revealing that either the interval between light transitions or the light duration is

inducing persistent effects on ψ, α, and τ. To determine whether the critical aspect of the light

cycle is the timing of the light/dark and dark/light transitions or the duration of light within each

cycle, we used skeleton photoperiods consisting of two 1-hour pulses of light, separated by

intervals of darkness, that mimic the timing of the light/dark transitions on full photoperiods of 16:8

and 8:16 (Figure 4A). Skeleton photoperiods have been used to assess the mechanism of

entrainment through seasonal changes in ψ (Pittendrigh and Minis, 1964), as well as to determine

the proximal effect on locomotor behavior duration (Pittendrigh and Daan, 1976a, 1976b). In order

to determine whether skeleton photoperiods can induce plasticity in α, τ, and ψ, we assessed

after-effects in locomotor behavioral rhythms in DD following 28 days of exposure to skeleton long

or short photoperiods in comparison to similar entrainment to full 16:8 and 8:16 photoperiods.

ψ exhibited main effects of both the duration of light in complete photoperiods

(Completeness, Figure 4B), and the timing of light transitions in skeleton photoperiods (Interval,

Figure 4B). The significant effect of the light duration of complete photoperiods indicates that there

is a parametric influence on the ψ of locomotor behavioral rhythms, in addition to the non-

parametric main effect of the interval between light transitions on skeleton photoperiods. In

contrast, we found a significant main effect only of the onset-offset interval of skeleton

photoperiods on α after-effects, suggesting that this effect is primarily non-parametric (Figure 4C).

We found that the complete and skeleton photoperiods had no significant after-effect effect on τ

(Figure 4D), however previous investigators have reported τ after-effects of complete and

skeleton photoperiods (Pittendrigh and Daan, 1976a). Thus we found that there are significant

parametric and non-parametric effects on ψ, non-parametric effects of onset-offset interval on α,

and non-significant after-effects on behavioral τ of both complete and skeleton photoperiods.

***Ex vivo* SCN after-effects**

Using *ex vivo* imaging of PER2::LUCIFERASE (PER2::LUC) rhythms following

entrainment to the same six light schedules described above, we investigated the effects of

photoperiod and T cycles on plasticity of the SCN pacemaker. We housed heterozygous

PER2::LUC animals in identical conditions to our complete photoperiod *in vivo* experiments,

including running wheels, and collected SCN slices from the animals after 14-28 days of exposure

to each of the light schedules described above. Slices were collected within 4 hours of lights off,

maintained in organotypic culture, and PER2::LUC luminescence recorded *ex vivo* for 5-7 days

(see Methods). Slice recordings were analyzed by detecting peak PER2::LUC expression in each

of a 102 x 102 grid of sub-cellular (~5 x 5 μm) regions of interest (ROIs) spread across the entire

SCN image. These PER2::LUC peak times were then used to calculate and map σ_ϕ (Figure 5A),

while the average PER2::LUC luminescence profile over time was used to calculate SCN τ.

Similar to the ψ of locomotor rhythms, we found that the ψ of SCN PER2::LUC rhythms

(relative to projected ZT12 of the previous light cycle) was significantly influenced by both T cycle

and photoperiod with a significant interaction between the two factors (Figure 5B). As T cycle

length increases, SCN ψ also becomes more advanced, and within each T cycle the long

photoperiod is more advanced than the short. Again, like the *in vivo* ψ, the *ex vivo* ψ featured a

narrowing difference between long and short as T cycle length increases. Similarly, mirroring *in*

vivo α, we found that the photoperiod modulated the degree of synchrony of PER2::LUC rhythms

within explanted SCN (σ_ϕ, Figure 5C), with increased phase dispersion in long photoperiods

compared to short. However, we also found a significant main effect of T cycle length on the SCN

σ_ϕ after-effect. This effect was driven by increased σ_ϕ in both T26 groups compared to both T24

groups (post-hoc multiple comparisons test between T cycle lengths, T24 vs. T26 adjusted p =

0.0329). There is no single consistent linear effect of T cycle on the *ex vivo* σ_ϕ after-effect. As

such, the pattern most resembled that of the example in Supplemental Figure 2A, but with the

T26 groups elevated.

SCN τ after-effects following T cycle entrainment have been a source of significant interest

in part because of the apparent disconnect between after-effects of T cycle (where *ex vivo* SCN

and behavioral measurements negatively correlate) and of photoperiod (where *ex vivo* SCN and

behavioral measurements agree). In our combined T cycle/photoperiod paradigm we found a

significant negative relationship between T cycle length and SCN τ length. Entrainment to short

T cycles *in vivo* resulted in lengthened SCN τ *ex vivo*, and *vice versa* as previously reported (Aton

et al., 2004; Azzi et al., 2017; Molyneux et al., 2008). Visually, there is an intriguing reversal in

trend direction of possible photoperiod effects between T24 and the two non-24 T cycles that

could indicate an interaction. In T24 cycles, the previously reported trend between long and short

photoperiod (shortened, behavior-like τ after-effect for long photoperiod) is present, while in T23

and T26 cycles, the trend between long and short is inverted. This relationship results in the

longest and shortest τ aligning with T23 Long and T26 Short, the same two groups highlighted

above (Figure 3) as overlap groups where the effects of T cycle and photoperiod are in line with

one another. However, no significant photoperiod effects or interactions were revealed by the

ANOVA, or by multiple post-hoc tests for differences within the T cycle groups, so these visual

trends require further testing.

***Ex vivo* SCN after-effects have characteristics similar to behavioral after-effects**

To investigate the relative influence of parametric versus non-parametric inputs on SCN

after-effects *ex vivo*, we collected slices from animals entrained to skeleton long and short

photoperiods (Figure 6A). Similar to *in vivo* ψ of the behavioral rhythm to skeleton photoperiods,

there was a significant effect of skeleton onset-offset interval on the ψ of the PER2::LUC rhythm

(Figure 6B). Unlike *in vivo*, the PER2::LUC ψ did not have a significant effect of photoperiod

completeness, although there was a trend towards a decrease in the difference in ψ between long

and short versions of skeleton photoperiods compared to the complete photoperiods. The σ_ϕ after-

effect was significantly influenced by the onset-offset interval (Figure 6C), similar to *in vivo*. The

σ_ϕ after-effect *ex vivo* was also significantly influenced by photoperiod completeness. However,

the mean difference in SCN σ_ϕ between the long and short versions of the skeleton photoperiods

was larger compared to that of the complete photoperiods. Though the interaction that this implies

was not statistically significant, it does suggest that skeleton interval is a principal driver of σ_ϕ

after-effects similar to *in vivo* α (Figure 4C). As with behavioral τ after-effects, we found no

significant effect of onset-offset interval or photoperiod completeness on the *ex vivo* τ after-effect,

again suggesting that the effects of photoperiod on τ after-effects are modest compared to T cycle

entrainment.

 DISCUSSION

Measurements in constant conditions reveal properties of the SCN and downstream

circadian outputs free of light input, and as such the study of after-effects is one means to

understand the plasticity of the SCN in response to seasonal light conditions. Understanding after-

effects is also useful *per se*, as the encoding of light information within the SCN allows seasonal

circadian properties to remain stable across ephemeral changes in the environment. Organisms

in the wild experience fluctuations in light exposure during the annual photoperiodic cycle because

of weather, nesting, hibernation, and other factors. Stable but flexible encoding of seasonal light

inputs within the circadian system would allow for consistency across these light exposure

fluctuations.

To improve our understanding of how after-effects are induced, we examined the relative

influences of T cycle length, photoperiod, and onset-offset interval on *in vivo* (α, τ) and *ex vivo*

(σ_ϕ, τ) after-effects, as well as on ψ (both *in vivo* and *ex vivo*). Using eight light cycle input groups,

including six T cycle/photoperiod combinations and two skeleton photoperiods, we tested which

specific components of light cycle input (light duration, onset-offset interval, repeated phase shifts)

are encoded by the SCN in a persistent manner by measuring after-effects in circadian behavior

and in SCN rhythms assayed *ex vivo*.

Comparison of *in vivo* and *ex vivo* responses

We found that ψ is strongly influenced by T cycle and photoperiod length, both *in vivo*

(Figure 2B) and *ex vivo* (Figure 5B), with a significant interaction between T cycles and

photoperiod in setting subsequent ψ in both cases. Using T24 skeleton photoperiods to assess

the influence of non-parametric (light onset-offset interval) and parametric (light interval

completeness) input, we found both input types significantly influence ψ *in vivo* (Figure 4B) while

observing significant influences of non-parametric input and a trend toward an influence of

parametric input *ex vivo* (Figure 6B). Determination of ψ has long been considered to be non-

parametric, with the exception of the continuity of light on full light cycles disambiguating onset

from offset on near symmetrical light cycles (e.g. near T24 12:12, Pittendrigh and Minis, 1964),

however our results suggest a more complex picture akin to the circadian surface entrainment

model of Roenneberg for *Neurospora* (Rémi et al., 2010; Roenneberg et al., 2010). Future

experiments may provide further information about the parametric and non-parametric influences

of T cycles and interval lengths through the use of skeleton photoperiods with non-24 hour T

cycles.

After-effects in α were strongly influenced by photoperiod length (Figure 2C) in a non-

parametric manner (Figure 4C). This result was mirrored by the significant main effect of

photoperiod on *ex vivo* SCN σ_ϕ (Figure 5C), which was also primarily non-parametric (Figure 6C).

One notable difference between *in vivo* α and *ex vivo* SCN σ_ϕ, however, was the general increase

in SCN σ_ϕ measured in both T26 photoperiods. The increase in the T26 Short group may be

related to the contracted α observed in LD during entrainment to T26 Long and Short (Figure 2A,

bottom row): Though the free-running α expands on the first day of DD, there is apparent negative

masking occurring in the early active phase of these mice during LD. Because slices were made

during the last LD cycle rather than on the first day of DD, the σ_ϕ could reflect that negatively-

masked pattern and imply that σ_ϕ is correlated with overt "net" α, rather than the underlying

assumed endogenous active phase length.

We found that behavioral τ after-effects are significantly influenced by both T cycle and

photoperiod length, but that the T cycle influence is of greater statistical significance and is the

source of a greater percentage of the overall variation (Figure 2D), suggesting that the T cycle

input to τ after-effects is more robust. Our results with T24 complete and skeleton photoperiods

in vivo support this notion, as we did not observe significant τ after-effects with either complete or

skeleton photoperiods (Figure 4D). Previous studies have described τ after-effects of complete

and skeleton photoperiods on behavioral τ in mice, but with modest differences between the two

skeleton photoperiod lengths (Pittendrigh and Daan, 1976a). While locomotor τ after-effects

reflected the length of the entraining T cycle, we observed negative correlations between the τ

after-effects of *ex vivo* SCN and the length of the entraining T cycle, consistent with previous

findings (Aton et al., 2004; Azzi et al., 2017; Molyneux et al., 2008). We did not detect any

statistically significant modulation of the T cycle effects on τ by varying photoperiod, but visually,

there are trend reversals in the expected direction of photoperiodic effects on τ in the non-24-hour

T cycle groups that suggest the possibility of interactions that we did not detect statistically (T24

Long vs. T23 Long, $p = 0.1513$; T24 Short vs. T26 Short, $p = 0.0584$; Sidak's Multiple

Comparisons, see further discussion below).

Differences between T cycle- and photoperiod-induced responses

Because α and τ after-effects have been observed together (shorter α, shorter τ; longer

α, longer τ), previous studies have attempted to explain both changes with a single underlying

mechanism (Beersma et al., 2017; Gu et al., 2016). We observed different patterns of responses

for after-effects in α and τ in photoperiod/T cycle combinations (Figure 2A, C-D) as well as in T24

photoperiods (Figure 4A, C-D), and correlation of changes in α and τ were specific to certain T

and photoperiod combinations (Figure 3). When measuring the correspondence of τ and α after-

effects *in vivo*, correlations were only observed in cases where T cycle and photoperiod effects

are expected to align (Figure 3). These results suggest that while the two after-effects may

frequently be observed together, their distinct patterning in response to particular T cycles and

photoperiods could be driven by disparate but conditionally convergent mechanisms.

Changes in SCN σ_ϕ remain a likely explanation for alterations in downstream locomotor

behavior duration, as an extended population-level high-firing phase of SCN neurons (for which

the PER2::LUC σ_ϕ is a proxy here) would logically underlie the regulation of timing activity or

inactivity (Ciarleglio et al., 2009, 2011; Inagaki et al., 2007; VanderLeest et al., 2007). Changes

in τ, however, shown here to be induced primarily through T cycle entrainment and likely the

requisite repeated phase shifts thereof, may be regulated in a different manner. Mechanisms

suggested recently by other groups (Azzi et al., 2014, 2017) surrounding the epigenetic changes

that may occur after extended entrainment to T cycles may help explain alterations in τ that we

observed.

Regional phase differences and their influence on after-effects

Regional phase differences within the SCN network have been observed in response to T

cycles and photoperiod, and have been proposed to drive τ after-effects (Azzi et al., 2017; Buijink

et al., 2016; Evans et al., 2013; Myung et al., 2015). In SCN explants from mice entrained to long

photoperiods, the ventral or ventromedial region SCN phase leads in some studies (Buijink et al.,

2016; Evans et al., 2013), but the reverse regional phase relationship was observed in another

(Myung et al., 2015). SCN τ shortening after entrainment to long photoperiods was observed to

correlate with ventral phase lead (Evans et al., 2013), with ventral phase lag (Myung et al., 2015),

or not be present (Buijink et al., 2016). Similarly, Azzi *et al.* (2017) found that following entrainment

to T cycles, there was an anticorrelation between SCN τ after-effect and the previous T cycle τ,

as expected from previous results (Aton et al., 2004; Molyneux et al., 2008). The same study also

found that shortened τ correlated with ventral phase lag, while lengthened τ correlated with ventral

phase lead – the opposite pattern previously found for photoperiodic τ after-effects by the same

group (Evans et al., 2013). Thus, there is no clear consensus in the literature as to whether SCN

regional phase differences mediate τ after-effects.

We also observed the anticorrelation of *ex vivo* SCN τ after T cycle entrainment *in vivo*,

as well as a variety of regional σ_ϕ across our experimental conditions that included examples of

ventral core phase lead in response to short T cycles (see T23 Long in Figure 6) similar to Azzi

et al. (2017), and examples of more ventromedial phase clustering similar to that of Buijink *et al.*

(2016) in response to long T cycles (see T26 Short in Figure 6). Differences in regional patterns

(ventral core vs. ventromedial) may be explained by slice position on the rostral caudal axis of the

SCN. Azzi et al. (2017) show that taking sections more caudal in the SCN changes the regional

phase pattern from ventral core grouping to a ventromedial grouping similar to Buijink (2016). We

speculate that we may not have controlled well enough for slice depth, resulting in our observation

of variations of both patterns in our data.

417   Conceptually, the anticorrelation of SCN *ex vivo* τ with *in vivo* τ after-effects of T cycles

presents a challenge. One interpretation of these results is that the observed *ex vivo* τ reflects a

deafferentation-induced reorganization of the SCN network, indicating extra-SCN input is key to

T cycle after-effects *in vivo* (Aton et al., 2004; Molyneux et al., 2008). Support for this view also

comes from genetic studies showing an anticorrelation between *in vivo* behavioral and *ex vivo*

SCN τ effects of genetic serotonergic de-afferentation of the SCN (Ciarleglio et al., 2014), and

lack of correlation of *in vivo* and *ex vivo* SCN period after-effects of genetic manipulations of SCN

gene expression (Mieda et al., 2015, 2016). In addition, part of the confound regarding previous

ex vivo SCN T-cycle after-effects appears to be a bias in the reporting of SCN regional rhythms

by the standard PER2::LUC reporter that more highly weights τ of the dorsal region (Azzi et al.,

2017). In any case, the lack of correspondence between *in vivo* and *ex vivo* T-cycle τ after-effects

in the SCN makes it difficult to attribute *ex vivo* findings of regional patterns or mechanisms to the

in vivo case, although the system can be used to study SCN *ex vivo* reorganization and its

mechanisms (Azzi et al., 2017).

SCN σ_ϕ and its relationship with α

The degree of SCN neuron σ_ϕ is a characteristic attribute of photoperiodic encoding, as it

represents a change in state of the SCN induced by the duration of the light interval of the daily

cycle and is maintained after transfer to constant conditions (Buijink et al., 2016; Ciarleglio et al.,

2011; Inagaki et al., 2007; VanderLeest et al., 2007). Our results suggest that the timing of

light/dark transitions is a key factor in influencing SCN σ_ϕ (Figure 6C) and *in vivo* α (Figure 4C),

although we also detected an effect of light duration on SCN σ_ϕ (Figure 6C). The spread in the

timing of neuronal rhythms allows for the daily duration of high firing activity of the SCN to be

broadened or contracted without required alteration to the waveforms of individual SCN neurons,

although this aspect is modulated by developmental inputs (Ciarleglio et al., 2011). As such, the

degree of σ_ϕ within the SCN is a potential target for artificial SCN manipulation – driving neurons

to adopt the σ_ϕ characteristic of a photoperiod, such as by pharmacologically, optogenetically, or

chemogenetically extending the high firing phase, would be expected to induce the behavioral α

characteristic of that particular photoperiod, and perhaps other photoperiod-dependent

responses, such as reproductive state and affective behaviors.

Because current methods for *in vivo* imaging are limiting, the measure of α after-effects

may be a useful proxy for inferring the waveform of the SCN. The integrity of this SCN waveform-

α connection depends on the correlation between the two. Our findings that α and *ex vivo* SCN

neuron σ_ϕ associate across multiple lighting conditions regardless of net daily phase shift do not

address causality, but strengthen the association between the two measurements. As the

broadened electrical waveform of the SCN is associated with the compressed behavioral α seen

in long photoperiods (and the contracted waveform with extended α in short photoperiods), the

spread in phase of SCN neurons and the corresponding widening of overall firing that occurs as

a result should account for the observed changes in α (Houben et al., 2009).

Changes to SCN network synchrony may have additional ramifications in the form of

altered responses to light input, potentially explaining previously observed larger phase shifts

after exposure to short days, and smaller phase shifts after exposure to long days (Refinetti,

2002). The role of SCN neuron σ_ϕ in mediating those altered responses and altered τ has been

explored computationally (Gu et al., 2016) and through altered GABA signaling (Farajnia et al.,

2014). In their analysis of after-effects on τ, Pittendrigh and Daan (1976b) suggested that this

change in circadian property may provide functional significance in terms of priming future

responses to light, but it remains to be seen whether any of these after-effects constitutes an

adaptive advantage or if they are simply a side effect of altered network synchrony.

464   **CONCLUSION**

Taken together, our results show that distinct components of light cycles can have distinct

effects on the circadian system, with (a) the timing of light/dark transitions in photoperiods driving

α plasticity; (b) combined photoperiod and T cycle length driving plasticity in behavioral τ and ψ

of both *in vivo* behaviors and *ex vivo* SCN gene expression; and (c) the light/dark transition interval

and long T cycle lengths affecting *ex vivo* SCN σ_ϕ. These results open the door to selective

manipulation of circadian plasticity in α or τ for further experimentation and for potential translation

to treat photoperiodic syndromes. The results shown here also demonstrate the utility of

deconvolving overlapping types of input when studying complex processes within the SCN. Our

results indicate that α and τ after-effects are possibly driven by distinct, though perhaps related,

mechanisms, and such information can be incorporated into experimental and computational

approaches to test the system in the future. With new techniques currently being established in

our field to measure SCN neuronal firing rate, calcium signaling, and activity levels with high

cellular specificity and extended timeframe, there will be an improved ability to characterize the

specific effect of photoperiodic influences on these circadian components.

ACKNOWLEDGMENTS

481 The authors would like to thank Maria Luísa Jabbur, Carl Johnson, Terry Page, and Chloe

482 Snider for their helpful comments and discussions. This work was supported by NIH R01

483 GM117650 to D.G.M., NIH F31 NS096813 to M.C.T., and NSF GRFP 0909667 to M.C.T.

AUTHOR CONTRIBUTIONS

486 M.C.T. and D.G.M conceived the experiments. M.C.T. performed the experiments. M.C.T. and

487 J.J.H. analyzed the data, M.C.T., J.J.H., and D.G.M. wrote the paper.

DECLARATION OF INTERESTS

489 The authors have no competing interests to report.

REFERENCES

Aton SJ, Block GD, Tei H, Yamazaki S and Herzog ED (2004) Plasticity of circadian behavior

and the suprachiasmatic nucleus following exposure to non-24-hour light cycles. *Journal of*

biological rhythms 19(3): 198–207.

Azzi A, Dallmann R, Casserly A, Rehrauer H, Patrignani A, Maier B, Kramer A and Brown SA

(2014) Circadian behavior is light-reprogrammed by plastic DNA methylation. *Nature*

neuroscience 17(3): 377–82.

Azzi A, Evans JA, Leise T, Myung J, Takumi T, Davidson AJ and Brown SA (2017) Network

Dynamics Mediate Circadian Clock Plasticity. *Neuron* 93(2): 441–450.

Bartness TJ and Wade GN (1984) Photoperiodic control of body weight and energy metabolism

in Syrian hamsters (Mesocricetus auratus): role of pineal gland, melatonin, gonads, and

diet. *Endocrinology* 114(2): 492–8.

Bartness TJ, Elliott JA and Goldman BD (1989) Control of torpor and body weight patterns by a

seasonal timer in Siberian hamsters. *The American journal of physiology* 257(1 Pt 2):

R142-9.

Basnet S, Merikanto I, Lahti T, Männistö S, Laatikainen T, Vartiainen E and Partonen T (2016)

Seasonal variations in mood and behavior associate with common chronic diseases and

symptoms in a population-based study. *Psychiatry research* 238: 181–188.

Beersma DGM, Gargar KA and Daan S (2017) Plasticity in the Period of the Circadian

Pacemaker Induced by Phase Dispersion of Its Constituent Cellular Clocks. *Journal of*

Biological Rhythms 32(3): 237–245.

Buijink MR, Almog A, Wit CB, Roethler O, Olde Engberink AHO, Meijer JH, Garlaschelli D,

Rohling JHT and Michel S (2016) Evidence for Weakened Intercellular Coupling in the

Mammalian Circadian Clock under Long Photoperiod. Yamazaki S (ed.) *PLOS ONE*

11(12): e0168954.

Ciarleglio CM, Gamble KL, Axley JC, Strauss BR, Cohen JY, Colwell CS and McMahon DG

(2009) Population Encoding by Circadian Clock Neurons Organizes Circadian Behavior.

Journal of Neuroscience 29(6): 1670–1676.

Ciarleglio CM, Axley JC, Strauss BR, Gamble KL and McMahon DG (2011) Perinatal

photoperiod imprints the circadian clock. *Nature Neuroscience* 14(1). Nature Publishing

Group: 25–27.

Ciarleglio CM, Resuehr HES, Axley JC, Deneris ES and McMahon DG (2014) Pet-1 Deficiency

Alters the Circadian Clock and Its Temporal Organization of Behavior. Mistlberger RE (ed.)

PLoS ONE 9(5): e97412.

Elliott JA and Goldman BD (1981) Seasonal Reproduction. In: Adler NT (ed.)
Neuroendocrinology of Reproduction. Boston, MA: Springer US, pp. 377–423.

Evans JA, Leise TL, Castanon-Cervantes O and Davidson AJ (2013) Dynamic interactions
mediated by nonredundant signaling mechanisms couple circadian clock neurons. *Neuron*
80(4): 973–983.

Farajnia S, van Westering TLE, Meijer JH and Michel S (2014) Seasonal induction of
GABAergic excitation in the central mammalian clock. *Proceedings of the National*
Academy of Sciences 111(26): 9627–9632.

Fernandez DC, Fogerson PM, Lazzerini Ospri L, Thomsen MB, Layne RM, Severin D, Zhan J,
Singer JH, Kirkwood A, Zhao H, Berson DM and Hattar S (2018) Light Affects Mood and
Learning through Distinct Retina-Brain Pathways. *Cell* 175(1): 71-84.e18.

Gu C, Rohling JHT, Liang X and Yang H (2016) Impact of dispersed coupling strength on the
free running periods of circadian rhythms. *Physical Review E* 93(3): 032414.

Gwinner E (1990) Circannual Rhythms in Bird Migration: Control of Temporal Patterns and
Interactions with Photoperiod. In: Gwinner E (ed.) *Bird Migration.* Berlin, Heidelberg:
Springer Berlin Heidelberg, pp. 257–268.

Houben T, Deboer T, van Oosterhout F and Meijer JH (2009) Correlation with Behavioral
Activity and Rest Implies Circadian Regulation by SCN Neuronal Activity Levels. *Journal of*
Biological Rhythms 24(6): 477–487.

Inagaki N, Honma S, Ono D, Tanahashi Y and Honma K (2007) Separate oscillating cell groups
in mouse suprachiasmatic nucleus couple photoperiodically to the onset and end of daily
activity. *Proc Natl Acad Sci USA* 104(18): 7664–7669.

Koorengevel KM, Beersma DG., den Boer J a and van den Hoofdakker RH (2003) Mood

regulation in seasonal affective disorder patients and healthy controls studied in forced

desynchrony. *Psychiatry Research* 117(1): 57–74.

Mickman CT, Stubblefield JS, Harrington ME and Nelson DE (2008) Photoperiod alters phase

difference between activity onset in vivo and mPer2::luc peak in vitro. *American Journal of*

Physiology-Regulatory, Integrative and Comparative Physiology 295(5): R1688–R1694.

Mieda M, Ono D, Hasegawa E, Okamoto H, Honma K ichi, Honma S and Sakurai T (2015)

Cellular clocks in AVP neurons of the scn are critical for interneuronal coupling regulating

circadian behavior rhythm. *Neuron* 85(5). Elsevier Inc.: 1103–1116.

Mieda M, Okamoto H and Sakurai T (2016) Manipulating the Cellular Circadian Period of

Arginine Vasopressin Neurons Alters the Behavioral Circadian Period. *Current Biology*.

Molyneux PC, Dahlgren MK and Harrington ME (2008) Circadian entrainment aftereffects in

suprachiasmatic nuclei and peripheral tissues in vitro. *Brain Research* 1228: 127–134.

Myung J, Hong S, DeWoskin D, De Schutter E, Forger DB and Takumi T (2015) GABA-

mediated repulsive coupling between circadian clock neurons in the SCN encodes

seasonal time. *Proceedings of the National Academy of Sciences* 112(29): E3920–E3929.

Oh E-Y, Ansell C, Nawaz H, Yang C-H, Wood PA and Hrushesky WJM (2010) Global breast

cancer seasonality. *Breast Cancer Research and Treatment* 123(1): 233–243.

Pittendrigh CS and Daan S (1976a) A functional analysis of circadian pacemakers in nocturnal

rodents (I). *Journal of Comparative Physiology A* 106(3). Springer-Verlag: 223–252.

Pittendrigh CS and Daan S (1976b) A functional analysis of circadian pacemakers in nocturnal

rodents (IV). *Journal of Comparative Physiology A* 106(3). Springer-Verlag: 291–331.

Pittendrigh CS and Minis DH (1964) The Entrainment of Circadian Oscillations by Light and

Their Role as Photoperiodic Clocks. *The American Naturalist* 98(902): 261–294.

Refinetti R (2002) Compression and expansion of circadian rhythm in mice under long and short

photoperiods. *Integrative Physiological & Behavioral Science* 37(2): 114–127.

Rémi J, Merrow M and Roenneberg T (2010) A Circadian Surface of Entrainment: Varying T, τ,

and Photoperiod in Neurospora crassa. *Journal of Biological Rhythms* 25(5): 318–328.

Roenneberg T, Rémi J and Merrow M (2010) Modeling a Circadian Surface. *Journal of*

Biological Rhythms 25(5): 340–349.

Schwartz WJ, Tavakoli-Nezhad M, Lambert CM, Weaver DR and de la Iglesia HO (2011)

Distinct patterns of Period gene expression in the suprachiasmatic nucleus underlie

circadian clock photoentrainment by advances or delays. *Proceedings of the National*

Academy of Sciences 108(41): 17219–17224.

Tackenberg MC and McMahon DG (2018) Photoperiodic Programming of the SCN and Its Role

in Photoperiodic Output. *Neural Plasticity* 2018: 1–9.

VanderLeest HT, Houben T, Michel S, Deboer T, Albus H, Vansteensel MJ, Block GD and

Meijer JH (2007) Seasonal Encoding by the Circadian Pacemaker of the SCN. *Current*

Biology 17(5): 468–473.

Wehr TA (2001) Photoperiodism in Humans and Other Primates: Evidence and Implications.

Journal of Biological Rhythms 16(4): 348–364.

Figure 1. Schematic of possible patterns of measurements across T23 L/S, T24 L/S, and T26 L/S.
A, a T-cycle dependent response where the measurement is graded depending on the T cycle
and photoperiod within the T-cycle has negligible effects. **B**, a photoperiod-dependent response
where light duty cycle dictates the measurement regardless of T cycle. **C**, a combination of
photoperiod and T cycle influences, with each of the 6 combinations having a characteristic
measurement level. Note that in all examples, the polarity of the measurement change is arbitrary
(e.g., T cycle response in A can increase or decrease with T cycle, photoperiod response in B
can have either Long or Short measure higher).These patterns are illustrations of individual and
combined main effects; possible interaction effects are not shown.

Figure 2. Behavioral after-effects of six photoperiod/T cycle combinations. **A**, representative actograms from long (left column) and short (right column) photoperiods combined with T23 (top row), T24 (middle row), and T26 (bottom row). **B**, ψ following exposure to the six light schedules. Values are relative to projected ZT12 (positive values advanced, negative values delayed). Interaction p = 0.0014 (7.242%), T Cycle Length p < 0.0001 (51.30%), Photoperiod Legth p < 0.0001 (20.97%). **C**, α after-effect across all six schedules as measured by automated α detection (see Methods). Interaction p = 0.4256 (2.081%), T Cycle Length p = 0.1648 (4.468%), Photoperiod Length p < 0.0001 (21.67%). **D**, period after-effect across the six schedules as measured by chi square periodogram. Interaction p = 0.7587 (0.2835%), T Cycle Length p < 0.0001 (56.17%), Photoperiod Length p = 0.0087 (3.778%). Two-way ANOVA results represent p value and percent variation (p < 0.0001 ****, p < 0.001 ***, p < 0.01 **, p < 0.05 *). T23 Long, n = 10; T23 Short, n = 8; T24 Long, n = 12; T24 Short, n = 12; T26 Long, n = 6; T26 Short, n = 14. T23 Long and Short groups are composed of a combination of two similar LD ratios (see Methods).

Figure 3. Within-individual correlations between τ and α after-effects following the given T-
cycle/photoperiod combination. T23 Long 0.8238 (p = 0.0034, **), T24 Long -0.06835 (p = 0.8328, ns),
T26 Long 0.004966 (p = 0.9926, ns), T23 Short -0.3147 (p = 0.4477, ns), T24 Short 0.2671 (p = 0.4013,
ns), T26 Short 0.5818 (p = 0.0291, *); Values reported are Pearson r and p value (p < 0.0001 ****, p <
0.001 ***, p < 0.01 **, p < 0.05 *).

Figure 4. Behavioral after-effects following exposure to skeleton long and short photoperiods compared to
T24 complete long and short photoperiods. **A**, representative actograms from long (top) and short (bottom)
skeleton photoperiods. **B**, ψ following exposure to skeleton long and short photoperiods. Values are relative
to projected ZT12 (positive values advanced, negative values delayed). Interaction $p = 0.3161$ (1.33%),
Completeness $p = 0.0003$ (21.24%), Onset-Offset Interval $p < 0.0001$ (29.32%). **C**, α after-effect of skeleton
long and short photoperiods. Interaction $p = 0.1861$ (4.332%), Completeness $p = 0.9858$ (0.0008%), Onset-
Offset Interval $p = 0.0376$ (11.15%). **D**, τ after-effect of skeleton long and short photoperiods. Interaction p
$= 0.7835$ (0.2063%), Completeness $p = 0.7006$ (0.4044%), Onset-Offset Interval $p = 0.0765$ (8.988%). Long
and Short complete photoperiod values in B-D are replotted from Figure 2. Two-way ANOVA results
represent p value and percent variation ($p < 0.0001$ ****, $p < 0.001$ ***, $p < 0.01$ **, $p < 0.05$ *). Complete
Long, n = 12; Complete Short, n = 12; Skeleton Long, n = 7, Skeleton Short, n = 6.

Figure 5. PER2::LUC after-effects following exposure to six photoperiod/T cycle combinations. **A**,
representative relative phase maps from long (left column) and short (right column) photoperiod combined
with T23 (top row), T24 (middle row), or T26 (bottom row). **B**, ψ of the PER2::LUC rhythm following
entrainment to the specified T cycle/photoperiod combination. Values represent the time of the first
PER2::LUC peak relative to the projected lights off for the first cultured cycle (projected ZT 12). Interaction
$p = 0.0052$ (8.185%), T Cycle Length $p < 0.0001$ (49.47%), Photoperiod Length $p < 0.0001$ (24.94%). **C**,
σ_ϕ (median absolute deviation of PER2::LUC peak times) across SCN slices from animals entrained to
each of the specified light schedules. Interaction $p = 0.7168$ (1.433%), T Cycle Length $p = 0.0353$ (15.89%),
Photoperiod Length $p = 0.0114$ (15.42%). **D**, τ of the PER2::LUC rhythm over 5 cycles *ex vivo*. Interaction
$p = 0.4104$ (4.109%), T Cycle Length $p = 0.0111$ (23.40%), Photoperiod Length $p = 0.4151$ (1.529%). Two-
way ANOVA results represent p value and percent variation ($p < 0.0001$ ****, $p < 0.001$ ***, $p < 0.01$ **, $p <$
0.05 *). T23 Long, n = 6; T23 Short, n = 4; T24 Long, n = 6; T24 Short, n = 8; T26 Long, n = 7; T26 Short,
n = 6. T23 Long is composed of a combination of two similar LD ratios (see Methods).

Figure 6. PER2::LUC after-effects following entrainment to skeleton long and short photoperiods compared to T24 complete long and short photoperiods. **A**, representative relative phase maps from skeleton long (top) and short (bottom) photoperiod. **B**, ψ of the PER2::LUC rhythm following entrainment to skeleton long and short photoperiods. Values represent the time of the first PER2::LUC peak relative to the projected lights off for the first cultured cycle (projected ZT 12). Interaction $p = 0.7514$ (0.1326%), Completeness $p = 0.1660$ (2.651%), Onset-Offset Interval $p < 0.0001$ (70.79%). **C**, σ_ϕ (median absolute deviation of PER2::LUC peak times) across SCN slices from animals entrained to skeleton long and short photoperiod. Interaction $p = 0.0654$ (7.695%), Completeness $p = 0.0318$ (10.77%), Onset-Offset Interval $p = 0.0003$ (37.26%). **D**, τ of the PER2::LUC rhythm over 5 cycles *ex vivo*. Interaction $p = 0.4867$ (2.277%), Completeness $p = 0.5608$ (1.587%), Onset-Offset Interval $p = 0.6959$ (0.7131%). Long and Short complete photoperiod values in B-D are replotted from Figure 5. Two-way ANOVA results represent p value and percent variation ($p < 0.0001$ ****, $p < 0.001$ ***, $p < 0.01$ **, $p < 0.05$ *). Complete Long, n = 6; Complete Short, n = 8; Skeleton Long, n =6, Skeleton Short, n = 5.

Supplementary Figures

Supplementary Figure 1. Mean period values for each interpeak interval for the first five cycles recorded *ex vivo*. Error bars represent standard error of the mean.

660   **Supplementary Figure 2**. Phase distribution values calculated each cycle for the first three
661   cycles *ex vivo*. Error bars represent standard error of the mean.