The synergistic actions of hydrolytic genes reveal the mechanism of *Trichoderma harzianum* for cellulose

degradation

Déborah Aires Almeida[1,2], Maria Augusta Crivelente Horta[1,2,3], Jaire Alves Ferreira Filho[1,2], Natália Faraj Murad[1]

and Anete Pereira de Souza[1,4,*]

[1]Center for Molecular Biology and Genetic Engineering (CBMEG), University of Campinas (UNICAMP),

Campinas, SP, Brazil

[2]Graduate Program in Genetics and Molecular Biology, Institute of Biology, UNICAMP, Campinas, SP, Brazil

[3]Holzforschung München, TUM School of Life Sciences Weihenstephan, Technical University of Munich, Freising,

85354, Germany

[4]Department of Plant Biology, Institute of Biology, UNICAMP, Campinas, SP, Brazil

*Corresponding author

Prof Anete Pereira de Souza

E-mail: anete@unicamp.br

ORCID

0000-0002-5095-5177, 0000-0003-2159-5296, 0000-0001-9690-7704, 0000-0001-7652-2567, 0000-0003-3831-

9829

Abstract

Bioprospecting genes and proteins related to plant biomass degradation is an attractive approach for the

identification of target genes for biotechnological purposes, especially genes with potential applications in the

biorefinery industry that can enhance second-generation ethanol production technology. *Trichoderma harzianum* is a

potential candidate for cellulolytic enzyme prospection and production. Herein, the enzymatic activities,

transcriptome, exoproteome, and coexpression networks of the *T. harzianum* strain CBMAI-0179 were examined

under biomass degradation conditions. We used RNA-Seq to identify differentially expressed genes (DEGs) and

carbohydrate-active enzyme (CAZyme) genes related to plant biomass degradation and compared them with the

genes of strains from congeneric species (*T. harzianum* IOC-3844 and *T. atroviride* CBMAI-0020). *T. harzianum*

CBMAI-0179 harbors strain- and treatment-specific CAZyme genes and transcription factors. We detected

important proteins related to biomass degradation, including β-glucosidases, endoglucanases, cellobiohydrolases,

lytic polysaccharide monooxygenases, endo-1,4-β-xylanases and β-mannanases, in the exoproteome under

cellulose growth conditions. Coexpression networks were constructed to explore the relationships among the genes

and corresponding secreted proteins that act synergistically for cellulose degradation. An enriched cluster with

degradative enzymes was described, and the subnetwork of CAZymes revealed strong correlations among the

secreted proteins (AA9, GH6, GH10, GH11 and CBM1) and differentially expressed CAZyme genes (GH45, GH7,

AA7 and GH1). Our results provide valuable information for future studies on the genetic regulation of plant cell

wall-degrading enzymes. This knowledge can be exploited for the improvement of enzymatic reactions in biomass

degradation for bioethanol production.

Keywords: *Trichoderma harzianum*, Cellulose, RNA-Seq, CAZymes, Exoproteome, Coexpression networks

Key points

Different biotechnological approaches were used to understand the mechanism of cellulose degradation of

Trichoderma spp.

T. harzianum CBMAI-0179 is a potential candidate for the production of cellulolytic enzymes.

Coexpression networks revealed genes and proteins acting synergistically for cellulose hydrolysis.

Introduction

The expanding worldwide demand for renewable and sustainable energy sources has increased the interest

in alternative energy sources, and the production of second-generation biofuels seems to be a viable option to

confront these issues (dos Santos Castro et al. 2014). Lignocellulosic biomass is the most abundant renewable

organic carbon resource on earth, consisting of three major polymers, cellulose, hemicellulose, and lignin (Bayer et

al. 2007; dos Santos Castro et al. 2014). However, due to its recalcitrant characteristics, degrading this complex

matrix requires a variety of enzymes acting in synergy. Interactions between different enzymes have been

investigated to identify optimal combinations and ratios of enzymes for efficient biomass degradation, which are

highly dependent on the properties of the lignocellulosic substrates and the surface structure of cellulose microfibrils

(van Dyk and Pletschke 2012; Wang et al. 2016). Microorganisms are considered natural producers of enzymes, and

many members of both bacteria and fungi have evolved to digest lignocellulose (Ahmed et al. 2017).

Filamentous fungi, including the genera *Trichoderma, Aspergillus*, *Penicillium* and *Neurospora*, produce

extracellular proteins capable of degrading plant cell walls and are widely used in the enzymatic industry (Miao et

al. 2015). Species in the filamentous ascomycete genus *Trichoderma* are among the most commonly isolated

saprotrophic fungi and are important from a biotechnological perspective (Horta et al. 2018). Most species from this

genus are used in agriculture as biocontrol agents due to their ability to antagonize plant-pathogenic fungi,

described as mycoparasites, and in industry as producers of hydrolytic enzymes (Druzhinina et al. 2011; Kubicek et

al. 2011). Within the *Trichoderma* genus, *T. reesei* is the most intensively studied species and a well-known

producer of cellulase and hemicellulase, which are widely employed in industry (Druzhinina and Kubicek 2017;

Martinez et al. 2008; Peterson and Nevalainen 2012). However, studies on *T. harzianum* strains have shown their

potential to produce different sets of enzymes that can degrade lignocellulosic biomass (Benoliel et al. 2013;

Delabona et al. 2012, 2013; Horta et al. 2014); therefore, *T. harzianum* strains are being investigated as potentially

valuable sources of new industrial cellulases (Rocha et al. 2016).

The high degree of genetic variation observed within *Trichoderma* species and strains (Al-Sadi et al. 2015;

Kubicek et al. 2019) makes it necessary to explore the genetic background of potential *Trichoderma* strains. In the

present study, we compared the transcriptome and exoproteome of the potential hydrolytic strain *T. harzianum*

CBMAI-0179 with those of *T. harzianum* IOC-3844 and *T. atroviride* CBMAI-0020. Coexpression networks

provided novel information about the synergistic action of the hydrolytic enzymes of *T. harzianum* CBMAI-

0179. Our findings provide insights into the genes/proteins that act synergistically in plant biomass conversion

and can be exploited to improve enzymatic hydrolysis, thereby increasing the efficiency of the saccharification

of lignocellulosic substrates for bioethanol production.

Materials and methods

Fungal strains, fermentation and enzymatic activities

The species originated from the Brazilian Collection of Environment and Industry Microorganisms

(CBMAI, Campinas, SP, Brazil) and the Institute Oswaldo Cruz (IOC, Rio de Janeiro, RJ, Brazil). *T. harzianum*

CBMAI-0179 (Th0179), *T. harzianum* IOC-3844 (Th3844) and *T. atroviride* CBMAI-0020 (Ta0020) strains were

cultivated on solid medium for 8 days at 37 °C to produce a sufficient number of spores as an inoculum for

fermentation, as described in a previous study (Horta et al. 2018). The fermentation process was initiated with the

inoculation of 10^7 spores/mL in an initial volume of 200 mL of Mandels Andreotti (MA) minimal medium (Mandels

and Andreotti 1978), and it was performed in biological triplicates using crystalline cellulose (Celuflok, São Paulo,

Brazil; degree of crystallinity, 0.72 g/g; composition, 0.857 g/g cellulose and 0.146 g/g hemicellulose) or glucose

(control condition) as the unique carbon source. The fermentation process continued for 96 h, and then, the

supernatant and RNA samples were collected (Horta et al. 2018).

Supernatants were collected to measure the enzymatic activity (Fig. S1) and determine the exoproteome

profile. Xylanase and β-glucosidase activities were determined using the methods described by Bailey and Poutanen

(1989) and Zhang et al. (2009), respectively. Cellulase activity was determined using the filter paper activity (FPA)

test according to Ghose (1987). Protein levels were measured based on the Bradford (1976) method.

RNA extraction

Mycelial samples from both conditions were extracted from Th0179, Th3844 and Ta0020, stored at -80 °C,

ground in liquid nitrogen using a mortar and pestle, and used for RNA extraction using the LiCl RNA extraction

protocol (Oliveira et al. 2015).

Library construction and RNA-Seq

RNA samples were quantified using a NanoDrop 8000 (Thermo Scientific, Wilmington, DE, USA). Paired-

end libraries were constructed using 1 µg of each RNA sample obtained from the mycelial samples and the TruSeq

RNA Sample Preparation Kit v2 (Illumina 2014) (Illumina Inc., San Diego, CA, USA) according to the

manufacturer's instructions. The expected target sizes were confirmed using a 2100 Bioanalyzer (Agilent

Technologies, Palo Alto, CA, USA) and the DNA 1000 Kit, and the libraries were quantified by qPCR using the

KAPA library quantification Kit for Illumina platforms (Kapa Biosystems, Wilmington, MA, USA). The average

insertion size was 260 bp. The sequencing was carried out on the HiSeq 2500 platform (Illumina, San Diego, CA,

USA) according to the manufacturer's specifications for paired-end reads of 150 bp.

Data sources

The reads were deposited into the Sequence Read Archive (SRA) database in the National Center for

Biotechnology Information (NCBI) under BioProject number PRJNA336221. The nucleotide and protein sequences

of *T. harzianum* T6776 (PRJNA252551) (Baroncelli et al. 2015) and *T. atroviride* IMI 206040 (PRJNA19867)

(Kubicek et al. 2011) were used as a reference for transcriptome mapping and downloaded from the NCBI database

(www.ncbi.nlm.nig.gov).

Phylogenetic analysis of* Trichoderma *spp.

Amplification of the internal transcribed spacer (ITS) region from the genomic DNA of the strains under

study was conducted via PCR. Multiple sequence alignment was performed using ClustalW (Thompson et al. 1994),

and a phylogenetic tree was created using the Molecular Evolutionary Genetics Analysis v7.0 (MEGA7) program

(Kumar et al. 2016). The maximum likelihood (ML) (Jones et al. 1992) method of inference was used based on the

Kimura two-parameter (K2P) model (Kimura 1980). We used 1,000 bootstrap replicates (Felsenstein 1985) for each

analysis. The trees were visualized and edited using the FigTree program v1.4.3

(http:/tree.bio.ed.ac.uk/software/figtree/). The phylogenetic tree showed a high phylogenetic proximity between

Th0179 and Th3844 (with bootstrap support of 92%), while Ta0020 shared closer affinity with *T. atroviride* CBS

142.95 (Fig. S2) (Rosolen et al. 2020).

Transcriptome analysis

The quality of the sequencing reads was assessed with FastQC v0.11.5 (Andrews 2010), and the trimming with a

sliding window of size 4, minimum quality of 15, and length filtering (minimal length of 36 bp) was performed with

Trimmomatic v0.36 (Bolger et al. 2014). The RNA-Seq data were analyzed using CLC Genomics Workbench

software (v6.5.2; CLC bio, Finlandsgade, Denmark) (CLC Genomics Workbench 2016). The reads were mapped

against the reference genomes of *T. harzianum* T6776 (Baroncelli et al. 2015) and *T. atroviride* IMI 206040

(Kubicek et al. 2011) using the following parameters: minimum length fraction = 0.5; minimum similarity fraction =

0.8; and maximum number of hits for a read = 10. For the paired settings, the parameters were minimum distance =

150 and maximum distance = 300, including the broken pairs. The gene expression values were normalized in

transcripts per million (TPM) (Conesa et al. 2016). To statistically analyze the differentially expressed genes

(DEGs), fold change ≥ 1.5 or ≤ -1.5 and p-value ≤ 0.05 were applied. Venn diagrams were constructed to compare

the genes with TPM expression values greater than zero under both conditions from all strains

(http://bioinformatics.psb.ugent.be/webtools/Venn/).

Transcriptome annotation and CAZyme determination

Sequences were functionally annotated according to the Gene Ontology (GO) terms (Ashburner et al. 2000)

with Blast2Go v4.1.9 (Conesa et al. 2005) using BLASTx-fast and a cutoff e-value of 10^{-6}. Information derived from

the CAZy database (Lombard et al. 2014) was downloaded (www.cazy.org) to locally build a CAZy database. The

protein sequences of *T. harzianum* T6776 and *T. atroviride* IMI 206040 were used as queries in basic local

alignment search tool (BLAST) searches against the locally built CAZy database. BLAST matches showing an e-

value less than 10^{-11}, identity greater than 30% and queries covering greater than 70% of the sequence length were

selected and classified according to the CAZyme catalytic group as follows: glycoside hydrolases (GHs),

carbohydrate-binding modules (CBMs), glycosyltransferases (GTs), carbohydrate esterases (CEs), auxiliary

activities (AAs) or polysaccharide lyases (PLs). CAZymes were also annotated according to their Enzyme

Commission (EC) number (Yamanishi et al. 2009) through the Braunschweig Enzyme Database (BRENDA)

(Schomburg et al. 2017) (www.brenda-enzymes.org) using BLASTp with an e-value cutoff of 10^{-10}, identity greater

than 30% and queries covering greater than 60% of the sequence length.

Coexpression networks

For the coexpression networks of Th0179, TPM expression values were used to perform the analyses. The

network was assembled by calculating Pearson's correlation coefficient for each pair of genes. Genes showing null

values for most of the replicates under different experimental conditions were excluded. The cluster analysis

procedure was performed with the heuristic cluster chiseling algorithm (HCCA), and the highest reciprocal rank

(HRR) (Mutwil et al. 2011; Mutwil et al. 2010) was used to empirically filter the edges, retaining edges with a HRR

$\leq$ 30. Cytoscape software v3.6.0 (Shannon et al. 2003) was used for network visualization.

Exoproteome analysis

The analysis of the exoproteome of Th0179 under both fermentative conditions was performed via liquid

chromatography tandem mass spectrometry (LC-MS/MS) using the data-independent method of acquisition MS^E as

described by Horta et al. (2018). The LC-MS/MS data were processed using ProteinLynx Global Server (PLGS)

v3.0.1 software (Waters, Milford, MA, USA), and the proteins were identified by comparison to the *Trichoderma*

sequence database available in the UniProt Knowledgebase (UniProtKB; https://www.uniprot.org/uniprot/).

BLASTp searches of the fasta sequences of the identified proteins were performed against the *T. harzianum* T6776

genome to identify the secreted proteins and compare them to the transcriptome.

RT-qPCR analysis

To verify the reliability and accuracy of the transcriptome data, reverse transcription-quantitative PCR (RT-

qPCR) was performed to validate the differential expression results. The RNeasy Mini Kit (Qiagen, Hilden,

Germany) was used for RNA extraction, and cDNA was synthesized using the QuantiTect Reverse Transcription Kit

(Qiagen, Hilden, Germany). Primers were designed using the Primer3Plus (Untergasser et al. 2007); the target

amplicon sizes ranged from 120-200 bp, with an optimal annealing temperature of 60 °C and an optimal primer

length of 20 bp.

Quantification of gene expression was performed by continuously monitoring SYBR Green fluorescence.

The reactions were performed in triplicate in a total volume of 6.22 μL. Each reaction contained 3.12 μL of SYBR

Green Supermix (Bio-Rad, Hercules, CA, USA), 1.0 μL of forward and reverse primers and 2.1 μL of diluted

cDNA. The reactions were assembled in 384-well plates. PCR amplification-based expression profiling of the

selected genes was performed using specific endogenous controls for each strain (Table S1). RT-qPCR was

conducted with the CFX384 Touch Real-Time PCR Detection System (Bio-Rad). Gene expression was calculated

via the delta-delta cycle threshold method (Livak and Schmittgen 2001). The obtained RT-qPCR results were

compared with the RNA-Seq results. The fold changes of the selected genes exhibited the same expression profiles

between the RT-qPCR and RNA-Seq analyses (Fig. S3).

Results

Transcriptome analysis of **Trichoderma** *spp. under cellulose growth conditions*

The present study represents the first deep genetic analysis of Th0179, describing and comparing the

transcriptome under two different growth conditions, cellulose and glucose, to identify the genes involved in plant

biomass degradation. Reads were mapped against reference genomes of *T. harzianum* (PRJNA252551) and *T.*

atroviride (PRJNA19867), generating 96.3, 111.8 and 133.3 million paired-end reads for Th0179, Th3844 and

Ta0020, respectively. To establish the degrees of similarity and difference in gene expression among strains and

between treatments, a principal component analysis (PCA) was performed using the *T. harzianum* T6776 genome as

a reference. The PCA results showed clustered groups with higher similarity between treatments than among strains.

The transcriptomes of Th0179 and Th3844 were more similar to each other than to the transcriptome of Ta0020

(Fig. 1a), showing that it is possible to capture differences among the three strains.

Venn diagrams of the genes exhibiting expression levels greater than zero were constructed based on the

similarities among the genes from all strains, showing strain-specific genes expressed under both conditions (Fig.

S4). The number of genes shared by Th0179 and Th3844 was higher under both conditions than that shared by

either Th0179 or Th3844 and Ta0020. Th0179 exhibited the highest number of unique expressed genes, with 374

and 168 unique genes under the cellulose and glucose growth conditions, respectively, in which major facilitator

superfamily (MFS) and ATP binding cassette (ABC) transporters and transcription factors were found (Table S2).

Among the genes that were upregulated in cellulose conditions relative to glucose conditions, 219 were identified

for Th0179, 281 were identified for Th3844, and 718 were identified for Ta0020 (Fig. 1b and Table S3).

CAZyme identification and distribution in **Trichoderma** *spp.*

The set of CAZymes was identified by mapping all of the protein sequences from *T. harzianum* T6776 and

T. atroviride IMI 206040 against the CAZy database. Based on the filtering criteria, a total of 631 proteins were

retained as CAZyme genes for *T. harzianum*, which corresponds to 5.5% of the total 11,498 proteins predicted for

this organism (Baroncelli et al. 2015), and 640 proteins were retained for *T. atroviride*, which corresponds to 5.4%

of the total 11,816 proteins predicted for this organism (Kubicek et al. 2011).

Considering the DEGs under cellulose growth conditions, we found 35, 78 and 31 differentially expressed

CAZyme genes for Th0179, Th3844 and Ta0020, respectively (Fig. 1b). The contents of the main CAZyme classes

(AA, GH, GT, CE and CBM) were identified (Fig. 1c), and the differences regarding the specific CAZyme families

for each strain are shown in Fig. 2. The classification of the CAZyme genes along with their enzyme activities, fold

change values, e-values, EC numbers and TPM expression values for Th0179, Th3844 and Ta0020 under cellulose

fermentative conditions are described in Table S4.

The expression levels of the main cellulase and hemicellulase families based on their principal enzyme

activity present in Th0179, Th3844 and Ta0020 were evaluated using the transcriptomic data (Fig. 3). GHs related

to cellulose degradation were found in Th3844, including two AA9 members with a CBM1 module for cellulose

binding. All GHs except GH12, including AA9/CBM1, were detected in Th0179. Only one gene belonging to the

GH5 family (TRIATDRAFT_81867 – 48.12 TPM) with cellulase activity was observed in Ta0020. The most highly

expressed genes were cellobiohydrolases from the GH6/CBM1 (THAR02_04414 – 299.66 TPM for Th0179 and

986.25 TPM for Th3844) and GH7/CBM1 families (THAR02_08897 – 369.92 TPM for Th0179; THAR02_03357

and THAR02_08897 – 1876.89 TPM for Th3844) (Fig. 3a).

In addition, families related to hemicellulose degradation were identified in Th3844, in which endo-1,4-β-

xylanases from the GH11 family had the greatest expression levels (THAR02_02147, THAR02_05896,

THAR02_08630 and THAR02_08858 – 1013.62 TPM). Endo-1,4-β-xylanase from the GH10 family was found in

both *T. harzianum* strains (THAR02_03271 – 47.58 TPM for Th0179 and 741.35 TPM for Th3844). Acetylxylan

esterases from the CE5/CBM1 family were only found in Th3844 (THAR02_01449 and THAR02_07663 – 227.56

TPM). We identified only one α-L-arabinofuranosidase from the GH54/CBM42 family (TRIATDRAFT_81098 –

18.07 TPM) in Ta0020, whereas for Th3844, three α-L-arabinofuranosidases from the GH54/CBM42 and GH62

families were identified (Fig. 3b). Despite the higher number of DEGs, the Ta0020 strain showed the lowest number

and expression levels of CAZyme genes related to cellulose and hemicellulose degradation.

Functional annotation of T. harzianum *CBMAI-0179 in the presence of cellulose*

The first functional annotation of the expressed genes under cellulose fermentative conditions for Th0179

was performed (Fig. 4). A total of 7,718 genes were annotated, which corresponds to 67.1% of the total genes

predicted for the reference strain *T. harzianum* T6776 (Baroncelli et al. 2015). Compared with Th3844, Th0179

presented 163 more genes related to catalytic activity, 53 more genes related to hydrolase activity, and 15 more

genes associated with TF activity but 10 fewer genes related to transmembrane transport, suggesting that these two

strains have developed different functional regulation. Ta0020 presented fewer genes related to any of the above

functions than Th3844 and Th0179 except for TF activity, for which Ta0020 had 53 more genes than Th0179.

Exoproteome and RNA-Seq data correlation of T. harzianum *CBMAI-0179*

Once the transcriptome was characterized, we analyzed the secreted proteins identified in the exoproteome

profile of Th0179. We identified 32 proteins in the cellulose aqueous extract, 12 only in the glucose aqueous extract,

and 20 under both conditions (Table S5). Among the 32 secreted proteins detected using cellulose as the carbon

source, the main CAZyme families observed were β-glucosidases (EC 3.2.1.21), endo-β-1,4-glucanases (EC

3.2.1.4), lytic polysaccharide monooxygenases (LPMOs) (EC 3.2.1.4), cellobiohydrolases (EC 3.2.1.91), endo-1,4-

β-xylanases (EC 3.2.1.8), and β-mannanases (EC 3.2.1.78). Two genes from the GH3 (THAR02_00656 and

THAR02_00890) and GH5/CBM1 (THAR02_04405 and THAR02_09719) families and one gene from the

AA9/CBM1 (THAR02_02134) and GH6/CBM1 (THAR02_04414) families corresponded to the main group of

secreted cellulases, whereas one gene from the GH10 (THAR02_03271) family and two genes from the GH11

(THAR02_02147 and THAR02_05896) family corresponded to the main group of hemicellulases detected in the

supernatant. In addition, a member of the GH5 cellulase family with β-mannanase activity (THAR02_03851) was

identified, and 8 other proteins were classified as uncharacterized proteins. We also identified hemicellulases in both

extracts, such as α-L-arabinofuranosidase B (EC 3.2.1.55) and xylan 1,4-β-xylosidase (EC 3.2.1.37).

Correlating the exoproteome with the transcriptome data under cellulose growth conditions, we observed

CAZyme genes showing high TPM values in the cellulose condition, including cellulose 1,4-β-cellobiosidase (EC

3.2.1.91) from the GH6/CBM1 family (299.66 TPM), cellulase (EC 3.2.1.4) from the AA9/CBM1 family (138.62

TPM) and endo-1,4-β-xylanase (EC 3.2.1.8) from the GH11/CBM1 family (119.56 TPM). Two uncharacterized

proteins (THAR02_02133 and THAR02_08479) with the CBM1 module of cellulose binding also showed increased

expression levels under the cellulose condition. In contrast, the THAR02_00656 gene, which displays β-glucosidase

(EC 3.2.1.21) activity, from the GH3 family had the lowest expression level (5.08 TPM) among the CAZyme genes,

indicating that genes with low expression levels are also important for functional secreted proteins (Alfaro et al.

2016).

Coexpression networks for **T. harzianum** *CBMAI-0179*

279   The coexpression network was assembled for Th0179 and included all genes obtained from the mapping

results using the *T. harzianum* T6776 genome as a reference under both conditions. The DEGs and CAZyme genes

were highlighted in the network (Fig. 5a), composed of a total of 11,104 nodes with 153,893 edges. We detected 219

genes corresponding to the DEGs under cellulose growth conditions and 367 DEGs under glucose growth

conditions. Differentially expressed CAZyme genes from both conditions were also identified in this network, with

35 and 28 CAZymes under cellulose and glucose growth conditions, respectively. The DEGs in glucose clustered on

the top while DEGs in cellulose clustered on the bottom part of the network, reflecting the different regulation of

degradative activities according to the substrate.

287   A subnetwork was generated based only on the secreted proteins that were present in the cellulose aqueous

extract and that had corresponding genes in the coexpression network (Fig. 5b). This subnetwork exclusively

represented the genes and their closest related genes that are correlated with the secreted proteins. It was composed

of 713 nodes and 6,124 edges, and includes the 32 genes that encode the secreted proteins. The CAZyme genes

found in this subnetwork included the GH1 family (THAR02_02251 and THAR02_05432) with β-glucosidase

activity, the GH7/CBM1 family (THAR02_08897) with cellulose 1,4-β-cellobiosidase activity, and the

GH45/CBM1 family (THAR02_02979) with cellulase activity. Despite the different functions of the correlated

genes, it is predicted that these genes participate in the genetic regulation of the detected CAZymes/proteins and are

important to the regulation of the hydrolytic system.

296   The cluster analysis classified 11,102 genes in 84 clusters (Table S6). We identified an enriched cluster

composed of 196 nodes and 2,125 edges (Fig. 5c and Table S6) containing the greatest number of CAZyme genes

among the clusters. Of the 12 CAZyme genes detected, 7 corresponded to the secreted proteins detected in the

cellulose aqueous extract, and 5 were differentially expressed. In addition, we identified 20 DEGs under cellulose

growth conditions and 119 uncharacterized proteins in this cluster, showing a strong correlation with the CAZyme

genes, which indicates unknown genes that are important to the degradation process. The CAZyme genes from the

cluster analysis were selected to generate a new subnetwork with their corresponding edges (Fig. 5d), showing linear

correlations among secreted proteins (AA9, GH6, GH10, GH11 and CBM1) and differentially expressed CAZyme

genes (GH45, GH7, AA7 and GH1).

Discussion

307   In this study, different biotechnological approaches were used in bioprospecting new and efficient enzymes

for possible applications in the enzymatic hydrolysis process. We performed the enzymatic activity, transcriptome,

exoproteome and coexpression network analyses of the Th0179 strain, which has potential for plant biomass

degradation under cellulose growth conditions, to gain insight into the genes and proteins produced that are

associated with cellulose hydrolysis. Furthermore, we compared the expression levels with *Trichoderma* spp. that

have been previously studied under the same conditions (Horta et al. 2018) to understand how the regulation varies

between hydrolytic strains (*T. harzianum* CBMAI-0179 and *T. harzianum* IOC-3844) and a common mycoparasitic

ancestor species (*T. atroviride* CBMAI-0020).

315   Through the transcriptome analysis, we identified the DEGs and the differentially expressed CAZyme

genes in all strains. Although the Ta0020 strain presented the highest number of identified DEGs, it was found to be

less efficient than other *Trichoderma* strains in degrading plant biomass. *T. atroviride* is mostly used as a biocontrol

agent and is among the best mycoparasitic fungus used in agriculture (Kubicek et al. 2011). Each strain presented a

different set of genes with different expression levels, which can be attributed to strain differences in the regulatory

mechanisms of hydrolysis. Additionally, differences regarding the carbon source metabolization were observed,

which plays an important role in the production of enzymes and promotes the expression of different sets of genes as

the fungus seeks to adapt to new environments (Horta et al. 2018). Even though all species are induced by cellulose

growth, *T. harzianum* strains showed high numbers of CAZyme genes with enhanced specificity for that type of

biomass.

325   Among the main CAZyme classes detected in all strains, GHs were well represented, with 21 genes in

Th0179 and twice that in Th3844. GHs compose an extremely important class in several metabolic routes in fungi,

including genes involved in cellulose and chitin degradation (Ferreira Filho et al. 2017). Here, we found the main

CAZyme families responsible for cellulose degradation, such as GH1, GH3, GH5, GH6, GH7, GH12, GH45 and an

LPMO from the AA9 family, and for hemicellulose degradation, such as GH10, GH11, GH54, GH62 and CE5

(Ferreira Filho et al. 2017; Sweeney and Xu 2012; Villares et al. 2017; Zhao et al. 2014). Within the GH class, an

important family expected in our data was GH18, which has chitinase activity; since members of the genus

Trichoderma (*T. harzianum* and *T. atroviride*) are capable of mycoparasitism, this class is directly related to the

biological control of these species (Binod et al. 2007; Martinez et al. 2008).

In comparing the two *T. harzianum* strains (Th0179 and Th3844), we observed that Th0179 presented

fewer CAZyme genes related to cellulose degradation than Th3844, with lower expression levels (Fig. 3a).

However, the measurement of the enzymatic activity from the culture supernatants after 96 h of growth showed

similar cellulase activity profiles during growth on cellulose, suggesting a greater potential of Th0179 to degrade

cellulose through the use of a more efficient set of enzymes. A similar profile was observed in *T. reesei*, the most

studied fungus within this genus and an important industrial producer of cellulolytic enzymes (Druzhinina et al.

2011); few CAZyme genes have been detected in its machinery, but it can reach the highest cellulolytic activity

(Druzhinina and Kubicek 2017; Horta et al. 2018; Manika et al. 2014). It is interesting to observe which proteins

found in the exoproteome of Th0179 may respond to this increased cellulase activity.

CAZyme genes related to cellulose degradation, including endo-β-1,4-glucanases, β-glucosidases, and

cellobiohydrolases, and CAZyme genes related to hemicellulose degradation, such as endo-1,4-β-xylanases, were

identified in the Th0179 transcriptome. In addition, we verified important secreted CAZymes that play important

roles in biomass degradation, such as β-glucosidases (A0A0F9XRC5 and A0A0F9XQT4), LPMO (A0A0F9XMI8),

cellulose 1,4-β-cellobiosidase (nonreducing end) (A0A0G0AEM7), cellulases (A0A0F9XG06 and A0A0F9WYH5),

endo-1,4-β-xylanases (A0A0F9Y0Y9, A0A0H3UCP8, and A0A0F9XXA4), and endo-1,4-β-mannosidase

(A0A0G0AGG8). One β-glucosidase from the GH3 family (THAR02_00656) was found in the exoproteome, which

had the lowest expression level among the CAZymes related to biomass degradation; the same pattern was observed

for *T. harzianum* IOC-3844 (Horta et al. 2018).

Horta et al. (2018) analyzed the exoproteomes of Th3844 and Ta0020 under equivalent conditions,

selecting a set of 80 proteins for a complete classification and analysis of expression levels based on their

transcriptomes. Comparison of the Th0179 and Th3844 exoproteomes indicated that both produced many of the

main CAZymes listed above; however, the THAR02_05896 gene, which encodes an endo-1,4-β-xylanase protein,

and the THAR02_03851 gene, which encodes a protein with endo-1,4-β-mannosidase activity, are exclusively

present in the Th0179 exoproteome. In the present study, when these exoproteomes were compared, we observed

that most of the proteins were upregulated in cellulose only for Th0179, including the AA9/CBM1 family

(THAR02_02134) that showed a 2.65-fold change (138.62 TPM). Among the strains, Ta0020 exhibited the lowest

number of secreted proteins related to cellulose and hemicellulose degradation. Thus, the exoproteome analysis

identified key enzymes that are fundamental for cellulose hydrolysis and act synergistically for efficient plant

biomass degradation.

The organization of the transcriptomic data into coexpression networks using graph theory allowed the

construction of gene interaction networks that were represented by nodes connected to edges (Azevedo et al. 2015;

Mutwil et al. 2010). Nodes represent the genes, and the edges represent the connections among these genes.

Correlations are determined based on the expression level of the genes pair by pair, indicating that genes spatially

closer to one another are more highly correlated than the genes that are farther apart. The coexpression subnetwork

revealed complex, specific relationships between CAZyme genes and genes involved in the production and secretion

of the detected proteins and is helpful for understanding the functions and regulation of genes. We identified an

enriched cluster with strong correlations among genes, CAZyme genes, secreted proteins and unknown genes in the

cluster analysis. Networks such as this one can be used as a platform to search for target genes or proteins in future

studies to comprehend the synergistic relationships between genes, their regulation and protein production, which is

very useful information for understanding the saccharification process.

Furthermore, through the functional annotation analysis of Th0179, we identified TFs and transporters,

which should be investigated in further studies to better understand the mechanisms by which these genes are

regulated. Most sugar transporters have yet to be characterized but play important roles in taking up mono- or

disaccharides into fungal cells after biomass degradation (de Gouvea et al. 2018; Peng et al. 2018). Fungal gene

expression is controlled at the transcriptional level (Liu et al. 2017), and its regulation affects the composition of

enzyme mixtures; accordingly, it is explored in several species because of its potential applications (Benocci et al.

2017). This further emphasizes the importance of deleting and/or overexpressing TFs that regulate specific genes

directly involved in plant biomass degradation (Liu et al. 2017).

In summary, the analyses of the transcriptome, exoproteome and coexpression networks revealed important

genes and enzymes that *T. harzianum* CBMAI-0179 uses to hydrolyze cellulose and that most likely act

synergistically to depolymerize polysaccharides. The results suggest great potential of this strain to degrade

cellulose and can contribute to the optimization of enzymatic cocktails for bioethanol production.

Conclusions

Bioprospecting new catalytic enzymes and improving technologies for the efficient enzymatic conversion

of plant biomass are required for advancing biofuel production. *T. harzianum* CBMAI-0179 is a novel potential

candidate producer of plant cell wall polysaccharide-degrading enzymes that can be biotechnologically exploited for

plant biomass degradation. The cellulase activity profile indicated high efficiency for the set of genes and proteins

produced and consequently the potential of this strain for cellulose degradation. A set of highly expressed CAZymes

and proteins that are strain- and treatment-specific was described in both the transcriptome and exoproteome

analyses. The coexpression network revealed the synergistic gene action to hydrolyze cellulose, in which the cluster

analysis revealed genes with strong correlations necessary for saccharification. Combined, these tools provide a

powerful approach for catalyst discovery and the selection of target genes for the heterologous expression of

proteins. In future studies, these tools can aid the selection of new species and the optimization of the production of

powerful enzymes for use in enzymatic cocktails for second-generation bioethanol production.

Declarations

Funding

This work was supported by grants from the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP

2015/09202-0; 2018/19660-4), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES,

Computational Biology Program) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

DAA received an MS fellowship from FAPESP (2017/17782-2) and CAPES – Computational Biology Program

(88882.160100/2017-01, 88887.336686/2019-00). MACH received a PD fellowship from FAPESP (2014/18856-1).

JAFF received a PhD fellowship from CNPq (170565/2017-3). NFM received a PhD fellowship from CNPq

(140909/2013-3), and APS is the recipient of a research fellowship from CNPq. The funding bodies played no role

in the design of the study, analysis, data interpretation and manuscript writing.

Conflict of interest

The authors declare that they have no conflict of interest.

Ethical approval

This article does not contain any studies with human participants or animals performed by any of the authors.

Consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and material

The datasets generated and/or analyzed during the current study are included in this published article and its

supplementary material. The reads have been deposited at the NCBI SRA and can be accessed under the BioProject

number PRJNA336221.

Authors' contributions

DAA and APS conceived and designed the study. DAA, MACH, JAFF and NFM performed the data analysis.

MACH designed and performed the fermentation experiments. DAA drafted the manuscript, which was critically

revised by MACH, JAFF and APS. All authors read and approved the final manuscript.

Acknowledgments

We would like to acknowledge the funding from Fundação de Amparo à Pesquisa do Estado de São Paulo

(FAPESP), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Computational Biology

Program) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). We thank the National

Institute of Metrology, Quality and Technology (INMETRO) for performing the proteomics analysis via LC-

MS/MS, the Brazilian Biorenewables National Laboratory (LNBR), Campinas – SP, for conducting the fermentation

References

Ahmed S, Mustafa G, Arshad M, Rajoka MI (2017) Fungal biomass protein production from *Trichoderma harzianum* using rice polishing. BioMed Res Int 2017:6232793. doi:10.1155/2017/6232793

Al-Sadi AM, Al-Oweisi FA, Edwards SG, Al-Nadabi H, Al-Fahdi AM (2015) Genetic analysis reveals diversity and genetic relationship among *Trichoderma* isolates from potting media, cultivated soil and uncultivated soil. BMC Microbiol 15(1):147. doi:10.1186/s12866-015-0483-8

Alfaro M, Castanera R, Lavin JL, Grigoriev IV, Oguiza JA, Ramirez L, Pisabarro AG (2016) Comparative and transcriptional analysis of the predicted secretome in the lignocellulose-degrading basidiomycete fungus *Pleurotus ostreatus*. Environ Microbiol 18(12):4710-4726. doi:10.1111/1462-2920.13360

Andrews S (2010) FastQC: a quality control tool for high throughput sequence data. http://www.bioinformatics.babraham.ac.uk/projects/fastqc. Accessed 25 March 2017

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G (2000) Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. Nat Genet 25(1):25-29. doi:10.1038/75556

Azevedo H, Bando SY, Bertonha FB, Moreira-Filho CA (2015) Redes de interação gênica e controle epigenético na transição saúde-doença. Rev Med 94(4):223. doi:10.11606/issn.1679-9836.v94i4p223-229

Bailey MJ, Poutanen K (1989) Production of xylanolytic enzymes by strains of *Aspergillus*. Appl Microbiol Biotechnol 30(1):5-10. doi:10.1007/bf00255989

Baroncelli R, Piaggeschi G, Fiorini L, Bertolini E, Zapparata A, Pe ME, Sarrocco S, Vannacci G (2015) Draft whole-genome sequence of the biocontrol agent *Trichoderma harzianum* T6776. Genome Announc 3(3):e00647-15. doi:10.1128/genomeA.00647-15

Bayer EA, Lamed R, Himmel ME (2007) The potential of cellulases and cellulosomes for cellulosic waste management. Current opinion in biotechnology 18(3):237-45. doi:10.1016/j.copbio.2007.04.004

Benocci T, Aguilar-Pontes MV, Zhou M, Seiboth B, de Vries RP (2017) Regulators of plant biomass degradation in ascomycetous fungi. Biotechnol Biofuels 10:152. doi:10.1186/s13068-017-0841-x

Benoliel B, Torres FA, de Moraes LM (2013) A novel promising *Trichoderma harzianum* strain for the production

of a cellulolytic complex using sugarcane bagasse in natura. Springerplus 2:656. doi:10.1186/2193-1801-2-

656

Binod P, Sukumaran RK, Shirke SV, Rajput JC, Pandey A (2007) Evaluation of fungal culture filtrate containing

chitinase as a biocontrol agent against *Helicoverpa armigera*. J Appl Microbiol 103(5):1845-1852.

doi:10.1111/j.1365-2672.2007.03428.x

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics

30(15):2114-2120. doi:10.1093/bioinformatics/btu170

Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing

the principle of protein-dye binding. Anal Biochem 72(1):248-254. doi:10.1006/abio.1976.9999

CLC Genomics Workbench (2016) Manual for CLC genomics workbench 6.5.2 Windows, Mac OS X and Linux

Denmark. QIAGEN (Aarhus A/S), Aarhus, Denmark

Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M, Robles M (2005) Blast2GO: a universal tool for annotation,

visualization and analysis in functional genomics research. Bioinformatics 21(18):3674-6.

doi:10.1093/bioinformatics/bti610

Conesa A, Madrigal P, Tarazona S, Gomez-Cabrero D, Cervera A, McPherson A, Szczesniak MW, Gaffney DJ, Elo

LL, Zhang X, Mortazavi A (2016) A survey of best practices for RNA-seq data analysis. Genome Biol

17(1):13. doi:10.1186/s13059-016-0881-8

de Gouvea PF, Bernardi AV, Gerolamo LE, de Souza Santos E, Riano-Pachon DM, Uyemura SA, Dinamarco TM

(2018) Transcriptome and secretome analysis of *Aspergillus fumigatus* in the presence of sugarcane

bagasse. BMC Genomics 19(1):232. doi:10.1186/s12864-018-4627-8

Delabona PDS, Cota J, Hoffmam ZB, Paixao DA, Farinas CS, Cairo JP, Lima DJ, Squina FM, Ruller R, Pradella JG

(2013) Understanding the cellulolytic system of *Trichoderma harzianum* P49P11 and enhancing

saccharification of pretreated sugarcane bagasse by supplementation with pectinase and alpha-L-

arabinofuranosidase. Bioresour Technol 131:500-507. doi:10.1016/j.biortech.2012.12.105

Delabona PDS, Farinas CS, da Silva MR, Azzoni SF, Pradella JG (2012) Use of a new *Trichoderma harzianum*

strain isolated from the Amazon rainforest with pretreated sugar cane bagasse for on-site cellulase

production. Bioresour Technol 107:517-521. doi:10.1016/j.biortech.2011.12.048

dos Santos Castro L, Pedersoli WR, Antonieto ACC, Steindorff AS, Silva-Rocha R, Martinez-Rossi NM, Rossi A,

Brown NA, Goldman GH, Faca VM, Persinoti GF, Silva RN (2014) Comparative metabolism of cellulose,

sophorose and glucose in *Trichoderma reesei* using high-throughput genomic and proteomic analyses.

Biotechnol Biofuels 7(1):41. doi:10.1186/1754-6834-7-41

Druzhinina IS, Kubicek CP (2017) Genetic engineering of *Trichoderma reesei* cellulases and their production.

Microb Biotechnol 10(6):1485-1499. doi:10.1111/1751-7915.12726

Druzhinina IS, Seidl-Seiboth V, Herrera-Estrella A, Horwitz BA, Kenerley CM, Monte E, Mukherjee PK, Zeilinger

S, Grigoriev IV, Kubicek CP (2011) *Trichoderma*: the genomics of opportunistic success. Nat Rev

Microbiol 9(10):749-759. doi:10.1038/nrmicro2637

Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. Evolution 39(4):783.

doi:10.2307/2408678

Ferreira Filho JA, Horta MAC, Beloti LL, dos Santos CA, de Souza AP (2017) Carbohydrate-active enzymes in

Trichoderma harzianum: a bioinformatic analysis bioprospecting for key enzymes for the biofuels industry.

BMC Genomics 18(1):779. doi:10.1186/s12864-017-4181-9

Ghose TK (1987) Measurement of cellulase activities. Pure Appl Chem 59(2):257-268.

doi:10.1351/pac198759020257

Horta MA, Vicentini R, Delabona PDS, Laborda P, Crucello A, Freitas S, Kuroshu RM, Polikarpov I, Pradella JG,

Souza AP (2014) Transcriptome profile of *Trichoderma harzianum* IOC-3844 induced by sugarcane

bagasse. PLoS One 9(2):e88689. doi:10.1371/journal.pone.0088689

Horta MAC, Filho JAF, Murad NF, de Oliveira Santos E, dos Santos CA, Mendes JS, Brandao MM, Azzoni SF, de

Souza AP (2018) Network of proteins, enzymes and genes linked to biomass degradation shared by

Trichoderma species. Sci Rep 8(1):1341. doi:10.1038/s41598-018-19671-w

Illumina (2014) TruSeq RNA, sample preparation v2 guide. Illumina, San Diego, US

Jones DT, Taylor WR, Thornton JM (1992) The rapid generation of mutation data matrices from protein sequences.

Comput Appl Biosci 8(3):275-282. doi:10.1093/bioinformatics/8.3.275

Kimura M (1980) A simple method for estimating evolutionary rates of base substitutions through comparative

studies of nucleotide sequences. J Mol Evol 16(2):111-120. doi:10.1007/BF01731581

Kubicek CP, Herrera-Estrella A, Seidl-Seiboth V, Martinez DA, Druzhinina IS, Thon M, Zeilinger S, Casas-Flores

S, Horwitz BA, Mukherjee PK, Mukherjee M, Kredics L, Alcaraz LD, Aerts A, Antal Z, Atanasova L,

Cervantes-Badillo MG, Challacombe J, Chertkov O, McCluskey K, Coulpier F, Deshpande N, von Döhren

H, Ebbole DJ, Esquivel-Naranjo EU, Fekete E, Flipphi M, Glaser F, Gómez-Rodríguez EY, Gruber S, Han

C, Henrissat B, Hermosa R, Hernández-Oñate M, Karaffa L, Kosti I, Le Crom S, Lindquist E, Lucas S,

Lübeck M, Lübeck PS, Margeot A, Metz B, Misra M, Nevalainen H, Omann M, Packer N, Perrone G,

Uresti-Rivera EE, Salamov A, Schmoll M, Seiboth B, Shapiro H, Sukno S, Tamayo-Ramos JA, Tisch D,

Wiest A, Wilkinson HH, Zhang M, Coutinho PM, Kenerley CM, Monte E, Baker SE, Grigoriev IV (2011)

Comparative genome sequence analysis underscores mycoparasitism as the ancestral life style of

Trichoderma. Genome Biol 12(4):R40. doi:10.1186/gb-2011-12-4-r40

Kubicek CP, Steindorff AS, Chenthamara K, Manganiello G, Henrissat B, Zhang J, Cai F, Kopchinskiy AG,

Kubicek EM, Kuo A, Baroncelli R, Sarrocco S, Noronha EF, Vannacci G, Shen Q, Grigoriev IV,

Druzhinina IS (2019) Evolution and comparative genomics of the most common Trichoderma species.

BMC Genomics 20(1):485. doi:10.1186/s12864-019-5680-7

Kumar S, Stecher G, Tamura K (2016) MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger

datasets. Mol Biol Evol 33(7):1870-1874. doi:10.1093/molbev/msw054

Liu R, Chen L, Jiang Y, Zou G, Zhou Z (2017) A novel transcription factor specifically regulates GH11 xylanase

genes in *Trichoderma reesei*. Biotechnol Biofuels 10:194. doi:10.1186/s13068-017-0878-x

Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the

2(-Delta Delta C(T)) method. Methods 25(4):402-408. doi:10.1006/meth.2001.1262

Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B (2014) The carbohydrate-active enzymes

database (CAZy) in 2013. Nucleic Acids Res 42(Database issue):D490-D495. doi:10.1093/nar/gkt1178

Mandels M, Andreotti RE (1978) Problems and challenges in the cellulose to cellulase fermentation. Process

Biochem 13:6-13

Manika S, Saju SS, Subhash C, Mukesh S, Pratibha S (2014) Comparative evaluation of cellulase activity in

Trichoderma harzianum and *Trichoderma reesei*. Afr J Microbiol Res 8(19):1939-1947.

doi:10.5897/ajmr2013.6488

Martinez D, Berka RM, Henrissat B, Saloheimo M, Arvas M, Baker SE, Chapman J, Chertkov O, Coutinho PM,

Cullen D, Danchin EG, Grigoriev IV, Harris P, Jackson M, Kubicek CP, Han CS, Ho I, Larrondo LF, de

Leon AL, Magnuson JK, Merino S, Misra M, Nelson B, Putnam N, Robbertse B, Salamov AA, Schmoll M,

Terry A, Thayer N, Westerholm-Parvinen A, Schoch CL, Yao J, Barabote R, Nelson MA, Detter C, Bruce

D, Kuske CR, Xie G, Richardson P, Rokhsar DS, Lucas SM, Rubin EM, Dunn-Coleman N, Ward M,

Brettin TS (2008) Genome sequencing and analysis of the biomass-degrading fungus *Trichoderma reesei*

(syn. *Hypocrea jecorina*). Nat Biotechnol 26(5):553-560. doi:10.1038/nbt1403

Miao Y, Liu D, Li G, Li P, Xu Y, Shen Q, Zhang R (2015) Genome-wide transcriptomic analysis of a superior

biomass-degrading strain of *A. fumigatus* revealed active lignocellulose-degrading genes. BMC Genomics

16(1):459. doi:10.1186/s12864-015-1658-2

Mutwil M, Klie S, Tohge T, Giorgi FM, Wilkins O, Campbell MM, Fernie AR, Usadel B, Nikoloski Z, Persson S

(2011) PlaNet: combined sequence and expression comparisons across plant networks derived from seven

species. Plant Cell 23(3):895-910. doi:10.1105/tpc.111.083667

Mutwil M, Usadel B, Schütte M, Loraine A, Ebenhöh O, Persson S (2010) Assembly of an interactive correlation

network for the Arabidopsis genome using a novel heuristic clustering algorithm. Plant physiology

152(1):29-43. doi:10.1104/pp.109.145318

Oliveira RR, Viana AJ, Reategui AC, Vincentz MG (2015) Short communication an efficient method for

simultaneous extraction of high-quality RNA and DNA from various plant tissues. Genet Mol Res

14(4):18828-18838. doi:10.4238/2015.December.28.32

Peng M, Aguilar-Pontes MV, de Vries RP, Mäkelä MR (2018) *In silico* analysis of putative sugar transporter genes

in *Aspergillus niger* using phylogeny and comparative transcriptomics. Front Microbiol 9:1045.

doi:10.3389/fmicb.2018.01045

Peterson R, Nevalainen H (2012) Trichoderma reesei RUT-C30 – thirty years of strain improvement. Microbiology

158(1):58-68. doi:10.1099/mic.0.054031-0

Rocha ALV, Maeda N, Pereira N, Kern FM, Elias L, Simister R, Steele-King C, Gomez LD, McQueen-Mason SJ

(2016) Characterization of the cellulolytic secretome of *Trichoderma harzianum* during growth on

sugarcane bagasse and analysis of the activity boosting effects of swollenin. Biotechnol Prog 32(2):327-

336. doi:10.1002/btpr.2217

Rosolen RR, Aono AH, Almeida DA, Ferreira Filho JA, Horta MAC, de Souza AP (2020) Comparative gene

coexpression networks analysis reveals different strategies of Trichoderma spp. associated with

XYR1 and CRE1 during cellulose degradation. bioRxiv:2020.05.02.074344.

doi:10.1101/2020.05.02.074344

Schomburg I, Jeske L, Ulbrich M, Placzek S, Chang A, Schomburg D (2017) The BRENDA enzyme information

system-from a database to an expert system. J Biotechnol 261:194-206. doi:10.1016/j.jbiotec.2017.04.020

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T (2003)

Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome

Res 13(11):2498-504. doi:10.1101/gr.1239303

Sweeney MD, Xu F (2012) Biomass converting enzymes as industrial biocatalysts for fuels and chemicals: recent

developments. Catalysts 2(2):244-263. doi:10.3390/catal2020244

Thompson JD, Higgins DG, Gibson TJ (1994) CLUSTAL W: improving the sensitivity of progressive multiple

sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice.

Nucleic Acids Res 22(22):4673-4680. doi:10.1093/nar/22.22.4673

Untergasser A, Nijveen H, Rao X, Bisseling T, Geurts R, Leunissen JA (2007) Primer3Plus, an enhanced web

interface to primer3. Nucleic Acids Res 35(Web Server issue):W71-W74. doi:10.1093/nar/gkm306

van Dyk JS, Pletschke BI (2012) A review of lignocellulose bioconversion using enzymatic hydrolysis and

synergistic cooperation between enzymes--factors affecting enzymes, conversion and synergy. Biotechnol

Adv 30(6):1458-1480. doi:10.1016/j.biotechadv.2012.03.002

Villares A, Moreau C, Bennati-Granier C, Garajova S, Foucat L, Falourd X, Saake B, Berrin JG, Cathala B (2017)

Lytic polysaccharide monooxygenases disrupt the cellulose fibers structure. Sci Rep 7:40262.

doi:10.1038/srep40262

Wang Y, Fan C, Hu H, Li Y, Sun D, Wang Y, Peng L (2016) Genetic modification of plant cell walls to enhance

biomass yield and biofuel production in bioenergy crops. Biotechnol Adv 34(5):997-1017.

doi:10.1016/j.biotechadv.2016.06.001

Yamanishi Y, Hattori M, Kotera M, Goto S, Kanehisa M (2009) E-zyme: predicting potential EC numbers from the

chemical transformation pattern of substrate-product pairs. Bioinformatics 25(12):i179-i186.

doi:10.1093/bioinformatics/btp223

Zhang YHP, Hong J, Ye X (2009) Cellulase Assays. In: Mielenz JR (ed) Biofuels: methods and protocols. Humana

608   Press, Totowa, NJ, pp 213-231

609   Zhao Z, Liu H, Wang C, Xu JR (2014) Correction: comparative analysis of fungal genomes reveals different plant

610   cell wall degrading capacity in fungi. BMC Genomics 15:6. doi:10.1186/1471-2164-15-6

Figure captions

Fig. 1 The transcriptome profiles, gene expression comparison and the main CAZyme classes identified for all

strains. PCA of transcriptome mapping according to species and growth conditions (CEL – cellulose and GLU –

glucose) using the *T. harzianum* T6776 genome as a reference (a), number of DEGs and differentially expressed

CAZyme genes in *T. harzianum* CBMAI-0179 (Th0179), *T. harzianum* IOC-3844 (Th3844), and *T. atroviride*

CBMAI-0020 (Ta0020) under cellulose growth conditions (b), the differentially expressed CAZyme classes

identified for each strain under cellulose growth conditions (c)

Fig. 2 Distribution of CAZyme families in *Trichoderma* spp. Classification and quantification of CAZyme families

in *T. harzianum* CBMAI-0179 (a), *T. harzianum* IOC-3844 (b), and *T. atroviride* CBMAI-0020 (c) under cellulose

growth conditions

Fig. 3 Evaluation of CAZyme family expression in *Trichoderma* spp. via RNA-Seq. Quantification of the

expression of the main families related to cellulose (a) and hemicellulose (b) degradation in TPM

Fig. 4 GO terms of *T. harzianum* CBMAI-0179 under cellulose growth conditions. The genes were annotated

according to the main GO terms: molecular function (a), biological process (b), and cellular component (c)

Fig. 5 Coexpression networks of *T. harzianum* CBMAI-0179. Complete coexpression network (a), the coexpression

subnetwork based on the exoproteome data (b), the enriched cluster analysis of the coexpression network (c), the

subnetwork of the CAZyme genes and the secreted proteins identified in the cluster analysis (d). Red squares

indicate DEGs under cellulose growth conditions, blue squares indicate DEGs under glucose growth conditions,

yellow triangles indicate CAZyme genes under cellulose growth conditions, light blue triangles indicate CAZyme

genes under glucose growth conditions and purple hexagons indicate the secreted proteins under cellulose growth

conditions

TH0179_CEL

TH3844_CEL

TA0020_CEL

a **Cellulase families**

b **Hemicellulase families**

Molecular Function

Biological Process

Cellular Component

a

b

c

GH6;CBM1

CBM1

GH7;CBM1

GH1

CBM1

GH10

AA9;CBM1

GH45;CBM1

GH11

GH11;CBM1

GH1

AA7

d

CBM1   CBM1

GH45;CBM1

GH7;CBM1

GH6;CBM1

GH11

GH10

GH11;CBM1

GH1

AA9;CBM1   GH1

AA7

■ DEGs in Cel   ■ DEGs in Glu   ⋎ CAZymes Cel   ▲ CAZymes Glu   ⬡ Secreted proteins