The synergistic actions of hydrolytic genes in coexpression networks reveal the potential

of *Trichoderma harzianum* for cellulose degradation

Déborah Aires Almeida[1,2], Maria Augusta Crivelente Horta[1,#], Jaire Alves Ferreira Filho[1,2],

Natália Faraj Murad[1] and Anete Pereira de Souza[1,3,*]

[1]Center for Molecular Biology and Genetic Engineering (CBMEG), University of Campinas

(UNICAMP), Campinas, SP, Brazil

[2]Graduate Program in Genetics and Molecular Biology, Institute of Biology, UNICAMP,

Campinas, SP, Brazil

[3]Department of Plant Biology, Institute of Biology, UNICAMP, Campinas, SP, Brazil

[#]Present Address: Holzforshung München, TUM School of Life Sciences Weihenstephan,

Technische Universität München, Freising, Germany

*Corresponding author

Profa Anete Pereira de Souza

Dept. de Biologia Vegetal, Universidade Estadual de Campinas, CEP 13083-875, Campinas,

São Paulo, Brazil

Tel.: +55-19-3521-1132

E-mail:anete@unicamp.br

Abstract

Background: Bioprospecting key genes and proteins related to plant biomass degradation is

an attractive approach for the identification of target genes for biotechnological purposes,

especially genes with potential applications in the biorefinery industry that can enhance

second-generation ethanol production technology. *Trichoderma harzianum* is a potential

candidate for cellulolytic enzyme production. Herein, the transcriptome, exoproteome,

enzymatic activities of extracts, and coexpression networks of the *T. harzianum* strain

CBMAI-0179 under biomass degradation conditions were examined.

Results: We used RNA-Seq to identify differentially expressed genes (DEGs) and

carbohydrate-active enzyme (CAZyme) genes related to plant biomass degradation and

compared them with genes of strains from congeneric species (*T. harzianum* IOC-3844 and

T. atroviride CBMAI-0020). *T. harzianum* CBMAI-0179 harbors species- and treatment-

specific CAZyme genes, transporters and transcription factors. Additionally, we detected

important proteins related to biomass degradation, including β-glucosidases, endoglucanases,

cellobiohydrolases, lytic polysaccharide monooxygenases (LPMOs), endo-1,4-β-xylanases

and β-mannanases, in the exoproteome under cellulose growth conditions. Coexpression

networks were constructed to explore the relationships among the genes with corresponding

secreted proteins that act synergistically for cellulose degradation. An enriched cluster with

degradative enzymes was described, and the subnetwork of CAZymes showed linear

correlations among secreted proteins (AA9, GH6, GH10, GH11 and CBM1) and

differentially expressed CAZyme genes (GH45, GH7, AA7 and GH1).

Conclusions: Our results provide valuable information for future studies on the genetic

regulation of plant cell wall-degrading enzymes. This knowledge can be exploited for the

improvement of enzymatic reactions to degrade plant biomass, which is useful for bioethanol

production.

Keywords: *Trichoderma harzianum*, Cellulose, RNA-Seq, CAZymes, Exoproteome,

Coexpression networks

Background

The expanding worldwide demand for renewable and sustainable energy sources has

increased the interest in alternative energy sources, and the production of second-generation

biofuels seems to be the most viable option to confront these issues [1, 2]. Lignocellulosic

biomass is the most abundant renewable organic carbon resource on earth, consisting of three

major polymers, cellulose, hemicellulose, and lignin [2]. However, due to its recalcitrant

characteristics that prevent enzyme access, degrading this complex matrix is still a major

challenge [3]. For the complete hydrolysis of lignocellulose, a variety of enzymes acting in

synergy are required, and much research has focused on this topic in recent decades [4].

Interactions between different enzymes have been investigated to identify optimal

combinations and ratios of enzymes for efficient biomass degradation, which are highly

dependent on the properties of the lignocellulosic substrates and the surface structure of

cellulose microfibrils [4, 5]. Due to their abundance in nature, microorganisms are considered

natural producers of enzymes, and many of them, including members of both bacteria and

fungi, have evolved to digest lignocellulose [6, 7]. The search for microorganisms that are

able to efficiently degrade lignocellulosic biomass is pivotal for the establishment of the

sustainable production of bioethanol [8].

Filamentous fungi, including the genera *Trichoderma, Aspergillus*, *Penicillium* and

Neurospora, produce extracellular proteins that act synergistically to degrade plant cell walls

and are widely used in the enzymatic industry [9]. Species in the filamentous ascomycete

genus *Trichoderma* are among the most commonly isolated saprotrophic fungi [10] and are

important from a biotechnological perspective [7]. *Trichoderma* species are widely used in

agriculture as biocontrol agents due their ability to antagonize plant-pathogenic fungi and in

industry as producers of plant cell wall-degrading enzymes [11-14]. In addition, *Trichoderma*

species are easily isolated from soil and decomposing organic matter [15]. Within the

Trichoderma genus, *T. reesei* is the most intensively studied species [16]. *T. reesei* is a well-

known producer of cellulase and hemicellulase, and due to the high effectiveness of the

synergistic cellulases in this species, it is widely employed in industry, and technologies for

its use and handling are based on seventy years of experience [8, 16-19]. However, studies on

T. harzianum strains have shown their potential to produce a set of enzymes that can degrade

lignocellulosic biomass [20-23]; therefore, *T. harzianum* strains are being investigated as

potentially valuable sources of industrial cellulases [6].

The identification of carbohydrate-active enzymes (CAZymes) that act synergistically

under biodegradation conditions [4] has the potential to improve the enzymatic hydrolysis

process by optimizing and reducing bioethanol costs. The CAZy database (www.cazy.org)

classifies CAZymes into six major groups: glycoside hydrolases (GHs), glycosyltransferases

(GTs), polysaccharide lyases (PLs), carbohydrate esterases (CEs), auxiliary activities (AAs),

and carbohydrate-binding modules (CBMs) [24]. CAZymes are extensively used for the

genetic classification of important hydrolytic enzymes [22, 25].

The conversion of cellulose to glucose involves the synergistic action of three

principal groups of enzymes: endo-β-1,4-glucanases (EC 3.2.1.4), β-glucosidases (EC

3.2.1.21), and cellobiohydrolases (EC 3.2.1.91/176) [20, 26]. For hemicellulose hydrolysis,

several enzymes are needed, such as endo-1,4-β-xylanases (EC 3.2.1.8), β-xylosidases (EC

3.2.1.37), β-mannanases (EC 3.2.1.78), arabinofuranosidases (EC 3.2.1.55), and acetylxylan

esterases (EC 3.1.1.72) [26, 27]. In addition, a number of auxiliary enzymes are involved in

this process, such as lytic polysaccharide monooxygenases (LPMOs), cellulose-induced

protein 1 and 2 (CIP1 and CIP2) and swollenin, which can increase the hydrolytic

performance of enzymatic cocktails used in industry for bioethanol production [6, 28-30].

As genetic variation occurs within species [31, 32], understanding and exploring the

genetic mechanisms of different *T. harzianum* strains can provide valuable information for

industrial applications. In the present study, we analyzed the enzymatic activity,

transcriptome and exoproteome of *T. harzianum* CBMAI-0179 and compared them with

those of other *Trichoderma* strains (*T. harzianum* IOC-3844 and *T. atroviride* CBMAI-

0020). RNA-Seq analysis was performed to construct coexpression networks, providing

novel information about the potential of this *T. harzianum* strain for biotechnological

applications. Our findings provide insights into the genes/proteins that act synergistically

in plant biomass conversion and can be exploited to improve enzymatic hydrolysis and

thereby increase the efficiency of the saccharification of lignocellulosic substrates for

bioethanol production.

Results

Transcriptome analysis of* Trichoderma *spp. under cellulose growth conditions

The present study represents the first deep genetic analysis of Th0179, describing and

comparing the transcriptome by RNA-Seq under two different growth conditions, cellulose

and glucose, to identify the genes involved in plant biomass degradation. Reads were mapped

against reference genomes of *T. harzianum* and *T. atroviride*, generating 96.3, 111.8 and

133.3 million paired-end reads for Th0179, Th3844 and Ta0020, respectively. To establish

the degrees of similarity and difference in gene expression among strains and between

treatments, a principal component analysis (PCA) was performed using the *T. harzianum*

T6776 genome as a reference. The PCA results showed clustered groups with higher

similarity between treatments than among strains. The transcriptomes of Th0179 and Th3844

were more similar to each other than to the transcriptome of Ta0020 (Fig. 1a), showing that it

is possible to capture differences among the three strains.

Venn diagrams of the genes exhibiting expression levels greater than zero were

constructed based on the similarities among the genes from all strains, thus showing the

species-specific genes expressed under both conditions (Additional file 4: Fig. S3). Through

the transcriptome analysis of the strains, we identified 11,250 genes exhibiting expression

levels greater than zero under cellulose growth conditions and 11,235 genes exhibiting

expression levels greater than zero under glucose growth conditions. The number of genes

shared by Th0179 and Th3844 was higher under both conditions than that shared by either

Th0179 or Th3844 and Ta0020. Th0179 exhibited the highest number of unique genes, with

374 and 168 unique genes under the cellulose and glucose growth conditions, respectively.

Among these unique genes under cellulose growth conditions, we found major facilitator

superfamily (MFS) transporters (THAR02_00234, THAR02_00911, THAR02_03251,

THAR02_03935, THAR02_07021, THAR02_07705 and THAR02_07942), an ATP-binding

cassette (ABC) transporter (THAR02_09958), a drug resistance protein (THAR02_04837), a

fungal specific transcription factor (TF) (THAR02_07743), and a C2H2 TF

(THAR02_11070). Among the genes that were upregulated in cellulose conditions relative to

glucose conditions, 219 were identified for Th0179, 281 were identified for Th3844, and 718

were identified for Ta0020 (Fig. 1b).

CAZyme identification and distribution in* Trichoderma *spp.

For CAZymes important in the degradation of biomass, the set of CAZymes was

identified by mapping all of the proteins of *T. harzianum* T6776 and *T. atroviride* IMI

206040 against the CAZy database using the BLASTp search tool. Based on the filtering

criteria, a total of 631 proteins were retained as CAZyme genes for *T. harzianum*, which

corresponds to 5.5% of the total 11,498 proteins predicted for this organism [33], and 640

proteins were retained for *T. atroviride*, which corresponds to 5.4% of the total 11,816

proteins predicted for this organism [34].

Considering the DEGs under cellulose growth conditions and using the established

CAZy database, we found 35, 78 and 31 differentially expressed CAZyme genes in our data

(Fig. 1b) for Th0179, Th3844 and Ta0020, respectively. We identified the main CAZyme

classes (AA, GH, GT, CE and CBM) and their contents for all strains (Fig. 1c). The Th3844

strain presented the highest number of classified genes from the AA family, a high number of

CBMs and twice the number of identified GHs found in the two other strains. Strain Ta0020

presented a higher number of genes from the GT family than Th0179, which exhibited a

higher number of classified genes from the CBM family than Ta0020. The differences

regarding the specific CAZyme families for each strain are shown in Fig. 2.

The GH group was the most represented class of enzymes present in all evaluated

strains. GHs are key enzymes for carbohydrate hydrolysis and include enzymes capable of

degrading cellulose [35-37], with many able to cleave glycosidic bonds between glucose

molecules. Another important family involved in degradation of the plant cell wall, the AA

family, was also identified in all strains [38]. The AA class currently harbors 9 families of

ligninolytic enzymes and 6 families of lytic polysaccharide mono-oxygenases that may not

act on carbohydrates. However, because lignin is invariably and intimately associated with

carbohydrates in the plant cell wall, these lignolytic enzymes cooperate with classical

polysaccharide depolymerases, named auxiliary enzymes [24, 38], that differentially

influence degradative activity among species.

Cellulose and hemicellulose degradative enzymes in* Trichoderma *spp.

173   In plant biomass degradation, a variety of enzymes working synergistically are

174 required for complete hydrolysis [4]. Under cellulose growth conditions, we found

175 upregulated families among the CAZymes responsible for cellulose degradation, including

176 endoglucanases (GH5, GH12, GH45 and CBM1), β-glucosidases (GH1 and GH3) and

177 cellobiohydrolases (GH6 and GH7), that play important roles in cellulose degradation. In

178 addition, among the CAZymes responsible for hemicellulose degradation, we identified endo-

179 1,4-β-xylanases (GH10, GH11 and CBM1), arabinofuranosidases (GH54, GH62 and

180 CBM42) and acetylxylan esterases (CE5 and CBM1). Additionally, we found the copper

181 enzymes LPMOs, classified as AAs in the family AA9 [39]; these are considered a

182 breakthrough in the enzymatic degradation of cellulose because they oxidatively cleave

183 glycosidic linkages that render substrates more susceptible to hydrolysis by conventional

184 cellulases [28]. The expression levels in TPM of the main cellulase and hemicellulase

185 families based on their principal enzyme activity present in Th0179, Th3844 and Ta0020

186 were evaluated using the RNA-Seq data (Fig. 3).

187   All of the GHs related to cellulose degradation were found in Th3844, including 2

188 AA9 members with a CBM1 module for cellulose binding. All GHs except GH12, including

189 AA9/CBM1, were detected in Th0179. Only one gene belonging to the GH5 family

190 (TRIATDRAFT_81867) with cellulase activity (EC 3.2.1.4) was observed in Ta0020 (48.12

191 TPM). The most highly expressed genes were cellobiohydrolases from the GH6/CBM1

192 (THAR02_04414 – 299.66 TPM for Th0179 and 986.25 TPM for Th3844) and GH7/CBM1

193 families (THAR02_08897 – 369.92 TPM for Th0179; THAR02_03357 and THAR02_08897

194 – 1876.89 TPM for Th3844) (Fig. 3a).

195   In addition, 5 families related to hemicellulose degradation were identified in Th3844,

196 in which endo-1,4-β-xylanases (EC 3.2.1.8) from the GH11 family had the greatest

197 expression levels (THAR02_02147, THAR02_05896, THAR02_08630 and THAR02_08858

– 1013.62 TPM). Endo-1,4-β-xylanase (EC 3.2.1.8) from the GH10 family was found in both

T. harzianum strains (THAR02_03271 – 47.58 TPM for Th0179 and 741.35 TPM for

Th3844). Acetylxylan esterases (EC 3.1.1.72) from the CE5/CBM1 family were found only

in Th3844 (THAR02_01449 and THAR02_07663 – 227.56 TPM). We identified only one α-

L-arabinofuranosidase (EC 3.2.1.55) from the GH54/CBM42 family (TRIATDRAFT_81098

– 18.07 TPM) in Ta0020, whereas for Th3844, three α-L-arabinofuranosidases from the

GH54/CBM42 and GH62 families were identified (Fig. 3b). The Ta0020 strain showed the

lowest number of genes and the lowest expression levels of the CAZyme families related to

cellulose and hemicellulose degradation. The classification of the CAZyme genes along with

their enzyme activities, fold change values, e-values, EC numbers and expression values

(TPM) for Th0179 under cellulose fermentative conditions are described in Table 1. The

corresponding information for the Th3844 and Ta0020 strains can be found in Additional file

5: Table S2.

[Insert Table 1 here]

Functional annotation of* T. harzianum *CBMAI-0179 in the presence of cellulose

The first functional annotation of the expressed sequences under cellulose

fermentative conditions for Th0179 was performed based on GO terms (Fig. 4). A total of

7,718 sequences were annotated, which corresponds to 67.1% of the total 11,498 proteins

predicted for this organism (*T. harzianum* T6776) [33]. Under the molecular function

category, catalytic activity was the main annotated function, with 3,350 identified sequences.

Other functions that were enriched under this category were hydrolase activity, nucleotide

binding and TF activity, with 1,111, 899 and 277 sequences, respectively. For the biological

process category, the main functions were metabolic process, with 3,607 sequences, and

cellular process, with 2,575 sequences. In addition, within the biological process category,

574 sequences were identified as associated with transmembrane transport, and 66 sequences

were identified as associated with regulation of catalytic activity. Compared with Th3844,

Th0179 presented 163 more sequences related to catalytic activity, 53 more sequences related

to hydrolase activity, and 15 more sequences associated with TF activity but 10 fewer

sequences related to transmembrane transport, suggesting that these two strains have

developed different functional regulation. Ta0020 presented fewer sequences related to any

of the above functions than Th3844 and Th0179 except for TF activity, for which Ta0020 had

53 more sequences than Th0179.

Exoproteome and RNA-Seq data correlation of* T. harzianum *CBMAI-0179

233   Once the transcriptome was characterized, we analyzed the secreted proteins

identified in the transcriptome profile of *T. harzianum* CBMAI-0179. A total of 64 proteins,

which had been secreted and were present in the culture medium after 96 h of fermentation,

were detected in the extracts. Of those, 32 proteins were present in the cellulose aqueous

extract (Table 2), 12 were present only in the glucose aqueous extract, and 20 were present

under both conditions (Additional file 6: Table S3). Among the 32 secreted proteins detected

using cellulose as the carbon source, the main CAZyme families observed exclusively in this

supernatant were among of the most important families for cellulose and hemicellulose

degradation, such as β-glucosidases (EC 3.2.1.21), endo-β-1,4-glucanases (EC 3.2.1.4),

LPMOs (EC 3.2.1.4), cellobiohydrolases (EC 3.2.1.91), endo-1,4-β-xylanases (EC 3.2.1.8),

and β-mannanases (EC 3.2.1.78). Two genes from the GH3 (THAR02_00656 and

THAR02_00890) and GH5/CBM1 (THAR02_04405 and THAR02_09719) families and one

gene from the AA9/CBM1 (THAR02_02134) and GH6/CBM1 (THAR02_04414) families

corresponded to the main group of secreted cellulases, whereas one gene from the GH10

(THAR02_03271) family and two genes from the GH11 (THAR02_02147 and

THAR02_05896) family corresponded to the main group of hemicellulases detected in the

supernatant. In addition, a member of the GH5 cellulase family with β-mannanase activity

(THAR02_03851) was identified. In addition to these secreted proteins, 8 other proteins were

classified as uncharacterized proteins. We also identified hemicellulose-degrading enzymes

in both extracts, such as α-L-arabinofuranosidase B (EC 3.2.1.55) and xylan 1,4-β-xylosidase

(EC 3.2.1.37).

[Insert Table 2 here]

Correlating the exoproteome data with the transcriptome data under cellulose growth

conditions, we observed the expression levels of genes that play important roles in plant

biomass degradation. CAZyme genes showing high TPM values in cellulose included one

cellulose 1,4-β-cellobiosidase (nonreducing end) (EC 3.2.1.91) from the GH6/CBM1 family

(299.66 TPM), one cellulase (EC 3.2.1.4) from the AA9/CBM1 family (138.62 TPM) and

one endo-1,4-β-xylanase (EC 3.2.1.8) from the GH11/CBM1 family (119.56 TPM). Two

uncharacterized proteins (THAR02_02133 and THAR02_08479) with the CBM1 module of

cellulose binding also showed increased expression levels under the cellulose condition. In

contrast, the THAR02_00656 gene, which displays β-glucosidase (EC 3.2.1.21) activity and

is from the GH3 family, had the lowest expression level (5.08 TPM) among the CAZyme

genes related to biomass degradation, indicating that genes with low expression levels are

also important for functional secreted proteins [40].

Coexpression networks for **T. harzianum** *CBMAI-0179*

The coexpression network was assembled for Th0179 and included all genes obtained

from the mapping results using the *T. harzianum* T6776 genome as a reference under both

conditions, and the DEGs and CAZyme genes were highlighted in the network (Fig. 5a). This

network, constructed based on the expression level data (TPM), was composed of a total of

11,104 nodes with 153,893 edges. We detected 219 genes corresponding to the DEGs under

cellulose growth conditions and 367 genes corresponding to the nodes under glucose growth

conditions. The respective CAZyme genes from both conditions were also identified in this

network, with 35 CAZymes under cellulose growth conditions and 28 CAZymes under

glucose growth conditions. The DEGs tended to cluster in the top (DEGs in glucose) and

bottom (DEGs in cellulose) parts of the network, reflecting the different regulation of

degradative activities according to substrate.

A subnetwork was generated based only on the secreted proteins that were present in

the cellulose aqueous extract and that had corresponding genes in the coexpression network

(Fig. 5b). This subnetwork exclusively represented the genes and their closest related genes

that are correlated to the proteins secreted. It was composed of 713 nodes and 6,124 edges,

including the 32 genes that encode the secreted proteins. We also identified 39 DEGs under

cellulose growth conditions, 8 DEGs under glucose growth conditions, 6 CAZyme genes

related to cellulose degradation and 1 CAZyme gene under glucose growth conditions in this

subnetwork. Among the CAZyme genes under cellulose growth conditions, the GH1 family

(THAR02_02251 and THAR02_05432) with β-glucosidase activity, the GH7/CBM1 family

(THAR02_08897) with cellulose 1,4-β-cellobiosidase activity, and the GH45/CBM1 family

(THAR02_02979) with cellulase activity were found in this subnetwork. Despite the different

functions of the related genes, it is predicted that these genes participate in the genetic

regulation of the detected CAZymes/proteins and are important to the regulation of the

hydrolytic system.

The cluster analysis classified 11,101 genes in 84 clusters (Additional file 7: Table

S4). We identified an enriched cluster composed of 196 nodes and 2,125 edges (Fig. 5c and

Additional file 7: Table S4) containing the greatest number of CAZyme genes among the

clusters. Of the 12 CAZyme genes detected, 7 corresponded to the secreted proteins detected

in the cellulose aqueous extract, and 5 were differentially expressed. In addition, we

identified 20 DEGs under cellulose growth conditions and 119 uncharacterized proteins in

this cluster, showing a strong correlation with the CAZyme genes, which indicates that the

unknown genes are important to the degradation process. The CAZyme genes from the

cluster analysis were selected to generate a new subnetwork with their corresponding edges

(Fig. 5d), showing linear correlations among secreted proteins (AA9, GH6, GH10, GH11 and

CBM1) and differentially expressed CAZyme genes (GH45, GH7, AA7 and GH1).

Discussion

Several studies, including transcriptomic and proteomic studies, have been performed

using filamentous fungi to bioprospect efficient catalysts for the development and

improvement of enzymatic cocktails to degrade plant biomass. Some fungi, including *T.*

reesei, *A. niger*, *A. nidulans* and *N. crassa* [9, 16, 17, 41-43], have been used for this purpose;

however, the search for new efficient strains is ongoing. Understanding the molecular

mechanisms by which filamentous fungi degrade plant biomass can improve the

saccharification process, a very important step in the production of second-generation ethanol

[42, 44]. The identification of new species and powerful enzymes can enhance the

technologies in the biofuel industry.

Trichoderma spp. are capable of degrading both fungal and plant cell wall materials

[45]. In this study, we chose to investigate species of the genus *Trichoderma* because this

genus is a common soil and wood-degrading fungi distributed worldwide [46-48], is easily

isolated from decomposing organic matter and soil [49], and harbors great potential to

produce enzymes that degrade plant biomass [10, 12, 22]. We selected two strains of *T.*

harzianum and one strain of *T. atroviride* to capture differences among strains. Different

biotechnological approaches were used in bioprospecting new and efficient enzymes for

possible applications in the enzymatic hydrolysis process. We performed transcriptome,

exoproteome and coexpression network analyses of a *T. harzianum* strain that has potential

for plant biomass degradation under cellulose growth conditions to gain insights into the

genes and proteins produced that are associated with cellulose hydrolysis. The analyses of

genetic expression together with the identified secreted proteins under biomass degradation

conditions allowed us to construct coexpression networks to investigate the relationships

among the genes. Furthermore, we compared the expression levels with *Trichoderma* spp.

that have been previously studied under the same conditions [10].

Through RNA-Seq, we identified the DEGs in all strains (Fig. 1b). Although Ta0020

presented the highest number of identified DEGs, this strain is less efficient than other

Trichoderma strains at degrading plant biomass. *T. atroviride* is mostly used as a biocontrol

agent and is among the best mycoparasitic fungus used in agriculture [34, 50, 51]. Although

the *T. atroviride* strain produced 31 differentially expressed CAZymes, it seemed to produce

less efficient enzymes with lower expression levels for plant biomass degradation than *T.*

harzianum, since it was observed to harbor only one gene with cellulase activity from the

GH5 family (TRIATDRAFT_81867) and only one gene with hemicellulase activity from the

GH54/CBM42 family (TRIATDRAFT_81098) (Fig. 3). In this analysis, each strain presented

a different set of genes with different expression levels, which can be attributed to strain

differences in the regulatory mechanisms of hydrolysis. Additionally, the carbon source,

which plays an important role in the production of enzymes, promotes the expression of

different sets of genes as the fungus seeks to adapt to new environments [10]. The *T.*

harzianum strains showed high numbers of CAZyme genes with enhanced specificity for

biomass degradation.

Among the main CAZyme classes detected in all strains, GHs were well represented,

with 21 genes in Th0179 and twice that in Th3844 (Fig. 1c). GHs compose an extremely

important class in several metabolic routes in fungi, including genes involved in cellulose and

chitin degradation [12, 36]. Here, we found the main CAZyme families responsible for

cellulose degradation, such as GH1, GH3, GH5, GH6, GH7, GH12, GH45 and an LPMO

from the AA9 family [12, 28]. The CAZyme families responsible for hemicellulose

degradation were as follows: GH10, GH11, GH54, GH62 and CE5 [12, 30, 52-54]. Within

the GH class, an important family expected in our data was GH18, which has chitinase

activity; this family was expected since members of the genus *Trichoderma* (such as *T.*

harzianum and *T. atroviride*) are capable of mycoparasitism and because this class is directly

related to the biological control of these species [17, 35, 55].

In comparing the two *T. harzianum* strains (Th0179 and Th3844), we observed that

Th0179 presented fewer CAZyme genes related to cellulose degradation than Th3844, with

lower expression levels (Fig. 3a). However, in measuring the enzymatic activity from the

culture supernatants after 96 h of growth, we found that both strains had similar cellulase

activity profiles during growth on cellulose (Additional file 1: Fig. S1), suggesting greater

potential of Th0179 to degrade cellulose. The detected enzymatic activity is related to

proteins secreted into the medium by the cells, and only the most stable proteins are detected

in this environment [10]. It is interesting to observe which proteins found in the exoproteome

of Th0179 may respond to this increased cellulase activity. A similar profile was observed in

T. reesei, the most studied fungus within this genus and an important industrial producer of

cellulolytic enzymes [13]: few CAZyme genes have been detected in its machinery, but it has

high cellulase activity [10, 19, 56].

The identification of LPMOs as a group of enzymes that accelerate the breakdown of

carbohydrate polymers, such as cellulose, by oxidative cleavage has been a breakthrough in

lignocellulose conversion research. The supplementation of AA9 enzymes in commercial

cocktails improves the hydrolysis of lignocellulose; they assist cellulases in attacking

crystalline substrate areas, resulting in rapid and relatively complete surface degradation [19].

CAZyme genes related to cellulose degradation, including endo-β-1,4-glucanases, β-

glucosidases, and cellobiohydrolases, and CAZyme genes related to hemicellulose

degradation, such as endo-1,4-β-xylanases, were identified in the Th0179 transcriptome

under cellulose growth conditions (Table 1). In addition, we verified important secreted

CAZymes that play important roles in biomass degradation (Table 2), such as β-glucosidases

(A0A0F9XRC5 and A0A0F9XQT4), an LPMO (A0A0F9XMI8), a cellulose 1,4-β-

cellobiosidase (nonreducing end) (A0A0G0AEM7), cellulases (A0A0F9XG06 and

A0A0F9WYH5), endo-1,4-β-xylanases (A0A0F9Y0Y9, A0A0H3UCP8, and

A0A0F9XXA4), and an endo-1,4-β-mannosidase (A0A0G0AGG8). One β-glucosidase from

the GH3 family (THAR02_00656) was found in the exoproteome, which had the lowest

expression level among the CAZymes related to biomass degradation; the same pattern was

observed for *T. harzianum* IOC-3844 [10]. All of these enzymes are well known and can be

used to improve enzymatic cocktails that can function in the degradation of specific

substrates [57], such as the Brazilian biomass sugarcane bagasse. For instance, it is known

that β-glucosidases are the rate-limiting enzymes in the degradation of cellulose [58].

These enzymes play a critical role among enzymes in enzymatic cocktails for biomass

degradation. Therefore, improving the activities of such enzymes can enhance the

efficiency of commercial enzymatic cocktails for bioethanol production [3].

Horta et al. [10] analyzed the exoproteome of Th3844 and Ta0020 under equivalent

conditions, selecting a set of 80 proteins for a complete classification and analysis of

expression levels based on their transcriptomes. Comparison of the Th0179 and Th3844

exoproteomes indicated that both produced many of the main CAZymes listed above;

however, the THAR02_05896 gene, which encodes an endo-1,4-β-xylanase protein, and the

THAR02_03851 gene, which encodes a protein with endo-1,4-β-mannosidase activity, were

not detected in the Th3844 secretome. In the present study, when these exoproteomes were

compared, we noticed that most of the proteins were differentially expressed in cellulose only

for Th0179, with an emphasis on the AA9/CBM1 family (THAR02_02134), showing a 2.65-

fold change (138.62 TPM). Among the strains, Ta0020 exhibited the lowest number of

secreted proteins related to cellulose and hemicellulose degradation. Thus, the exoproteome

analysis identified key enzymes that may be fundamental for cellulose hydrolysis and that act

synergistically for efficient plant biomass degradation.

The organization of the transcriptomic data into coexpression networks using graph

theory allowed the construction of gene interaction networks that were represented by nodes

connected to edges [59]. Nodes represent the genes, and the edges represent the connections

among these genes. Correlations are determined based on the expression level of the genes

pair by pair, indicating that genes spatially closer to one another are more highly correlated

than the genes that are farther apart. The coexpression subnetwork (Fig. 5b) revealed

complex, specific relationships between CAZyme genes and genes involved in the production

and secretion of the detected proteins and is helpful for understanding the functions and

regulation of genes. Networks such as this one can be used as a platform to search for target

genes or proteins in future studies to comprehend the synergistic relationships between genes

and their regulation, which is very useful information for understanding the saccharification

process. We identified an enriched cluster with strong correlations among genes, CAZyme

genes, secreted proteins and unknown genes in the cluster analysis.

Furthermore, through the functional annotation analysis of Th0179, we identified TFs

and transporters, which should be investigated in further studies to better understand the

mechanisms by which these genes are regulated. Most sugar transporters have yet to be

characterized [1] but play important roles in taking up mono- or disaccharides into fungal

cells after biomass degradation [1, 60]. Fungal gene expression is controlled at the

transcriptional level [61], and its regulation affects the composition of enzyme mixtures;

accordingly, it is studied in several species because of its potential applications [62]. This

further emphasizes the importance of deleting and/or overexpressing TFs that regulate

specific genes directly involved in plant biomass degradation [61].

In summary, the analyses of the enzymatic activity of cellulase, the transcriptome, the

exoproteome and the coexpression networks revealed important enzymes that *T. harzianum*

CBMAI-0179 uses to hydrolyze cellulose and that most likely act synergistically to

depolymerize polysaccharides. The results suggest great potential of this strain to degrade

cellulose and can contribute to the optimization of enzymatic cocktails for bioethanol

production.

Conclusions

Bioprospecting new catalytic enzymes and improving technologies for the efficient

enzymatic conversion of plant biomass are required for advancing biofuel production. *T.*

harzianum CBMAI-0179 is a novel potential candidate producer of plant cell wall

polysaccharide-degrading enzymes that can be biotechnologically exploited for plant biomass

degradation. The cellulase activity profile indicated high efficiency and high potential of this

strain for cellulose degradation. A set of highly expressed CAZymes and proteins that are

species- and treatment-specific was observed in both the transcriptome and exoproteome

analyses. The coexpression network revealed coexpressed genes and CAZymes that act

synergistically to hydrolyze cellulose. In addition, the cluster analysis revealed genes with

strong correlations that are necessary for saccharification. Combined, these tools provide a

powerful approach for protein discovery and the selection of target genes through the

heterologous expression of proteins. In future studies, these tools can aid the selection of new

species and the optimization of the production of powerful enzymes for use in enzymatic

cocktails for second-generation bioethanol production.

Methods

Fungal strains, fermentation and enzymatic activities

The species originated from the Brazilian Collection of Environment and Industry

Microorganisms (CBMAI), located on CPQBA/UNICAMP, in Campinas, Brazil. *T.*

harzianum CBMAI-0179 (Th0179), *T. harzianum* IOC-3844 (Th3844) and *T. atroviride*

CBMAI-0020 (Ta0020) strains were grown as described in a previous work on solid medium

to produce sufficient spores for the fermentation process, which was performed in biological

triplicates using crystalline cellulose (Celuflok, São Paulo, Brazil; degree of crystallinity,

0.72 g/g; composition, 0.857 g/g cellulose and 0.146 g/g hemicellulose) or glucose as the

carbon source [10]. Glucose was used as a control in all experimental conditions.

Supernatants were collected to measure enzymatic activity and determine the exoproteome

profile.

Xylanase and β-glucosidase activities were determined using the methods described

by Bailey and Poutanen [63] and Zhang et al. [64], respectively. Cellulase activity was

determined using the filter paper activity (FPA) test according to Ghose [65]. Protein levels

were measured based on the Bradford [66] method. The enzymatic activities and protein

contents in culture supernatants were determined in a previous work and are shown in

Additional file 1: Fig. S1.

RNA extraction

Mycelial samples from cellulose and glucose conditions were extracted from Th0179,

Th3844 and Ta0020 after 96 h of fermentation, stored at -80 °C, ground in liquid nitrogen

using a mortar and pestle, and used for RNA extraction using the LiCl RNA extraction

protocol according to the method reported by Oliveira et al. [67].

Data sources

The nucleotide and protein sequences of *T. harzianum* T6776 (PRJNA252551) and *T.*

atroviride IMI 206040 (PRJNA19867) used as reference for transcriptome assembly were

downloaded from the NCBI database (www.ncbi.nlm.nig.gov).

Library construction and RNA-Seq

RNA samples were quantified using a NanoDrop 8000 (Thermo Scientific,

Wilmington, DE, USA). The libraries were constructed using 1 µg of each RNA sample

obtained from the mycelial samples and the TruSeq RNA Sample Preparation Kit v2 [68]

(Illumina Inc., San Diego, CA, USA) according to the manufacturer's instructions. The

expected target sizes were confirmed using a 2100 Bioanalyzer (Agilent Technologies, Palo

Alto, CA, USA) and the DNA 1000 Kit, and the libraries were quantified by qPCR using the

KAPA library quantification Kit for Illumina platforms (Kapa Biosystems, Wilmington, MA,

USA). The average insertion size was 260 bp. A total of 18 biological triplicate samples were

multiplexed with different adapters and organized in different lanes of the flow cell for high-

throughput sequencing. The sequencing was carried out on the HiSeq 2500 platform

(Illumina, San Diego, CA, USA) according to the manufacturer's specifications for paired-

end reads of 150 bp.

Transcriptome assembly and mapping

After sequencing was completed, the data were transferred to a local high-

performance computing server at the Center for Molecular Biology and Genetic Engineering

(CBMEG, University of Campinas, Campinas, Brazil). FastQC v0.11.5 [69] was used to

visually assess the quality of the sequencing reads. Removal of the remaining adapter

sequences and quality trimming with a sliding window of size 4, minimum quality of 15, and

length filtering (minimal length of 36 bp) was performed with Trimmomatic v0.36 [70].

The RNA-Seq data were analyzed using CLC Genomics Workbench software (v6.5.2;

CLC bio, Finlandsgade, Denmark) [71]. The reads were mapped against the reference

genomes of *T. harzianum* T6776 [33] and *T. atroviride* IMI 206040 [34] using the following

parameters: minimum length fraction = 0.5; minimum similarity fraction = 0.8; and

maximum number of hits for a read = 10. For the paired settings, the parameters were

minimum distance = 150 and maximum distance = 300, including the broken pairs counting

scheme. The gene expression values were expressed in reads per kilobase of exon model per

million mapped reads (RPKM), and the normalized value for each sample was calculated in

transcripts per million (TPM) [72]. To statistically analyze the differentially expressed genes

(DEGs), the following parameters were used: fold change greater than or equal to 1.5 or

lower than or equal to -1.5 and p-value lower than 0.05.

Gene comparisons between species

Venn diagrams were constructed to compare the genes with TPM expression values

greater than zero under both conditions from all species using

http://bioinformatics.psb.ugent.be/webtools/Venn/.

Transcriptome annotation and CAZyme determination

Sequences were functionally annotated according to the Gene Ontology (GO) terms

[73] with Blast2Go v4.1.9 [74] using BLASTx-fast and a cutoff e-value of 10^{-6}. Information

derived from the CAZy database [24] was downloaded (www.cazy.org) to locally build a

CAZy database (2017). The protein sequences of *T. harzianum* T6776 and *T. atroviride* IMI

206040 were used as queries in basic local alignment search tool (BLAST) searches against

the locally built CAZy BLAST database. BLAST matches showing an e-value less than 10^{-11},

identity greater than 30% and queries covering greater than 70% of the sequence length were

selected and classified according to the CAZyme catalytic group as GHs, CBMs, GTs, CEs,

AAs or PLs and their respective CAZyme families.

CAZymes were also annotated according to Enzyme Commission (EC) number [75]

through BRENDA (Braunschweig Enzyme Database) [76] (www.brenda-enzymes.org), using

BLASTp with an e-value cutoff of 10^{-10}, identity greater than 30% and queries covering

greater than 60% of the sequence length.

Coexpression networks

The coexpression networks for Th0179 were assembled from the mapped RNA-Seq

data against the genome of *T. harzianum* T6776 using biological triplicates. The network was

assembled by calculating Pearson's correlation coefficient for each pair of genes. Genes

showing null values for most of the replicates under different experimental conditions were

excluded to decrease noise and to remove residuals from the analysis. The highest reciprocal

rank (HRR) method proposed by Mutwil et al. [77] was used to empirically filter the edges,

retaining edges with an HRR less than or equal to 30. Thus, only edges representing the

strongest correlations were selected. Cytoscape software v3.6.0 [78] was used for data

analysis and network construction. The cluster analysis procedure was performed with the

Heuristic Cluster Chiseling Algorithm (HCCA) [77].

Exoproteome analysis

The analysis of the exoproteome of Th0179 under both fermentative conditions was

performed via liquid chromatography tandem mass spectrometry (LC-MS/MS) using the

data-independent method of acquisition MS^E as described by Horta et al. [10] The LC-

MS/MS data were processed using ProteinLynx Global Server (PLGS) v3.0.1 software

(Waters, Milford, MA, USA), and the proteins in the processed files (.pkl) were identified by

comparison to the *Trichoderma* sequence database available in the UniProt Knowledgebase

(UniProtKB; https://www.uniprot.org/uniprot/). BLASTp searches of the fasta sequences of

the identified proteins were performed against the *T. harzianum* T6776 genome to identify

the secreted proteins and compare them to the transcriptome.

RT-qPCR analysis

To validate the expression profiles of the transcriptome analysis, reverse

transcription-quantitative PCR (RT-qPCR) was performed for selected genes. The RNeasy

Mini Kit (Qiagen) was used for RNA extraction, and cDNA was synthesized using the

QuantiTect Reverse Transcription Kit (Qiagen, Germany) according to the manufacturer's

instructions. DEGs with a high number of reads and TPM were selected and are shown in

Additional file 2: Table S1, together with the primers and annealing temperatures. Primers

were synthesized using the Primer3Plus web interface [79] with a fusion temperature between

58 and 60 °C and amplicon sizes between 120 and 200 bp.

Quantification of gene expression was performed by continuously monitoring SYBR

Green fluorescence. The reactions were performed in triplicate in a total volume of 6.22 μL.

Each reaction contained 3.12 μL of SYBR Green Supermix (Bio-Rad, USA), 1.0 μL of

forward and reverse primers and 2.1 μL of diluted cDNA. The reactions were assembled in

384-well plates. PCR amplification-based expression profiling of the selected genes was

performed using specific endogenous controls for each strain, which are described in

Additional file 2: Table S4. RT-qPCR was conducted with the CFX384 Touch Real-Time

PCR Detection System (Bio-Rad). The real-time PCR program was as follows: initial

denaturation at 95 °C for 10 min, followed by 40 cycles of 15 sec at 95 °C and 60 sec at 60

°C. Gene expression was calculated via the delta-delta cycle threshold method [80]. The

obtained RT-qPCR results were compared with the RNA-Seq results from the generated

assemblies. The selected genes exhibited the same expression profiles between the RT-qPCR

and RNA-Seq analyses (Additional file 3: Fig. S2).

Additional files

Additional file 1: Fig. S1. Enzymatic activity. Enzymatic activities (UI mL^{-1}) of β-

glucosidase (a), cellulase (b), and xylanase (c) and protein contents (d) in the culture

supernatants of *Trichoderma* spp. measured after 96 h of growth. Each bar represents the

mean and standard deviation of biological triplicates.

Additional file 2: Table S1. Sequences of the primers from the endogenous controls and

DEGs in RNA-Seq data selected for RT-qPCR.

Additional file 3: Fig. S2. RNA-Seq analysis validation. The obtained RT-qPCR results

were compared with the RNA-Seq results for transcriptome analysis validation of

Trichoderma spp.

Additional file 4: Fig. S3. Venn diagrams. Venn diagrams of the genes identified in

Trichoderma spp. with expression levels higher than zero under cellulose (a) and glucose (b)

growth conditions using the *T. harzianum* T6776 genome as a reference.

Additional file 5: Table S2. Classification of the CAZyme genes under cellulose conditions

for *T. harzianum* IOC-3844 and *T. atroviride* CBMAI-0020.

Additional file 6: Table S3. Proteins identified in both the transcriptome and exoproteome of

T. harzianum CBMAI-0179 under both glucose and cellulose growth conditions.

Additional file 7: Table S4. Genes identified in the cluster analysis from the main

coexpression network of *T. harzianum* CBMAI-0179.

List of abbreviations

AA: Auxiliary enzymes; **BLAST:** Basic Local Alignment Search Tool; **bp:** Base pair;

BRENDA: Braunschweig Enzyme Database; **CAZymes:** Carbohydrate-active enzymes;

CBM: Carbohydrate-binding module; **cDNA:** Complementary DNA; **CE:** Carbohydrate

esterases; **CEL:** Cellulose; **DEGs:** Differentially expressed genes; **EC:** Enzyme commission

number; **FPA:** Filter paper activity; **GH:** Glycoside hydrolases; **GLU:** Glucose; **GO:** Gene

Ontology; **GT:** Glycosyltransferases; **HCCA:** Heuristic Cluster Chiseling Algorithm; **HRR:**

Highest reciprocal rank; **kb:** Kilobases; **LPMO:** Lytic polysaccharides monooxygenase;

MA2: Malt extract agar, 2% w/w; **Mb:** Megabase; **mL:** Milliliter; **mRNA:** Messenger RNA;

PCA: Principal component analysis; **PDA:** Potato dextrose agar; **PL:** Polysaccharide lyases;

RNA: Ribonucleic acid; **RNA-Seq:** RNA sequencing; **RPKM:** Reads per kilobase of exon

model per million mapped reads; **RT-qPCR:** Real-time quantitative PCR; **Ta0020:**

Trichoderma atroviride CBMAI-0020; **TFs:** Transcription factors; **Th0179:** *Trichoderma*

harzianum CBMAI-0179; **Th3844:** *Trichoderma harzianum* IOC-3844; **TPM:** Transcripts

per million; **UI:** International Unit; **µl:** Microliter

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The datasets generated and/or analyzed during this study are included in this published article

and Additional files 1, 2, 3, 4, 5, 6 and 7. The reads have been deposited at the NCBI

Sequence Read Archive and can be accessed under the BioProject number PRJNA336221.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by grants from the Fundação de Amparo à Pesquisa do Estado de

São Paulo (FAPESP 2015/09202-0, 2018/19660-4), Coordenação de Aperfeiçoamento de

Pessoal de Nível Superior (CAPES, Computational Biology Program) and Conselho Nacional

de Desenvolvimento Científico e Tecnológico (CNPq). DAA received an MS fellowship

from FAPESP (2017/17782-2) and CAPES – Computational Biology Program

(88882.160100/2017-01, 88887.336686/2019-00). MACH received a PD fellowship from

FAPESP (2018/18856-1). JAFF received a PhD fellowship from CNPq (170565/2017-3).

NFM received a PD fellowship from CNPq and CAPES, Computational Biology Program,

and APS is the recipient of a research fellowship from CNPq.

Authors' contributions

DAA and APS conceived and designed the study. DAA, MACH, JAFF and NFM performed

the data analysis. DAA drafted the manuscript, which was critically revised by MACH, JAFF

and APS. All authors read and approved the final manuscript.

Acknowledgements

We would like to acknowledge the funding from Fundação de Amparo à Pesquisa do Estado

de São Paulo (FAPESP 2015/09202-0), Coordenação de Aperfeiçoamento de Pessoal de

Nível Superior (CAPES, Computational Biology Program) and Conselho Nacional de

Desenvolvimento Científico e Tecnológico (CNPq). We thank the National Institute of

Metrology, Quality and Technology (INMETRO) for performing the proteomics analysis via

LC-MS/MS, the Brazilian Biorenewables National Laboratory (LNBR), Campinas – SP, for

conducting the fermentation experiments and the Center of Molecular Biology and Genetic

Engineering (CBMEG) at the University of Campinas, SP, for use of the center and

laboratory space.

References

1.   de Gouvêa PF, Bernardi AV, Gerolamo LE, Santos EDS, Riaño-Pachón DM,

Uyemura SA, et al. Transcriptome and secretome analysis of *Aspergillus fumigatus* in

the presence of sugarcane bagasse. BMC Genomics. 2018;19:232.

2.   Castro LDS, Pedersoli WR, Antoniêto ACC, Steindorff AS, Silva-Rocha R, Martinez-

Rossi NM, et al. Comparative metabolism of cellulose, sophorose and glucose in

Trichoderma reeseiusing high-throughput genomic and proteomic analyses.

Biotechnol Biofuels. 2014;7:41.

3.   Santos CA, Morais MAB, Terrett OM, Lyczakowski JJ, Zanphorlin LM, Ferreira-

Filho JA, et al. An engineered GH1 β-glucosidase displays enhanced glucose

tolerance and increased sugar release from lignocellulosic materials. Sci Rep.

2019;9:4903.

4.   van Dyk JS, Pletschke BI. A review of lignocellulose bioconversion using enzymatic

hydrolysis and synergistic cooperation between enzymes-factors affecting enzymes,

conversion and synergy. Biotechnol Adv. 2012;30:1458-80.

5.   Wang Y, Fan C, Hu H, Li Y, Sun D, Wang Y, et al. Genetic modification of plant cell

walls to enhance biomass yield and biofuel production in bioenergy crops. Biotechnol

Adv. 2016;34:997-1017.

6.   Rocha VAL, Maeda RN, Pereira N, Kern MF, Elias L, Simister R, et al.

Characterization of the cellulolytic secretome of *Trichoderma harzianum* during

growth on sugarcane bagasse and analysis of the activity boosting effects of

swollenin. Biotechnol Prog. 2016;32:327-36.

7.   Ahmed S, Mustafa G, Arshad M, Rajoka MI. Fungal biomass protein production from

Trichoderma harzianum using rice polishing. BioMed Res Int. 2017;2017:6232793.

8.   de Souza WR. Microbial degradation of lignocellulosic biomass**.** In: Chandel AK, da

Silva SS, editors. Sustainable degradation of lignocellulosic biomass - techniques,

applications and commercialization. London, UK: IntechOpen; 2013. p. 207-46.

9.   Miao Y, Liu D, Li G, Li P, Xu Y, Shen Q, et al. Genome-wide transcriptomic analysis

of a superior biomass-degrading strain of *A. fumigatus* revealed active lignocellulose-

degrading genes. BMC Genomics. 2015;16:459.

10.   Horta MAC, Filho JAF, Murad NF, Santos EDO, dos Santos CA, Mendes JS, et al.

Network of proteins, enzymes and genes linked to biomass degradation shared by

Trichoderma species. Sci Rep. 2018;8:1341.

11.   Kumar M, Ashraf S. Role of *Trichoderma* spp. as a biocontrol agent of fungal plant

693   pathogens**.** In: Kumar V, Kumar M, Sharma S, Prasad R, editors. Probiotics and plant

694   health. Singapore: Springer Singapore; 2017. p. 497-506.

695   12.   Filho JAF, Horta MAC, Beloti LL, dos Santos CA, de Souza AP. Carbohydrate-active

696   enzymes in *Trichoderma harzianum*: a bioinformatic analysis bioprospecting for key

697   enzymes for the biofuels industry. BMC Genomics. 2017;18:779.

698   13.   Druzhinina IS, Seidl-Seiboth V, Herrera-Estrella A, Horwitz BA, Kenerley CM,

699   Monte E, et al. *Trichoderma*: the genomics of opportunistic success. Nat Rev

700   Microbiol. 2011;9:749-59.

701   14.   Mukherjee PK, Horwitz BA, Herrera-Estrella A, Schmoll M, Kenerley CM.

702   *Trichoderma* research in the genome era. Annu Rev Phytopathol. 2013;51:105-29.

703   15.   Ghildiyal A, Pandey A. Isolation of cold tolerant antifungal strains of *Trichoderma*

704   sp. from glacial sites of Indian Himalayan region. Res J Microbiol. 2008;3:559-64.

705   16.   Peterson R, Nevalainen H. *Trichoderma reesei* RUT-C30 – thirty years of strain

706   improvement. Microbiology. 2012;158:58-68.

707   17.   Martinez D, Berka RM, Henrissat B, Saloheimo M, Arvas M, Baker SE, et al.

708   Genome sequencing and analysis of the biomass-degrading fungus *Trichoderma*

709   *reesei* (syn. *Hypocrea jecorina*). Nat Biotechnol. 2008;26:553-60.

710   18.   Margolles-Clark E, Ihnen M, Penttilä M. Expression patterns of ten hemicellulase

711   genes of the filamentous fungus *Trichoderma reesei* on various carbon sources. J

712   Biotechnol. 1997;57:167-79.

713   19.   Druzhinina IS, Kubicek CP. Genetic engineering of *Trichoderma reesei* cellulases and

714   their production. Microb Biotechnol. 2017;10:1485-99.

20. Benoliel B, Torres FAG, de Moraes LMP. A novel promising *Trichoderma*

harzianum strain for the production of a cellulolytic complex using sugarcane bagasse

in natura. SpringerPlus. 2013;2:656.

21. Delabona PDS, Farinas CS, da Silva MR, Azzoni SF, Pradella JGDC. Use of a new

Trichoderma harzianum strain isolated from the Amazon rainforest with pretreated

sugar cane bagasse for on-site cellulase production. Bioresour Technol.

2012;107:517-21.

22. Horta MAC, Vicentini R, Delabona PDS, Laborda P, Crucello A, Freitas S, et al.

Transcriptome profile of *Trichoderma harzianum* IOC-3844 induced by sugarcane

bagasse. PLoS One. 2014;9:e88689.

23. Delabona PDS, Cota J, Hoffmam ZB, Paixão DAA, Farinas CS, Cairo JPLF, et al.

Understanding the cellulolytic system of *Trichoderma harzianum* P49P11 and

enhancing saccharification of pretreated sugarcane bagasse by supplementation with

pectinase and α-l-arabinofuranosidase. Bioresour Technol. 2013;131:500-7.

24. Lombard V, Ramulu HG, Drula E, Coutinho PM, Henrissat B. The carbohydrate-

active enzymes database (CAZy) in 2013. Nucleic Acids Res. 2014;42:D490-5.

25. Cairo JPLF, Leonardo FC, Alvarez TM, Ribeiro DA, Büchli F, Costa-Leonardo AM,

et al. Functional characterization and target discovery of glycoside hydrolases from

the digestome of the lower termite *Coptotermes gestroi*. Biotechnol Biofuels.

2011;4:50.

26. Montella S, Ventorino V, Lombard V, Henrissat B, Pepe O, Faraco V. Discovery of

genes coding for carbohydrate-active enzyme by metagenomic analysis of

lignocellulosic biomasses. Sci Rep. 2017;7:42623.

27. Suwannarangsee S, Bunterngsook B, Arnthong J, Paemanee A, Thamchaipenet A,

Eurwilaichitr L, et al. Optimisation of synergistic biomass-degrading enzyme systems

for efficient rice straw hydrolysis using an experimental mixture design. Bioresour

Technol. 2012;119:252-61.

28.   Villares A, Moreau C, Bennati-Granier C, Garajova S, Foucat L, Falourd X, et al.

Lytic polysaccharide monooxygenases disrupt the cellulose fibers structure. Sci Rep.

2017;7:40262.

29.   Bennati-Granier C, Garajova S, Champion C, Grisel S, Haon M, Zhou S, et al.

Substrate specificity and regioselectivity of fungal AA9 lytic polysaccharide

monooxygenases secreted by *Podospora anserina*. Biotechnol Biofuels. 2015;8:90.

30.   Bischof RH, Ramoni J, Seiboth B. Cellulases and beyond: the first 70 years of the

enzyme producer *Trichoderma reesei*. Microb Cell Fact. 2016;15:106.

31.   Ellegren H, Galtier N. Determinants of genetic diversity. Nat Rev Genet.

2016;17:422-33.

32.   Al-Sadi AM, Al-Oweisi FA, Edwards SG, Al-Nadabi H, Al-Fahdi AM. Genetic

analysis reveals diversity and genetic relationship among *Trichoderma* isolates from

potting media, cultivated soil and uncultivated soil. BMC Microbiology. 2015;15:147.

33.   Baroncelli R, Piaggeschi G, Fiorini L, Bertolini E, Zapparata A, Pè ME, et al. Draft

whole-genome sequence of the biocontrol agent *Trichoderma harzianum* T6776.

Genome Announc. 2015;3:e00647-15.

34.   Kubicek CP, Herrera-Estrella A, Seidl-Seiboth V, Martinez DA, Druzhinina IS, Thon

M, et al. Comparative genome sequence analysis underscores mycoparasitism as the

ancestral life style of *Trichoderma*. Genome Biol. 2011;12:R40.

35.   Limón MC, Chacón MR, Mejías R, Delgado-Jarana J, Rincón AM, Codón AC, et al.

Increased antifungal and chitinase specific activities of *Trichoderma harzianum*

CECT 2413 by addition of a cellulose binding domain. Appl Microbiol Biotechnol.

2004;64:675-85.

36.   Pellegrini VOA, Serpa VI, Godoy AS, Camilo CM, Bernardes A, Rezende CA, et al.

766   Recombinant *Trichoderma harzianum* endoglucanase I (Cel7B) is a highly acidic and

767   promiscuous carbohydrate-active enzyme. Appl Microbiol Biotechnol. 2015;99:9591-

768   604.

769   37.   Valadares F, Gonçalves TA, Gonçalves DSPO, Segato F, Romanel E, Milagres AMF,

770   et al. Exploring glycoside hydrolases and accessory proteins from wood decay fungi

771   to enhance sugarcane bagasse saccharification. Biotechnol Biofuels. 2016;9:110.

772   38.   Levasseur A, Drula E, Lombard V, Coutinho PM, Henrissat B. Expansion of the

773   enzymatic repertoire of the CAZy database to integrate auxiliary redox enzymes.

774   Biotechnol Biofuels. 2013;6:41.

775   39.   Johansen KS. Lytic polysaccharide monooxygenases: the microbial power tool for

776   lignocellulose degradation. Trends Plant Sci. 2016;21:926-36.

777   40.   Alfaro M, Castanera R, Lavín JL, Grigoriev IV, Oguiza JA, Ramírez L, et al.

778   Comparative and transcriptional analysis of the predicted secretome in the

779   lignocellulose-degrading basidiomycete fungus *Pleurotus ostreatus*. Environ

780   Microbiol. 2016;18:4710-26.

781   41.   Saykhedkar S, Ray A, Ayoubi-Canaan P, Hartson SD, Prade R, Mort AJ. A time

782   course analysis of the extracellular proteome of *Aspergillus nidulans* growing on

783   sorghum stover. Biotechnol Biofuels. 2012;5:52.

784   42.   Borin GP, Sanchez CC, de Santana ES, Zanini GK, dos Santos RAC, Pontes ADO, et

785   al. Comparative transcriptome analysis reveals different strategies for degradation of

786   steam-exploded sugarcane bagasse by *Aspergillus niger* and *Trichoderma reesei*.

787   BMC Genomics. 2017;18:501.

788   43.   Borin GP, Sanchez CC, de Souza AP, de Santana ES, de Souza AT, Leme AFP, et al.

789   Comparative secretome analysis of *Trichoderma reesei* and *Aspergillus niger* during

790   growth on sugarcane biomass. PLoS One. 2015;10:e0129275.

791   44.   Vicentini R, Bottcher A, Brito MDS, dos Santos AB, Creste S, Landell MG, et al.

792   Large-scale transcriptome analysis of two sugarcane genotypes contrasting for lignin

793   content. PLoS One. 2015;10:e0134909.

794   45.   Aranda-Martinez A, Lenfant N, Escudero N, Zavala-Gonzalez EA, Henrissat B,

795   Lopez-Llorca LV. CAZyme content of *Pochonia chlamydosporia* reflects that chitin

796   and chitosan modification are involved in nematode parasitism. Environ Microbiol.

797   2016;18:4200-15.

798   46.   Leelavathi MS, Vani L, Reena P. Antimicrobial activity of *Trichoderma harzianum*

799   against bacteria and fungi. Int J Curr Microbiol App Sci. 2014;3:96-103.

800   47.   Contreras-Cornejo HA, Macías-Rodríguez L, Cortés-Penagos C, López-Bucio J.

801   *Trichoderma virens*, a plant beneficial fungus, enhances biomass production and

802   promotes lateral root growth through an auxin-dependent mechanism in *Arabidopsis*.

803   Plant Physiol. 2009;149:1579-92.

804   48.   Jang S, Kwon SL, Lee H, Jang Y, Park MS, Lim YW, et al. New report of three

805   unrecorded species in *Trichoderma harzianum* species complex in Korea.

806   Mycobiology. 2018;46:177-84.

807   49.   Sharma PK, Gothalwal R. *Trichoderma*: a potent fungus as biological control agent.

808   In: Singh JS, Seneviratne G, editors. Agro-environmental sustainability: volume 1:

809   managing crop health. Cham, Switzerland: Springer International Publishing; 2017. p.

810   113-25.

811   50.   Xie B-B, Qin Q-L, Shi M, Chen L-L, Shu Y-L, Luo Y, et al. Comparative genomics

812   provide insights into evolution of *Trichoderma* nutrition style. Genome Biol Evol.

813   2014;6:379-90.

814   51.   Brunner K, Zeilinger S, Ciliento R, Woo SL, Lorito M, Kubicek CP, et al.

815   Improvement of the fungal biocontrol agent *Trichoderma* atroviride to enhance both

816   antagonism and induction of plant systemic disease resistance. Appl Environ

817   Microbiol. 2005;71:3959.

818   52.   Zhao Z, Liu H, Wang C, Xu J-R. Correction: comparative analysis of fungal genomes

819   reveals different plant cell wall degrading capacity in fungi. BMC Genomics.

820   2014;15:274.

821   53.   Sweeney MD, Xu F. Biomass converting enzymes as industrial biocatalysts for fuels

822   and chemicals: recent developments. Catalysts. 2012;2:244-63.

823   54.   Javier PFI, Óscar G, Sanz-Aparicio J, Díaz P. Xylanases: molecular properties and

824   applications**.** In: Polaina J, MacCabe AP, editors. Industrial enzymes: structure,

825   function and applications. Dordrecht, Netherlands: Springer Netherlands; 2007. p. 65-

826   82.

827   55.   Binod P, Sukumaran RK, Shirke SV, Rajput JC, Pandey A. Evaluation of fungal

828   culture filtrate containing chitinase as a biocontrol agent against *Helicoverpa*

829   *armigera*. J Appl Microbiol. 2007;103:1845-52.

830   56.   Manika S, Saju S, Subhash C, Mukesh S, Sharma P. Comparative evaluation of

831   cellulase activity in *Trichoderma harzianum* and *Trichoderma reesei*. Afr J Microbiol

832   Res. 2014;8:1939-47.

833   57.   Lopes AM, Filho EXF, Moreira LRS. An update on enzymatic cocktails for

834   lignocellulose breakdown. J Appl Microbiol. 2018;125:632-45.

835   58.   Zang X, Liu M, Fan Y, Xu J, Xu X, Li H. The structural and functional contributions

836   of β-glucosidase-producing microbial communities to cellulose degradation in

837   composting. Biotechnol Biofuels. 2018;11:51.

838   59.   Azevedo H, Bando S, Bertonha F, Moreira-Filho CA. Redes de interação gênica e

839   controle epigenético na transição saúde-doença. Rev Med. 2015;94:223-9.

840   60.   Peng M, Aguilar-Pontes MV, de Vries RP, Mäkelä MR. In silico analysis of putative

841   sugar transporter genes in *Aspergillus niger* using phylogeny and comparative

842   transcriptomics. Front Microbiol. 2018;9:1045.

843   61.   Liu R, Chen L, Jiang Y, Zou G, Zhou Z. A novel transcription factor specifically

844   regulates GH11 xylanase genes in *Trichoderma reesei*. Biotechnol Biofuels.

845   2017;10:194.

846   62.   Benocci T, Aguilar-Pontes MV, Zhou M, Seiboth B, de Vries RP. Regulators of plant

847   biomass degradation in *Ascomycetous fungi*. Biotechnol Biofuels. 2017;10:152.

848   63.   Bailey MJ, Poutanen K. Production of xylanolytic enzymes by strains of *Aspergillus*.

849   Appl Microbiol Biotechnol. 1989;30:5-10.

850   64.   Zhang YHP, Hong J, Ye X. Cellulase assays. In: Mielenz JR, editors. Biofuels:

851   methods and protocols. Totowa, NJ: Humana Press; 2009. p. 213-31.

852   65.   Ghose TK. Measurement of cellulase activities. Pure Appl Chem. 1987;59:257-68.

853   66.   Bradford MM. A rapid and sensitive method for the quantitation of microgram

854   quantities of protein utilizing the principle of protein-dye binding. Anal Biochem.

855   1976;72:248-54.

856   67.   Oliveira RR, Viana AJC, Reátegui ACE, Vincentz MGA. Short communication an

857   efficient method for simultaneous extraction of high-quality RNA and DNA from

858   various plant tissues. Genet Mol Res. 2015;14:18828-38.

859   68.   Illumina. TruSeq RNA, sample preparation v2 guide. San Diego, US: Illumina; 2014.

860   69.   Andrews S. FastQC: a quality control tool for high throughput sequence data.

861   http://www.bioinformatics.babraham.ac.uk/projects/fastqc (2010). Accessed 25 Mar

862   2017.

863   70.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

864   sequence data. Bioinformatics. 2014;30:2114-20.

865   71.   CLC Genomics Workbench. Manual for CLC genomics workbench 6.5.2 Windows,

866   Mac OS X and Linux Denmark. Aarhus, Denmark: QIAGEN (Aarhus A/S); 2016.

867   72.   Conesa A, Madrigal P, Tarazona S, Gomez-Cabrero D, Cervera A, McPherson A, et

868   al. A survey of best practices for RNA-seq data analysis. Genome Biol. 2016;17:13.

869   73.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene

870   ontology: tool for the unification of biology. The gene ontology consortium. Nat

871   Genet. 2000;25:25-9.

872   74.   Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M, Robles M. Blast2GO: a

873   universal tool for annotation, visualization and analysis in functional genomics

874   research. Bioinformatics. 2005;21:3674-6.

875   75.   Yamanishi Y, Hattori M, Kotera M, Goto S, Kanehisa M. E-zyme: predicting

876   potential EC numbers from the chemical transformation pattern of substrate-product

877   pairs. Bioinformatics. 2009;25:i179-86.

878   76.   Schomburg I, Jeske L, Ulbrich M, Placzek S, Chang A, Schomburg D. The BRENDA

879   enzyme information system–from a database to an expert system. J Biotechnol.

880   2017;261:194-206.

881   77.   Mutwil M, Klie S, Tohge T, Giorgi FM, Wilkins O, Campbell MM, et al. PlaNet:

882   combined sequence and expression comparisons across plant networks derived from

883   seven species. Plant Cell. 2011;23:895-910.

78. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a

software environment for integrated models of biomolecular interaction networks.

Genome Res. 2003;13:2498-504.

79. Untergasser A, Nijveen H, Rao X, Bisseling T, Geurts R, Leunissen JAM.

Primer3Plus, an enhanced web interface to primer3. Nucleic Acids Res.

2007;35:W71-4.

80. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time

quantitative PCR and the 2(-delta delta C(T)) method. Methods. 2001;25:402-8.

Figure legends

Fig. 1. The transcriptome profiles, gene expression comparison and the main CAZyme

classes identified for all strains. PCA of transcriptome mapping according to species and

growth conditions (CEL – cellulose and GLU – glucose) using the *T. harzianum* T6776

genome as a reference (a), number of DEGs and differentially expressed CAZyme genes in *T.*

harzianum CBMAI-0179 (Th0179), *T. harzianum* IOC-3844 (Th3844), and *T. atroviride*

CBMAI-0020 (Ta0020) under cellulose growth conditions (b), the differentially expressed

CAZyme classes identified for each strain under cellulose growth conditions (c).

Fig. 2. Distribution of CAZyme families in *Trichoderma* spp. Classification and

quantification of CAZyme families in *T. harzianum* CBMAI-0179 (a), *T. harzianum* IOC-

3844 (b), and *T. atroviride* CBMAI-0020 (c) under cellulose growth conditions.

Fig. 3. Evaluation of CAZyme family expression in *Trichoderma* spp. via RNA-Seq.

Quantification of the expression of the main families related to cellulose (a) and

hemicellulose (b) degradation in TPM.

Fig. 4. GO terms of *T. harzianum* CBMAI-0179 under cellulose growth conditions. The

sequences were annotated according to the main GO terms: molecular function (a), biological

process (b), and cellular component (c).

Fig. 5. Coexpression networks of *T. harzianum* CBMAI-0179. Complete coexpression

network (a), the coexpression subnetwork based on the exoproteome data (b), the enriched

cluster analysis of the coexpression network (c), the subnetwork of the CAZyme genes and

the secreted proteins identified in the cluster analysis (d). Red squares indicate DEGs under

cellulose growth conditions, blue squares indicate DEGs under glucose growth conditions,

yellow triangles indicate CAZyme genes under cellulose growth conditions, light blue

triangles indicate CAZyme genes under glucose growth conditions and purple hexagons

indicate the secreted proteins under cellulose growth conditions.

Tables

Table 1. Classification of the CAZyme genes under cellulose growth conditions for *T.*

harzianum CBMAI-0179

Gene ID	Protein product	Fold change	e-value	CAZy classification	Enzyme activity	EC number	Cellulose TPM	Glucose TPM
THAR02 _00068	KKP07860.1	1.87	1.00E-143	GH18 CBM1	chitinase	3.2.1.14	42.13	22.14
THAR02 _00890	KKP07011.1	2.10	4.00E-15	GH3	beta-glucosidase	3.2.1.21	68.64	32.01
THAR02 _01434	KKP06476.1	1.75	0	GH16	endo-1,3(4)-beta-glucanase	3.2.1.6	341.65	191.83
THAR02 _01911	KKP05955.1	1.64	2.00E-53	GT4	1-acyl-sn-glycerol-3-phosphate acyltransferase	-	94.76	56.86
THAR02 _02132	KKP05758.1	2.89	0	GH3	beta-glucosidase	3.2.1.21	16.39	5.56
THAR02 _02133	KKP05759.1	2.27	3.00E-160	CBM1	cellulase	3.2.1.4	47.86	20.67
THAR02 _02134	KKP05760.1	2.60	0	AA9 CBM1	cellulase	3.2.1.4	138.62	52.35
THAR02 _02251	KKP05610.1	2.21	0	GH1	beta-glucosidase	3.2.1.21	88.05	39.02
THAR02 _02560	KKP05371.1	1.97	2.00E-29	GH18	uncharacterized protein	-	151.83	75.55
THAR02	KKP04958.1	2.05	3.00E-114	GH45 CBM1	cellulase	3.2.1.4	52.97	25.32

_02979

THAR02 _03008	KKP04907.1	2.02	7.00E-30	GH55	glucan 1,3-beta-glucosidase	3.2.1.58	28.88	14.01
THAR02 _03217	KKP04674.1	1.52	8.00E-14	GH17	hypothetical protein THAR02_03217	-	71.68	46.17
THAR02 _03271	KKP04658.1	2.25	5.00E-137	GH10	endo-1,4-beta-xylanase	3.2.1.8	47.58	20.75
THAR02 _03302	KKP04612.1	1.85	4.00E-95	GH16	glucan endo-1,3-beta-D-glucosidase	3.2.1.39	212.11	112.64
THAR02 _04021	KKP03872.1	2.11	2.00E-84	GH72	1,3-beta-glucanosyltransferase	2.4.1.-	178.64	82.96
THAR02 _04405	KKP03485.1	2.09	0	GH5 CBM1	cellulase	3.2.1.4	39.92	18.77
THAR02 _04414	KKP03494.1	1.52	0	GH6 CBM1	cellulose 1,4-beta-cellobiosidase (nonreducing end)	3.2.1.91	299.66	192.97
THAR02 _05432	KKP02477.1	2.64	0	GH1	beta-glucosidase	3.2.1.21	125.51	46.73
THAR02 _05677	KKP02215.1	40.48	2.00E-127	AA7	UDP-N-acetylmuramate dehydrogenase	-	20.74	0.50
THAR02 _07531	KKP00372.1	2.05	0	GH20	beta-N-acetylhexosaminidase	3.2.1.52	1925.74	920.50
THAR02 _07716	KKP00192.1	1.98	4.00E-92	GH64 CBM6	glucanase B	-	271.18	134.28
THAR02 _07777	KKP00125.1	1.79	0	GH18 CBM1	chitinase	3.2.1.14	108.01	59.24
THAR02 _07958	KKO99924.1	2.60	3.00E-162	GH27	alpha-galactosidase	3.2.1.22	22.70	8.58
THAR02 _08897	KKO99004.1	1.73	0	GH7 CBM1	cellulose 1,4-beta-cellobiosidase (reducing end)	3.2.1.176	369.92	209.24
THAR02 _10108	KKO97789.1	1.51	4.00E-29	CE9	glucosamine-6-phosphate isomerase	-	3705.87	2411.20
THAR02 _10110	KKO97791.1	1.64	0	CE9	N-acetylglucosamine-6-phosphate deacetylase	3.5.1.25	6362.40	3809.76
THAR02 _10273	KKO97625.1	1.62	0	GH17	glucan endo-1,3-beta-D-glucosidase	3.2.1.39	95.61	57.93

923 **Table 2.** Proteins identified in both the transcriptome and exoproteome of *T. harzianum*

924 CBMAI-0179 grown in cellulose

Gene ID	Accession number	Protein name	e-value	CAZy classification	EC number	Cellulose TPM	Glucose TPM
THAR02_00377	G0RX84	Predicted protein	3.00E-15	-	-	1113.19	1128.71
THAR02_00656	A0A0F9XRC5	Beta-glucosidase	0	GH3	3.2.1.21	5.08	2.55
THAR02_00890	A0A0F9XQT4	Beta-glucosidase	0	GH3	3.2.1.21	68.64	32.01
THAR02_01069	G9MX73	Glycoside hydrolase family 64 protein	1.00E-64	GH64	-	17.67	25.95
THAR02_01434	A0A0F9XP75	Uncharacterized protein	0	GH16	3.2.1.6	341.65	191.83
THAR02_01982	A0A0F9Y1F6	Beta-galactosidase	0	GH35	3.2.1.23	10.08	6.85
THAR02_02133	A0A0G0AME2	Uncharacterized protein	0	CBM1	-	47.86	20.67
THAR02_02134	A0A0F9XMI8	Cellulase	0	AA9 CBM1	3.2.1.4	138.62	52.35
THAR02_02147	A0A0F9Y0Y9	Endo-1,4-beta-xylanase	0	GH11 CBM1	3.2.1.8	119.56	86.24
THAR02_02289	A0A0F9Y0G5	Cel74a	0	CBM1	-	22.53	24.27
THAR02_03210	A0A0F9ZXC9	WSC domain-containing protein	0	AA5_1	-	33.57	38.75
THAR02_03271	A0A0F9XXA4	Beta-xylanase	0	GH10	3.2.1.8	47.58	20.75
THAR02_03624	G0RX52	Extracellular metalloproteinase (Fungalysin)	0	CBM50	3.4.24.-	1.02	0.31
THAR02_03851	A0A0G0AGG8	Mannan endo-1,4-β-mannosidase	0	GH5 CBM1	3.2.1.78	67.98	57.54
THAR02_04062	A0A0F9XH17	Uncharacterized protein	2.00E-170	-	-	21.80	18.38
THAR02_04344	G9MY63	Glycoside hydrolase family 71 protein	0	GH71 CBM24	3.2.1.59	3.82	3.87
THAR02_04405	A0A0F9XG06	Cellulase	0	GH5 CBM1	3.2.1.4	39.92	18.77
THAR02_04414	A0A0G0AEM7	Cellulose 1,4-β-cellobiosidase (nonreducing end)	0	GH6 CBM1	3.2.1.91	299.66	192.97
THAR02_04626	G9NK86	Glycoside hydrolase family 92 protein	0	GH92	-	1.62	0.36
THAR02_04782	A0A024HVI0	Chitinase 18-5 (Fragment)	6.00E-80	GH18	-	21.24	16.93
THAR02_05380	A0A0F9XQN9	Uncharacterized protein	3.00E-171	-	-	102.99	81.78
THAR02_05501	G0R911	Glycoside hydrolase family 92	0	GH92	-	3.88	2.03
THAR02_05896	A0A0H3UCP8	Endo-1,4-beta-xylanase	8.00E-156	GH11	3.2.1.8	33.19	27.46
THAR02_06252	A0A0F9XN06	Murein transglycosylase	0	GH71 CBM24	3.2.1.59	81.33	131.74
THAR02_07321	A0A0F9X7S7	Uncharacterized protein	0	-	-	33.34	42.55
THAR02_07975	G0RXE3	Predicted protein	0	-	-	8.39	15.64
THAR02_08235	A0A0F9ZHA7	Chitinase 3	0	GH18	3.2.1.96	62.16	62.00
THAR02_08478	A0A0G0A296	Uncharacterized protein	0	GH30_7	3.2.1.-	6.37	2.67
THAR02_08479	A0A0F9X463	Uncharacterized protein	0	CBM1	-	28.08	15.80
THAR02_09247	A0A0F9ZZN6	Uncharacterized protein	0	CBM43	-	142.27	133.87
THAR02_09257	E2PTX8	Endochitinase 42 (Fragment)	6.00E-36	GH18	-	8.28	8.50

| THAR02
_09719 | A0A0F9WYH5 | Cellulase | 0 | GH5
CBM1 | 3.2.1.4 | 18.31 | 12.14 |

a

Principal Component Analysis

b

c

TH0179_CEL

TH3844_CEL

TA0020_CEL

Molecular Function

- binding
- catalytic activity
- hydrolase activity
- nucleotide binding
- transcription factor activity

Biological Process

- cellular process
- cellular response to stimulus
- metabolic process
- regulation of catalytic activity
- single-organism process
- transmembrane transport

Cellular Component

- cell
- cell part
- extracellular region
- membrane
- membrane part
- organelle
- organelle part

a

b

c

CBM1

GH6;CBM1

GH7;CBM1

GH1

CBM1

GH10

GH45;CBM1

AA9;CBM1

GH11

GH11;CBM1

GH1

AA9

d

CBM1

CBM1

GH45;CBM1

GH7;CBM1

GH6;CBM1

GH11

GH11;CBM1

GH10

GH1

AA9;CBM1

GH1

AA9

| ■ DEGs in Cel | ■ DEGs in Glu | ⋎ CAZymes Cel | ▲ CAZymes Glu | ⬢ Secreted proteins |