Cardiac Microlesions Form During Severe Bacteremic *Enterococcus faecalis* Infection

Armand O. Brown[1], Kavindra V. Singh[2], Melissa R. Cruz[1], Karan Gautam Kaval[1], Liezl E. Francisco[3], Barbara E. Murray[1,2], and Danielle A. Garsin[1*]

[1]Department of Microbiology and Molecular Genetics, The University of Texas Health Science Center at Houston, Houston TX, USA

[2] Division of Infectious Diseases, Department of Internal Medicine, The University of Texas Health Science Center at Houston, Houston, TX, USA

[3] Department of Biochemistry and Structural Biology, The University of Texas Health Science Center at San Antonio, San Antonio, TX, USA

*Corresponding author
Danielle.A.Garsin@uth.tmc.edu
713-500-5454

Running Title: Cardiac microlesions formed by *E. faecalis*

Abstract

Enterococcus faecalis is a significant cause of hospital-acquired bacteremia. Herein, the discovery is reported that cardiac microlesions form during severe bacteremic *E. faecalis* infection in mice. The cardiac microlesions were identical in appearance to those formed by *Streptococcus pneumoniae* during invasive pneumococcal disease (IPD). However, *E. faecalis* does not encode the virulence determinants implicated in pneumococcal microlesion formation. Rather, disulfide bond forming protein DsbA was found to be required for *E. faecalis* virulence in a *C. elegans* model and was necessary for efficient cardiac microlesion formation. Furthermore, *E. faecalis* promoted cardiomyocyte necroptotic cell death at sites of microlesion formation. Additionally, loss of DsbA caused an increase in pro-inflammatory cytokines unlike the wild type strain, which suppressed the immune response. In conclusion, we establish that *E. faecalis* is capable of forming cardiac microlesions and identify features of both the bacterium and the host response that are mechanistically involved.

KEYWORDS: Enterococcus faecalis, cardiac, microlesions, sepsis, bacteremia, infection, virulence, thioredoxin, disulfide bond, necroptosis

ABSTRACT WORD COUNT: 147

TEXT WORD COUNT: 3,478

Introduction

Enterococcus faecalis is a Gram-positive bacterium that normally co-exists with the host as a harmless commensal. However, it is capable of causing a variety of infections such as infective endocarditis (IE), urinary tract infection (UTI), bacteremia, peritonitis, prosthetic joint infection, and endophthalmitis. A common complication associated with enterococcal bacteremia is endocarditis, an infection that has a 1-year mortality rate of approximately 29% despite antibiotic therapy [6]. Enterococci are the third most common cause of endocarditis in North America [8] and, in patients with this infection, heart failure has been observed to occur with increased frequency [9-15]. Infective endocarditis involves the formation of biofilms on heart valves and inner chambers that develop into complex masses called vegetations [16]. Thus, heart infections associated with *E. faecalis* have traditionally been characterized as occurring on the surfaces of the heart, rather than in the deeper myocardial tissue, which would require active invasion.

Interestingly, another important Gram-positive pathogen, *Streptococcus pneumoniae,* was recently shown to invade the heart tissue and form cardiac microlesions during invasive pneumococcal disease (IPD) [17]. Importantly, pneumococcal microlesions were shown to be associated with altered cardiac electrophysiology, heart failure and scar formation during convalescence, which is thought to predispose individuals to major adverse cardiac events (MACE) [17, 18]. Furthermore, cardiac microlesions form following vascular endothelial cell invasion into the myocardial tissue through the interactions of pneumococcal Choline binding protein A (CbpA) and pneumococcal teichoic acid associated phosphorylcholine (ChoP) with host laminin receptor (LR) and platelet activating factor receptor (PAFr), respectively. Cardiomyocyte death is mediated by exposure to the pneumococcal secreted cholesterol-dependent pore forming toxin, pneumolysin [17, 18], and hydrogen peroxide [19, 20]. Since the original observation of

pneumococcal microlesions in 2014, similar observations were seen during blood stream

infections with *Francisella tularensis* [21], and *Mycobacterium avium* [22] , however the

mechanisms involved were not characterized.

In this report, we show that cardiac microlesions also form during invasive *E. faecalis* infection.

The observed cardiac microlesions were identical in appearance and size to *S. pneumoniae*

mediated cardiac microlesions, were diffuse across the myocardium, and characterized by an

absence of immune cell infiltration. Furthermore, an enterococcal determinant required for

microlesion formation, DsbA, was identified. Finally, *E. faecalis* infection of cardiomyocytes was

discovered to cause necroptosis and suppress the immune response.

Materials and Methods

C. elegans **survival assay**

E. faecalis strains were tested in *C. elegans* for survival as previously described [44]. The data is representative of 3 independent experiments. Nematode survival was plotted with the log-rank (Mantel-Cox) method using Prism 7 statistical software (GraphPad).

Bacterial strains and growth analysis

All *E. faecalis* strains used in this project were used in previously published studies [23] and are listed in Table S1. For growth analysis, overnight cultures were diluted to an OD_{600nm} of 0.05 in brain heart infusion (BHI) agar medium (Difco) in a volume of 200µl / well in a 96-well plate. The 96-well plate containing the diluted cultures were placed in a prewarmed 37ºC BioTek Cytation 5 plate reader, and OD_{600nm} measurements were taken every 10 minutes for 15 hours.

Mouse peritonitis and cardiac microlesion model

The mouse peritonitis model was performed as described previously [45]. The method for identifying and evaluating cardiac microlesion formation was conducted as described previously [17]. Briefly, female, ~4 – 6-week old (~25g), ICR outbred mice were obtained from Envigo (Indianapolis, Indiana) and infected intraperitoneally with 7×10^8 CFUs of *E. faecalis*. Mouse hearts were obtained 24-48 hours post-infection, formalin fixed and sent to The University of Texas Health Science Center Histopathology Laboratory for paraffin embedding, sectioning, and hematoxylin and eosin H&E) staining. Paraffin embedded mouse hearts were first bisected and sectioned longitudinally. 20 sections were cut per block, and every 5th section H&E stained in order to measure the number and relative distribution of the microlesions. Cardiac microlesions were averaged across sections, and the differences between mouse groups compared using

Student's two-tailed t-test. Experiments were performed in accordance with protocols approved

by the Animal Welfare Committee of the University of Texas Health Science Center at Houston

AWC-19-0058.

Necroptosis inhibition assay

HL-1 cardiomyocytes were seeded in 24-well plates at a concentration of 5×10^5 cells/ml. 24 hours

later, the cells were washed three times with Dulbecco's (1X) PBS, followed by pretreatment with

antibiotic free Claycomb medium supplemented with 10 μM of Necrostatin-1 (Sigma Aldrich),

GSK'872 (RD Systems), and Necrosulfonamaide (Millipore-Sigma) for 1 hour, followed by

infection at an MOI of 10 for 4 hours, following which the supernatant was collected, and

cytotoxicity in each well accessed using the CytoTox-ONE™ Homogeneous Membrane Integrity

Assay Kit (Promega). Each experiment was conducted in triplicate, on 3 separate occasions, and

averaged values between groups compared using *Student's* two-tailed t-test.

Annexin V apoptosis and necrosis assay.

The annexin V apoptosis and necrosis assay was conducted using RealTime-Glo™ Annexin V

Apoptosis and Necrosis Assay Kit (Promega). The kit contains near-equimolar ratios of two

annexin V fusion proteins (Annexin V-LgBiT and Annexin V-SmBit) containing complementary

subunits of NanoBiT® Luciferase that, when combined following exposure to phosphatidyl serine,

results in an emission of a luminescent signal. The kit also contains a cell-impermeant

profluorescent DNA dye, which detects necrosis. Briefly, HL-1 cardiomyocytes (Sigma-Aldrich)

were seeded in a 96 well plate. 24 hours later, the cells were infected with *E. faecalis* at an MOI

= 50, in the presence of kit reagents. Luminescent and fluorescent readings were obtained every

30 minutes for up to 6 hours.

Invasion assay

SVEC4-10 Endothelial Cells (ATCC) were seeded in 24 well plates at a cell density of 1.5×10^5 cells/ml. After 24 hours, the cells were washed three times with PBS and infected with *E. faecalis* at a MOI of 10 for 4 hours in antibiotic free Claycomb medium. After the incubation, the cells were washed with Dulbecco's (1X) PBS three times and treated with trypsin, diluted in PBS, and plated on BHI agar plates for attachment quantification. To measure intracellular numbers of *E. faecalis*, cells that were washed three times with Dulbecco's (1X) PBS 4 h.p.i., then treated with Dulbecco's Modified Eagle's Medium (ATCC) supplemented with 15 µg/ml vancomycin and 150 µg/ml gentamicin. 24 hours later, the wells were washed three times with Dulbecco's (1X) PBS, lysed using 0.1% Triton-X-100, and diluted and plated on BHI agar plates for intracellular quantification. The ratio of invasion to attachment for a given enterococcal strain was used to calculate the percent of invasion. The experiments were conducted on 3 independent occasions and were performed using triplicate wells for each strain. Attachment and invasion values were averaged from each experiment, and statistically analyzed using *Student's* two-tailed t-test.

Cytotoxicity and inflammatory cytokine panel microtiter plate reader assay

HL-1 cardiomyocytes were obtained from Sigma-Aldrich and SVEC4-10 Endothelial Cells were obtained from ATCC. HL-1 cardiomyocytes (Sigma-Aldrich) were seeded into 24-well plates at a concentration of 5×10^5 cells/ml. 24 hours later, the cells were washed three times with Dulbecco's (1X) PBS, and infected with *E. faecalis* at an MOI of 10 for 4 hours in antibiotic free Claycomb medium. Supernatant was collected, and cytotoxicity assayed using CytoTox-ONE™ Homogeneous Membrane Integrity Assay Kit (Promega). In addition to measuring cytotoxicity in HL-1 cardiomyocytes infected with *E. faecalis* 4 h.p.i, inflammatory chemokines and cytokines were analyzed using a Mouse Cytokine ELISA Plate Array III Colorimetric Assay (Signosis). The

data represent an average of at least 3 independent experiments for each isogenic mutant tested

and were analyzed using *Student's* two-tailed t-test.

Immunofluorescence microscopy

Immunofluorescence microcopy was conducted as described previously [17]. Briefly, paraffin

embedded sections were treated with 1:100 rabbit Anti-mouse MLKL (phospho S345) antibody

[EPR9515(2)] in order to detect phosphorylated MLKL (Abcam), and 1:100 rat anti-mouse MLKL

antibody (Millipore-Sigma) for MLKL detection followed by treatment with 1:1000 chicken anti-

rabbit IgG (H+L) cross-adsorbed secondary Alexa Fluor 594 antibody (ThermoFisher), and 1:1000

goat anti-rat IgG (H+L) secondary FITC antibody (ThermoFisher) respectively. A rabbit anti-

Enterococcus faecalis antibody (Abcam) was used at a concentration of 1:1000 to detect *E.*

faecalis within cardiac microlesions followed by treatment with 1:1000 goat anti-rabbit IgG (H&L),

FITC conjugated antibody (Invitrogen). All stained sections were counterstained with 1:1000 DAPI

(ThermoFisher) and mounted with Hardset™ Antifade Mounting Medium. Visualization of

fluorescence was obtained using standard excitation and emission wavelengths for Alexa 594,

FITC, and DAPI on an Olympus FV3000RS high-speed, high-sensitivity, inverted laser scanning

confocal microscope.

Generation of rabbit anti-DsbA antisera

His-tagged recombinant DsbA was overexpressed in Rosetta-2(DE3) *E. coli* (Sigma-Aldrich) and

isolated by affinity purification via HisPur Nickel-NTA Resin (ThermoFisher) packed columns

followed by further purification and buffer exchange using Fast Protein Liquid Chromatography

(FPLC) to obtain highly pure samples of recombinant DsbA. Cocalico Biologicals (Reamstown,

PA) generated anti-DsbA antisera by immunizing rabbits with the recombinant protein.

E. faecalis whole cell lysate and Western blot

An overnight culture of *E. faecalis* grown in BHI medium was centrifuged, washed with BHI and then diluted 1/100 in 5 ml of BHI. The culture was subsequently incubated at 37ºC for an additional 3.5 hours to obtain a culture of *E. faecalis* in exponential growth. The culture was spun down and the supernatant removed. Following sonication of the pellet, the sample was centrifuged at 10,000 G for 5 minutes and supernatant collected for electrophoresis. Following electrophoresis and transfer of protein to a PVDF membrane, the membrane was incubated overnight at 4ºC in a 5% milk solution containing a 1:1000 dilution of rabbit anti-DsbA antisera followed by incubation with a Goat anti-rabbit IgG HRP (ThermoFisher) secondary antibody at a dilution of 1:10,000 at room temperature for 30 minutes prior to development with a chemiluminescent kit (ThermoFisher).

Results

Cardiac microlesions form during severe bacteremic *E. faecalis* infection. In previous work,

we implicated a previously uncharacterized *E. faecalis* thioredoxin known as DsbA (disulfide bond

forming protein A) as a requirement for the stability of EntV, a secreted bacteriocin that we

determined inhibits hyphal morphogenesis of the human fungal pathogen *Candida albicans* [23,

24]. To test if these proteins had any effects on the fitness and pathogenicity of *E. faecalis*, we

infected *Caenorhabditis elegans* with *ΔdsbA* and *ΔentV E. faecalis* mutants and measured

survival. As shown in Figure 1A, loss of *dsbA* significantly attenuated mortality in infected

nematodes. Note that the *ΔdsbA* mutant did not manifest different growth kinetics when grown in

culture medium compared to wild type, ruling out a general growth defect (Figure S1). Infection

with the complement, *ΔdsbA::dsbA*, partially rescued the *ΔdsbA* phenotype. The partial

complementation was attributed to the DsbA protein being produced at lower levels in the

complemented strain (Figure 1B). No significant difference in nematode survival was observed

following infection with either the *ΔentV* mutant or its complement (Fig. 1A). Considering that EntV

is the only known substrate of *E. faecalis* DsbA, the results suggest that DsbA has additional

targets that affect pathogenicity.

Based on these results, we wanted to know if loss of DsbA also affected virulence in a vertebrate

animal model of *E. faecalis* infection. Therefore, the mutants were tested in a mouse model in

which animals were injected intra-peritoneally with *E. faecalis* resulting in generalized bacteremia

leading to mortality. Significant survival differences between the mice infected with the different

strains were not observed. However, various organs were collected in order to determine if *E.

faecalis* infection induced any strain-specific, pathological effects. To our surprise, cardiac

microlesions were discovered within the myocardium that were devoid of immune cell infiltrate

(Fig. 2 A, B). The microlesions appeared to form within roughly the same time frame (~24 h.p.i.)

as cardiac microlesions that have previously been reported to occur during invasive infection with

S. pneumoniae, the only other pathogen known to cause cardiac microlesions to date [17].

To verify that the observed microlesions were the result of infection by *E. faecalis*,

immunofluorescent staining on paraffin embedded cardiac tissue sections from mice infected with

E. faecalis with a polyclonal anti-enterococcal IgG antibody was performed and revealed the

presence of dense aggregates of *E. faecalis* within the microlesions (Fig. 2C). The presence of

bacterial DNA was further verified by DNA counterstaining with DAPI that revealed an

accumulation of non-cardiomyocyte nuclear DNA within microlesions (Fig. 2D). To determine the

contribution of DsbA toward microlesion formation, cardiac microlesions found on hematoxylin

and eosin stained paraffin embedded cardiac sections were quantified following infection with

ΔdsbA. A significant attenuation in microlesion formation during infection with *ΔdsbA* was

observed relative to wild type, while no significant difference was observed in mice infected with

the *dsbA* complement (Fig. 2E). Furthermore, when the sizes of the cardiac microlesions were

compared, the wild-type lesions were significantly larger than those found in the *ΔdsbA* infected

mice (Fig. 2F). Importantly, the size of the microlesions appeared to correlate with level of *dsbA*

expression observed in Western blots of *E. faecalis* whole cell lysates (Fig. 1B), with the

ΔdsbA::dsbA strain manifesting an in-between phenotype to that of the *ΔdsbA* and wild-type

strains, consistent with partial complementation. In conclusion, these data demonstrate that

cardiac microlesion formation occurs during systematic infection with *E. faecalis* and is dependent

on DsbA.

Necroptosis contributes to cardiomyocyte cell death within microlesions. In previous

studies of pneumococcal cardiac microlesion formation, apoptosis was found to be associated

with microlesion formation in mice (22). A subsequent study determined that cardiomyocyte cell

death occurs following exposure to pneumolysin (*ply*) and pneumococcal derived hydrogen

peroxide (*spxB*) [20]. Based on these observations and the fact that *E. faecalis* lacks *ply* and *spxB*

homologs, we sought to determine whether *E. faecalis* could promote cell death of

cardiomyocytes and identify the mechanism. First, HL-1 cardiomyocytes were infected with *E.*

faecalis and subsequently lactate dehydrogenase (LDH) release into the cell culture supernatant

was measured to assess cell death. From these experiments, it was concluded that *E. faecalis*

induces cell death of cardiomyocytes. Furthermore, infection with a *ΔdsbA* mutant resulted in a

significant decrease in cardiomyocyte cell death (Fig. 3B), suggesting DsbA is required to

modulate this process. Since cardiomyocyte cell death was observed following exposure to *E.*

faecalis, we wanted to identify the mechanism. To test this question, an annexin V apoptosis and

necrosis assay was performed to determine if *E. faecalis* promotes apoptosis or necrosis during

cardiomyocyte infection. The signals for both apoptosis and necrosis increased concomitantly

with time (Figure S2), suggesting that cell death is occurring by an apoptosis-independent

mechanism (see Materials and Methods). One apoptosis-independent cell death pathway is

necroptosis, and this mechanism of cell death is consistent with data published by investigators

studying *S. pnuemoniae* microlesions, which showed that markers of necroptosis could be

detected within the myocardium of acutely ill mice [17] and non-human primates [18]. Importantly,

necroptosis is considered a key cell death pathway in cardiomyocytes during ischemia-

reperfusion injury and acute coronary syndrome [25-27]. To test the hypothesis that

cardiomyocytes undergo necroptosis following exposure to *E. faecalis*, we infected HL-1

cardiomyocytes with this bacterium in the presence of various inhibitors of necroptosis and

observed a significant suppression of cell death (Fig. 3C). Additionally, staining for markers of

necroptosis [28], such as mixed lineage kinase domain-like pseudokinase (MLKL) and

phosphorylated MLKL, on cardiac tissue sections obtained from infected mice showed

concentrated signals within the microlesions (Fig 3D). Despite the ability of *E. faecalis* to promote death of the cardiomyocytes following exposure, antibiotic protection assays indicated that the bacteria were unable to invade cardiomyocytes relative to endothelial cells (Fig. 3A). No difference was observed in either the ability to promote necroptosis or cell invasion following exposure to *ΔdsbA* relative to wild-type (Figure S3). Taken together, these results suggest that during bacteremia, *E. faecalis* is able to translocate across the vascular endothelium in order to enter the myocardial tissue. Once there, *E. faecalis* is unable to invade cardiomyocytes, but can form microcolonies that promote necroptotic cell death.

Cardiomyocyte inflammatory cytokine response is suppressed following exposure to *E. faecalis*. Cardiac microlesions are characterized by bacteria filled vacuoles that are largely devoid of immune cell infiltrate [17, 29]. Previous efforts to explain the lack of an immune response to pneumococcal microlesions have yielded interesting insights. For example, *S. pneumoniae* within microlesions are thought to subvert the host response through a biofilm-mediated mechanism involving the pneumolysin mediated killing of resident macrophages, thus creating a more permissive environment for pneumococcal growth and survival [19]. There are also examples of *E. faecalis* subverting the immune response. In previous work, it was shown that *E. faecalis* can survive within mouse peritoneal macrophages for extended periods of time enabling the bacteria to replicate and spread [30]. In more recent work, *E. faecalis* was shown to activate Phosphatidylinositol 3-Kinase signaling that inhibits apoptosis [31]. *E. faecalis* can also resist phagosome acidification and autophagy within macrophages [32]. Because *E. faecalis* does not encode a pneumolysin homolog, nor does it secrete toxins known to kill macrophages or other immune cells, we wanted to examine the immune response to *E. faecalis* within the heart by measuring various inflammatory cytokines and chemokines. Experiments assessing cardiomyocyte inflammatory chemokine and cytokine levels showed that wild-type *E. faecalis* promotes an immune quiescent response, while loss of *dsbA* leads to a more inflammatory

immune response (Fig. 4A). Interestingly, EGF, M1P-1a, GM-CSF, IL-6, G-CSF, and INFr were

significantly upregulated during infection with *ΔdsbA* relative to the wild-type infected

cardiomyocytes. Thus, these results implicate the contribution of specific host signaling pathways

involved in pathogen clearance that are normally suppressed in the presence of DsbA during

infection, as modeled in Figure 4B. Collectively, these data suggest that *E. faecalis* inhibits a

vigorous cardiomyocyte immune response in a manner similar to that of *S. pneumoniae* despite

lacking the components that *S. pneumoniae* uses during immune subversion. Future studies will

be focused on understanding the cardiomyocyte immune response during enterococcal infection,

and how DsbA contributes.

Discussion

In this work, we demonstrated for the first time that cardiac microlesions form during severe *E. faecalis* bacteremic infection of mice. Importantly, microlesions occurred despite *E. faecalis* not encoding homologs of the pneumococcal virulence determinants implicated in microlesion formation. These data suggest that *E. faecalis* may be expressing functionally redundant proteins that serve to promote similar interactions to *S. pneumoniae* in mediating microlesion formation, such as CbpA and pneumolysin, which facilitate vascular endothelial cell translocation and cardiomyocyte cell death, respectively. Indeed, while *E. faecalis* does not encode a CbpA homolog, it does encode other proteins capable of binding host extracellular matrix proteins. These include EfbA, which binds fibronectin [33], ACE, which binds laminin receptor [34], and Ebp pili that bind fibrinogen [35].

Interestingly, we did discover one factor, DsbA, that contributes to microlesion formation and inhibition of the cardiomyocyte inflammatory immune response, as modeled in Figure 4B. Dsb proteins work to catalyze disulfide bond formation in proteins and have been best studied in Gram-negative bacteria where they catalyze the oxidative folding of proteins in the periplasm [36, 37]. While Dsb proteins and oxidative protein folding have established importance in the biology of certain Gram-positive bacteria, like the Actinomyces, Dsb proteins are not well studied in Firmicutes like *E. faecalis* and have generally not been considered important [36]. Most of these species secrete few proteins containing cysteines and some, notably *S. pneumoniae*, do not encode any thiol-disulfide oxidoreductases [38, 39]. A deletion of the only DsbA-like protein in *S. aureus* resulted in no significant phenotypes including no effects on pathogenicity [40]. Thus, it will be of interest to further investigate the role of Dsb proteins in these species.

Our prior work established EntV as a DsbA substrate, currently the only one identified [23].

However, loss of EntV did not reduce pathogenicity in the worm model (Figure 1A), and exposure

to EntV, even at very high concentrations, was non-toxic to mammalian cells [24]. Thus, we do

not favor EntV as being the source of DsbA's effects. A primary future direction will be to identify

additional substrates of DsbA and to determining which contribute to microlesion formation.

Pneumococcal ChoP interactions with host PAFr were determined to be a prerequisite for

vascular endothelial cell invasion during IPD. In light of our observations of *E. faecalis* invasion of

vascular endothelial cells despite an absence of this wall teichoic acid structure, it is possible that

another host receptor is required for endothelial cell invasion. Interestingly, recent studies focused

on bacterial cell wall entry and signaling in eukaryotic cells have identified a PAFr-independent

mechanism of tissue invasion that is actin-dependent and engenders Rac1, Cdc42, and

phosphatidylinositol 3-kinase (PI3K) signaling leading to cell wall internalization via a

macropinoctytosis-like pathway [41]. Whether such a pathway contributes to microlesion

formation is unknown. In the future, it will be of interest to study the mechanisms of tissue invasion

and the host signaling pathways involved.

Cardiac microlesion formation is becoming increasingly appreciated as a risk factor during acute

infection [42], and this study establishes that *E. faecalis* is another etiological agent capable of

causing this pathology in a vertebrate animal model. Considering that nosocomial derived

enterococcal infections are increasingly problematic among the elderly and immunocompromised

patients [4-7, 43], the development of successful treatments will require identifying all disease

manifestations as well as the bacterial and the host factors involved.

369
370
371

Table S1. Strains used in this study

Enterococcus faecalis Strains				
Strain	**Genotype**	**Characteristics**	**Parent**	**Source**
OG1RF	Wild-type	Gel$^+$ Spr$^+$ Rf^R Fa^R	OG1	[46]
AOBD1	*ΔdsbA*	Gel$^+$ Spr$^+$ Rf^R Fa^R	OG1	[23]
AOBD2	*ΔdsbA::dsbA*	Gel$^+$ Spr$^+$ Rf^R Fa^R	OG1	[23]

Figure Legends

Figure 1. *dsbA* deficient *E. faecalis* show reduced virulence in a *C. elegans* survival assay. **(A)** L4 stage *C. elegans* were infected with *E. faecalis*. This figure is representative of 3 independent experiments with 30 worms/group. ****Asterisk indicate a $P < 0.0001$ and were analyzed by the log-rank (Mantel-Cox). **(B)** Western blot performed by SDS-PAGE separation of *E. faecalis* whole cell lysates followed by hybridization of DsbA using rabbit anti-DsbA antisera. Image is representative of 3 independent Western blot experiments.

Figure 2. Cardiac microlesions form during invasive *E. faecalis* infection and are attenuated by loss of *dsbA*. Cardiac microlesions were observed using light microscopy of hematoxylin and eosin (H&E) stained paraffin embedded cardiac sections at 40X magnification **(A)**, and 100X magnification **(B)**. Immunofluorescent detection of *E. faecalis* was imaged using a 20X objective at 3X zoom (total of 60X magnification), a FITC conjugated anti-enterococcal antibody **(C)**, and DAPI for DNA staining **(D)**. Total microlesion counts **(E)**, and size measurements **(F)**, were obtained by quantification using light microscopy of H&E stained paraffin embedded cardiac sections from mice infected for 24-48 hours with *E. faecalis* (OG1RF n=14, Δ*dsbA* n=14, Δ*dsbA::dsbA* n=14). The data were averaged from 3 independent experiments, and analyzed using *Student's* two-tailed t-test (*** indicate a $P < 0.0001$).

Figure 3. Necroptosis contributes to cardiotoxicity within cardiac microlesions. **(A)** Comparison of OG1RF invasion rates into HL-1 cardiomyocytes and SVEC endothelial cells. The graph represents data from 3 independent experiments that were analyzed using *Student's* two-

tailed t-test (** indicate a *P* < 0.01). **(B)** *E. faecalis* induced cell death in HL-1 cardiomyocytes 4

hours post-infection (h.p.i.) as determined by measuring lactate dehydrogenase. The data

represent an average of 3 independent experiments and were analyzed using *Student's* two-tailed

t-test (* indicate a *P* < 0.05). **(C)** *E. faecalis* induced cell death in HL-1 cardiomyocytes 4 h.p.i. and

was inhibited in the presence of inhibitors of necroptosis. The data represent an average of 3

independent experiments and were analyzed using *Student's* two-tailed t-test (**** indicate a *P* <

0.0001). **(D)** Immunofluorescent microscopy of paraffin embedded cardiac cross sections at 20X

were performed to detect MLKL using FITC conjugated anti-MLKL antibodies, and Alexa 594

conjugated anti-phosphorylated MLKL antibodies at sites of microlesion formation, while DAPI

was used to stain nuclei.

Figure 4. Wild-type *E. faecalis* suppresses the cardiomyocyte inflammatory response *in-*

***vitro*. (A)** HL-1 cardiomyocytes exposed to *E. faecalis* promote a suppressed inflammatory

response while Δ*dsbA* infected cardiomyocytes yield a more inflammatory response. Data was

obtained from the fold-expression change analysis (relative to uninfected controls) of at least 3

independent experiments for each strain. **(B)** Model of *E. faecalis* mediated suppression of the

host cardiomyocyte immune response and effect on microlesion size.

Supplemental Figure 1. *E. faecalis* growth curve. *E. faecalis* strains were grown for 15 hours

in BHI and absorbance (OD_{600nm}) measured every 10 minutes. This figure is representative of 3

independent experiments.

Supplemental Figure 2. Cardiomyocytes infected with *E. faecalis* die through an apoptosis

independent mechanism. HL-1 cardiomyocytes were infected with *E. faecalis* at an MOI=50 for

4 hours in the presence of annexin V fusion proteins containing complimentary subunits of

NanoBiT® Luciferase combined with a cell-impermeant profluorescent DNA dye to detect

necrosis. A concomitant increase in both luminescent (apoptosis) **(A)** and fluorescent (necrosis)

(B) signals are indicative of cell death through an apoptosis independent mechanism. The graphs

represent an average of 3 independent experiments.

Supplemental Figure 3. Loss of *dsbA* does has no effect on endothelial cell invasion or

necroptosis induced cell death of cardiomyocytes. (A) Comparison of WT and *ΔdsbA* invasion

rates into HL-1 cardiomyocytes and SVEC endothelial cells showed no significant difference. The

graph represents data from 3 independent experiments analyzed using *Student's* two-tailed t-test

(NS denotes no significance). **(B)** Comparison of WT and *ΔdsbA* in HL-1 cardiomyocytes 4 h.p.i.

in the presence of inhibitors of necroptosis *in-vitro* was determined by measuring lactate

dehydrogenase. No significant difference was observed between necroptosis inhibitor conditions

tested. The data represent an average of 3 independent experiments and were analyzed using

Student's two-tailed t-test (* indicate a $P < 0.05$).

References

1. Lebreton F, Willems RJL, Gilmore MS. Enterococcus Diversity, Origins in Nature, and Gut Colonization. In: Gilmore MS, Clewell DB, Ike Y, Shankar N, eds. Enterococci: From Commensals to Leading Causes of Drug Resistant Infection. Boston, **2014**.

2. Armstrong-Evans M, Litt M, McArthur MA, et al. Control of transmission of vancomycin-resistant Enterococcus faecium in a long-term-care facility. Infect Control Hosp Epidemiol **1999**; 20:312-7.

3. Goh HMS, Yong MHA, Chong KKL, Kline KA. Model systems for the study of Enterococcal colonization and infection. Virulence **2017**; 8:1525-62.

4. Malani PN KC, Zervis MJ. Enterococcal disease, epidemiology and treatment. Washington, D.C.: ASM Press, **2002** The Enterococci: Pathogenesis, Molecular Biology and Antibiotic Resistance).

5. Wisplinghoff H, Bischoff T, Tallent SM, Seifert H, Wenzel RP, Edmond MB. Nosocomial bloodstream infections in US hospitals: analysis of 24,179 cases from a prospective nationwide surveillance study. Clin Infect Dis **2004**; 39:309-17.

6. Chirouze C, Athan E, Alla F, et al. Enterococcal endocarditis in the beginning of the 21st century: analysis from the International Collaboration on Endocarditis-Prospective Cohort Study. Clin Microbiol Infect **2013**; 19:1140-7.

7. McDonald JR, Olaison L, Anderson DJ, et al. Enterococcal endocarditis: 107 cases from the international collaboration on endocarditis merged database. Am J Med **2005**; 118:759-66.

8. Agudelo Higuita NI, Huycke MM. Enterococcal Disease, Epidemiology, and Implications for Treatment. In: Gilmore MS, Clewell DB, Ike Y, Shankar N, eds. Enterococci: From Commensals to Leading Causes of Drug Resistant Infection. Boston, **2014**.

9. Cabell CH, Jollis JG, Peterson GE, et al. Changing patient characteristics and the effect on mortality in endocarditis. Arch Intern Med **2002**; 162:90-4.

10. Garvey GJ, Neu HC. Infective endocarditis--an evolving disease. A review of endocarditis at

the Columbia-Presbyterian Medical Center, 1968-1973. Medicine (Baltimore) **1978**; 57:105-27.

11. Geraci JE, Martin WJ. Antibiotic therapy of bacterial endocarditis. VI. Subacute enterococcal

endocarditis; clinical, pathologic and therapeutic consideration of 33 cases. Circulation **1954**;

10:173-94.

12. Jawetz E, Sonne M. Penicillin-streptomycin treatment of enterococcal endocarditis. A re-

evaluation. N Engl J Med **1966**; 274:710-5.

13. Lerner PI, Weinstein L. Infective endocarditis in the antibiotic era. N Engl J Med **1966**;

274:388-93 concl.

14. Olaison L, Schadewitz K, Swedish Society of Infectious Diseases Quality Assurance Study

Group for E. Enterococcal endocarditis in Sweden, 1995-1999: can shorter therapy with

aminoglycosides be used? Clin Infect Dis **2002**; 34:159-66.

15. Pelletier LL, Jr., Petersdorf RG. Infective endocarditis: a review of 125 cases from the

University of Washington Hospitals, 1963-72. Medicine (Baltimore) **1977**; 56:287-313.

16. Garsin DA, Frank KL, Silanpaa J, et al. Pathogenesis and Models of Enterococcal Infection.

In: Gilmore MS, Clewell DB, Ike Y, Shankar N, eds. Enterococci: From Commensals to Leading

Causes of Drug Resistant Infection. Boston, **2014**.

17. Brown AO, Mann B, Gao G, et al. Streptococcus pneumoniae translocates into the

myocardium and forms unique microlesions that disrupt cardiac function. PLoS Pathog **2014**;

10:e1004383.

18. Reyes LF, Restrepo MI, Hinojosa CA, et al. Severe Pneumococcal Pneumonia Causes Acute

Cardiac Toxicity and Subsequent Cardiac Remodeling. Am J Respir Crit Care Med **2017**;

196:609-20.

19. Shenoy AT, Brissac T, Gilley RP, et al. Streptococcus pneumoniae in the heart subvert the

host response through biofilm-mediated resident macrophage killing. PLoS Pathog **2017**;

13:e1006582.

20. Brissac T, Shenoy AT, Patterson LA, Orihuela CJ. Cell Invasion and Pyruvate Oxidase-

Derived H2O2 Are Critical for Streptococcus pneumoniae-Mediated Cardiomyocyte Killing. Infect

Immun **2018**; 86.

21. Makara MA, Hoang KV, Ganesan LP, et al. Cardiac Electrical and Structural Changes During

Bacterial Infection: An Instructive Model to Study Cardiac Dysfunction in Sepsis. J Am Heart

Assoc **2016**; 5.

22. Headley CA, Gerberick A, Mehta S, et al. Nontuberculous mycobacterium M. avium infection

predisposes aged mice to cardiac abnormalities and inflammation. Aging Cell **2019**; 18:e12926.

23. Armand O. Brown CEG, Melissa R. Cruz, Michael C. Lorenz, Danielle A. Garsin. Antifungal

activity of the Enterococcus faecalis peptide EntV requires protease cleavage and disulfide bond

formation. Mbio **2019**; 10:1-13.

24. Graham CE, Cruz MR, Garsin DA, Lorenz MC. Enterococcus faecalis bacteriocin EntV inhibits

hyphal morphogenesis, biofilm formation, and virulence of Candida albicans. Proc Natl Acad Sci

U S A **2017**; 114:4507-12.

25. Zhang T, Zhang Y, Cui M, et al. CaMKII is a RIP3 substrate mediating ischemia- and oxidative

stress-induced myocardial necroptosis. Nat Med **2016**; 22:175-82.

26. Linkermann A, Hackl MJ, Kunzendorf U, Walczak H, Krautwald S, Jevnikar AM. Necroptosis

in immunity and ischemia-reperfusion injury. Am J Transplant **2013**; 13:2797-804.

27. Linkermann A, Brasen JH, Darding M, et al. Two independent pathways of regulated necrosis

mediate ischemia-reperfusion injury. Proc Natl Acad Sci U S A **2013**; 110:12024-9.

28. Kitur K, Wachtel S, Brown A, et al. Necroptosis Promotes Staphylococcus aureus Clearance

by Inhibiting Excessive Inflammatory Signaling. Cell Rep **2016**; 16:2219-30.

29. Gilley RP, Gonzalez-Juarbe N, Shenoy AT, et al. Infiltrated Macrophages Die of Pneumolysin-

Mediated Necroptosis following Pneumococcal Myocardial Invasion. Infect Immun **2016**; 84:1457-

69.

30. Gentry-Weeks CR, Karkhoff-Schweizer R, Pikis A, Estay M, Keith JM. Survival of

Enterococcus faecalis in mouse peritoneal macrophages. Infect Immun **1999**; 67:2160-5.

31. Zou J, Shankar N. Enterococcus faecalis infection activates phosphatidylinositol 3-kinase

signaling to block apoptotic cell death in macrophages. Infect Immun **2014**; 82:5132-42.

32. Zou J, Shankar N. The opportunistic pathogen Enterococcus faecalis resists phagosome

acidification and autophagy to promote intracellular survival in macrophages. Cell Microbiol **2016**;

18:831-43.

33. Singh KV, La Rosa SL, Somarajan SR, Roh JH, Murray BE. The fibronectin-binding protein

EfbA contributes to pathogenesis and protects against infective endocarditis caused by

Enterococcus faecalis. Infect Immun **2015**; 83:4487-94.

34. Nallapareddy SR, Qin X, Weinstock GM, Hook M, Murray BE. Enterococcus faecalis adhesin,

ace, mediates attachment to extracellular matrix proteins collagen type IV and laminin as well as

collagen type I. Infect Immun **2000**; 68:5218-24.

35. .Nallapareddy SR, Singh KV, Sillanpaa J, et al. Endocarditis and biofilm-associated pili of

Enterococcus faecalis. J Clin Invest **2006**; 116:2799-807.

36. Reardon-Robinson ME, Ton-That H. Disulfide-Bond-Forming Pathways in Gram-Positive

Bacteria. J Bacteriol **2015**; 198:746-54.

37. Landeta C, Boyd D, Beckwith J. Disulfide bond formation in prokaryotes. Nat Microbiol **2018**;

3:270-80.

38. Daniels R, Mellroth P, Bernsel A, et al. Disulfide bond formation and cysteine exclusion in

gram-positive bacteria. J Biol Chem **2010**; 285:3300-9.

39. Dutton RJ, Boyd D, Berkmen M, Beckwith J. Bacterial species exhibit diversity in their

mechanisms and capacity for protein disulfide bond formation. Proc Natl Acad Sci U S A **2008**;

105:11933-8.

40. Dumoulin A, Grauschopf U, Bischoff M, Thony-Meyer L, Berger-Bachi B. Staphylococcus

aureus DsbA is a membrane-bound lipoprotein with thiol-disulfide oxidoreductase activity. Arch

Microbiol **2005**; 184:117-28.

41. Loh LN, Gao G, Tuomanen EI. Dissecting Bacterial Cell Wall Entry and Signaling in Eukaryotic

Cells: an Actin-Dependent Pathway Parallels Platelet-Activating Factor Receptor-Mediated

Endocytosis. MBio **2017**; 8.

42. Musher DM, Abers MS, Corrales-Medina VF. Acute Infection and Myocardial Infarction. N

Engl J Med **2019**; 380:171-6.

43. Weiner LM, Webb AK, Limbago B, et al. Antimicrobial-Resistant Pathogens Associated With

Healthcare-Associated Infections: Summary of Data Reported to the National Healthcare Safety

Network at the Centers for Disease Control and Prevention, 2011-2014. Infect Control Hosp

Epidemiol **2016**; 37:1288-301.

44. Garsin DA, Sifri CD, Mylonakis E, et al. A simple model host for identifying Gram-positive

virulence factors. Proc Natl Acad Sci U S A **2001**; 98:10892-7.

45. Singh KV, Qin X, Weinstock GM, Murray BE. Generation and testing of mutants of

Enterococcus faecalis in a mouse peritonitis model. J Infect Dis **1998**; 178:1416-20.

46. Murray BE, Singh KV, Ross RP, Heath JD, Dunny GM, Weinstock GM. Generation of

restriction map of Enterococcus faecalis OG1 and investigation of growth requirements and

regions encoding biosynthetic function. J Bacteriol **1993**; 175:5216-23.

FOOTNOTE PAGE

Conflict of Interest statement

There are no conflicts of interest.

Funding statement

This work was supported by the National Institute of Dental and Craniofacial Research of the

National Institutes of Health under [grant number R01DE027608 to D. A. G.] and [grant number

F32DE027580 to A. O. B.]

Meetings

This work has not been presented.

Corresponding author contact information

Danielle A. Garsin, Ph.D.

6431 Fannin St.

MSB 1.168

Houston, TX 77030

Telephone: (713) 500-5454

Fax: (713) 500-5499

Email: Danielle.A.Garsin@uth.tmc.edu

A.

	OG1RF	ΔdsbA
IL-12		
Rantes		
VEGF		
Leptin		
EGF		
FGFb		
MCP-1		
M1P-1a		
IGF-1		
IL-3		
Resistin		
IL-10		
b-NGF		
IL-1a		
PDGF-BB		
IL-2		
IL-13		
IL-4		
CXCL1		
GM-CSF		
IL-6		
G-CSF		
IFNr		
CCL21		
CCL11		
TNF-a		
IL-1b		
CXCL10		
SCF		
IL-17a		
IL-21		
IL-22		

B.

OG1RF   ΔdsbA

- Increased immune suppression
- Increased cell death
- Increased microlesion size

- Decreased immune suppression
- Decreased cell death
- Decreased microlesion size