Cardiac Microlesions Form During Severe Bacteremic *Enterococcus faecalis* Infection

Armand O. Brown[1], Kavindra V. Singh[2], Melissa R. Cruz[1], Karan Gautam Kaval[1], Liezl E. Francisco[3], Barbara E. Murray[1,2], and Danielle A. Garsin[1*]

[1]Department of Microbiology and Molecular Genetics, The University of Texas Health Science Center at Houston, Houston TX, USA

[2] Division of Infectious Diseases, Department of Internal Medicine, The University of Texas Health Science Center at Houston, Houston, TX, USA

[3] Department of Biochemistry and Structural Biology, The University of Texas Health Science Center at San Antonio, San Antonio, TX, USA

*Corresponding author
Danielle.A.Garsin@uth.tmc.edu
713-500-5454

Running Title: Cardiac microlesions formed by *E. faecalis*

Abstract

Enterococcus faecalis is a significant cause of hospital-acquired bacteremia. Herein, the discovery is reported that cardiac microlesions form during severe bacteremic *E. faecalis* infection in mice. The cardiac microlesions were identical in appearance to those formed by *Streptococcus pneumoniae* during invasive pneumococcal disease (IPD). However, *E. faecalis* does not encode the virulence determinants implicated in pneumococcal microlesion formation. Rather, disulfide bond forming protein DsbA was found to be required for *E. faecalis* virulence in a *C. elegans* model and was necessary for efficient cardiac microlesion formation. Furthermore, *E. faecalis* promoted cardiomyocyte apoptotic and necroptotic cell death at sites of microlesion formation. Additionally, loss of DsbA caused an increase in pro-inflammatory cytokines unlike the wild-type strain, which suppressed the immune response. In conclusion, we establish that *E. faecalis* is capable of forming cardiac microlesions and identify features of both the bacterium and the host response that are mechanistically involved.

SUMMARY: This work presents the observation of cardiac microlesion formation during severe blood stream infection with *Enterococcus faecalis* in mice. Moreover, we identify the contribution of a novel enterococcal virulence determinant in modulating microlesion formation and the host immune response.

KEYWORDS: Enterococcus faecalis, cardiac, microlesions, sepsis, bacteremia, infection, virulence, thioredoxin, disulfide bond, necroptosis

ABSTRACT WORD COUNT: 149

TEXT WORD COUNT: 3,496

Introduction

Enterococcus faecalis is a Gram-positive bacterium that normally co-exists with the host as a harmless commensal. However, it is capable of causing a variety of infections such as infective endocarditis (IE), urinary tract infection (UTI), bacteremia, peritonitis, prosthetic joint infection, and endophthalmitis. A common complication associated with enterococcal bacteremia is endocarditis, an infection that has a 1-year mortality rate of approximately 29% despite antibiotic therapy [1]. Enterococci are the third most common cause of endocarditis in North America [2] and, in patients with this infection, heart failure has been observed to occur with increased frequency [3-9]. Endocarditis only involves infection of the surfaces of the heart - the formation of biofilms on heart valves and inner chambers that develop into complex masses called vegetations [10]. *E. faecalis* invasion of the deeper myocardial tissue to cause cardiac microlesions has not previously been described.

Interestingly, another Gram-positive pathogen, *Streptococcus pneumoniae,* was recently shown to invade the heart tissue during invasive pneumococcal disease (IPD) [11]. Importantly, pneumococcal microlesions were determined to be associated with altered cardiac electrophysiology, heart failure and scar formation during convalescence, which is thought to predispose individuals to major adverse cardiac events (MACE) [11, 12]. Furthermore, cardiac microlesions form following vascular endothelial cell invasion into the myocardial tissue through the interactions of pneumococcal Choline binding protein A (CbpA) and pneumococcal teichoic acid associated phosphorylcholine (ChoP) with host laminin receptor (LR) and platelet activating factor receptor (PAFr), respectively. Cardiomyocyte death is mediated by exposure to the pneumococcal secreted cholesterol-dependent pore forming toxin, pneumolysin [11, 12], and hydrogen peroxide [13, 14]. Since the original observation of pneumococcal microlesions in 2014,

similar observations were seen during blood stream infections with *Francisella tularensis* [15], and

Mycobacterium avium [16], however the mechanisms involved were not characterized.

In this report, we show that cardiac microlesions also form during invasive *E. faecalis* infection.

The observed cardiac microlesions were identical in appearance and size to those formed by *S.*

pneumoniae, were diffuse across the myocardium, and characterized by an absence of immune

cell infiltration and cell death by apoptosis and necroptosis. Furthermore, an enterococcal

determinant required for microlesion formation, DsbA, was identified. DsbA was additionally

shown to enhance cell death and promote suppression of the immune response during *E. faecalis*

infection of a cardiomyocyte cell line.

Materials and Methods

C. elegans survival assay

E. faecalis strains were tested in *C. elegans* for survival as previously described [17]. The data is representative of 3 independent experiments. Nematode survival was plotted with the log-rank (Mantel-Cox) method using Prism 7 statistical software (GraphPad).

Bacterial strains and growth analysis

All *E. faecalis* strains used in this project are listed in Table S1. For growth analysis, overnight cultures were diluted to an OD_{600nm} of 0.05 in brain heart infusion (BHI) medium (Difco) or serum [18, 19] in a volume of 200µl / well in a 96-well plate. The 96-well plate containing the diluted cultures were placed in a prewarmed 37ºC BioTek Cytation 5 plate reader, and OD_{600nm} measurements were taken every 10 minutes for 16 hours.

Mouse peritonitis and cardiac microlesion model

The mouse peritonitis model was performed as described previously [20]. The method for identifying and evaluating cardiac microlesion formation was conducted as described previously [11]. Briefly, female, ~4 – 6-week old (~25g), ICR outbred mice were obtained from Envigo (Indianapolis, Indiana) and infected intraperitoneally with 3.8 - 4.2 x10^8 CFUs of *E. faecalis*. Mouse hearts were harvested 24-48 hours post-infection at the point when the animals showed signs of severe sepsis indicated by ruffled fur, labored breathing and little movement. The hearts were formalin fixed and sent to The University of Texas Health Science Center Histopathology Laboratory for paraffin embedding, sectioning, and hematoxylin and eosin H&E) staining. Paraffin embedded mouse hearts were first bisected and sectioned longitudinally. 20 sections were cut per block, and every 5th section H&E stained in order to measure the number and relative

distribution of the microlesions. Cardiac microlesions were averaged across sections, and the

differences between mouse groups compared using *Student's* two-tailed t-test. Experiments were

performed in accordance with protocols approved by the Animal Welfare Committee of the

University of Texas Health Science Center at Houston AWC-19-0058.

Invasion assay

SVEC4-10 Endothelial Cells (ATCC) were seeded in 24 well plates at a cell density of $1.5x10^5$

cells/ml. After 24 hours, the cells were washed three times with PBS and infected with *E. faecalis*

at a MOI of 10 for 4 hours in antibiotic free Dulbecco's Modified Eagle's Medium (ATCC). The

cells were then washed with Dulbecco's (1X) PBS three times, treated with trypsin, diluted in PBS,

and plated for quantification. To measure intracellular numbers of *E. faecalis*, cells that were

washed three times with Dulbecco's (1X) PBS 4 h.p.i., then treated with DMEM (ATCC)

supplemented with 15 µg/ml vancomycin and 150 µg/ml gentamicin. 24 hours later, the wells were

washed three times with Dulbecco's (1X) PBS, lysed using 0.1% Triton-X-100, and diluted and

plated for intracellular quantification. The ratio of invasion to attachment was used to calculate the

percent of invasion. The experiments were conducted on 3 independent occasions and were

performed using triplicate wells for each strain. Attachment and invasion values were averaged

from each experiment, and statistically analyzed using *Student's* two-tailed t-test.

Cytotoxicity and inflammatory cytokine panel microtiter plate reader assay

HL-1 cardiomyocytes were obtained from Sigma-Aldrich and SVEC4-10 Endothelial Cells were

obtained from ATCC. HL-1 cardiomyocytes (Sigma-Aldrich) were seeded into 24-well plates at a

concentration of $5x10^5$ cells/ml. 24 hours later, the cells were washed three times with Dulbecco's

(1X) PBS, and infected with *E. faecalis* at a MOI of 10 for 4 hours in antibiotic free Claycomb

medium. Supernatant was collected, and cytotoxicity assayed using CytoTox-ONE™

Homogeneous Membrane Integrity Assay Kit (Promega). For HL-1 infected cardiomyocytes,

inflammatory chemokines and cytokines were additionally analyzed using a Mouse Cytokine

ELISA Plate Array III Colorimetric Assay (Signosis). The data represent an average of at least 3

independent experiments for each strain and were analyzed using *Student's* two-tailed t-test.

Immunofluorescence microscopy

Immunofluorescence microscopy was conducted as described previously [11]. Briefly, paraffin

embedded sections were treated with 1:100 rabbit Anti-mouse MLKL (phospho S345) antibody

[EPR9515(2)] in order to detect phosphorylated MLKL (Abcam), and 1:100 rat anti-mouse MLKL

antibody (Millipore-Sigma) for MLKL detection followed by treatment with 1:1000 chicken anti-

rabbit IgG (H+L) cross-adsorbed secondary Alexa Fluor 594 antibody (ThermoFisher), and 1:1000

goat anti-rat IgG (H+L) secondary FITC antibody (ThermoFisher) respectively. A rabbit anti-

Enterococcus faecalis antibody (Abcam) was used at a concentration of 1:1000 to detect *E.*

faecalis followed by treatment with 1:1000 goat anti-rabbit IgG (H&L), FITC conjugated antibody

(Invitrogen). A rat anti-PARP (MAB600) antibody (R&DSystems) was used at a concentration of

1:100 followed by treatment with 1:1000 goat anti-rat IgG (H+L) cross-adsorbed secondary Alexa

Fluor 594 antibody (ThermoFisher) to detect PARP. All stained sections were counterstained with

1:1000 DAPI (ThermoFisher) and mounted with Hardset™ Antifade Mounting Medium.

Visualization of fluorescence was obtained using standard excitation and emission wavelengths

for Alexa 594, FITC, and DAPI on an Olympus FV3000RS high-speed, high-sensitivity, inverted

laser scanning confocal microscope.

Generation of rabbit anti-DsbA antisera

His-tagged recombinant DsbA was overexpressed in Rosetta-2(DE3) *E. coli* (Sigma-Aldrich) and

isolated by affinity purification via HisPur Nickel-NTA Resin (ThermoFisher) packed columns

followed by Fast Protein Liquid Chromatography (FPLC) to obtain highly pure samples of recombinant DsbA. Cocalico Biologicals (Reamstown, PA) generated anti-DsbA antisera by immunizing rabbits with the recombinant protein.

E. faecalis whole cell lysate and Western blot

A 5 ml culture of *E. faecalis* grown in BHI medium at 37ºC for 15-16 hours followed by a 1:10 dilution in 5 ml of fresh BHI. After 3.5 hours to achieve exponential growth, the culture was centrifuged at 6,010 x G and the supernatant removed. The pellet was resuspended in 1ml of buffer (10 mM Tris/HCl pH 7.5 and 200 mM NaCl) containing cOmplete™, EDTA-free Protease Inhibitor Cocktail (Roche). Following sonication of the pellet on ice, the sample was centrifuged at 10,000 x G for 5 minutes and supernatant collected for electrophoresis. Total protein in the whole cell lysate was measured using the *DC* Protein Assay (BioRad). Following the loading of 10µg of cell lysate, electrophoresis and transfer of protein to a PVDF membrane, the membrane was incubated overnight at 4ºC in a 5% milk solution containing a 1:1000 dilution of rabbit anti-DsbA antisera followed by incubation with a Goat anti-rabbit IgG HRP (ThermoFisher) secondary antibody at a dilution of 1:10,000 at room temperature for 30 minutes prior to development with a chemiluminescent kit (ThermoFisher).

Results

Cardiac microlesions form during severe bacteremic *E. faecalis* infection. In previous work, we implicated a previously uncharacterized *E. faecalis* thioredoxin known as DsbA (disulfide bond forming protein A) as a requirement for the stability of EntV, a secreted bacteriocin that inhibits hyphal morphogenesis of the human fungal pathogen *Candida albicans* [21, 22]. To test if these proteins had any effects on the fitness and pathogenicity of *E. faecalis*, we infected *Caenorhabditis elegans* with Δ*dsbA* and Δ*entV E. faecalis* mutants and measured survival. As shown in Figure 1A, loss of *dsbA* significantly attenuated mortality in infected nematodes. Note that the Δ*dsbA* mutant did not manifest different growth kinetics when grown in culture medium compared to wild-type, ruling out a general growth defect (Figure S1A). Infection with the complement, Δ*dsbA::dsbA*, partially rescued the Δ*dsbA* phenotype. The partial complementation was attributed to the DsbA protein being produced at lower levels in the complemented strain (Figure 1B and S2). No significant difference in nematode survival was observed following infection with either the Δ*entV* mutant or its complement (Figure 1A). Considering that EntV is the only known substrate of *E. faecalis* DsbA, the results suggest that DsbA has additional targets that affect pathogenicity.

Based on these results, we wanted to know if loss of DsbA also affected virulence in a vertebrate animal model of *E. faecalis* infection. Therefore, the mutants were tested in a mouse model in which animals were injected intra-peritoneally with *E. faecalis* resulting in generalized bacteremia leading to mortality. Significant survival differences between the mice infected with the different strains were subtle, but significant attenuation by the *dsbA* mutant was observed when data from three separate experiments were combined to generate an n close to 20 for each group (Figure S3). To further study the contribution of *dsbA*, various organs were collected in order to determine

if *E. faecalis* infection induced any strain-specific, pathological effects. To our surprise, cardiac microlesions were discovered within the myocardium that were devoid of immune cell infiltrate regardless of which *E. faecalis* strain was used (Figures 2A, B and S4). The microlesions appeared to form within roughly the same time frame (~24 h.p.i.) as cardiac microlesions that have previously been reported to occur during invasive infection with *S. pneumoniae* [11].

To verify that the observed microlesions were the result of infection by *E. faecalis*, immunofluorescent staining on paraffin embedded cardiac tissue sections from mice infected with *E. faecalis* with a polyclonal anti-enterococcal IgG antibody was performed and revealed the presence of dense aggregates of *E. faecalis* within the microlesions (Figure 2C). The presence of bacterial DNA was further verified by counterstaining with DAPI, revealing an accumulation of non-cardiomyocyte nuclear DNA within microlesions (Figure 2D). To determine the contribution of DsbA, cardiac microlesions found on hematoxylin and eosin stained paraffin embedded cardiac sections were quantified following infection with Δ*dsbA*. A significant attenuation in microlesion formation during infection with Δ*dsbA* was observed relative to wild-type, while no significant difference was observed in mice infected with the *dsbA* complement (Figure 2E). Furthermore, when the sizes of the cardiac microlesions were compared, the wild-type lesions were significantly larger than those found in the Δ*dsbA* infected mice (Figure 2F). Importantly, the size of the microlesions appeared to correlate with level of *dsbA* expression observed in Western blots of *E. faecalis* whole cell lysates (Figure 1B), with the Δ*dsbA*::*dsbA* strain manifesting an in-between phenotype to that of the Δ*dsbA* and wild-type strains, consistent with partial complementation. It is possible that less microlesion formation occurred with the Δ*dsbA* strain due poor growth in the blood, but we do not favor this model as in vitro growth curves in serum showed no defect (Figure S1B). In conclusion, these data demonstrate that cardiac microlesion formation occurs during systematic infection with *E. faecalis* and is dependent on DsbA.

Apoptosis and necroptosis contribute to cardiomyocyte cell death within microlesions. In previous studies of pneumococcal cardiac microlesion formation, apoptosis was found to be associated with microlesion formation in mice [11]. A subsequent study determined that cardiomyocyte cell death occurs following exposure to pneumolysin (*ply*) and pneumococcal pyruvate oxidase (*spxB*) derived hydrogen peroxide [14]. Based on these observations and the fact that *E. faecalis* lacks *ply* and *spxB* homologs, we sought to determine whether *E. faecalis* could promote cell death of cardiomyocytes and identify the mechanism. To determine if apoptosis contributes to *E. faecalis* microlesion formation, we performed immunofluorescent microscopy on heart sections of infected mice to detect PARP [Poly(ADP-ribose) Polymerase], an enzyme that is proteolytically cleaved by the effector Caspase-3, and is one hallmark event of apoptosis. In agreement with previous reports, markers of apoptosis were observed to be concentrated at sites of microlesion formation (Figure 3A and S5).

In addition to apoptosis, we hypothesized that necroptosis also contributes to *E. faecalis* mediated cardiac microlesion formation. Necroptosis is a programmed form of necrotic cell death that is generally inflammatory, and has been shown to contribute to the formation of *S. pneumoniae* microlesions within the myocardium of acutely ill mice and non-human primates [12]. Importantly, necroptosis is considered a key cell death pathway in cardiomyocytes during ischemia-reperfusion injury and acute coronary syndrome [23-25]. To test the hypothesis that necroptosis contributes to the formation of *E. faecalis* microlesions, we stained for markers of necroptosis [26], such as mixed lineage kinase domain-like pseudokinase (MLKL) and phosphorylated MLKL, on cardiac tissue sections obtained from infected mice. As a result, we detected concentrated signals within the microlesions (Fig 3B and Figure S6) that suggest that necroptosis does indeed contribute to *E. faecalis* cardiac microlesion formation.

To further look at the effects of *E. faecalis* and the *dsbA* mutant on heart cells, we infected HL-1

cardiomyocytes and subsequently measured lactate dehydrogenase (LDH) release into the cell

culture supernatant to assess cell death. As shown in Figure 3C, cell death of the cardiomyocytes

is significantly decreased following exposure to the *dsbA* mutant compared to wild-type

suggesting that DsbA contributes to cardiac cell death (Figure 3C). While *E. faecalis* promoted

death of the cardiomyocytes, antibiotic protection assays indicated that both the wild-type and

dsbA mutant bacteria were unable to invade cardiomyocytes relative to endothelial cells (Figure

3D). Taken together, these results suggest that during bacteremia, *E. faecalis* is able to

translocate across the vascular endothelium in order to enter the myocardial tissue. Once there,

E. faecalis is unable to invade cardiomyocytes, but can form microcolonies that promote

necroptosis in addition to apoptosis. Whether other cell death pathways additionally contribute

will be fodder for future investigations.

Cardiomyocyte inflammatory cytokine response is suppressed following exposure to *E.*

***faecalis*.** Cardiac microlesions are characterized by bacteria filled vacuoles that are largely devoid

of immune cell infiltrate [11, 27]. Previous efforts to explain the lack of an immune response to

pneumococcal microlesions have yielded interesting insights. For example, *S. pneumoniae* within

microlesions are thought to subvert the host response through a biofilm-mediated mechanism

involving the pneumolysin mediated killing of resident macrophages, thus creating a more

permissive environment for pneumococcal growth and survival [13]. There are also examples of

E. faecalis subverting the immune response. In previous work, it was shown that *E. faecalis* can

survive within mouse peritoneal macrophages for extended periods of time [28] and can also resist

phagosome acidification and autophagy [29]. Additionally, *E. faecalis* was shown to activate

Phosphatidylinositol 3-Kinase signaling that inhibits apoptosis [30]. Because *E. faecalis* does not

encode a pneumolysin homolog, nor does it secrete toxins known to kill macrophages or other

immune cells, we wanted to examine the immune response to *E. faecalis* within cardiomyocytes.

Therefore, following infection with *E. faecalis*, inflammatory chemokine and cytokine levels were

measured. Wild-type *E. faecalis* was discovered to promote an immune quiescent response, while

the *dsbA* mutant caused a more inflammatory immune response (Figure 4A and Table S2).

Specifically, infection of cardiomyocytes with wild-type increased the expression of two pro-

inflammatory cytokines IL-4 and b-NGF cytokines [31, 32]. Modest increases were also observed

in VEGF, an important determinant of sepsis morbidity and mortality [33], and GM-CSF that

serves as an immunomodulatory cytokine [34]. In contrast, cardiomyocytes infected with Δ*dsbA*

led to increases in Rantes, IFNr, MIP-1a, and IL-12, with modest increases also observed in Leptin

and G-CSF. With the exception of G-CSF, an immunomodulatory cytokine with roles in cardiac

cell generation and repair [35, 36], all of the other aforementioned cytokines are pro-inflammatory

[37-40]. These results implicate the contribution of specific host signaling pathways that are

normally suppressed in the presence of DsbA during infection, as modeled in Figure 4B.

Collectively, these data suggest that *E. faecalis* is capable of suppressing the host cardiomyocyte

immune response.

Discussion

In this work, we demonstrated for the first time that cardiac microlesions form during severe *E. faecalis* bacteremic infection of mice. Importantly, microlesions occurred despite *E. faecalis* not encoding homologs of the pneumococcal virulence determinants implicated in microlesion formation. These data suggest that *E. faecalis* may be expressing functionally redundant proteins that serve to promote similar interactions to *S. pneumoniae* in mediating microlesion formation, such as CbpA and pneumolysin, which facilitate vascular endothelial cell translocation and cardiomyocyte cell death, respectively. Indeed, while *E. faecalis* does not encode a CbpA homolog, it does encode other proteins capable of binding host extracellular matrix proteins. These include EfbA, which binds fibronectin [41], ACE, which binds laminin receptor [42], and Ebp pili that bind fibrinogen [43].

Interestingly, we did discover one factor, DsbA, that contributes to microlesion formation and inhibits the cardiomyocyte inflammatory immune response, as modeled in Figure 4B. Dsb proteins work to catalyze disulfide bond formation in proteins and have been best studied in Gram-negative bacteria [44, 45]. While Dsb proteins have established importance in the biology of certain Gram-positive bacteria, Dsb proteins are not well studied in Firmicutes like *E. faecalis* and have generally not been considered important [44]. Most of these species secrete few proteins containing cysteines and some, notably *S. pneumoniae*, do not encode any thiol-disulfide oxidoreductases [46, 47]. Thus, it will be of interest to further investigate the role of Dsb proteins in these species.

Our prior work established EntV as a DsbA substrate, currently the only one identified [21]. However, loss of EntV did not reduce pathogenicity in the worm model (Figure 1A), and exposure to EntV, even at very high concentrations, was non-toxic to mammalian cells [22]. As such, we do

not favor EntV as being the source of DsbA's effects. While the specific *E. faecalis* proteins

involved in modulating the host immune response are unknown, we speculate that the

Fibronectin-binding protein (Efb) and Choline binding protein may contribute to cardiac

microlesion formation by serving as functionally redundant proteins to pneumococcal CbpA, a

virulence determinant known to be integral to microlesion formation [11]. Moreover, both proteins

contain 5 cysteines and are expressed extracellularly. Thus, our future research efforts will be

geared toward identifying additional substrates of DsbA and to determining their contribution to

microlesion formation.

Pneumococcal ChoP interactions with host PAFr were determined to be a prerequisite for

vascular endothelial cell invasion during IPD. In light of our observations of *E. faecalis* invasion of

vascular endothelial cells despite an absence of this wall teichoic acid structure, it is possible that

another host receptor is required for endothelial cell invasion. Interestingly, recent studies focused

on bacterial cell wall entry and signaling in eukaryotic cells have identified a PAFr-independent

mechanism of tissue invasion that is actin-dependent and engenders Rac1, Cdc42, and

phosphatidylinositol 3-kinase (PI3K) signaling leading to cell wall internalization via a

macropinoctytosis-like pathway [48]. Whether such a pathway contributes to microlesion

formation is unknown. In the future, it will be of interest to study the mechanisms of tissue invasion

and the host signaling pathways involved.

Although the role of cardiac microlesion formation in humans during invasive disease remains

unclear, cardiac microlesion formation is becoming increasingly appreciated as a risk factor [49].

To our knowledge, no one has investigated whether cardiac microlesions form in humans during

enterococcal infection. We speculate that formation occurs and contributes to cardiac

abnormalities or death in patients with enterococcal infective endocarditis or sepsis [1, 3-9]. This

study builds on our understanding of cardiac microlesions by establishing *E. faecalis* as another

etiological agent capable of causing them in a vertebrate animal model.　Considering that *E.*

faecalis microlesion pathology appears identical to that caused by *S. pneumoniae*, it is likely that

scar formation occurs following resolution of infection, potentially predisposing individuals to future

adverse cardiac events [11]. Taken together, these data highlight the need for further investigation

in order to understand the impact of microlesion formation on the heart and mortality during

invasive *E. faecalis* infection.

Table S1. Strains used in this study
343

Enterococcus faecalis Strains				
Strain	**Genotype**	**Characteristics**	**Parent**	**Source**
OG1RF	Wild-type	Gel$^+$ Spr$^+$ Rf^R Fa^R	OG1	[50]
AOBD1	Δ*dsbA*	Gel$^+$ Spr$^+$ Rf^R Fa^R	OG1	[21]
AOBD2	Δ*dsbA::dsbA*	Gel$^+$ Spr$^+$ Rf^R Fa^R	OG1	[21]

344
345
346
347 **Table S2. Cytokine Profiling Expression Data**
348
349

Cytokine Profiling - Fold Change Data for 3 independent Experiments
4 hour infection with an MOI=10

	Exp. 1 OG1RF	Exp. 2 OG1RF	Exp. 3 OG1RF	Average	Std. Dev
TNF-a	-7%	7%	-2%	-1%	7%
IGF-1	2%	-7%	4%	0%	6%
VEGF	23%	4%	-2%	8%	13%
IL-6	7%	5%	1%	4%	3%
FGFb	11%	-28%	0%	-5%	20%
IFNr	5%	3%	-11%	-1%	9%
EGF	-10%	33%	2%	8%	23%
Leptin	0%	-16%	-1%	-6%	9%
IL-1a	-7%	-14%	4%	-6%	9%
IL-1b	-8%	-22%	18%	-4%	20%
G-CSF	1%	-22%	2%	-6%	14%
GM-CSF	-10%	19%	14%	7%	16%
MCP-1	-7%	-8%	11%	-1%	11%
M1P-1a	3%	3%	-17%	-4%	12%
SCF	-15%	5%	-3%	-5%	10%
Rantes	-13%	19%	-7%	0%	17%
PDGF-BB	-16%	8%	7%	0%	13%
b-NGF	-20%	12%	17%	3%	20%
IL-17a	-22%	-4%	9%	-5%	16%
IL-2	-9%	-10%	-5%	-8%	3%
IL-4	12%	-1%	9%	7%	7%
IL-10	11%	-13%	-3%	-2%	12%
Resistin	-13%	-3%	-6%	-8%	5%
IL-12	-3%	0%	-17%	-7%	9%
CCL11	-9%	-25%	3%	-10%	14%
CCL21	-17%	0%	-5%	-7%	9%
IL-3	8%	3%	5%	5%	2%
IL-13	-7%	-7%	10%	-1%	10%
IL-21	-24%	-14%	1%	-12%	13%
IL-22	-18%	-19%	-12%	-16%	4%
CXCL10	-12%	-10%	2%	-7%	8%
CXCL1	-5%	-4%	-13%	-7%	5%

*OG1RF ordered from Largest to smallest

	OG1RF	ΔdsbA
EGF	8%	3%
VEGF	8%	1%
GM-CSF	7%	1%
IL-4	7%	-12%
IL-3	5%	-1%
IL-6	4%	2%
b-NGF	3%	-8%
PDGF-BB	0%	-2%
IGF-1	0%	-6%
Rantes	0%	4%
TNF-a	-1%	-2%
IFNr	-1%	4%
IL-13	-1%	-3%
MCP-1	-1%	-2%
IL-10	-2%	-2%
M1P-1a	-4%	3%
IL-1b	-4%	-13%
SCF	-5%	-7%
FGFb	-5%	-8%
IL-17a	-5%	-14%
IL-1a	-6%	-7%
Leptin	-6%	0%
G-CSF	-6%	2%
CXCL10	-7%	-11%
IL-12	-7%	1%
CXCL1	-7%	-3%
CCL21	-7%	-9%
Resistin	-8%	-10%
IL-2	-8%	-11%
CCL11	-10%	-12%
IL-21	-12%	-9%
IL-22	-16%	-5%

	Exp. 1 ΔdsbA	Exp. 2 ΔdsbA	Exp. 3 ΔdsbA	Average	Std. Dev
TNF-a	-2%	5%	-8%	-2%	6%
IGF-1	-15%	-7%	3%	-6%	9%
VEGF	2%	1%	0%	1%	1%
IL-6	0%	13%	-8%	2%	10%
FGFb	-7%	-15%	-2%	-8%	7%
IFNr	16%	7%	-12%	4%	14%
EGF	-8%	31%	-12%	3%	24%
Leptin	13%	-8%	-4%	0%	11%
IL-1a	-17%	-10%	7%	-7%	12%
IL-1b	-20%	-19%	-1%	-13%	11%
G-CSF	-12%	14%	3%	2%	13%
GM-CSF	-18%	18%	2%	1%	18%
MCP-1	-20%	-9%	23%	-2%	22%
M1P-1a	-6%	3%	13%	3%	10%
SCF	-10%	-20%	10%	-7%	15%
Rantes	-3%	20%	-4%	4%	14%
PDGF-BB	-23%	15%	4%	-2%	20%
b-NGF	-16%	-10%	1%	-8%	9%
IL-17a	-45%	-1%	4%	-14%	27%
IL-2	-35%	-1%	1%	-11%	20%
IL-4	-17%	-21%	2%	-12%	13%
IL-10	0%	-2%	-4%	-2%	2%
Resistin	-19%	-6%	-6%	-10%	7%
IL-12	-13%	14%	1%	1%	14%
CCL11	-13%	-30%	7%	-12%	18%
CCL21	-33%	4%	3%	-9%	21%
IL-3	-8%	2%	3%	-1%	6%
IL-13	-25%	-3%	19%	-3%	22%
IL-21	-23%	-10%	6%	-9%	15%
IL-22	-26%	-7%	18%	-5%	22%
CXCL10	-14%	-17%	-2%	-11%	8%
CXCL1	-12%	-15%	17%	-3%	17%

Figure Legends

Figure 1. *dsbA* deficient *E. faecalis* show reduced virulence in a *C. elegans* survival assay. **(A)** L4 stage *C. elegans* were infected with *E. faecalis*. This figure is representative of 3 independent experiments with 30 worms/group. ****Asterisks indicate a *P* < 0.0001, ** a P < 0.01 when compared to OG1RF following pairwise analysis by log-rank (Mantel-Cox). **(B)** Western blot performed by SDS-PAGE separation of *E. faecalis* whole cell lysates followed by hybridization of DsbA using rabbit anti-DsbA antisera. Equal loading is ensured by visualization of a non-specific band at 40 kDa. Image is representative of 3 independent Western blot experiments.

Figure 2. Cardiac microlesions form during invasive *E. faecalis* infection and are attenuated by loss of *dsbA*. Cardiac microlesions were observed using light microscopy of hematoxylin and eosin (H&E) stained paraffin embedded cardiac sections at 40X magnification **(A)**, and 100X magnification **(B)**. Immunofluorescent detection of wild-type *E. faecalis* was imaged using a 20X objective at 3X zoom (total of 60X magnification), a FITC conjugated anti-enterococcal antibody **(C)**, and DAPI for DNA staining **(D)**. Total microlesion counts **(E)**, and size measurements **(F)**, were obtained by quantification using light microscopy of H&E stained paraffin embedded cardiac sections from mice infected for 24-48 hours with *E. faecalis* (OG1RF n=14, Δ*dsbA* n=14*,* Δ*dsbA*::*dsbA* n=14). The data were averaged from 3 independent experiments and analyzed using *Student's* two-tailed t-test (* indicate a *P* < 0.05, ** *P* < 0.01, *** a *P* < 0.001).

Figure 3. Necroptosis contributes to cardiotoxicity within cardiac microlesions. **(A)** Immunofluorescent microscopy of paraffin embedded cardiac cross sections at 20X were performed to detect *E. faecalis* using FITC conjugated anti-faecalis antibodies, and apoptosis using Alexa 594 conjugated anti-PARP antibodies at sites of microlesion formation, while DAPI

was used to stain nuclei. **(B)** Immunofluorescent microscopy of paraffin embedded cardiac cross

sections at 20X were performed to detect MLKL using FITC conjugated anti-MLKL antibodies, and

Alexa 594 conjugated anti-phosphorylated MLKL antibodies at sites of microlesion formation,

while DAPI was used to stain nuclei. **(C)** *E. faecalis* induced cell death in HL-1 cardiomyocytes 4

hours post-infection (h.p.i.) as determined by measuring lactate dehydrogenase. The data

represent an average of 3 independent experiments and were analyzed using *Student's* two-tailed

t-test (* indicate a *P* < 0.05). **(D)** Comparison of OG1RF and Δ*dsbA* invasion rates into HL-1

cardiomyocytes and SVEC endothelial cells. The graph represents data from 3 independent

experiments that were analyzed using *Student's* two-tailed t-test.

Figure 4. Wild-type *E. faecalis* suppresses the cardiomyocyte inflammatory response *in-

***vitro*. (A)** HL-1 cardiomyocytes exposed to *E. faecalis* promote a suppressed inflammatory

response while Δ*dsbA* infected cardiomyocytes yield a more inflammatory response. Data was

obtained from the fold-expression change analysis (relative to uninfected controls) of at least 3

independent experiments for each strain. **(B)** Model of *E. faecalis* mediated suppression of the

host cardiomyocyte immune response and effect on microlesion size. Wild-type *E. faecalis* is

postulated to downregulate the cardiomyocyte immune response more effectively than the *dsbA*

mutant resulting in more cell death and increased lesion size. However, both strains cause cell

death by apoptosis and necroptosis and inhibit immune cell infiltration.

Supplemental Figure 1. *E. faecalis* growth curve. (A) *E. faecalis* strains were grown for 15

hours in BHI and absorbance (OD_{600nm}) measured every 10 minutes for 16 hours. **(B)** *E. faecalis*

grown in serum and absorbance (OD_{600nm}) measured every 10 minutes for 16 hours. Figures are

representative of 3 independent experiments.

Supplemental Figure 2. Coomassie stained SDS-PAGE gel of *E. faecalis* whole cell lysates separated by electrophoresis. (A) SDS-PAGE separation of *E. faecalis* whole cell lysates was stained with Coomassie to verify that each well received 10 µg / lane of whole cell lysate. Coomassie stained gel is the loading control for the Western blot in Figure 1B.

Supplemental Figure 3. DsbA is required for *E. faecalis* pathogenesis. (A) Mice (n=17-20) challenged with 3.8 – 4.2 x 10^8 CFUs of *E. faecalis* in a mouse model of peritonitis were assessed for survival over 96 hours. Mice infected with Δ*dsbA E. faecalis* lived significantly longer (*P* = 0.0051) than mice challenged with wild-type or the complemented strain (no significant difference). Survival curve was calculated from 3 independent experiments using log-rank analysis.

Supplemental Figure 4. (A) 40X magnification image of cardiac microlesion formation in mice infected with Δ*dsbA.* **(B)** 100X image of cardiac microlesion in Figure S4A. **(C)** 40X magnification image of cardiac microlesion formation in mice infected with Δ*dsbA*::*dsbA*. **(D)** 100X image of cardiac microlesion in Figure S4C.

Supplemental Figure 5. Necroptosis contributes to cardiotoxicity within cardiac microlesions. **(A)** Immunofluorescent microscopy of paraffin embedded cardiac cross sections using a 20X objective at 4X Zoom (80X total magnification) were performed to detect MLKL using FITC conjugated anti-MLKL antibodies, and Alexa 594 conjugated anti-phosphorylated MLKL antibodies at sites of microlesion formation, while DAPI was used to stain nuclei.

Supplemental Figure 6. Apoptosis contributes to cell death within cardiac microlesions. **(A)** Immunofluorescent microscopy of paraffin embedded cardiac cross sections using a 20X

objective at 4X Zoom (80X total magnification) were performed to detect *E. faecalis* using FITC

conjugated anti-*E. faecalis* antibodies, and Alexa 594 conjugated anti-PARP antibodies at sites of

microlesion formation, while DAPI was used to stain nuclei.

References

1. Chirouze C, Athan E, Alla F, et al. Enterococcal endocarditis in the beginning of the 21st

century: analysis from the International Collaboration on Endocarditis-Prospective Cohort Study.

Clin Microbiol Infect 2013; 19:1140-7.

2. Agudelo Higuita NI, Huycke MM. Enterococcal Disease, Epidemiology, and Implications for

Treatment. In: Gilmore MS, Clewell DB, Ike Y, Shankar N, eds. Enterococci: From Commensals

to Leading Causes of Drug Resistant Infection. Boston, 2014.

3. Cabell CH, Jollis JG, Peterson GE, et al. Changing patient characteristics and the effect on

mortality in endocarditis. Arch Intern Med 2002; 162:90-4.

4. Garvey GJ, Neu HC. Infective endocarditis--an evolving disease. A review of endocarditis at

the Columbia-Presbyterian Medical Center, 1968-1973. Medicine (Baltimore) 1978; 57:105-27.

5. Geraci JE, Martin WJ. Antibiotic therapy of bacterial endocarditis. VI. Subacute enterococcal

endocarditis; clinical, pathologic and therapeutic consideration of 33 cases. Circulation 1954;

10:173-94.

6. Jawetz E, Sonne M. Penicillin-streptomycin treatment of enterococcal endocarditis. A re-

evaluation. N Engl J Med 1966; 274:710-5.

7. Lerner PI, Weinstein L. Infective endocarditis in the antibiotic era. N Engl J Med 1966; 274:388-

93 concl.

8. Olaison L, Schadewitz K, Swedish Society of Infectious Diseases Quality Assurance Study

Group for E. Enterococcal endocarditis in Sweden, 1995-1999: can shorter therapy with

aminoglycosides be used? Clin Infect Dis 2002; 34:159-66.

9. Pelletier LL, Jr., Petersdorf RG. Infective endocarditis: a review of 125 cases from the University

of Washington Hospitals, 1963-72. Medicine (Baltimore) 1977; 56:287-313.

10. Garsin DA, Frank KL, Silanpaa J, et al. Pathogenesis and Models of Enterococcal Infection.

In: Gilmore MS, Clewell DB, Ike Y, Shankar N, eds. Enterococci: From Commensals to Leading

Causes of Drug Resistant Infection. Boston, 2014.

11. Brown AO, Mann B, Gao G, et al. Streptococcus pneumoniae translocates into the

myocardium and forms unique microlesions that disrupt cardiac function. PLoS Pathog 2014;

10:e1004383.

12. Reyes LF, Restrepo MI, Hinojosa CA, et al. Severe Pneumococcal Pneumonia Causes Acute

Cardiac Toxicity and Subsequent Cardiac Remodeling. Am J Respir Crit Care Med 2017;

196:609-20.

13. Shenoy AT, Brissac T, Gilley RP, et al. Streptococcus pneumoniae in the heart subvert the

host response through biofilm-mediated resident macrophage killing. PLoS Pathog 2017;

13:e1006582.

14. Brissac T, Shenoy AT, Patterson LA, Orihuela CJ. Cell Invasion and Pyruvate Oxidase-

Derived H_2O_2 Are Critical for Streptococcus pneumoniae-Mediated Cardiomyocyte Killing. Infect

Immun 2018; 86.

15. Makara MA, Hoang KV, Ganesan LP, et al. Cardiac Electrical and Structural Changes During

Bacterial Infection: An Instructive Model to Study Cardiac Dysfunction in Sepsis. J Am Heart

Assoc 2016; 5.

16. Headley CA, Gerberick A, Mehta S, et al. Nontuberculous mycobacterium M. avium infection

predisposes aged mice to cardiac abnormalities and inflammation. Aging Cell 2019; 18:e12926.

17. Garsin DA, Sifri CD, Mylonakis E, et al. A simple model host for identifying Gram-positive

virulence factors. Proc Natl Acad Sci U S A 2001; 98:10892-7.

18. Nallapareddy SR, Murray BE. Role played by serum, a biological cue, in the adherence of

Enterococcus faecalis to extracellular matrix proteins, collagen, fibrinogen, and fibronectin. J

Infect Dis 2008; 197:1728-36.

19. Teng F, Singh KV, Bourgogne A, Zeng J, Murray BE. Further characterization of the epa gene

cluster and Epa polysaccharides of Enterococcus faecalis. Infect Immun 2009; 77:3759-67.

20. Singh KV, Qin X, Weinstock GM, Murray BE. Generation and testing of mutants of

Enterococcus faecalis in a mouse peritonitis model. J Infect Dis 1998; 178:1416-20.

21. Armand O. Brown CEG, Melissa R. Cruz, Michael C. Lorenz, Danielle A. Garsin. Antifungal

activity of the Enterococcus faecalis peptide EntV requires protease cleavage and disulfide bond

formation. Mbio 2019; 10:1-13.

22. Graham CE, Cruz MR, Garsin DA, Lorenz MC. Enterococcus faecalis bacteriocin EntV inhibits

hyphal morphogenesis, biofilm formation, and virulence of Candida albicans. Proc Natl Acad Sci

U S A 2017; 114:4507-12.

23. Zhang T, Zhang Y, Cui M, et al. CaMKII is a RIP3 substrate mediating ischemia- and oxidative

stress-induced myocardial necroptosis. Nat Med 2016; 22:175-82.

24. Linkermann A, Hackl MJ, Kunzendorf U, Walczak H, Krautwald S, Jevnikar AM. Necroptosis

in immunity and ischemia-reperfusion injury. Am J Transplant 2013; 13:2797-804.

25. Linkermann A, Brasen JH, Darding M, et al. Two independent pathways of regulated necrosis

mediate ischemia-reperfusion injury. Proc Natl Acad Sci U S A 2013; 110:12024-9.

26. Kitur K, Wachtel S, Brown A, et al. Necroptosis Promotes Staphylococcus aureus Clearance

by Inhibiting Excessive Inflammatory Signaling. Cell Rep 2016; 16:2219-30.

27. Gilley RP, Gonzalez-Juarbe N, Shenoy AT, et al. Infiltrated Macrophages Die of Pneumolysin-

Mediated Necroptosis following Pneumococcal Myocardial Invasion. Infect Immun 2016; 84:1457-

69.

28. Gentry-Weeks CR, Karkhoff-Schweizer R, Pikis A, Estay M, Keith JM. Survival of

Enterococcus faecalis in mouse peritoneal macrophages. Infect Immun 1999; 67:2160-5.

29. Zou J, Shankar N. The opportunistic pathogen Enterococcus faecalis resists phagosome

acidification and autophagy to promote intracellular survival in macrophages. Cell Microbiol 2016;

18:831-43.

30. Zou J, Shankar N. Enterococcus faecalis infection activates phosphatidylinositol 3-kinase

signaling to block apoptotic cell death in macrophages. Infect Immun 2014; 82:5132-42.

31. Huang XL, Wang YJ, Yan JW, et al. Role of anti-inflammatory cytokines IL-4 and IL-13 in

systemic sclerosis. Inflamm Res 2015; 64:151-9.

32. Minnone G, De Benedetti F, Bracci-Laudiero L. NGF and Its Receptors in the Regulation of

Inflammatory Response. Int J Mol Sci 2017; 18.

33. Yano K, Liaw PC, Mullington JM, et al. Vascular endothelial growth factor is an important

determinant of sepsis morbidity and mortality. J Exp Med 2006; 203:1447-58.

34. Bhattacharya P, Thiruppathi M, Elshabrawy HA, Alharshawi K, Kumar P, Prabhakar BS. GM-

CSF: An immune modulatory cytokine that can suppress autoimmunity. Cytokine 2015; 75:261-

71.

35. Shimoji K, Yuasa S, Onizuka T, et al. G-CSF promotes the proliferation of developing

cardiomyocytes in vivo and in derivation from ESCs and iPSCs. Cell Stem Cell 2010; 6:227-37.

36. Martins A, Han J, Kim SO. The multifaceted effects of granulocyte colony-stimulating factor in

immunomodulation and potential roles in intestinal immune homeostasis. IUBMB Life 2010;

62:611-7.

37. Lee SH, Kwon JY, Kim SY, Jung K, Cho ML. Interferon-gamma regulates inflammatory cell

death by targeting necroptosis in experimental autoimmune arthritis. Sci Rep 2017; 7:10133.

38. Kothapalli R, Nyland SB, Kusmartseva I, Bailey RD, McKeown TM, Loughran TP, Jr.

Constitutive production of proinflammatory cytokines RANTES, MIP-1beta and IL-18

characterizes LGL leukemia. Int J Oncol 2005; 26:529-35.

39. La Cava A. Leptin in inflammation and autoimmunity. Cytokine 2017; 98:51-8.

40. Trinchieri G. Interleukin-12: a proinflammatory cytokine with immunoregulatory functions that

bridge innate resistance and antigen-specific adaptive immunity. Annu Rev Immunol 1995;

13:251-76.

41. Singh KV, La Rosa SL, Somarajan SR, Roh JH, Murray BE. The fibronectin-binding protein

EfbA contributes to pathogenesis and protects against infective endocarditis caused by

Enterococcus faecalis. Infect Immun 2015; 83:4487-94.

42. Nallapareddy SR, Qin X, Weinstock GM, Hook M, Murray BE. Enterococcus faecalis adhesin,

ace, mediates attachment to extracellular matrix proteins collagen type IV and laminin as well as

collagen type I. Infect Immun 2000; 68:5218-24.

43. .Nallapareddy SR, Singh KV, Sillanpaa J, et al. Endocarditis and biofilm-associated pili of

Enterococcus faecalis. J Clin Invest 2006; 116:2799-807.

44. Reardon-Robinson ME, Ton-That H. Disulfide-Bond-Forming Pathways in Gram-Positive

Bacteria. J Bacteriol 2015; 198:746-54.

45. Landeta C, Boyd D, Beckwith J. Disulfide bond formation in prokaryotes. Nat Microbiol 2018;

3:270-80.

46. Daniels R, Mellroth P, Bernsel A, et al. Disulfide bond formation and cysteine exclusion in

gram-positive bacteria. J Biol Chem 2010; 285:3300-9.

47. Dutton RJ, Boyd D, Berkmen M, Beckwith J. Bacterial species exhibit diversity in their

mechanisms and capacity for protein disulfide bond formation. Proc Natl Acad Sci U S A 2008;

105:11933-8.

48. Loh LN, Gao G, Tuomanen EI. Dissecting Bacterial Cell Wall Entry and Signaling in Eukaryotic

Cells: an Actin-Dependent Pathway Parallels Platelet-Activating Factor Receptor-Mediated

Endocytosis. MBio 2017; 8.

49. Musher DM, Abers MS, Corrales-Medina VF. Acute Infection and Myocardial Infarction. N

Engl J Med 2019; 380:171-6.

50. Murray BE, Singh KV, Ross RP, Heath JD, Dunny GM, Weinstock GM. Generation of

restriction map of Enterococcus faecalis OG1 and investigation of growth requirements and

regions encoding biosynthetic function. J Bacteriol 1993; 175:5216-23.

FOOTNOTE PAGE

Conflict of Interest statement

There are no conflicts of interest.

Funding statement

This work was supported by the National Institute of Dental and Craniofacial Research of the

National Institutes of Health under [grant number R01DE027608 to D. A. G.] and [grant number

F32DE027580 to A. O. B.]

Meetings

This work has not been presented.

Corresponding author contact information

Danielle A. Garsin, Ph.D.

6431 Fannin St.

MSB 1.168

Houston, TX 77030

Telephone: (713) 500-5454

Fax: (713) 500-5499

Email: Danielle.A.Garsin@uth.tmc.edu

Figure 1

Figure 2

Figure 3

Figure 4