Cardiac Microlesions Form During Severe Bacteremic *Enterococcus faecalis* Infection

Armand O. Brown[1], Kavindra V. Singh[2], Melissa R. Cruz[1], Karan Gautam Kaval[1], Liezl E. Francisco[3], Barbara E. Murray[1,2], and Danielle A. Garsin[1*]

[1]Department of Microbiology and Molecular Genetics, The University of Texas Health Science Center at Houston, Houston TX, USA

[2] Division of Infectious Diseases, Department of Internal Medicine, The University of Texas Health Science Center at Houston, Houston, TX, USA

[3] Department of Biochemistry and Structural Biology, The University of Texas Health Science Center at San Antonio, San Antonio, TX, USA

*Corresponding author

Email: Danielle.A.Garsin@uth.tmc.edu

Abstract

Enterococci, such as *Enterococcus faecalis,* are a significant cause of hospital-acquired bacteremia and mortality among the elderly and patients with comorbid conditions in the healthcare setting. In this study, the novel discovery is made that cardiac microlesions form during severe bacteremic *E. faecalis* infection in mice. The cardiac microlesions were identical in appearance to those previously observed to form by *Streptococcus pneumoniae* during invasive pneumococcal disease (IPD) in mice, non-human primates, and humans. However, *E. faecalis* does not encode the virulence determinants implicated in pneumococcal microlesion formation, suggesting that formation is through an alternative pathway. In previous work, we showed that *E. faecalis* DsbA is required for the post-translational disulfide bond formation and stability of EntV, a secreted *E. faecalis* bacteriocin. Here, the role of DsbA in *E. faecalis* pathogenicity was explored. DsbA was discovered to be required for *E. faecalis* virulence in a *C. elegans* model and, importantly, was necessary for efficient cardiac microlesion formation. By investigating the cardiac host response, it was discovered that *E. faecalis* promotes cardiomyocyte necroptotic cell death at sites of microlesion formation. Additionally, by examining the cytokines induced by exposure of *E. faecalis* to cardiomyocytes, it was revealed that loss of DsbA caused a pro-inflammatory response unlike the wild type strain, which suppressed the immune response. In conclusion, we establish that *E. faecalis* is capable of forming cardiac microlesions and identify features of both the bacterial and host responses that are mechanistically involved.

Significance Statement

E. faecalis is a bacterium capable of causing serious disease in humans, including endocarditis,

an infection on the surface of the heart valves with a high mortality rate. In this work, we document

for the first time that *E. faecalis* is capable of also causing an invasive infection of the heart,

forming cardiac microlesions in a mouse model of severe bloodstream infection. Additionally, a

contributing bacterial factor is identified and the immune response of heart cells to infection with

E. faecalis is characterized. Overall, this work makes a significant contribution to the field of

infectious diseases by identifying a new enterococcal disease pathology, cardiac microlesion

formation, which was previously known to occur only with IPD.

Introduction

Enterococcus faecalis is a Gram-positive bacterium that normally co-exists with the host as a harmless commensal. However, it is capable of causing a variety of infections such as infective endocarditis (IE), urinary tract infection (UTI), bacteremia, peritonitis, prosthetic joint infection, and endophthalmitis. These infections can become life-threatening if left untreated (1, 2). Furthermore, immunocompromised individuals and those with underlying disease are at an increased risk (3-5). Enterococcal infections are increasingly characterized by the presence of antimicrobial resistance such as vancomycin resistant enterococci (VRE). Due to their intrinsic ability to survive various environmental habitats (3), the bacterium poses a significant threat in hospitals and long-term care facilities where VRE has increased access to immunocompromised patients through infection of open wounds, or through the use of contaminated medical devices (6, 7). As such, enterococci are a common nosocomial pathogen, having been observed to be the etiological agent responsible for 14% of all hospital-acquired infections (HAIs) in the US between 2011–2014 (8). Therefore, it is no surprise that enterococci are a significant cause of hospital-acquired bacteremia and mortality among the elderly and patients with comorbid conditions in the healthcare setting (9-12).

Enterococcal infections often lead to other clinical manifestations. A common complication associated with bacteremia is endocarditis, an infection that has a 1-year mortality rate of approximately 29% despite antibiotic therapy (11). Enterococci are the third most common cause of endocarditis in North America (5) and, in patients with this infection, heart failure has been observed to occur with increased frequency (13-19). Infective endocarditis involves the formation of biofilms on heart valves and inner chambers that develop into complex masses called vegetations (2). Thus, heart infections associated with *E. faecalis* have traditionally been

characterized as occurring on the surfaces of the heart, rather than in the deeper myocardial

tissue, which would require active invasion by the bacterium.

Interestingly, another important Gram-positive pathogen, *Streptococcus pneumoniae,* was

recently shown to invade the heart tissue and form cardiac microlesions during invasive

pneumococcal disease (IPD) (20). The microlesions occurred in mice, non-human primates, and

humans with IPD and were shown to be associated with altered cardiac electrophysiology, heart

failure in mice, and scar formation during convalescence (20). Scar formation following

microlesion development is thought to predispose individuals to major adverse cardiac events

(MACE) (21). Moreover, cardiac microlesions were shown to form following vascular endothelial

cell invasion into the myocardial tissue, followed by subsequent interactions between

pneumococcal Choline binding protein A (CbpA) and pneumococcal teichoic acid associated

phosphorylcholine (ChoP) with host laminin receptor (LR) and platelet activating factor receptor

(PAFr), respectively. Furthermore, cardiomyocyte death is mediated by exposure to the

pneumococcal secreted cholesterol-dependent pore forming toxin, pneumolysin (20, 21), and

hydrogen peroxide (22, 23). As such, cardiac microlesions have been proposed to be a

contributory factor to the onset of MACE observed in older patients with IPD, or in convalescent

patients previously hospitalized with IPD (20, 24, 25). Importantly, invasive infection is now

appreciated as a risk factor for myocardial infarction, and the risk appears to be more pronounced

depending on the severity of the infection (26, 27). Since the original observation of pneumococcal

microlesions in 2014, no other organism has been observed to cause such pathology.

In this report, we show that cardiac microlesions also form during invasive *E. faecalis* infection.

The observed cardiac microlesions are identical in appearance and size to *S. pneumoniae*

mediated cardiac microlesions, are diffuse across the myocardium, and are characterized by an

absence of immune cell infiltration. Furthermore, we identify a mode of cardiomyocyte cell death,

and an enterococcal determinant required for microlesion formation. Our findings suggest a

previously unrecognized pathological aspect of enterococcal infection that may be a contributing

factor to the increased rate of heart failure and mortality observed in patients with endocarditis

(13-19). Importantly, because *E. faecalis* lacks homologs of the virulence determinants previously

implicated in pneumococcal microlesion formation, the mechanism of microlesion formation for *E.*

faecalis is likely through an alternative pathway.

Results

Cardiac microlesions form during severe bacteremic *E. faecalis* infection. In previous work, we implicated a previously uncharacterized *E. faecalis* thioredoxin known as DsbA (disulfide bond forming protein A) as a requirement for the stability of EntV, a secreted bacteriocin that we determined inhibits hyphal morphogenesis of the human fungal pathogen *Candida albicans* (28, 29). To test if these proteins had any effects on the fitness and pathogenicity of *E. faecalis*, we infected *Caenorhabditis elegans* with *ΔdsbA* and *ΔentV E. faecalis* mutants and measured survival. As shown in Figure 1A, loss of *dsbA* significantly attenuated mortality in infected nematodes. Note that the *ΔdsbA* mutant did not manifest different growth kinetics when grown in culture medium compared to wild type, ruling out a general growth defect (Figure S1). Infection with the complement, *ΔdsbA::dsbA*, partially rescued the *ΔdsbA* phenotype. The partial complementation was attributed to the DsbA protein being produced at lower levels in the complemented strain (Figure 1B). No significant difference in nematode survival was observed following infection with either the *ΔentV* mutant or its complement (Fig. 1A). Considering that EntV is the only known substrate of *E. faecalis* DsbA, the results suggest that DsbA has additional targets that affect pathogenicity.

Based on these results, we wanted to know if loss of DsbA also affected virulence in a vertebrate animal model of *E. faecalis* infection. Therefore, the mutants were tested in a mouse model in which animals were injected intra-peritoneally with *E. faecalis* resulting in generalized bacteremia leading to mortality. Significant survival differences between the mice infected with the different strains were not observed. However, various organs were collected in order to determine if *E. faecalis* infection induced any strain-specific, pathological effects. To our surprise, cardiac microlesions were discovered within the myocardium that were devoid of immune cell infiltrate

(Fig. 2 A, B). The microlesions appeared to form within roughly the same time frame (~24 h.p.i.)

as cardiac microlesions that have previously been reported to occur during invasive infection with

S. pneumoniae, the only other pathogen known to cause cardiac microlesions to date (20).

S. pneumoniae mediated microlesions have been shown to form in mice, non-human primates,

and humans, and were correlated with heart failure in mouse models of infection (20). Because

E. faecalis lacks homologs of the pneumococcal virulence determinants identified to promote

pneumococcal cardiac microlesion formation, such as choline binding protein A (CbpA) (20), the

cholesterol-dependent pore forming toxin pneumolysin (20), and hydrogen peroxide (30), we

decided to study this phenomenon more closely. To verify that the observed microlesions were

the result of infection by *E. faecalis*, immunofluorescent staining on paraffin embedded cardiac

tissue sections from mice infected with *E. faecalis* with a polyclonal anti-enterococcal IgG antibody

was performed and revealed the presence of dense aggregates of *E. faecalis* within the

microlesions (Fig. 2C). The presence of bacterial DNA was further verified by DNA counterstaining

with DAPI that revealed an accumulation of non-cardiomyocyte nuclear DNA within microlesions

(Fig. 2D). To determine the contribution of DsbA toward microlesion formation, cardiac

microlesions found on hematoxylin and eosin stained paraffin embedded cardiac sections were

quantified following infection with Δ*dsbA*. From these experiments, a significant attenuation in

microlesion formation during infection with Δ*dsbA* was observed relative to wild type, while no

significant difference was observed in mice infected with the *dsbA* complement (Fig. 2E).

Furthermore, when the sizes of the cardiac microlesions were compared, the wild-type lesions

were significantly larger than those found in the Δ*dsbA* infected mice (Fig. 2F). Importantly, the

size of the microlesions appeared to correlate with level of *dsbA* expression observed in Western

blots of *E. faecalis* whole cell lysates (Fig. 1B), with the Δ*dsbA::dsbA* strain manifesting an in-

between phenotype to that of the Δ*dsbA* and wild-type strains, consistent with partial

complementation. In conclusion, these data demonstrate that cardiac microlesion formation

occurs during systematic infection with *E. faecalis* and is dependent on DsbA. Thus, for the first

time, loss of DsbA in *E. faecalis* is shown to affect pathogenicity in animal models of infection,

both invertebrate (*C. elegans*) and vertebrate (mice).

Necroptosis contributes to cardiomyocyte cell death within microlesions. In previous

studies of pneumococcal cardiac microlesion formation, apoptosis was found to be associated

with microlesion formation in mice (22). A subsequent study determined that cardiomyocyte cell

death occurs following exposure to pneumolysin (*ply*) and pneumococcal derived hydrogen

peroxide (*spxB*) (23). Based on these observations and the fact that *E. faecalis* lacks *ply* and *spxB*

homologs, we sought to determine whether *E. faecalis* could promote cell death of

cardiomyocytes and identify the mechanism. First, HL-1 cardiomyocytes were infected with *E.*

faecalis and subsequently lactate dehydrogenase (LDH) release into the cell culture supernatant

was measured to assess cell death. From these experiments, it was concluded that *E. faecalis*

induces cell death of cardiomyocytes. Furthermore, infection with a *ΔdsbA* mutant resulted in a

significant decrease in cardiomyocyte cell death (Fig. 3B), suggesting DsbA is required to

modulate this process. Since cardiomyocyte cell death was observed following exposure to *E.*

faecalis, we wanted to identify the mechanism. To test this question, an annexin V apoptosis and

necrosis assay was performed to determine if *E. faecalis* promotes apoptosis or necrosis during

cardiomyocyte infection. The signals for both apoptosis and necrosis increased concomitantly

with time (Figure S2), and this read-out means that cell death is occurring by an apoptosis-

independent mechanism (see Materials and Methods). One apoptosis-independent cell death

pathway is necroptosis, and this mechanism of cell death is consistent with data published by

investigators studying *S. pnuemoniae* microlesions, which showed that markers of necroptosis

could be detected within the myocardium of acutely ill mice (20) and non-human primates (21).

Importantly, necroptosis is considered a key cell death pathway in cardiomyocytes during

ischemia-reperfusion injury and acute coronary syndrome (31-33). To test the hypothesis that

cardiomyocytes undergo necroptosis following exposure to *E. faecalis*, we infected HL-1

cardiomyocytes with this bacterium in the presence of various inhibitors of necroptosis and

observed a significant suppression of cell death (Fig. 3C). Additionally, staining for markers of

necroptosis (34), such as mixed lineage kinase domain-like pseudokinase (MLKL) and

phosphorylated MLKL, on cardiac tissue sections obtained from infected mice showed

concentrated signals within the microlesions (Fig 3D). Despite the ability of *E. faecalis* to promote

death of the cardiomyocytes following exposure, antibiotic protection assays indicated that the

bacteria were unable to invade cardiomyocytes relative to endothelial cells (Fig. 3A). No difference

was observed in either the ability to promote necroptosis or cell invasion following exposure to

ΔdsbA relative to wild-type (Figure S3). Taken together, these results suggest that during

bacteremia, *E. faecalis* is able to translocate across the vascular endothelium in order to enter

the myocardial tissue. Once there, *E. faecalis* is unable to invade cardiomyocytes, but can form

microcolonies that promote necroptotic cell death.

Cardiomyocyte inflammatory cytokine response is suppressed following exposure to *E.

faecalis.** Cardiac microlesions are characterized by bacteria filled vacuoles that are largely devoid

of immune cell infiltrate (20, 30). Therefore, this pathology was distinguished from myocardial

abscesses, commonly caused by *Staphylococcus aureus*, which are suppurative infections

characterized by a robust immune cell infiltrate within the cardiac tissue (20, 35). Previous efforts

to explain the lack of an immune response to pneumococcal microlesions have yielded interesting

insights. For example, *S. pneumoniae* within microlesions are thought to subvert the host

response through a biofilm-mediated mechanism involving the pneumolysin mediated killing of

resident macrophages, thus creating a more permissive environment for pneumococcal growth

and survival (22). There are also examples of *E. faecalis* subverting the immune response. In

previous work, it was shown that *E. faecalis* can survive within mouse peritoneal macrophages

for extended periods of time enabling the bacteria to replicate and spread (36). In more recent

work, *E. faecalis* was shown to activate Phosphatidylinositol 3-Kinase signaling that inhibits

apoptosis (37). *E. faecalis* can also resist phagosome acidification and autophagy within

macrophages (38). Because *E. faecalis* does not encode a pneumolysin homolog, nor does it

secrete toxins known to kill macrophages or other immune cells, we wanted to examine the

immune response to *E. faecalis* within the heart by measuring various inflammatory cytokines and

chemokines. Experiments assessing cardiomyocyte inflammatory chemokine and cytokine levels

showed that wild-type *E. faecalis* promotes an immune quiescent response, while loss of *dsbA*

leads to a more inflammatory immune response (Fig. 4A). Interestingly, EGF, M1P-1a, GM-CSF,

IL-6, G-CSF, and INFr were significantly upregulated during infection with *ΔdsbA* relative to the

wild-type infected cardiomyocytes. Thus, these results implicate the contribution of specific host

signaling pathways involved in pathogen clearance that are normally suppressed in the presence

of DsbA during infection, as modeled in Figure 4B. Collectively, these data suggest that *E. faecalis*

inhibits a vigorous cardiomyocyte immune response in a manner similar to that of *S. pneumoniae*

despite lacking the components that *S. pneumoniae* uses during immune subversion. Future

studies will be focused on understanding the cardiomyocyte immune response during

enterococcal infection, and how DsbA contributes.

Discussion

In this work, we demonstrated for the first time that cardiac microlesions form during severe *E. faecalis* bacteremic infection of mice. Importantly, microlesions occur despite *E. faecalis* not encoding homologs of the pneumococcal virulence determinants implicated in microlesion formation. These data suggest that *E. faecalis* may be expressing functionally redundant proteins that serve to promote similar interactions to *S. pneumoniae* in mediating microlesion formation, such as CbpA and pneumolysin, which facilitate vascular endothelial cell translocation and cardiomyocyte cell death, respectively. Indeed, while *E. faecalis* does not encode a CbpA homolog, it does encode other proteins capable of binding host extracellular matrix proteins. These include EfbA, which binds fibronectin (39), ACE, which binds laminin receptor (40), and Ebp pili that bind fibrinogen (41).

Interestingly, we did discover one factor, DsbA, that contributes to microlesion formation and inhibition of the cardiomyocyte inflammatory immune response, as modeled in Figure 4B. Dsb proteins work to catalyze disulfide bond formation in proteins and have been best studied in Gram-negative bacteria where they catalyze the oxidative folding of proteins in the periplasm (42, 43). In Gram-positive bacteria, oxidative protein folding of proteins targeted for the cell envelope or secretion is thought to occur in the exoplasm, the area immediately outside of the cytoplasmic membrane but before the peptidoglycan layer (42). While Dsb proteins and oxidative protein folding have established importance in the biology of certain Gram-positive bacteria, like the Actinomyces, Dsb proteins are not well studied in Firmicutes like *E. faecalis* and have generally not been considered important (42). Most of these species secrete few proteins containing cysteines and some, notably *S. pneumoniae*, do not encode any thiol-disulfide oxidoreductases (44, 45). A deletion of the only DsbA-like protein in *S. aureus* resulted in no significant phenotypes

including no effects on pathogenicity (46). However, the discovery in this work that loss of DsbA

in *E. faecalis* profoundly affected pathogenicity in the *C. elegans* model and reduced cardiac

microlesion formation in the mouse sepsis model suggests that the lack of Dsb-related

phenotypes in *S. aureus* does not necessarily apply to all Firmicutes. It will be of interest to further

investigate the role of Dsb proteins in these species.

Because of the well-known role of Dsbs in the folding of secreted and cell envelope associated

proteins, we hypothesize that the effects on pathogenicity is the result of disruption in the stability

of unknown DsbA substrate(s). Our prior work established EntV as a DsbA substrate, currently

the only one identified (28). However, loss of EntV did not reduce pathogenicity in the worm model

(Figure 1A), and exposure to EntV, even at very high concentrations, was non-toxic to mammalian

cells (29). Thus, we do not favor EntV as being the source of DsbA's effects. A primary future

direction will be identifying additional substrates of DsbA and determining which contribute to

microlesion formation. Not only will this provide insight into the developmental requirements for

this cardiac pathology that promotes cardiomyocyte necroptosis, it will also impact understanding

of *E. faecalis* pathogenesis and the biology of disulfide bond forming proteins in bacteria.

Pneumococcal ChoP interactions with host PAFr were determined to be a prerequisite for

vascular endothelial cell invasion during IPD. In *S. pneumoniae*, the main wall teichoic acid chain

repeat is composed of 2-acetamido-4-amino-2,4,6-trideoxygalactose, glucose, ribitol-phosphate

and two GalNAc moieties, each decorated with phosphorylcholine. This cell wall teichoic acid

structure is considered rare and is thought to be exclusive to *S. pneumoniae* (47, 48). In light of

our observations of *E. faecalis* invasion of vascular endothelial cells despite an absence of this

wall teichoic acid structure, it is possible that another host receptor is required for endothelial cell

invasion. Interestingly, recent studies focused on bacterial cell wall entry and signaling in

eukaryotic cells have identified a PAFr-independent mechanism of tissue invasion that is actin-

dependent and engenders Rac1, Cdc42, and phosphatidylinositol 3-kinase (PI3K) signaling

leading to cell wall internalization via a macropinoctytosis-like pathway (49). Whether such a

pathway contributes to microlesion formation is unknown. In the future, it will be of interest to study

the mechanisms of tissue invasion and the host signaling pathways involved.

Cardiac microlesion formation is becoming increasingly appreciated as a risk factor during acute

infection (25), and this study establishes that *E. faecalis* is another etiological agent capable of

causing this pathology in a vertebrate animal model. There is an increased occurrence of heart

failure observed in patients with endocarditis (13-19), and we hypothesize that invasive infection

resulting in microlesions additionally occurs and is a contributing factor. It will be important to

investigate whether *E. faecalis* microlesions occur in humans and whether they contribute to

adverse cardiac events. As nosocomial derived enterococcal infections are increasingly

problematic among the elderly and immunocompromised patients (8-12), the development of

successful treatments will require identifying all disease manifestations as well as the bacterial

and the host factors involved.

317
318
319

Methods and Materials

C. elegans **survival assay**

E. faecalis strains were tested in *C. elegans* for survival as previously described (50). Briefly, *C. elegans* strain N2 was maintained and propagated on *Escherichia coli* strain OP50 with standard techniques. 30 *C. elegans* L4 nematodes were transferred from a lawn of *Escherichia coli* OP50 to a lawn of *E. faecalis* to be tested, incubated at 25°C, and examined at ≈24-hr intervals with a dissecting microscope for viability. Worms were considered dead when they did not respond to touch with a platinum wire pick. The data is representative of 3 independent experiments. Nematode survival was plotted with the log-rank (Mantel-Cox) method using Prism 7 statistical software (GraphPad).

Bacterial strains and growth analysis

All *E. faecalis* strains used in this project were used in previously published studies (28) and are listed in Table S1. In a 96 well plate, overnight cultures (grown ~15 hours) of OG1RF strains were diluted to an OD_{600nm} of 0.05 in brain heart infusion (BHI) agar medium (Difco) in a volume of 200µl / well in a 96-well plate. The 96-well plate containing the diluted cultures were then placed in a prewarmed 37ºC BioTek Cytation 5 plate reader, and OD_{600nm} measurements were taken every 10 minutes for 15 hours. Prior to measurement, the plate was gently agitated for 5 seconds.

Mouse peritonitis and cardiac microlesion model

The mouse peritonitis model was performed as described previously (51). The method for identifying and evaluating cardiac microlesion formation was conducted as described previously (20). Briefly, female, ~4 – 6-week old (~25g), ICR outbred mice were obtained from Envigo (Indianapolis, Indiana) and were infected intraperitoneally with 7×10^8 CFUs of *E. faecalis*. Mouse

hearts were obtained 24-48 hours post-infection, formalin fixed and sent to The University of

Texas Health Science Center Histopathology Laboratory for paraffin embedding, sectioning, and

hematoxylin and eosin H&E) staining. Paraffin embedded mouse hearts were first bisected and

sectioned longitudinally. 20 sections were cut per block, and every 5th section H&E stained in

order to measure the number and relative distribution of cardiac microlesions. Cardiac

microlesions were averaged across sections, and the differences between mouse groups

compared using *Student's* two-tailed t-test. Microlesions size measurements were also obtained

by measuring cardiac microlesion size on each section, and then averaging the values across

sections. ? Blinded reading? If so, might be worthwhile mentioning

Necroptosis inhibition assay

HL-1 cardiomyocytes were seeded in 24-well plates at a concentration of 5×10^5 cells/ml. 24 hours

later, the cells were washed three times with Dulbecco's (1X) PBS, followed by pretreatment with

antibiotic free Claycomb medium supplemented with 10 µM of Necrostatin-1 (Sigma Aldrich),

GSK'872 (RD Systems), and Necrosulfonamaide (Millipore-Sigma) for 1 hour, followed by

infection at an MOI of 10 for 4 hours, following which the supernatant was collected, and

cytotoxicity in each well accessed using the CytoTox-ONE™ Homogeneous Membrane Integrity

Assay Kit (Promega). Each experiment was conducted in triplicate, on 3 separate occasions, and

averaged values between groups compared using using *Student's* two-tailed t-test.

Annexin V apoptosis and necrosis assay.

The annexin V apoptosis and necrosis assay was conducted using RealTime-Glo™ Annexin V

Apoptosis and Necrosis Assay Kit (Promega). The kit contains near-equimolar ratios of two

annexin V fusion proteins (Annexin V-LgBiT and Annexin V-SmBit) containing complementary

subunits of NanoBiT® Luciferase that, when combined following exposure to phosphatidyl serine,

results in an emission of a luminescent signal. The kit also contains a cell-impermeant

profluorescent DNA dye, which detects necrosis. Briefly, HL-1 cardiomyocytes (Sigma-Aldrich)

were seeded in a 96 well plate. 24 hours later, the cells were infected with *E. faecalis* at an MOI

= 50, in the presence of kit reagents. Luminescent and fluorescent readings were obtained every

30 minutes for up to 6 hours.

Invasion assay

SVEC4-10 Endothelial Cells (ATCC) were seeded in 24 well plates at a cell density of 1.5×10^5

cells/ml. After 24 hours, the cells were washed three times with PBS, and infected with *E. faecalis*

at a MOI of 10 for 4 hours in antibiotic free Claycomb medium. After the incubation, the cells were

washed with Dulbecco's (1X) PBS three times and treated with trypsin, diluted in PBS, and plated

on BHI agar plates for attachment quantification. To measure intracellular numbers of *E. faecalis*,

cells that were washed three times with Dulbecco's (1X) PBS 4 h.p.i., then treated with Dulbecco's

Modified Eagle's Medium (ATCC) supplemented with 15 µg/ml vancomycin and 150 µg/ml

gentamicin. 24 hours later, the wells were washed three times with Dulbecco's (1X) PBS, lysed

using 0.1% Triton-X-100, and diluted and plated on BHI agar plates for intracellular quantification.

The ratio of invasion to attachment for a given enterococcal strain was used to calculate the

percent of invasion. The experiments were conducted on 3 independent occasions and were

performed using triplicate wells for each strain. Attachment and invasion values were averaged

from each experiment, and statistically analyzed using *Student's* two-tailed t-test.

Cytotoxicity and inflammatory cytokine panel microtiter plate reader assay

HL-1 cardiomyocytes were obtained from Sigma-Aldrich and SVEC4-10 Endothelial Cells were

obtained from ATCC. HL-1 cardiomyocytes (Sigma-Aldrich) were seeded into 24-well plates at a

concentration of 5×10^5 cells/ml. 24 hours later, the cells were washed three times with Dulbecco's

(1X) PBS, and infected with *E. faecalis* at an MOI of 10 for 4 hours in antibiotic free Claycomb

medium. Supernatants were collected and centrifuged for 5 minutes at 10,000 G. Supernatant

was collected, and cytotoxicity assayed using CytoTox-ONE™ Homogeneous Membrane

Integrity Assay Kit (Promega). In addition to measuring cytotoxicity in HL-1 cardiomyocytes

infected with *E. faecalis* 4 h.p.i, inflammatory chemokines and cytokines were analyzed using a

Mouse Cytokine ELISA Plate Array III Colorimetric Assay (Signosis). The data represent an

average of at least 3 independent experiments for each isogenic mutant tested and were analyzed

using *Student's* two-tailed t-test.

Immunofluorescence microscopy

Immunofluorescence microcopy was conducted as described previously (20). Briefly, paraffin

embedded sections were treated with 1:100 rabbit Anti-mouse MLKL (phospho S345) antibody

[EPR9515(2)] in order to detect phosphorylated MLKL (Abcam), and 1:100 rat anti-mouse MLKL

antibody (Millipore-Sigma) for MLKL detection followed by treatment with 1:1000 chicken anti-

rabbit IgG (H+L) cross-adsorbed secondary Alexa Fluor 594 antibody (ThermoFisher), and 1:1000

goat anti-rat IgG (H+L) Secondary FITC Antibody (ThermoFisher) respectively. A rabbit anti-

Enterococcus faecalis antibody (Abcam) was used at a concentration of 1:1000 to detect *E.*

faecalis within cardiac microlesions followed by treatment with 1:1000 goat anti-rabbit IgG (H&L),

F(ab)'2 Fragment-affinity purified fluorescein (FITC) conjugate antibody (Invitrogen). All stained

sections were counterstained with 1:1000 DAPI (ThermoFisher) and mounted with Hardset™

Antifade Mounting Medium. Visualization of fluorescence was obtained using standard excitation

and emission wavelengths for Alexa 594, FITC, and DAPI on an Olympus FV3000RS high-speed

high-sensitivity inverted laser scanning confocal microscope.

Generation of rabbit anti-DsbA antisera

His-tagged recombinant DsbA was overexpressed in Rosetta-2(DE3) *E. coli* (Sigma-Aldrich) and

isolated by affinity purification via HisPur Nickel-NTA Resin (ThermoFisher) packed columns

followed by further purification and buffer exchange using Fast Protein Liquid Chromatography

(FPLC) to obtain highly pure samples of recombinant DsbA. The recovered recombinant protein

was then shipped to Cocalico Biologicals (Reamstown, PA) for generation of anti-DsbA antisera

through the immunization of rabbits. Throughout the immunization schedule, regular test bleeds

were performed and tested by Western blot using *E. faecalis* whole cell lysates and recombinant

DsbA in order to ensure the specificity of the rabbit anti-DsbA antisera.

***E. faecalis* whole cell lysate and Western blot**

E. faecalis was grown overnight in 5 ml of BHI. The following day (~16 hours later), the media

was centrifuged at 6,738 G for 10 min at 4ºC. The pellet was resuspended in 5 ml of BHI, and

diluted 10X into fresh BHI media. The culture was subsequently incubated at 37ºC for an

additional 3.5 hours to obtain a culture of *E. faecalis* in exponential growth. The culture was then

spun down again at 6,738 G and supernatant removed. The pellet was then resuspended in 1ml

of ZAP buffer (10 mM Tris, 200 mM NaCl, pH 7.5) containing 1X protease inhibitor cocktail

(Roche) sonicated on ice with a max amplitude of 30%, and alternating on/off cycles consisting of

1 minute on, and 10 seconds off for a total of 5 minutes. Following sonication, the sample was

centrifuged at 10,000 G for 5 minutes and supernatant collected. The amount of protein in the

supernatant was quantified using a DC protein modified Bradford assay (BIORAD), and 10µg of

lysate was loaded in each lane of a 12% SDS-PAGE gel. Following electrophoresis and transfer

of protein to a PVDF membrane, the membrane was incubated overnight at 4ºC in a 5% milk

solution containing a 1:1000 dilution of rabbit anti-DsbA antisera followed by incubation with a

Goat anti-rabbit IgG HRP (ThermoFisher) secondary antibody at a dilution of 1:10,000 at room

temperature for 30 minutes prior to exposure.

445
446
447 **Acknowledgements**

448 This work was supported by the National Institute of Dental and Craniofacial Research of the

449 National Institutes of Health under award numbers R01DE027608 to D. A. G. and F32DE027580

450 to A. O. B.

451
452
453
454 **Table S1. Strains used in this study**

Enterococcus faecalis Strains				
Strain	**Genotype**	**Characteristics**	**Parent**	**Source**
OG1RF	Wild-type	Gel$^+$ Spr$^+$ Rf^R Fa^R	OG1	(52)
AOBD1	*ΔdsbA*	Gel$^+$ Spr$^+$ Rf^R Fa^R	OG1	(28)
AOBD2	*ΔdsbA::dsbA*	Gel$^+$ Spr$^+$ Rf^R Fa^R	OG1	(28)

493
494
Figure 1. *dsbA* **deficient** *E. faecalis* **show reduced virulence in a** *C. elegans* **survival**
assay. **(A)** L4 stage *C. elegans* were infected with *E. faecalis*. This figure is representative
of 3 independent experiments with 30 worms/group. ****Asterisk indicate a *P* < 0.0001 and
were analyzed by the log-rank (Mantel-Cox). **(B)** Western blot performed by SDS-PAGE
separation of *E. faecalis* whole cell lysates followed by hybridization of DsbA using rabbit
anti-DsbA antisera. Image is representative of 3 independent Western blot experiments.

Figure 2. **Cardiac microlesions form during invasive *E. faecalis* infection and are attenuated by loss of *dsbA***. Cardiac microlesions were observed using light microscopy of hematoxylin and eosin (H&E) stained paraffin embedded cardiac sections at 40X magnification **(A)**, and 100X magnification **(B)**. Immunofluorescent detection of *E. faecalis* was imaged using a 20X objective at 3X zoom (total of 60X magnification), a FITC conjugated anti-enterococcal antibody **(C)**, and DAPI for DNA staining **(D)**. Total microlesion counts **(E)**, and size measurements **(F)**, were obtained by quantification using light microscopy of H&E stained paraffin embedded cardiac sections from mice infected for 24-48 hours with *E. faecalis* (OG1RF n=14, Δ*dsbA* n=14*,* Δ*dsbA::dsbA* n=14). The data were averaged from 3 independent experiments, and analyzed using *Student's* two-tailed t-test (*** indicate a *P* < 0.0001).

Figure 3. Necroptosis contributes to cardiotoxicity within cardiac microlesions. **(A)** Comparison of OG1RF invasion rates into HL-1 cardiomyocytes and SVEC endothelial cells. The graph represents data from 3 independent experiments that were analyzed using *Student's* two-tailed t-test (** indicate a *P* < 0.01). **(B)** *E. faecalis* induced cell death in HL-1 cardiomyocytes 4 hours post-infection (h.p.i.) as determined by measuring lactate dehydrogenase. The data represent an average of 3 independent experiments and were analyzed using *Student's* two-tailed t-test (* indicate a *P* < 0.05). **(C)** *E. faecalis* induced cell death in HL-1 cardiomyocytes 4 h.p.i. and was inhibited in the presence of inhibitors of necroptosis. The data represent an average of 3 independent experiments and were analyzed using *Student's* two-tailed t-test (**** indicate a *P* < 0.0001). **(D)** Immunofluorescent microscopy of paraffin embedded cardiac cross sections at 20X were performed to detect MLKL using FITC conjugated anti-MLKL antibodies, and Alexa 594 conjugated anti-phosphorylated MLKL antibodies at sites of microlesion formation, while DAPI was used to stain nuclei.

Figure 4. Wild-type *E. faecalis* suppresses the cardiomyocyte inflammatory response *in-vitro*. (A) HL-1 cardiomyocytes exposed to *E. faecalis* promote a suppressed inflammatory response while Δ*dsbA* infected cardiomyocytes yield a more inflammatory response. Data was obtained from the fold-expression change analysis (relative to uninfected controls) of at least 3 independent experiments for each strain. **(B)** Model of *E. faecalis* mediated suppression of the host cardiomyocyte immune response and effect on microlesion size.

Supplemental Figure 1. *E. faecalis* **growth curve.** *E. faecalis* strains were grown for 15 hours in BHI and absorbance (OD_{600nm}) measured every 10 minutes. This figure is representative of 3 independent experiments.

Supplemental Figure 2. Cardiomyocytes infected with *E. faecalis* die through an apoptosis independent mechanism. HL-1 cardiomyocytes were infected with *E. faecalis* at an MOI=50 for 4 hours in the presence of annexin V fusion proteins containing complimentary subunits of NanoBiT® Luciferase combined with a cell-impermeant profluorescent DNA dye to detect necrosis. A concomitant increase in both luminescent (apoptosis) **(A)** and fluorescent (necrosis) **(B)** signals are indicative of cell death through an apoptosis independent mechanism. The graphs represent an average of 3 independent experiments.

Supplemental Figure 3. Loss of *dsbA* does has no effect on endothelial cell invasion or necroptosis induced cell death of cardiomyocytes. (A) Comparison of WT and *ΔdsbA* invasion rates into HL-1 cardiomyocytes and SVEC endothelial cells showed no significant difference. The graph represents data from 3 independent experiments analyzed using *Student's* two-tailed t-test (NS denotes no significance). **(B)** Comparison of WT and *ΔdsbA* in HL-1 cardiomyocytes 4 h.p.i. in the presence of inhibitors of necroptosis *in-vitro* was determined by measuring lactate dehydrogenase. No significant difference was observed between necroptosis inhibitor conditions tested. The data represent an average of 3 independent experiments and were analyzed using *Student's* two-tailed t-test (* indicate a *P* < 0.05).

References

1. K. L. Frank *et al.*, Use of recombinase-based in vivo expression technology to characterize Enterococcus faecalis gene expression during infection identifies in vivo-expressed antisense RNAs and implicates the protease Eep in pathogenesis. *Infect Immun* **80**, 539-549 (2012).

2. D. A. Garsin *et al.*, "Pathogenesis and Models of Enterococcal Infection" in Enterococci: From Commensals to Leading Causes of Drug Resistant Infection*,* M. S. Gilmore, D. B. Clewell, Y. Ike, N. Shankar, Eds. (Boston, 2014).

3. F. Lebreton, R. J. L. Willems, M. S. Gilmore, "Enterococcus Diversity, Origins in Nature, and Gut Colonization" in Enterococci: From Commensals to Leading Causes of Drug Resistant Infection*,* M. S. Gilmore, D. B. Clewell, Y. Ike, N. Shankar, Eds. (Boston, 2014).

4. H. S. Gold, Vancomycin-resistant enterococci: mechanisms and clinical observations. *Clin Infect Dis* **33**, 210-219 (2001).

5. N. I. Agudelo Higuita, M. M. Huycke, "Enterococcal Disease, Epidemiology, and Implications for Treatment" in Enterococci: From Commensals to Leading Causes of Drug Resistant Infection*,* M. S. Gilmore, D. B. Clewell, Y. Ike, N. Shankar, Eds. (Boston, 2014).

6. M. Armstrong-Evans *et al.*, Control of transmission of vancomycin-resistant Enterococcus faecium in a long-term-care facility. *Infect Control Hosp Epidemiol* **20**, 312-317 (1999).

7. H. M. S. Goh, M. H. A. Yong, K. K. L. Chong, K. A. Kline, Model systems for the study of Enterococcal colonization and infection. *Virulence* **8**, 1525-1562 (2017).

8. L. M. Weiner *et al.*, Antimicrobial-Resistant Pathogens Associated With Healthcare-Associated Infections: Summary of Data Reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2011-2014. *Infect Control Hosp Epidemiol* **37**, 1288-1301 (2016).

703   9.   K. C. Malani PN, Zervis MJ, *Enterococcal disease, epidemiology and treatment*, The

704   Enterococci: Pathogenesis, Molecular Biology and Antibiotic Resistance (ASM Press,

705   Washington, D.C., 2002).

706   10.   H. Wisplinghoff *et al.*, Nosocomial bloodstream infections in US hospitals: analysis of

707   24,179 cases from a prospective nationwide surveillance study. *Clin Infect Dis* **39**, 309-317

708   (2004).

709   11.   C. Chirouze *et al.*, Enterococcal endocarditis in the beginning of the 21st century: analysis

710   from the International Collaboration on Endocarditis-Prospective Cohort Study. *Clin*

711   *Microbiol Infect* **19**, 1140-1147 (2013).

712   12.   J. R. McDonald *et al.*, Enterococcal endocarditis: 107 cases from the international

713   collaboration on endocarditis merged database. *Am J Med* **118**, 759-766 (2005).

714   13.   C. H. Cabell *et al.*, Changing patient characteristics and the effect on mortality in

715   endocarditis. *Arch Intern Med* **162**, 90-94 (2002).

716   14.   G. J. Garvey, H. C. Neu, Infective endocarditis--an evolving disease. A review of

717   endocarditis at the Columbia-Presbyterian Medical Center, 1968-1973. *Medicine*

718   *(Baltimore)* **57**, 105-127 (1978).

719   15.   J. E. Geraci, W. J. Martin, Antibiotic therapy of bacterial endocarditis. VI. Subacute

720   enterococcal endocarditis; clinical, pathologic and therapeutic consideration of 33 cases.

721   *Circulation* **10**, 173-194 (1954).

722   16.   E. Jawetz, M. Sonne, Penicillin-streptomycin treatment of enterococcal endocarditis. A re-

723   evaluation. *N Engl J Med* **274**, 710-715 (1966).

724   17.   P. I. Lerner, L. Weinstein, Infective endocarditis in the antibiotic era. *N Engl J Med* **274**,

725   388-393 concl (1966).

726   18.   L. Olaison, K. Schadewitz, E. Swedish Society of Infectious Diseases Quality Assurance

727   Study Group for, Enterococcal endocarditis in Sweden, 1995-1999: can shorter therapy

728   with aminoglycosides be used? *Clin Infect Dis* **34**, 159-166 (2002).

729   19.   L. L. Pelletier, Jr., R. G. Petersdorf, Infective endocarditis: a review of 125 cases from the

730   University of Washington Hospitals, 1963-72. *Medicine (Baltimore)* **56**, 287-313 (1977).

731   20.   A. O. Brown *et al.*, Streptococcus pneumoniae translocates into the myocardium and forms

732   unique microlesions that disrupt cardiac function. *PLoS Pathog* **10**, e1004383 (2014).

733   21.   L. F. Reyes *et al.*, Severe Pneumococcal Pneumonia Causes Acute Cardiac Toxicity and

734   Subsequent Cardiac Remodeling. *Am J Respir Crit Care Med* **196**, 609-620 (2017).

735   22.   A. T. Shenoy *et al.*, Streptococcus pneumoniae in the heart subvert the host response

736   through biofilm-mediated resident macrophage killing. *PLoS Pathog* **13**, e1006582 (2017).

737   23.   T. Brissac, A. T. Shenoy, L. A. Patterson, C. J. Orihuela, Cell Invasion and Pyruvate

738   Oxidase-Derived H2O2 Are Critical for Streptococcus pneumoniae-Mediated

739   Cardiomyocyte Killing. *Infect Immun* **86** (2018).

740   24.   A. T. Shenoy, C. J. Orihuela, Anatomical site-specific contributions of pneumococcal

741   virulence determinants. *Pneumonia (Nathan)* **8** (2016).

742   25.   D. M. Musher, M. S. Abers, V. F. Corrales-Medina, Acute Infection and Myocardial

743   Infarction. *N Engl J Med* **380**, 171-176 (2019).

744   26.   V. F. Corrales-Medina *et al.*, Association between hospitalization for pneumonia and

745   subsequent risk of cardiovascular disease. *JAMA* **313**, 264-274 (2015).

746   27.   S. R. Jafarzadeh, B. S. Thomas, D. K. Warren, J. Gill, V. J. Fraser, Longitudinal Study of

747   the Effects of Bacteremia and Sepsis on 5-year Risk of Cardiovascular Events. *Clin Infect*

748   *Dis* **63**, 495-500 (2016).

28. C. E. G. Armand O. Brown, Melissa R. Cruz, Michael C. Lorenz, Danielle A. Garsin,

Antifungal activity of the Enterococcus faecalis peptide EntV requires protease cleavage

and disulfide bond formation. *Mbio* **10**, 1-13 (2019).

29. C. E. Graham, M. R. Cruz, D. A. Garsin, M. C. Lorenz, Enterococcus faecalis bacteriocin

EntV inhibits hyphal morphogenesis, biofilm formation, and virulence of Candida albicans.

Proc Natl Acad Sci U S A **114**, 4507-4512 (2017).

30. R. P. Gilley *et al.*, Infiltrated Macrophages Die of Pneumolysin-Mediated Necroptosis

following Pneumococcal Myocardial Invasion. *Infect Immun* **84**, 1457-1469 (2016).

31. T. Zhang *et al.*, CaMKII is a RIP3 substrate mediating ischemia- and oxidative stress-

induced myocardial necroptosis. *Nat Med* **22**, 175-182 (2016).

32. A. Linkermann *et al.*, Necroptosis in immunity and ischemia-reperfusion injury. *Am J

Transplant* **13**, 2797-2804 (2013).

33. A. Linkermann *et al.*, Two independent pathways of regulated necrosis mediate ischemia-

reperfusion injury. *Proc Natl Acad Sci U S A* **110**, 12024-12029 (2013).

34. K. Kitur *et al.*, Necroptosis Promotes Staphylococcus aureus Clearance by Inhibiting

Excessive Inflammatory Signaling. *Cell Rep* **16**, 2219-2230 (2016).

35. F. J. Ramos Tuarez, M. A. Law, "Cardiac Abscess" in StatPearls. (Treasure Island (FL),

2019).

36. C. R. Gentry-Weeks, R. Karkhoff-Schweizer, A. Pikis, M. Estay, J. M. Keith, Survival of

Enterococcus faecalis in mouse peritoneal macrophages. *Infect Immun* **67**, 2160-2165

(1999).

37. J. Zou, N. Shankar, Enterococcus faecalis infection activates phosphatidylinositol 3-kinase

signaling to block apoptotic cell death in macrophages. *Infect Immun* **82**, 5132-5142

(2014).

38.   J. Zou, N. Shankar, The opportunistic pathogen Enterococcus faecalis resists phagosome

774   acidification and autophagy to promote intracellular survival in macrophages. *Cell Microbiol*

775   **18**, 831-843 (2016).

776   39.   K. V. Singh, S. L. La Rosa, S. R. Somarajan, J. H. Roh, B. E. Murray, The fibronectin-

777   binding protein EfbA contributes to pathogenesis and protects against infective

778   endocarditis caused by Enterococcus faecalis. *Infect Immun* **83**, 4487-4494 (2015).

779   40.   S. R. Nallapareddy, X. Qin, G. M. Weinstock, M. Hook, B. E. Murray, Enterococcus faecalis

780   adhesin, ace, mediates attachment to extracellular matrix proteins collagen type IV and

781   laminin as well as collagen type I. *Infect Immun* **68**, 5218-5224 (2000).

782   41.   S. R. Nallapareddy *et al.*, Endocarditis and biofilm-associated pili of Enterococcus faecalis.

783   *J Clin Invest* **116**, 2799-2807 (2006).

784   42.   M. E. Reardon-Robinson, H. Ton-That, Disulfide-Bond-Forming Pathways in Gram-

785   Positive Bacteria. *J Bacteriol* **198**, 746-754 (2015).

786   43.   C. Landeta, D. Boyd, J. Beckwith, Disulfide bond formation in prokaryotes. *Nat Microbiol*

787   **3**, 270-280 (2018).

788   44.   R. Daniels *et al.*, Disulfide bond formation and cysteine exclusion in gram-positive bacteria.

789   *J Biol Chem* **285**, 3300-3309 (2010).

790   45.   R. J. Dutton, D. Boyd, M. Berkmen, J. Beckwith, Bacterial species exhibit diversity in their

791   mechanisms and capacity for protein disulfide bond formation. *Proc Natl Acad Sci U S A*

792   **105**, 11933-11938 (2008).

793   46.   A. Dumoulin, U. Grauschopf, M. Bischoff, L. Thony-Meyer, B. Berger-Bachi,

794   Staphylococcus aureus DsbA is a membrane-bound lipoprotein with thiol-disulfide

795   oxidoreductase activity. *Arch Microbiol* **184**, 117-128 (2005).

47.   D. Denapaite, R. Bruckner, R. Hakenbeck, W. Vollmer, Biosynthesis of teichoic acids in
Streptococcus pneumoniae and closely related species: lessons from genomes. *Microb*
Drug Resist **18**, 344-358 (2012).

48.   W. Fischer, T. Behr, R. Hartmann, J. Peter-Katalinic, H. Egge, Teichoic acid and
lipoteichoic acid of Streptococcus pneumoniae possess identical chain structures. A
reinvestigation of teichoid acid (C polysaccharide). *Eur J Biochem* **215**, 851-857 (1993).

49.   L. N. Loh, G. Gao, E. I. Tuomanen, Dissecting Bacterial Cell Wall Entry and Signaling in
Eukaryotic Cells: an Actin-Dependent Pathway Parallels Platelet-Activating Factor
Receptor-Mediated Endocytosis. *MBio* **8** (2017).

50.   D. A. Garsin *et al.*, A simple model host for identifying Gram-positive virulence factors. *Proc*
Natl Acad Sci U S A **98**, 10892-10897 (2001).

51.   K. V. Singh, X. Qin, G. M. Weinstock, B. E. Murray, Generation and testing of mutants of
Enterococcus faecalis in a mouse peritonitis model. *J Infect Dis* **178**, 1416-1420 (1998).

52.   B. E. Murray *et al.*, Generation of restriction map of Enterococcus faecalis OG1 and
investigation of growth requirements and regions encoding biosynthetic function. *J*
Bacteriol **175**, 5216-5223 (1993).