1 Bioluminescence dynamics in single germinating bacterial spores reveal metabolic heterogeneity

2 Zak Frentz and Jonathan Dworkin*

3 Department of Microbiology and Immunology, College of Physicians and Surgeons, Columbia

4 University, New York, NY 10032

6 * Corresponding author,

Abstract

Spore-forming bacteria modulate their metabolic rate by over 5 orders of magnitude as they transition between dormant spores and vegetative cells, and thus represent an extreme case of phenotypic variation. During environmental changes in nutrient availability, clonal populations of spore-forming bacteria exhibit individual differences in cell fate, timing of phenotypic transitions, and gene expression. One potential source of this variability is metabolic heterogeneity, but this has not yet been measured, as existing single-cell methods are not easily applicable to spores due to their small size and strong autofluorescence. Here, we use the bacterial bioluminescence system and a highly sensitive microscope to measure metabolic dynamics in thousands of *B. subtilis* spores as they germinate. We observe and quantitate large variations in the bioluminescence dynamics across individual spores that can be decomposed into contributions from variability in germination timing, the amount of endogenously produced luminescence substrate, and the intracellular reducing power. This work shows that quantitative measurement of spore metabolism is possible and thus it opens venues for future study of the thermodynamic nature of dormant states.

Introduction

Some bacteria respond to nutrient depletion by differentiating from growing, vegetative cells into spores that can survive extreme temperatures, radiation, and long periods without nutrients, while retaining the ability to germinate and reinitiate growth in the presence of sufficient nutrients [1-3]. One of the most remarkable aspects of spore-forming bacteria is the extreme reduction of their metabolic rate, by at least 5 orders of magnitude, during sporulation [4]. Metabolic dormancy and resistance allow spores to survive for long times in harsh environments, which typically fluctuate. Dynamically sensing the environment requires energy expenditure [5], and thus is likely difficult for spores given their low or nonexistent metabolism. Therefore, sporulating populations must adapt to fluctuating environments through other strategies, such as phenotypic diversification, or "bet-hedging" [6, 7].

Significant heterogeneity is observed in populations during both the sporulation and germination transitions. For example, after nutrient limitation, only a portion of vegetative cells commit to sporulation, while other cells either lyse or survive in the vegetative state by using secondary metabolites [8]. The time at which a vegetative cell sporulates varies significantly within a population, and this time correlates with the time at which the ensuing spore germinates and in its ability to resume growth in various nutrients [9]. Spores germinate in reponse to low concentrations of nutrients at variable times [7], and this variability is partially explained by the number of germinant receptors in each spore [10]. Even in the absence of germinants, a small fraction of spores germinates spontaneously [11, 12]. These phenomena of phenotypic heterogeneity have been linked to variations in gene expression [8, 9, 12, 13]. However, the remarkable dynamics of metabolism during sporulation and germination have not been measured in individual cells, due to lack of a suitable technique.

The dynamics of energy metabolism can be measured in single vegetative bacteria using several *in vivo* methods, most of which are based on fluorescence [14]. These are difficult to apply to spores due to their strong autofluorescence, primarily from calcium dipicolinate, NADH, and

flavins [15, 16]. The intracellular redox potential drives all pathways of energy metabolism other

than fermentation [17, 18], and has been measured *in vivo* in single bacterial cells using a custom-

fabricated microelectrode coupled to an optical trap [19]. However, this technique requires a

specialized device, operates in anaerobic conditions, and only measures a small number of cells.

An alternative approach uses the bacterial luminescence system [20]. An enzyme,

luciferase, encoded by the *luxAB* genes, catalyzes the light-emitting oxidation of a substrate,

luciferin, which can be supplied exogenously or synthesized endogenously by expression of the

luxCDE genes. The oxidation reaction's rate depends on the reduced form of the electron

transporter flavin mononucleotide ($FMNH_2$), whose concentration is determined by the

intracellular reducing power [21]. Thus, the rate of light production directly relates to the redox

potential. In addition, since the emission of light occurs without external illumination,

bioluminescent techniques can be considered "zero background", and therefore offer improved

sensitivity as compared to fluorescent techniques. Here we describe a microscope designed to

capture the dynamics of bioluminescence emitted by single cells, which we used to non-invasively

assay metabolic activity of single germinating spores of *Bacillus subtilis*.

Results

Bioluminescence microscopy of single spores

To date, the bioluminescence dynamics of sporulating and germinating cells have been measured

only in populations, typically using a photomultiplier tube for photon counting [22, 23]. A *B. subtilis*

strain that expresses the *lux* genes under a sporulation-specific promoter (P_{sspB}) produces spores

that contain a sufficient number of luciferase enzymes to emit detectable light only upon

germination [23]. Prior to germination, spores do not emit detectable light, presumably due to their

dormant metabolic state.

The feasibility of measuring bioluminescence dynamics in single germinating spores is

determined by the rate of photon emission, which must be sufficiently high to be resolved over

measurement noise. This rate increases with higher expression of the *luxABCDE* genes, as has

recently been achieved in *B. subtilis* by modifying the ribosomal binding sites upstream of each

gene [24]. The rate of bioluminescent photon emission at the population level has been reported

in relative light units, or RLUs [24, 25], which are defined differently for various devices. RLUs can

be converted to units of photons per time per cell by calibration with a luminescent standard [26],

although this calibration has not been included in the published results. As a rough guide,

microscopy of individual metabolically active *Escherichia coli*, *Synechoccocus elongatus*, *Vibrio*

fischeri, and *Vibrio harveyi* cells expressing the lux system typically measures ~60 photons min^{-1}

cell^{-1} [27-29]. Our preliminary analysis suggested that a microscope utilizing recent improvements

in CCD technology could resolve bioluminescence signals as dim as 1 photons min^{-1} spore^{-1} (SI

Appendix).

We therefore constructed a custom microscope for measuring the dynamics of

bioluminescence in single spores (Fig. 1A, Materials and Methods). The microscope includes an

optical path for bright field imaging, which is used to localize spores and measure their refractility,

as described below. The setup consists of an inverted microscope with a high-NA 63X objective

and a cooled, back-illuminated CCD camera. We used 10 min exposure times, sufficient to

resolve germination dynamics while achieving sensitivity to permit measurements from single

germinating spores. Over a period of 10 min, Brownian motion displaces a spore by a root-mean-

square displacement of approximately 50 μm. To prevent such displacements, we trapped the

spores by using hydrophobic glass coverslips. With this method, the displacement of spores over

10 min periods was below the optical resolution of the bright field images (<160 nm). The sample

was held in a perfusion chamber, which dynamically controlled the chemical environment (Fig.

1B) , and the temperature at the sample was controlled to ±0.05 ℃ by an air-to-air heat exchanger.

We initially investigated the dynamics of bioluminescence emitted by single *B. subtilis*

spores as they germinated in response to L-alanine. These spores (JDB3780) expressed the

luxABCDE genes under the control of a sporulation-specific promoter. A field of approximately

900 spores was imaged in bright field and luminescence every 10 min for 1.5 h (Fig. 1C). Prior to

the induction of germination, the dormant spores appeared dark, with high contrast, and produced

no resolvable luminescence signal (Figs. 1D and E). L-alanine was added to the perfusion media

at a concentration of 100 μM, and the ensuing germination response was recorded for 15 h.

After an initial lag following the addition of L-alanine, most spores showed a significant

decrease of contrast in bright field, and many produced luminescence well above the noise

background (Figs. 1F and G). The predominant source of measurement noise was sensor read

noise; we found that its contribution could be reduced significantly by applying a temporal low-

pass filter with a cutoff period of 36 min (SI Appendix). Sensor read noise has a Gaussian

distribution, and can therefore produce estimated photon counts that are negative [30]. We

determined that the technique had a sensitivity limit of 0.4 photons min^{-1} spore^{-1} (SI Appendix).

The dynamics of individual spores in the bright field and luminescence images were

analyzed, resulting in time series of bright field contrast and estimated luminescent photon counts

for individual spores (Figs. 1H and I). Rapid loss of contrast after the addition of L-alanine was

evident in most spores, indicating an early event in germination, namely, the release of calcium

dipicolinate from the spore [31]. Spores differed significantly in their time of germination (Fig. 1H)

and in their bioluminescence signals (Fig. 1I).

Bioluminescence dynamics are synchronized with germination time

In order to compare the measurements in the microscope to an established method, we used a

96-well plate reader to measure the population-averaged bioluminescence dynamics of spores

germinating in response to the addition of L-alanine at 5 different concentrations (1 μM, 10 μM,

100 μM, 1 mM, and 10 mM) (Fig. 2A). Time-lapse experiments were performed in the microscope

setup at the same L-alanine concentrations, and the population-averaged bioluminescence

dynamics were calculated (Fig. 2B). The dynamics as measured by both methods shared some

features: Signal intensity generally increased with increasing L-alanine concentration, from total

darkness at 1 μM, to a peak level at 1 mM, then decreased slightly at 10 mM (Fig. 2C). The L-

alanine concentration also affected the temporal patterns of the dynamics: In the measurements

performed with the microscope, a peak in the bioluminescence signal was evident approximately

1 h after germination with 10 mM, 1 mM, and 100 μM L-alanine, but this peak was not observed

following germination with 10 μM L-alanine (Fig. 2B). Although much less pronounced, this early

peak was also apparent in the plate reader measurements after germination with 10 mM and 1

mM L-alanine; with 100 μM and 10 μM L-alanine, no peak was resolved, but a "shoulder" at 40

min was apparent in both conditions (Fig. 2A). The primary difference in the dynamics measured

by the two methods was the much higher intensity of the early peak measured in the microscope.

We observed similar differences caused by varying the spore density in the plate reader

experiments (Fig. S1), which are likely due to chemical interactions between germinating spores.

This effect can explain the differences observed between the dynamics measured in the plate

reader, at a density of 3×10^7 spores ml^{-1}, and in the microscope, at a density of less than 10^5

spores ml^{-1}.

The observed population-averaged bioluminescence dynamics arise from temporal

fluctuations in individual spores as well as variability in the population, and these two sources can

be separated using the single-spore technique. In particular, individual spores germinate at

different times, with large variability observed at low L-alanine concentrations [32]. With

automated image analysis, we identified the time τ_g of rapid contrast loss in each germinating

spore (SI Appendix), and interpreted τ_g as the time at which each spore started germination. We

considered the statistics of germination dynamics by calculating the fraction of the population that

had not yet germinated by time t: This is the tail distribution function of germination time, $P(\tau_g > t)$

(Fig. 3A). At L-alanine concentrations of 100 μM and higher, most spores germinated within 1 h

of induction, and over 99% of spores germinated within 13 h. In contrast, spores germinated much

more slowly in response to the addition of 10 μM L-alanine, with only half of all spores germinating

within 13 h. For the 1 μM L-alanine concentration, we did not observe any germination for spores

up to 82 h after induction.

The distributions of germination time τ_g reveal significant variability in the population of

spores, especially evident in the 10 μM L-alanine condition. To investigate the relationship

between this variability and the bioluminescence dynamics, we considered the bioluminescence

of each cell as a function of time elapsed since germination: $\phi_g(t) = \phi(t-\tau_g)$, defined for all spores

that germinated during the observation interval. Note that the germination time τ_g was determined

exclusively by the bright field images, through a procedure that was independent of the

bioluminescence dynamics (SI Appendix). The population averages of $\phi_g(t)$ in each germination

condition indicate significant synchronization of the bioluminescence dynamics with the time of

germination, as evidenced by the qualitatively similar temporal patterns in the 4 conditions for

which germination was observed (Fig. 3B). In particular, we observe remarkable alignment in the

timing of a primary peak at $t \approx 40$ min, the appearance of a secondary peak between $t = 2$ h and

$t = 3$ h, and similar profiles of signal attenuation at later times, possibly due to degradation of the

lux enzymes (Fig. S2).

Bioluminescence varies significantly across spores

Aligning the dynamics to the time of germination τ_g accounted for the occurrence and timing of

germination as a source of the variability observed in the bioluminescence dynamics. Significant

variability remained in the aligned dynamics $\phi_g(t)$ (Fig. 4A). We analyzed the variability of $\phi_g(t)$

over spores to answer two questions: First, what are the temporal correlations of the

bioluminescence dynamics in individual spores? Second, how much of the variability in

bioluminescence is due to variations in the factors that affect the reaction chemistry, namely, the

redox potential, luciferin concentration, and luciferase copy number?

To analyze temporal correlations in the bioluminescence dynamics, we calculated the

temporal covariance matrix $C(s,t)$ from the photon counts ϕ_g at times s and t (SI Appendix). This

calculation was performed over all germinating spores from the 4 concentrations of L-alanine for

which germination was observed. The covariance matrix was analyzed using principal component

analysis (PCA): PCA applied to time series in this manner is often referred to as empirical

orthogonal function analysis [33]. The first principal component explained 30% of the variance

over individuals, while additional components each explained only a few percent of the total

variance (Fig. S3). Therefore, we focused on the pattern of correlations described by the first

component, which had temporal structure similar to the population-averaged dynamics (Fig. 4B).

This similarity indicates that the dynamics in most spores were well-described by the mean

dynamics multiplied by a coefficient specific to each spore, which we call its amplitude. This simple

mode of variation was useful for quantifying the contributions from different sources, as described

below.

Bioluminescence variability arises from variations in redox potential and luciferin

concentration

Besides the timing of germination, several factors could contribute to variability in the

bioluminescence dynamics. The redox potential and concentration of luciferin in each cell are two

obvious candidates. To investigate these, we performed an experiment with fumarate added to

the germination buffer, in order to drive the citric acid cycle and thus increase the intracellular

redox potential. In a second experiment, decanal was added as an exogenous luciferin. In a third

experiment, both fumarate and decanal were added to the germination buffer. Spores germinated

in response to the addition of 1 mM L-alanine. We calculated the population-averaged

bioluminescence dynamics and performed PCA for each data set independently (Figs. 4C-E).

In the presence of fumarate, the population-averaged bioluminescence dynamics $\langle \phi_g(t) \rangle$

exhibited 2 distinct peaks, approximately 40 min and 3 h after germination (Fig. 4C). These

dynamics appear to be similar to those observed in germination buffer without fumarate (Fig. 4B),

with two major differences: The addition of fumarate increased the intensity of the second peak,

and strongly attenuated the spore-to-spore variability during the first peak, as indicated by the

disappearance of the first peak from the first principal component. The addition of exogenous

luciferin (decanal), both with and without fumarate, resulted in population-averaged

bioluminescence dynamics with a single peak, occurring approximately 40 min after germination,

with intensity approximately 10-fold higher than the initial peak observed in the absence of

exogenous luciferin (Figs. 4D and 4E).

In all conditions, PCA revealed a single significant component of variation (Fig. S3), and

in each condition, this mode's eigenvector was strongly correlated with the average dynamics

(Figs. 4B-E). Thus the primary mode of variation in all conditions was variability in amplitude. This

simple pattern facilitates the quantitation of variability, since the contribution of each spore's signal

to the total variability can be summarized by a single variable: the signal amplitude. Thus we

calculated for each germinating spore the total number of photons counted during the 12 h

following germination, n_g.

The mean value of n_g varied significantly in different conditions. Therefore, to compare

variability in n_g across conditions, we considered the ratio of the standard deviation to the mean,

$\eta = \sigma_{n_g} \langle n_g \rangle^{-1}$, also known as the coefficient of variation. We also considered the asymmetry in the

distribution of n_g by calculating its skew, γ. The estimated statistics are summarized in Table 1.

Variability was high for spores germinating in the absence of exogenous factors (η=160%), and

the distribution of n_g had strong positive skew (γ=3.3). The addition of fumarate significantly

decreased the coefficient of variation and the skew (η=125%, γ=1.4). The addition of decanal

reduced the variability and skew more strongly (η=55%, γ=0.6). The addition of both fumarate

and decanal had the largest effect (η=39%, γ=0.32). We conclude that the large amount of

variability in bioluminescence signal observed in spores germinating in the absence of exogenous

factors was predominantly due to variability in both intracellular luciferin concentration and redox

potential.

The bioluminescence dynamics following germination are influenced by metabolic

changes that occur early in germination, including the initiation of electron transport [23, 34]. In

addition to inducing germination, L-alanine may also fuel energy metabolism [9]. To investigate

this possibility, we considered the photon counts n_g from spores germinating in response to 4

concentrations of L-alanine, in germination buffer (Fig. S4). The population-average $\langle n_g \rangle$

increased 2-fold as the L-alanine concentration increased from 10 µM to 1 mM, then decreased

at 10 mM. The variation in amplitude with L-alanine concentration arose primarily from a

subpopulation of the brightest spores (Fig. S5). The weak, non-monotonic dependence of the

bioluminescence signal on L-alanine concentration indicates that exogenous L-alanine does not

significantly drive energy metabolism in germinating spores.

Discussion

The spore is an extreme phenotypic state that allows an organism to survive extended periods of

starvation, drought, and radiation. Metabolism decreases to extremely low levels during

sporulation, then rapidly increases during germination. Laboratory studies of populations

undergoing the sporulation and germination transitions have also revealed large amounts of

variability over individuals, suggesting that phenotypic diversification in natural spore-forming

populations is a bet-hedging strategy for surviving in fluctuating environments. It is therefore

important to study metabolic dynamics in individual spores.

No previously described methods for quantifying metabolic dynamics in single cells are

applicable to *Bacillus* spores. Therefore, we developed a microscopic technique for this purpose,

using a bioluminescent reporter for the intracellular redox potential. In this paper, we measured

bioluminescence dynamics from individual spores germinating in response to L-alanine, which is

both a germinant and a potential nutrient. Comparing the bioluminescence dynamics measured

in individuals to the population average, we exposed significant variability in the population. A

major source of variability was shown to be associated with differences in the timing of

germination. We could remove this contribution from the bioluminescence dynamics by

simultaneously using bright field microscopy to precisely determine the time of germination of

each spore. The bioluminescence dynamics could then be analyzed for its temporal correlations.

The analysis indicated that the primary mode of variation was associated with the amplitude of

the bioluminescence signal produced by each spore.

By performing experiments at different concentrations of L-alanine, we were able to show

that it was not a major source of variations in bioluminescence amplitude, suggesting that

metabolism in germinating spores was driven not by L-alanine, but rather by endogenous energy

sources stored during sporulation. Through the addition of exogenous nutrients (fumarate) and

bioluminescence substrate (luciferin), we were able to decompose the variability in

bioluminescence amplitude into two main sources: the intracellular redox potential and luciferin

concentration. Both of these sources relate to energy metabolism, as the redox potential drives

ATP synthesis, and the biosynthesis of luciferin depends on both ATP and NADPH [35]. Our

results show that we can begin to understand the metabolic transitions between the growing and

dormant states at the level of single cells.

We foresee two important extensions of the current work. First, one can study the

variability of metabolic dynamics in fluctuating environments. Vegetative cells can be tracked as

they sporulate in response to experimentally imposed starvation, and later germinate and outgrow

in response to nutrients [9]. By simultaneously measuring the bioluminescence dynamics in this

type of experiment, we can study the correlations between spores' timing of sporulation, timing of

germination, ability to outgrow, and metabolic state. Environments also fluctuate due to biotic

interactions. Bacterial spores contain high concentrations of calcium dipicolinate, a compound

that affects resistance and dormancy, but also induces germination in dormant spores [36].

Because spores release large amounts of calcium dipicolinate during germination [37], the local

concentration of spores can affect their germination behavior. Such interactions may explain the

differences we observed in the bioluminescence dynamics with respect to spore density (Figs. 2A

and B, Fig. S1). Interactions between spores and their effects on metabolism could be studied

systematically by analyzing the spatial statistics of germination timing and bioluminescence.

A second extension is to study the thermodynamic nature of dormant states. The

maintenance of order in living matter seems to require energy expenditure, in order to counteract

spontaneously occurring degradation [38, 39]. Therefore, a central question arises: What are the

energetic limits for survival? This question has primarily been studied in microbial communities

from low-energy environments, including deep sediments, permafrost, and subglacial lakes [40-

42]. Metabolic activity in such environments, averaged over the community, has been found to be

reduced by 5 to 6 orders of magnitude compared to the state of growth when nutrients are

supplied [40, 41]. These low metabolic rates are approximately equal to the energetic costs of

repairing molecular damage to proteins and DNA at the corresponding temperatures [41, 42].

However, these analyses included contributions from cells with metabolic activity that was

potentially significantly lower than the population average, including dormant spores [43].

Therefore, spores likely represent the most extreme case of metabolic dormancy in nature.

Most studies of metabolism in dormant spores have not detected activity [44-47].

Extraodinarily sensitive measurements of metabolism in populations of dormant spores using

radiochemical labelling of metabolites did detect activity, reduced by over 5 orders of magnitude

compared to the vegetative state [4]. However, the detected metabolism could have been due to

a rare subpopulation of germinating spores [4, 11, 12], and therefore can serve only as an upper

bound on the rate of metabolic activity in dormant spores. By resolving individual spontaneous

germination events, bioluminescence microscopy could achieve unprecedented sensitivity. The

intensity of bioluminescence is linear in the concentration of $FMNH_2$ at low concentrations [48],

so its utility as a metabolic reporter should extend to cells with very low metabolic rates, such as

dormant spores. A measurement on a single spore at a single time point with signal-to-noise ratio

ξ can be averaged over N individuals and T time points to produce a bulk measurement with

signal-to-noise ratio $\xi(NT)^{1/2}$. For our method applied to germinating spores, ξ is ~20 (SI

Appendix). This suggests that bioluminescence measurements could achieve sensitivity

concomitant with a 10^5-fold reduction in signal compared to germinating spores by measuring, for

example, a population of 10^4 dormant spores over an interval of 18 d, which is feasible using the

setup described here.

A possible mechanism for long-term survival without any metabolic activity is the formation

of a thermodynamically metastable state. The glassy state is an example of metastability that has

been invoked to explain spore dormancy [49, 50], as macromolecules in a glass change

conformation extremely slowly [51]. Therefore, a glassy environment could in principle reduce the

rate of molecular damage to proteins and DNA, while also reducing the activity of metabolic

enzymes and thus causing dormancy. Direct characterization of glassiness in spores is difficult,

as the internal structure of the spore is highly heterogeneous [52]. Instead, one can study a feature

of spores from the perspective of metastability, namely, the statistics of spontaneous germination

and its dependence on environmental conditions such as temperature. This approach should be

possible with our technique, which would also quantify possible differences in metabolism

between spores germinating spontaneously and those germinating in response to nutrients.

Acknowledgments

We thank members of the Dworkin and S. Leibler labs for helpful discussions. ZF was supported

by a grant from Simons Foundation to S. Leibler through Rockefeller University Grant 345430 and

JD was supported by NIH GM122146.

Materials and Methods

Strains

JDB3780 (168 *trpC2 sacA::*P*~sspB~-luxABCDE::cm*) was constructed by transforming genomic DNA

from RL4921 (PY79 *sacA::*P*~sspB~-luxABCDE::cm*) into JDB1772 (168 *trpC2).*

Sporulation protocol

Spores were prepared in Difco sporulation media (DSM). Single colonies grown on LB agar were

used to inoculate 3 ml cultures in test tubes. Cultures were grown with shaking at 30 ℃ for 3 h.

30 µl aliquots were used to inoculate 30 ml cultures in baffled flasks. After growing with shaking

at 30 ℃ for 36 h, spores were purified by spinning at 5,400 g for 10 min at 4 ℃ and resuspending

in sterile water. Cold water washes were repeated 4 times, yielding samples that were

predominantly refractile spores (>90%). Samples were stored at 4 ℃.

Plate reader assay

To assay bioluminescence dynamics in response to various concentrations of L-alanine, dormant

spores (JDB3780) were suspended in 25 mM Tris-HCl buffer at pH 8.2, at a density of 3×10^7

spores ml^{-1}. 200 µl of spore suspension was pipetted into wells of a 96-microwell plate (Nunc

236108). In order to avoid both signal contamination from neighboring wells as well as the

influence of thermal gradients, wells were filled only in the plate interior, in a checkerboard pattern.

2 µl of L-alanine solutions at concentrations between 1 M and 1 mM were added to the appropriate

wells. Each germinant condition was prepared in 3 replicate wells. Three wells with 200 µl of

buffer were included as blanks. The plate was sealed with a gas permeable, partially adhesive

film (Uniscience C20613201), and positioned within the plate reader (Perkin Elmer Wallac Victor^2

1420-011). Luminescence measurements were made without an optical filter, normal aperture

setting, and 10 s integration time. The temperature was maintained at 30 ± 1 ℃. Between

luminescence readings, the plate was shaken orbitally for 10 s with a 2 mm diameter at the slow

speed setting. The plate reader assay of bioluminescence in the presence of chloramphenicol

was performed using a similar procedure, with spores of JDB3785 suspended at a density of

1.2×10^8 spores ml^{-1} and germinated with 1 mM L-alanine.

Reagents and Media

All germination experiments were conducted in germination buffer (25 mM Tris-HCl, pH 8.2).

Decanal was dissolved at a concentration of 15 mM in a 1:1 solution of methanol and water.

Microscopy

<u>Sample preparation</u>

All sample preparation was conducted with sterile technique at the bench.

Adhesion of dormant and germinating spores to glass coverslips was improved by

rendering the glass surface hydrophobic. Coverslips were treated with vapor deposition of

hexamethyldisilazane (Alfa Aesar L16519AE) in a sealed chamber for 1 h, followed by baking at

80 ℃ for 1 h.

Agarose pads were prepared by pipetting 700 µl of molten 4% low-melt agarose (Fisher

BP165-25) between two 22 mm diameter glass coverslips. After 1 h at room temperature, the

agarose pad was exposed by removing the top coverslip, and dried for 5 min.

Prior to spotting the pad, dormant spores were washed twice with water (5400 g, 10 min,

4 ℃). 2 µl of the spore suspension was pipetted onto the center of the dried pad, and the spot

was allowed to completely evaporate (~10 min). The pad was then separated from the bottom

coverslip, and inverted onto a hydrophobic glass coverslip (30 mm diameter, #1.5 thickness,

Bioptechs). The coverslip-bonded pad was left to evaporate for 30 min, then assembled into the

bottom of the perfusion chamber (Bioptechs, ICD with heated lid, gas and media perfusion).

Molten 4% low-melt agarose was added to the exposed portion of the coverslip, and allowed to

gel for 30 min. Creating a contiguous layer of agarose in this manner was found to attenuate

coverslip bounce resulting from the perfusion flow.

Perfusion

A peristaltic pump (Cole-Parmer, Masterflex L/S digital miniflex, dual-channel) was used to draw

media from a reservoir bottle, through sterile tubing (Cole-Parmer, C-Flex #13, i.d. 0.8 mm), into

the perfusion chamber, and finally into a waste reservoir bottle. Tubing was connected with

standard Luer locks, and the bottle openings were wrapped tightly with Parafilm and covered in

aluminum foil. At the beginning of each experiment and whenever the medium was changed (for

example, when germinant was added), the flow was primed at 1 ml min^{-1}. Otherwise, the flow rate

was 50 µl min^{-1}.

Prior to conducting each imaging experiment, the perfusion tubing was sterilized by

flowing 100 ml 20% bleach, followed by 100 ml isopropanol, followed by 5 subsequent bottles

containing 250 ml sterile, purified water (Milli-Q). The perfusion chamber was cleaned by rinsing

with 20% bleach, isopropanol, and distilled water, then air dried.

Imaging

The imaging setup consisted of an inverted optical microscope (Zeiss Axiovert 135), a 63X 1.4

NA oil immersion DIC objective (Zeiss 1113-108) operated without the Wollaston prism, and a

sensitive, back-illuminated, cooled CCD camera (Princeton Instruments, PIXIS 1024B).

A collimated, 445 nm LED (Thorlabs, Solis-445C) was used as external illumination for

bright field imaging. For acquisition of bright field images, the LED was pulsed for 50 ms at low

current (40 mA), and was otherwise fully off.

Autofocus was maintained via a pupil obscuration method applied to an infrared (780 nm)

beam reflected from the glass-media interface (Applied Scientific Instrumentation, CRISP). The

defocus detector controlled the focal position through a DC servo motor connected to the

mechanical focusing shaft of the microscope (Applied Scientific Instrumentation, MFC-2000).

Luminescence images were acquired with 10 min exposure time. The readout was

performed at 100 kHz (read noise 3.56 e$^-$), high gain (1.00 e$^-$ ADU^{-1}), and with 2x2 binning. The

CCD was cooled to -65 ℃ with a combination of thermoelectric and low-vibration liquid cooling

(Princeton Instruments, CoolCUBE II). At this temperature, the dark current was measured at 10^{-3}

e$^-$ pixel^{-1} s^{-1}.

The microscope and camera were enclosed inside a box made of styrofoam-lined black

plastic. Exterior edges and corners of the box were covered with opaque black fabric and taped.

A removable styrofoam-lined plastic panel provided access to the microscope and sample holder,

and was held in place during data acquisition with Velcro straps and metal L-brackets. The

temperature within the box was controlled by a thermoelectric device (TE Technology, AC-220),

coupled to a PID feedback device (TE Technology, TC-720) monitoring a thermistor placed as

close as possible to the perfusion chamber. The interior temperature of the box was maintained

at 30±0.05 ℃. The entire imaging apparatus was supported by a vibration-damped optical table,

situated in a temperature-controlled room (28±0.1 ℃).

Image analysis and statistics

See SI Appendix.

Table

media	#spores	mean: [photons] $\langle n_g \rangle$	coefficient of variation: η	skew: γ
germination buffer only	11,882	1,300±19	161% [154%, 169%]	3.326±0.022
germ. buffer +0.4% fumarate	1,976	2,862±81	125% [117%, 137%]	1.449±0.055
germ. buffer +30 µM decanal	928	18,600±340	55% [52%, 59%]	0.56±0.080
germ. buffer +30 µM decanal +0.4% fumarate	284	14,000±320	39% [35%, 43%]	0.32±0.14

Table 1. Bioluminescence signal amplitude statistics. Columns denote media, the number of spores analyzed, and the statistics of n_g, the photon counts per spore integrated over the 12 h after germination. The ± signs indicate 1 standard error. The bracketed interval in the coefficient of variation column is the 95% confidence interval.

Figure Legends

Fig. 1. Schematic of bioluminescence microscope and perfusion chamber with example images and time series. (A) Microscope schematic showing 3 light paths: bright field (blue), luminescence (green), and autofocus infrared (red). For bright field imaging (blue path), the external illumination is focused near the bottom of the agarose pad (yellow slab), where sample cells are sandwiched against a glass coverslip. The sample emits luminescence (green path) in all directions. A temperature-controlled, light-tight box (thick black lines) encloses the microscope. **(B)** The perfusion chamber flows medium around the agarose pad (yellow) that traps sample cells against a glass coverslip on the chamber bottom, sealed by an O-ring (orange). **(C)** Bright field image of P$_{sspB}$-*luxABCDE* spores, before the addition of germinant (time 0 h). Highly refractile dormant spores appear dark, *i.e.,* high contrast. The field of view contains approximately 900

spores. Scale bar: 20 µm. **(D,E)** The subregion bounded by the green box is shown in panels (D)

(bright field) and (E) (luminescence), showing the lack of luminescence signal from dormant

spores. **(F,G)** The same region is depicted 3.5 h after the addition of 100 µM L-alanine, showing

spores' typical loss of bright field contrast upon germination (F), and the corresponding increase

in luminescence signal (G) that occurs in some cells after germination. **(H,I)** Time series of bright

field contrast (H) and luminescent photon counts (I) of the 5 representative spores outlined in (D)-

(G). The rapid loss of contrast shown in panel (H) defines the time of germination, τ_g, for each

spore. Colored cell outlines are dilated by 1 pixel for visualization. Panel (D) scale bar: 5 µm.

Fig. 2. Bioluminescence dynamics of spore populations germinating in response to L-

alanine. (A) The bioluminescence emitted by P$_{sspB}$-*luxABCDE* spores germinating in a range of

L-alanine concentrations was assayed in a 96-well plate reader. Each well contained

approximately 6×10^6 spores. Lines depict the mean signal over 3 replicates; the standard

deviation was negligible on this scale. **(B)** The bioluminescence signal $\phi(t)$ was measured in

individual spores germinating in response to various concentrations of L-alanine added at time 0

h. The mean signal over all spores (>600 individuals per concentration) is plotted at each time

point. **(C)** The bioluminescence signals in (A) and (B) were integrated over the period between 0

and 12 h after the addition of alanine to calculate the population-averaged photon counts n,

plotted as a function of alanine concentration.

Fig. 3. Time of germination varies in populations and synchronizes with bioluminescence

dynamics. Dormant spores were exposed to 5 concentrations of L-alanine at time 0 h. **(A)** The

fraction of spores that had not yet germinated by time t, P(τ_g>t), is plotted on a linear scale (left)

and a log scale (right). An offset is added to avoid divergence of zero values on the log scale. **(B)**

The luminescence signal of each spore as a function of time elapsed since the time of germination

is $\phi_g(t) = \phi(t-\tau_g)$, defined for spores that germinated during the acquisition interval. The population

average of $\phi_g(t)$ is plotted for each concentration of L-alanine. Note that no spores germinated in

1 µM L-alanine and so no curve for that condition appears in (B).

Fig. 4. Variability in bioluminescence dynamics over spores. (A) Bioluminescence as a

function of time since germination is analyzed in terms of variability over individuals. Data include

all spores that germinated in response to L-alanine at 4 concentrations (Table 1). The time series

of $\phi_g(t)$ for a randomly chosen subset of 500 cells are plotted as thin black lines. The solid red line

indicates the median over the 11,000 cells comprising the data set, and the dashed red lines

indicate the 16% and 84% quartiles. (**B-E**) Principal component analysis was performed on time

series $\phi_g(t)$ from 4 experimental conditions: germination buffer only (B), 0.4% sodium fumarate

(C), 30 µM decanal (D), and 30 µM decanal plus 0.4% sodium fumarate (E). The dark gray line

shows the population average $\langle \phi_g(t) \rangle$ for each condition. The first principal component for each

condition explained 30%, 55%, 68%, and 58% of the total variance, respectively. The red line

shows the loadings of the first principle component, *i.e.*, $\lambda_1^{1/2} w_1(t)$, where λ_1 is the largest

eigenvalue of the covariance matrix, and w_1 is the first principle component, scaled to unit norm.

The average and loadings are plotted on separate axes. The first principal component and the

population average were highly correlated in all conditions: defining $r = \mathrm{cor}(w_1(t), \langle \phi_g(t) \rangle)$, $r_{\mathrm{buffer-}}$

$_{\mathrm{only}} = 84\%$, $r_{\mathrm{fumarate}} = 97\%$, $r_{\mathrm{decanal}} = 89\%$, $r_{\mathrm{decanal+fumarate}} = 99\%$ (all *p*-values $< 2.2 \times 10^{-16}$).

Figures

Figure 1

Figure 2

Figure 3

 Figure 4

References

1. Setlow, P., *Spores of Bacillus subtilis: their resistance to and killing by radiation, heat and chemicals.* Journal of Applied Microbiology, 2006. **101**(3): p. 514-525.

2. Graham-Smith, G.S., *The longevity of dry spores of B. anthracis.* Journal of Hygiene, 1930. **30**(2): p. 213-215.

3. Sneath, P.H.A., *Longevity of micro-organisms.* Nature, 1962. **195**(4842): p. 643-646.

4. Desser, H. and E. Broda, *Radiochemical investigation of the respiration of spores of Bacillus cereus.* Archiv Für Mikrobiologie, 1969. **65**(1): p. 76-86.

5. Mehta, P. and D.J. Schwab, *Energetic costs of cellular computation.* Proceedings of the National Academy of Sciences of the United States of America, 2012. **109**(44): p. 17978-17982.

6. Xue, B.K., P. Sartori, and S. Leibler, *Environment-to-phenotype mapping and adaptation strategies in varying environments.* Proceedings of the National Academy of Sciences of the United States of America, 2019. **116**(28): p. 13847-13855.

7. Yi, X., J. Liu, J.R. Faeder, and P. Setlow, *Synergism between different germinant receptors in the germination of Bacillus subtilis spores.* Journal of Bacteriology, 2011. **193**(18): p. 4664-4671.

8. Veening, J.-W., E.J. Stewart, T.W. Berngruber, F. Taddei, O.P. Kuipers, and L.W. Hamoen, *Bet-hedging and epigenetic inheritance in bacterial cell development.* Proceedings of the National Academy of Sciences of the United States of America, 2008. **105**(11): p. 4393-4398.

9. Mutlu, A., S. Trauth, M. Ziesack, K. Nagler, J.P. Bergeest, K. Rohr, N. Becker, T. Höfer, and I.B. Bischofs, *Phenotypic memory in Bacillus subtilis links dormancy entry and exit by a spore quantity-quality tradeoff.* Nature Communications, 2018. **9**.

10.   Zhang, J.-Q., K.K. Griffiths, A. Cowan, P. Setlow, and J. Yu, *Expression level of Bacillus*

subtilis germinant receptors determines the average rate but not the heterogeneity of

spore germination. Journal of Bacteriology, 2013. **195**(8): p. 1735-1740.

11.   Powell, J.F. and R.E. Strange, *Biochemical changes occurring during the germination of*

bacterial spores. Biochemical Journal, 1953. **54**(2): p. 205-209.

12.   Sturm, A. and J. Dworkin, *Phenotypic diversity as a mechanism to exit cellular dormancy.*

Curr Biol, 2015. **25**(17): p. 2272-2277.

13.   de Jong, I.G., J.-W. Veening, and O.P. Kuipers, *Heterochronic phosphorelay gene*

expression as a source of heterogeneity in Bacillus subtilis spore formation. Journal of

Bacteriology, 2010. **192**(8): p. 2053-2067.

14.   Takhaveev, V. and M. Heinemann, *Metabolic heterogeneity in clonal microbial*

populations. Current Opinion in Microbiology, 2018. **45**: p. 30-38.

15.   Laflamme, C., D. Verreault, S. Lavigne, L. Trudel, J. Ho, and C. Duchaine,

Autofluorescence as a viability marker for detection of bacterial spores. Frontiers in

Bioscience, 2005. **10**: p. 1647-1653.

16.   Sarasanandarajah, S., J. Kunnil, B.V. Bronk, and L. Reinisch, *Two-dimensional*

multiwavelength fluorescence spectra of dipicolinic acid and calcium dipicolinate. Applied

Optics, 2005. **44**(7): p. 1182-1187.

17.   Jelen, B.I., D. Giovannelli, and P.G. Falkowski, *The role of microbial electron transfer in*

the coevolution of the biosphere and geosphere. The Annual Review of Microbiology,

2016. **70**: p. 45-62.

18.   Kim, B.H. and G.M. Gadd, *Bacterial Physiology and Metabolism*. 2008, Cambridge:

Cambridge University Press.

19.   Gross, B.J. and M.Y. El-Naggar, *A combined electrochemical and optical trapping platform*

for measuring single cell respiration rates at electrode interfaces. Review of Scientific

Instruments, 2015. **86**(6).

20.   Hastings, J.W., *Chemistries and colors of bioluminescent reactions: A review.* Gene, 1996.

173(1): p. 5-11.

21.   Koga, K., T. Harada, H. Shimizu, and K. Tanaka, *Bacterial luciferase activity and the*

intracellular redox pool in Escherichia coli. Molecular Genetics and Genomics, 2005.

274(2): p. 180-188.

22.   Carmi, O.A., G.S.A.B. Stewart, S. Ulitzur, and J. Kuhn, *Use of bacterial luciferase to*

establish a promoter probe vehicle capable of nondestructive real-time analysis of gene

expression in Bacillus spp. Journal of Bacteriology, 1987. **169**(5): p. 2165-2170.

23.   Hill, P.J., L. Hall, D.A. Vinicombe, C.J. Soper, P. Setlow, W.M. Waites, S. Denyer, and

G.S.A.B. Stewart, *Bioluminescence and spores as biological indicators of inimical*

processes. Journal of Applied Bacteriology, 1994. **76**: p. S129-S134.

24.   Schmalisch, M., E. Maiques, L. Nikolov, A.H. Camp, B. Chevreux, A. Muffler, S.

Rodriguez, J. Perkins, and R. Losick, *Small genes under sporulation control in the Bacillus*

subtilis genome. Journal of Bacteriology, 2010. **192**(20): p. 5402-5412.

25.   Ciarciaglini, G., P.J. Hill, K. Davies, P.J. McClure, D. Kilsby, M.H. Brown, and P.J. Coote,

Germination-induced bioluminescence, a route to determine the inhibitory effect of a

combination preservation treatment on bacterial spores. Applied and Environmental

Microbiology, 2000. **66**(9): p. 3735-3742.

26.   Hastings, J.W. and G. Weber, *Total quantum flux of isotropic sources.* Journal of the

Optical Society of America, 1963. **53**(12): p. 1410-1415.

27.   Gregor, C., K.C. Gwosch, S.J. Sahl, and S.W. Hell, *Strongly enhanced bacterial*

bioluminescence with the ilux operon for single-cell imaging. Proceedings of the National

Academy of Sciences of the United States of America, 2018. **115**(5): p. 962-967.

28.   Mihalcescu, I., W. Hsing, and S. Leibler, *Resilient circadian oscillator revealed in individual*

cyanobacteria. Nature, 2004. **430**(6995): p. 81-85.

29.   Phiefer, C.B., R.J. Palmer, and D.C. White, *Comparison of relative photon flux from single*
581   *cells of the bioluminescent marine bacteria Vibrio fischeri and Vibrio harveyi using photon-*
582   *counting microscopy.* Luminescence, 1999. **14**(3): p. 147-151.

583   30.   McLean, I.S., *Electronic Imaging in Astronomy: Detectors and Instrumentation.* 2008:
584   Springer Science & Business Media.

585   31.   Zhang, P., L. Kong, G. Wang, M. Scotland, S. Ghosh, B. Setlow, P. Setlow, and Y.-Q. Li,
586   *Analysis of the slow germination of multiple individual superdormant Bacillus subtilis*
587   *spores using multifocus Raman microspectroscopy and differential interference contrast*
588   *microscopy.* Journal of Applied Microbiology, 2012. **112**(3): p. 526-536.

589   32.   Cabrera-Martinez, R.-M., F. Tovar-Rojo, V.R. Vepachedu, and P. Setlow, *Effects of*
590   *overexpression of nutrient receptors on germination of spores of Bacillus subtilis.* Journal
591   of Bacteriology, 2003. **185**(8): p. 2457-2464.

592   33.   Weare, B.C., A.R. Navato, and R.E. Newell, *Empirical orthogonal analysis of Pacific sea*
593   *surface temperatures.* Journal of Physical Oceanography, 1976. **6**(5): p. 671-678.

594   34.   Scott, I.R. and D.J. Ellar, *Metabolism and the triggering of germination of Bacillus*
595   *megaterium. Use of L-[3H]alanine and tritiated water to detect metabolism.* Biochemical
596   Journal, 1978. **174**(2): p. 635-640.

597   35.   Soly, R.R. and E.A. Meighen, *Identification of the acyl transfer site of fatty acyl-protein*
598   *synthetase from bioluminescent bacteria.* Journal of Molecular Biology, 1991. **219**(1): p.
599   69-77.

600   36.   Paidhungat, M., K. Ragkousi, and P. Setlow, *Genetic requirements for induction of*
601   *germination of spores of Bacillus subtilis by Ca2+-dipicolinate.* Journal of Bacteriology,
602   2001. **183**(16): p. 4886-4893.

603   37.   Huang, S.-S., D. Chen, P.L. Pelczar, V.R. Vepachedu, P. Setlow, and Y.-Q. Li, *Levels of*
604   *Ca(2+)-dipicolinic acid in individual Bacillus spores determined using microfluidic Raman*
605   *tweezers.* Journal of Bacteriology, 2007. **189**(13): p. 4681-4687.

38. Schrödinger, E., *What is life? The physical aspect of the living cell and mind.* 1944,
Cambridge: Cambridge University Press.

39. Hoehler, T.M., *Biological energy requirements as quantitative boundary conditions for life*
in the subsurface. Geobiology, 2004. **2**(4): p. 205-215.

40. Morita, R.Y., *Bacteria in Oligotrophic Environments.* 1997, New York: Chapman & Hall.

41. Price, P.B. and T. Sowers, *Temperature dependence of metabolic rates for microbial*
growth, maintenance, and survival. Proceedings of the National Academy of Sciences of
the United States of America, 2004. **101**(13): p. 4631-4636.

42. Lever, M.A., K.L. Rogers, K.G. Lloyd, J. Overmann, B. Schink, R.K. Thauer, T.M. Hoehler,
and B.B. Jørgensen, *Life under extreme energy limitation: a synthesis of laboratory- and*
field-based investigations. FEMS Microbiology Reviews, 2015. **39**(5): p. 688-728.

43. Hoehler, T.M. and B.B. Jørgensen, *Microbial life under extreme energy limitation.* Nature
Reviews Microbiology, 2013. **11**(2): p. 83-94.

44. Nelson, D.L. and A. Kornberg, *Biochemical studies of bacterial sporulation and*
germination. XIX. Phosphate metabolism during sporulation. The Journal of Biological
Chemistry, 1970. **245**(5): p. 1137-1145.

45. Swerdlow, B.M., B. Setlow, and P. Setlow, *Levels of H+ and other monovalent cations in*
dormant and germinating spores of Bacillus megaterium. Journal of Bacteriology, 1981.
148(1): p. 20-29.

46. Singh, R.P., B. Setlow, and P. Setlow, *Levels of small molecules and enzymes in the*
mother cell compartment and the forespore of sporulating Bacillus megaterium. Journal of
Bacteriology, 1977. **130**(3): p. 1130-1138.

47. Ghosh, S., G. Korza, M. Maciejewski, and P. Setlow, *Analysis of metabolism in dormant*
spores of Bacillus species by 31P nuclear magnetic resonance analysis of low-molecular-
weight compounds. Journal of Bacteriology, 2015. **197**(5): p. 992-1001.

48.   Matheson, I.B.C. and J. Lee, *Kinetics of bacterial bioluminescence and the fluorescent*

transient. Photochemistry and Photobiology, 1983. **38**(2): p. 231-240.

49.   Sapru, V. and T.P. Labuza, *Glassy state in bacterial spores predicted by polymer glass-*

transition theory. Journal of Food Science, 1993. **58**(2): p. 445-448.

50.   Ablett, S., A.H. Darke, P.J. Lillford, and D.R. Martin, *Glass formation and dormancy in*

bacterial spores. International Journal of Food Science and Technology, 1999. **34**(1): p.

59-69.

51.   Hagen, S.J., J. Hofrichter, and W.A. Eaton, *Protein reaction kinetics in a room-temperature*

glass. Science, 1995. **269**(5226): p. 959-962.

52.   Sunde, E.P., P. Setlow, L. Hederstedt, and B. Halle, *The physical state of water in bacterial*

spores. Proceedings of the National Academy of Sciences of the United States of America,

2009. **106**(46): p. 19334-19339.

Supporting Materials and Methods

Sensitivity limits of bioluminescence microscopy

Sensor read noise, dark current, and high energy particles such as cosmic rays generate a noise background for imaging. The signal measured from a steadily emitting source accumulates with increasing exposure time t. Noise due to dark current also accumulates with t, but modern cooled CCDs have very low dark current, so that sensor read noise dominates for t up to about 1 h. The number of high energy particle artifacts per image also increases with t. These artifacts can be identified automatically [1], so that pixels affected by these artifacts can be removed from subsequent analysis. From a practical perspective, t must be short enough to ensure that the majority of pixels are not affected by these artifacts. The flux of high energy particles from cosmic rays is about 1 cm^{-2} min^{-1} [1]. In our preliminary experiments, artifacts from high energy particles limited t to 10 min.

After removing artifacts due to high energy particles, imaging noise is dominated by sensor readout noise. We consider the following parameters: the average rate of photon emission per spore, ψ; the average number of pixels in the luminescence image of a spore, N; the RMS readout noise per pixel σ_r; the exposure time t, and the collection efficiency of the imaging setup s. Note that the collection efficiency s is the ratio of the number of emitted photons to the number of electrons measured by the sensor. It includes geometric effects, $i.e.$, losses due to the collection geometry, reflections from optical surfaces, and the quantum yield of the sensor.

The signal to noise ratio for each pixel in the image of the spore, χ_p, is the number of electrons per pixel per exposure arising from emitted luminescence, divided by the magnitude of the readout noise: $\chi_p = s\psi t N^{-1}\sigma_r^{-1}$. The signal from each spore is the sum over the signals from the pixels comprising its image. The readout noise is independent for each pixel, so the signal to noise ratio per spore, χ, equals $\chi_p N^{1/2} = s\psi t N^{-1/2}\sigma_r^{-1}$. The sensitivity limit of the technique is taken to be the rate of photon emission per spore for which the signal to noise ratio χ is equal to 1: $\psi_{min} = s^{-1}t^{-1}N^{1/2}\sigma_r$.

Optimizing the sensitivity of the technique amounts to maximizing collection efficiency s, minimizing the number of pixels per spore n, and minimizing the sensor read noise σ_r. Practically, s is less than 1/2 electrons per photon, since at least half of photons are emitted away from the collection objective. The numerical aperture of the objective should be as high as possible, to maximize collection of the remaining photons. The quantum efficiency of back-illuminated CCDs in the spectral range of bacterial bioluminescence (~490 nm) typically exceeds 0.9 electrons per photon. Accounting for these effects, we estimated that we could achieve a value of s as high as 0.3 electrons per photon. For the camera we used in the microscope, the parameters were as follows: $s = 0.3$ e$^-$ photon^{-1} (estimated), $t = 10$ min, $N = 10$ pixels, $\sigma_r = 3.6$ e$^-$, which determines the sensitivity limit $\psi_{min} \approx 3.8$ photons min^{-1}. Published results typically report the number of measured photons per time ($\gamma^{-1}s\psi$, where γ is the gain, expressed in electrons per photon), rather than the number emitted per time (ψ), and so we introduce a new symbol for the former rate: $\Phi = \gamma^{-1}s\psi$, and estimate the sensitivity limit $\Phi_{min} = \gamma^{-1}s\psi_{min} \approx 1$ photons min^{-1} ($\gamma = 1$ e$^-$ photon^{-1} for our setup).

We analyzed the signal to noise ratio of our measurements on a per-spore basis. The noise background was quantified by considering the statistics of luminescence signal from the reported experiments in regions of the image that did not contain spores. These regions were defined to each include the same number of pixels as a real spore. Twelve thousand such regions were analyzed, and the standard deviation of the estimated photon counts from each region was calculated: $\sigma = 3.5$ photons min^{-1}. The bioluminescence dynamics per spore $\Phi(t)$ varied over time and across spores. Detection of the signal from a particular spore is primarily determined by that spore's maximal signal. For the purposes of determining a single signal to noise ratio that would describe the typical sensitivity of the technique, we calculated the population-average of the temporal maximum bioluminescence signal:

$$\phi^* = \langle \max_t \phi(t) \rangle$$

The signal to noise ratio is given by $\chi = \Phi^* \sigma^{-1}$. Its values were $\chi = 4.2$ (germination buffer only), 5.1 (0.4% fumarate), 22 (30 µM decanal), and 18 (30 µM decanal + 0.4% fumarate).

Image analysis

<u>Background subtraction</u>

Spatial inhomogeneities were observed in long exposure images, even those acquired in complete darkness without a luminescent sample. These inhomogeneities were found to be nearly constant over time and were treated as a form of background. To remove this background, a sequence of at least 20 images was acquired in darkness prior to each experiment, and the temporal median of this sequence, *i.e.*, the estimated background, was subtracted from each luminescence image subsequently acquired during the experiment.

<u>Removal of artifacts due to high energy particles</u>

Each luminescence image included several small regions of extremely bright pixels, due to high energy particles, such as cosmic rays, ionizing the CCD. In order to prevent these artifacts from corrupting the luminescence signal, we developed a routine to automatically detect pixels that appeared to be affected by such events. Pixels affected by artifacts had a particularly identifying feature: extreme brightness, that appeared rapidly in time and with very high spatial contrast. Temporal and spatial high-pass filters were applied to the images, and the resulting images were segmented, using a threshold chosen to separate the vast majority (>99%) of affected pixels from unaffected pixels. The affected regions were grown by 2 pixels to avoid any remaining signal contamination. The mean number of artifacts was 2.0 min^{-1}, and the mean number of pixels affected by each artifact was 28.

<u>Spore identification and segmentation</u>

Spores were identified by determining the maximally stable extremal regions of the bright field image, using the VLFeat software library [2, 3]. The parameters of the method were tuned manually to identify >99% of spores, with a negligible false identification rate (<0.1%).

Determination of Germination Time τ_g

The bright field contrast of each spore was determined by calculating the spatial gradient of each bright field image, ∇I_{BF}. The magnitude of the gradient was averaged over the pixels p corresponding to each spore to calculate its contrast:

$$C(t) = \sum_p \|\nabla I_{BF}(t)\|^2$$

The dynamics of contrast of germinating spores typically appeared as noisy step function, declining from a high value prior to germination to a low value after. The time of germination τ_g was determined for each spore by fitting the measured contrast with a regression tree having a single split, using the rpart package for R, v. 4.1-15. The automatically determined τ_g was found to exactly match the value determined by manually considering the time-lapse images in 100 spores selected randomly from 4 data sets.

Statistical analysis

Temporal filtering

Camera read noise and photon counting noise (shot noise) corrupt the counts of bioluminescence photons. Each of the noise sources is expected to have zero temporal autocorrelation, while the bioluminescence dynamics are expected to have significant autocorrelation. Defining the photon counts within each pixel p as $X_p(t)$, we expect the power spectral density of X_p to be different for pixels located within the image of a spore vs. pixels located outside (background). To text this, we estimated the power spectral density of $X_p(t)$ over 110,000 pixels within the image of spores ("signal": $P_S(\omega)$) and over 200,000 pixels not within the image of spores (background: $P_{BG}(\omega)$).

The power spectral density of the background was flat, while that of the signal was flat above $\omega \approx 0.84$ h^{-1}. In order to filter the background noise without significantly blurring the signal dynamics, we applied a low-pass filter (sinc filter), with a cutoff period of 36 min, to the bioluminescence signal measured from each pixel. For each spore, the sum of these counts over all pixels inside its bright field image was calculated for each time point, resulting in $\Phi(t)$.

Principal component analysis

The covariance matrix $C(s,t)$ was calculated over all germinating spores as:

$$C(s,t) = N^{-1} \sum_{i=1}^{N} \left(\phi_g^i(s) - \langle \phi_g(s) \rangle \right) \left(\phi_g^i(t) - \langle \phi_g(t) \rangle \right)$$

where $\phi_g^i(t)$ is the estimated rate of photons counted from spore i at time t, and N is the number of spores in the data set. The times s and t took values corresponding to 87 time points: 12 before germination (2 h), and 75 following germination (12.3 h). Principal component analysis was performed on the covariance matrix $C(s,t)$, resulting in the eigenvalues λ_i and the eigenvectors $w_i(t)$. The eigenvalues have units (photons min^{-1})^2 and the eigenvectors are unitless and have norm 1:

$$\sum_{j=1}^{87} \left(w_i(t_j) \right)^2 = 1$$

The temporal pattern of variation explained by mode i is quantified by $\lambda_i^{1/2} w_i(t)$, as is plotted in Figs. 4B-E for $i = 1$.

Statistics of n_g

The number of photons counted per spore was calculated as

$$n_g = \sum_{t \geq 0}^{t \leq 12 \text{ h}} \phi_g(t) \Delta t$$

where $\Delta t = 10$ min is the exposure time.

For Table 1, the estimates of the skew of n_g and its standard error were calculated for each data set using method 3 of Joanes & Gill [4]. The confidence interval of the coefficient of variation was estimated using the formula of McKay [5].

Figure S1. Bioluminescence dynamics depend on spore concentration. The population-averaged bioluminescence signal $\langle\phi(t)\rangle$ was measured in a 96-well plate reader, at a range of spore densities, and normalized by the number of spores per well to facilitate comparison between densities. Spores were induced to germinate with 1 mM L-alanine. Lines depict the mean signal over 3 replicates and the shaded region encloses ±1 standard deviation. Spore density in the experiments performed with the microscope was less than 10^5 ml^{-1}.

Figure S2. Bioluminescence signal decays exponentially after 4 h post-germination. The population-averaged bioluminescence dynamics of germinating spores $\langle \phi_g(t) \rangle$, depicted by the black line, exhibit exponential decline, beginning at 4 h post-germination. The rate of exponential decline r was estimated in each of 4 conditions: $r_{\text{buffer-only}} = 22\%$ h^{-1}, $r_{\text{fumarate}} = 22\%$ h^{-1}, $r_{\text{decanal}} = 8\%$ h^{-1}, $r_{\text{decanal+fumarate}} = 12\%$ h^{-1}. Estimated fits are depicted by red lines.

Figure S3. Variance explained by principal components. Principal component analysis was performed on the bioluminescent photon counts $\phi_g(t)$ for each of 4 experimental conditions, as described in the Results section. The variance explained by principal component number k is λ_k, and the fraction of variance explained is $\lambda_k(\Sigma_i \lambda_i)^{-1}$. The data sets included 87 time points and therefore 87 principal components; only the first 25 components are included in the plots.

Figure S4. Bioluminescent photons counted per germinating spore. The bioluminescence signal as a function of time since germination, $\phi_g(t) = \phi(t-\tau_g)$, was integrated from 0 to 12 h post-germination to calculate the number of photons counted per germinating spore, n_g. The average of n_g over all germinating spores is plotted against L-alanine concentration.

Figure S5. Quantile-quantile plot of bioluminescent photons counted per spore, germinating in response to 4 concentrations of L-alanine. Quantiles of the number of photons emitted per germinating spore during the 12 h post-germination, n_g, were determined for 4 concentrations of L-alanine. Quantiles were also calculated for the data from all 4 L-alanine concentrations, weighted for equal contribution from each condition. The quantiles in each condition are plotted on the vertical axis, as a function of the corresponding quantile in all conditions. Quantiles were determined in increments of 5%, starting at 5% and ending at 95%.

References

1. Groom, D., *Cosmic rays and other nonsense in astronomical CCD imagers.* Experimental Astronomy, 2002. **14**(1): p. 45-55.
2. Matas, J., O. Chum, M. Urban, and T. Pajdla, *Robust wide-baseline stereo from maximally stable extremal regions.* Image and Vision Computing, 2004. **22**(10): p. 761-767.
3. Vedaldi, A.F., B., *VLFeat: an open and portable library of computer vision algorithms*. 2008: http://www.vlfeat.org/.
4. Joanes, D.N. and C.A. Gill, *Comparing measures of sample skewness and kurtosis.* Journal of the Royal Statistical Society Series D (The Statistician), 1998. **47**(1): p. 183-189.
5. McKay, A.T., *Distribution of the coefficient of variation and the extended "t" distribution.* Journal of the Royal Statistical Society, 1932. **95**(4): p. 695-698.