A-to-I RNA editing in honeybees shows signals of adaptation and convergent evolution

Yuange Duan[1,2], Shengqian Dou[1], Hagit T. Porath[3], Jiaxing Huang[4], Eli Eisenberg[5,*], and Jian Lu[1,6,*]

1. State Key Laboratory of Protein and Plant Gene Research, Center for Bioinformatics, School of Life Sciences, Peking University, Beijing, 100871, China
2. Academy for Advanced Interdisciplinary Studies, Peking University, Beijing, 100871, China
3. The Mina and Everard Goodman Faculty of Life Sciences, Bar Ilan University, Ramat-Gan 52900, Israel
4. Key Laboratory for Insect-Pollinator Biology of the Ministry of Agriculture and Rural Affairs, Institute of Apicultural Research, Chinese Academy of Agricultural Sciences, Beijing, 100093, China
5. Raymond and Beverly Sackler School of Physics and Astronomy and Sagol School of Neuroscience, Tel Aviv University, Tel Aviv 6997801, Israel
6. Lead Contact

*Correspondence:
Jian Lu
Email: luj@pku.edu.cn

Eli Eisenberg
Email: elieis@post.tau.ac.il

SUMMARY

Social insects exhibit extensive phenotypic diversities among the genetically similar individuals, suggesting a role for the epigenetic regulations beyond the genome level. The ADAR-mediated adenosine-to-inosine (A-to-I) RNA editing, an evolutionarily conserved mechanism, facilitates adaptive evolution by expanding proteomic diversities. Here, we characterize the A-to-I RNA editome of honeybees (*Apis mellifera*), identifying 407 high-confidence A-to-I editing sites. Editing is most abundant in the heads, and shows signatures for positive selection. Editing behavior differs between foragers and nurses, suggesting a role for editing in caste differentiation. Although only five sites are conserved between bees and flies, an unexpectedly large number of genes exhibit editing in both species, albeit at different locations, including the nonsynonymous auto-editing of *Adar*. This convergent evolution, where the same target genes independently acquire recoding events in distant diverged clades, together with the signals of adaptation observed in honeybees alone, further supports the notion of recoding being adaptive.

Keywords: A-to-I RNA editing; Honeybee; Caste-dependent editing; Adaptation; Convergent evolution.

INTRODUCTION

Adenosine (A) to inosine (A-to-I) RNA editing, catalyzed by enzymes of the ADAR (adenosine deaminase acting on RNA) family (Savva et al. 2012b), is an evolutionarily conserved mechanism that expands RNA diversity at the co-transcriptional or post-transcriptional level in metazoans (Bass 2002; Nishikura 2010; Eisenberg and Levanon 2018). Two catalytically-active *Adar* genes are encoded in most metazoans. However, insects have lost ADAR1 and encode only a single *Adar* gene (Keegan et al. 2011). Due to the structural similarity between inosine (I) and guanosine (G), I is generally believed to be recognized as G in many cellular processes such as mRNA splicing (Rueter et al. 1999; Flomen et al. 2004; Jin et al. 2007; Lev-Maor et al. 2007), microRNA (miRNA) biogenesis or target recognition (Liang and Landweber 2007; Borchert et al. 2009; Alon et al. 2012), and mRNA translation (Basilio et al. 1962; Licht et al. 2019). Therefore, an A-to-I RNA editing usually has a similar effect as an A-to-G DNA substitution. A-to-I editing plays essential roles in many biological and physiological processes(Keegan et al. 2001; Nishikura 2010), and dysregulation of A-to-I editing might be associated with cancer, autoimmune disorders, or other human diseases (Gallo et al. 2017).

During the past two decades A-to-I RNA editing sites have been systematically characterized in various metazoan species (Ramaswami and Li 2016). The majority of RNA editing sites are located in clusters in non-coding regions of humans (Athanasiadis et al. 2004; Blow et al. 2004; Kim et al. 2004; Levanon et al. 2004; Picardi et al. 2017), monkeys (Chen et al. 2014; Yang et al. 2015), mice (Neeman et al. 2006; Danecek et al. 2012), worms (Morse and Bass 1999; Zhao et al. 2015; Goldstein et al. 2017), corals (Porath et al. 2017b) and many other species (Porath et al. 2017a). In most species, only a minute fraction of the edits resides within the coding sequence. Notable exceptions are *Drosophila* (Graveley et al. 2011; Rodriguez et al. 2012; St Laurent et al. 2013; Mazloomian and Meyer 2015; Yu et al. 2016; Buchumenski et al. 2017; Duan et al. 2017; Zhang et al. 2017) and cephalopod species (Alon et al. 2015; Liscovitch-Brauer et al. 2017). Despite the deep conservation of A-to-I editing mechanism, the target landscapes of editing have considerably evolved during metazoan evolution. Only one editing site is known to be conserved across virtually all mammals, *Drosophila*, and cephalopods (Porath et al. 2019), and it thus seems that RNA editing modulates the diversity of

the transcriptomes and proteomes in a lineage-specific manner.

The forces driving the evolution of A-to-I editing across species at the macro-evolutional

scale are not well understood. RNA editing was hypothesized to facilitate adaptive evolution

by increasing proteomic diversities temporally or spatially, in a manner more flexible than

genomic mutations (Gommans et al. 2009; Nishikura 2010; Klironomos et al. 2013; Rosenthal

2015; Nishikura 2016). However, in most species studied the ratio of nonsynonymous *(N)* to

synonymous (*S*) editing sites (*N/S*) is lower than expected for random sites, suggesting that

recoding events may be overall non-adaptive (Xu and Zhang 2014). Here too *Drosophila* and

cephalopods stand out as exceptions, exhibiting high *N/S* ratios indicating positive selection of

recoding. Furthermore, hundreds of recoding sites were shown to be conserved across the

Drosophila lineage (Graveley et al. 2011; Rodriguez et al. 2012; St Laurent et al. 2013;

Mazloomian and Meyer 2015; Yu et al. 2016; Buchumenski et al. 2017; Duan et al. 2017; Zhang

et al. 2017), and thousands in the behaviorally sophisticated cephalopods (Garrett and

Rosenthal 2012; Alon et al. 2015; Liscovitch-Brauer et al. 2017), supporting the functional

importance of these editing events. However, in only a few examples was the advantageous

effect conferred by RNA editing explicitly demonstrated (see for example (Garrett and

Rosenthal 2012)), and even it is not yet clear to what extent editing is indeed utilized for

proteome diversification.

Social insects, including bees and ants, show extensive phenotypic plasticity. The

morphologically and behaviorally differentiated social castes such as queens, workers, and

drones have the same set of diploid or haploid genome. The social insects provide us with model

system to study how phenotypic diversity is regulated (Page et al. 2012; Yan et al. 2014). RNA

editing is well suited to contribute to the behavioral variation among genetically similar

individuals. Indeed, differential editing was demonstrated for a few sites in leaf-cutting ants

and in worker bumblebees (Li et al. 2014; Porath et al. 2019).

Here we wish to study the contribution of RNA editing to caste differentiation in honeybee

(*Apis mellifera*), an important pollinator and a model for complicated social behaviors in insects

(Page et al. 2012). Unlike bumblebees, honeybees have a sharp task-specialization, with distinct

worker castes. Genetic mapping demonstrated that the phenotypic plasticity in honeybees is

associated with complex epistatic and pleiotropic genetic networks that influence reproductive

regulation and foraging behaviors (Page et al. 2012). We wish to check whether A-to-I RNA

editing may contribute to the proteomic diversity underlying this phenotypic plasticity. We

investigate the A-to-I RNA editome in different tissues of four honeybee drone individuals and

detect over four hundred A-to-I sites in multiple tissues. Editing is enriched in the head, and

exhibit signs for positive selection and a particularly high *N/S* ratio. We show editing is elevated

in foragers compared to nurses. Five editing sites are conserved between honeybee, bumblebee

and *Drosophila*. Interestingly, we find a significantly high number of cases where the same

gene in edited in both bees and flies, even if not at the same position. One example is the auto-

editing of *Adar* mRNA in bees and flies, which might play an auto-regulatory role in the two

clades. This finding may further support possible convergent evolution and adaptation.

RESULTS

Editome of honeybee

We deep-sequenced genomic DNA and RNA from the head, thorax, and abdomen for each

of four individual drones (12 RNA-seq samples) (Figure 1A). The transcriptome libraries of

drones 1 and 2 were constructed by selecting polyA tailed mRNAs while the libraries of drones

3 and 4 were constructed by using the Ribo-Zero kit to deplete ribosomal RNAs (Transparent

Methods). The male drone is haploid, simplifying identification of its genomic SNPs. DNA-

Seq reads were mapped to the reference genome to identify 826,890-858,949 SNPs in each of

the four drones, and 1,303,225 unique sites combined (Transparent Methods; Table S1). Median

sequencing coverage at the SNP sites was 39, and in 96.6% of the sites all DNA reads supported

the SNP (Supplemental Fig S1). The identified SNPs were then used to produce a masked

genome version where the reference allele is substituted by the individual-specific alleles

identified for each drone, to facilitate accurate detection of RNA editing sites (Figure 1B;

Transparent Methods).

For each of the 12 RNA-Seq samples, 9.2-21.5 million reads were uniquely mapped to the

reference genome, and similar numbers of reads were mapped to the masked genome sequences

(Transparent Methods; Table S2). Analyzing pooled data for each of the three tissues separately,

we identified (FDR=0.05) 376 (84.3%) A-to-G sites among 446 variations in heads, 106 (66.7%)

A-to-G sites among 159 variations in thoraxes, and 137 (67.5%) A-to-G sites among 203

variations in abdomens (Figure S2). Combined, we obtained 407 (80.1%) unique A-to-G sites

among 508 variations (Figure 1C). These 407 sites are regarded as A-to-I RNA editing sites

(Table S3, Information of the candidate editing sites identified in this study, Related to Figure

1). The nucleotide context around these putative editing sites is consistent with the known

ADAR binding motif (Figure S3). The median DNA coverage (over the 407 A-to-I editing sites)

was 25 reads, sufficient to exclude possible SNPs (Figure S4).

To improve mapping accuracy, we discarded variants in repeat regions, and required a

variation site to be found by mapping to both the reference genome and the masked genome

(Transparent Methods). To test whether these criteria are too stringent, we checked the variation

sites discarded by the two filtering steps. First, we looked at the variation sites located in the

repeat regions. Without any filter, we found 19,573 unique variation sites that overlapped with

repeat regions, only 3,694 (19%) of which were A-to-G variants. Following a binomial test and

multiple testing correction (to remove random sequencing errors), 1,277 variation sites were

maintained, only 324 (25%) of them A-to-G variations. Thus, filtering repetitive regions does

contribute appreciably to the precision of our detection. Consistently, hyper-editing analysis did

not reveal multiple sites in repetitive regions (Transparent Methods).

As for the masked genome filter, only 38% (98/256) of the variation sites found by

mapping to the masked genome but not to the reference genome are A-to-G mismatches.

Following binomial test for sequencing error and multiple testing correction for P values, only

41 variation sites were maintained, none of them was an A-to-G variation. Conversely, only 31%

(82/268) variation sites found by mapping to the reference genome but not to the masked

genome are A-to-G mismatches. Of these 268 variation sites, 82 were maintained after multiple

testing correction, 20 of them were A-to-G variations. Thus, most of the sites supported by only

one method of mapping are likely not due to A-to-I RNA editing.

Among the 407 editing sites we identified, 199 sites are located in gene regions, including

111 nonsynonymous (*N*) sites, 9 synonymous (*S*) sites, and several sites in UTRs and introns

(Figure 1D). The other 208 sites are annotated as intergenic. As expected from the different

library construction strategies, the fraction of sites in coding regions is higher and the intergenic

fraction is lower for sites observed in drones 1 and 2 compared with drones 3 and 4 ($23.2 \pm 2.6\%$

in CDS in drones 1 and 2 versus $14.5 \pm 5.1\%$ in CDS in drones 3 and 4; $54.9 \pm 2.2\%$ in intergenic

regions in drones 1 and 2 versus 70.5 ± 5.4% in intergenic regions in drones 3 and 4). The

number of sites for which editing is observed in each sample (per-sample editing level > 0)

varies considerably (40-278 sites per sample; Table S4). The partial overlap between these sets

of sites is mostly due to sites being undetected at a given sample due to low per-sample coverage

(Figure S5).

Interestingly, editing seems to be enriched in heads. The number of detected sites (Figure

S6), the editing levels (Figure 1E) and the fraction of coding sites (Figure S7), are all higher in

heads. Clustering the samples by their editing profile, the three tissue-types form distinct

clusters (Figure 1F). Consistently, *Adar* expression is also higher in heads than other tissues

(Figure S8). Looking at specific sites, 78 sites (out of 378 sites with sufficient coverage, see

Transparent Methods) are differentially-edited in the head, compared to non-head tissues. In all

of these 78 sites, editing is higher in the head (Table S3, Information of the candidate editing

sites identified in this study, Related to Figure 1).

Signals of adaptation

The nonsynonymous to synonymous (N/S) ratio for sites in the coding region is $111/9 = 12.3$

(Table S4), compared to a ratio of 2.26 observed for random adenosine to guanosine

substitutions (see Transparent Methods). This strongly suggests that the nonsynonymous editing

events in honeybees are overall adaptive. Notably, the N/S is exceptionally high, even compared

to the ratios we previously found in the brains of *Drosophila* (Duan et al. 2017) or in

cephalopods (Alon et al. 2015; Liscovitch-Brauer et al. 2017). Looking at each sample separately,

at pooled tissue data, or focusing on sites that appear in at least two individuals per tissue all

show the N/S ratio to be especially high in heads (Figure 2A and Table S4). Furthermore, the

editing level at nonsynonymous sites is significantly higher than synonymous sites editing

levels in heads of honeybees (Figure 2B).

A previous study in cephalopods proposed that the coleoids massively edit their RNAs to

diversify the transcriptome at the cost of constraining the evolution of genomic sequences

(Liscovitch-Brauer et al. 2017). Maintaining beneficial editing requires genomic conservation

of the genomic sequence encoding for the dsRNA structures that allow ADAR to bind and

deaminate the adenosine. To test this, we first confirmed that the editing sites in honeybees are

enriched in hairpin structures (Figure 2C). Then, we demonstrated that SNPs are depleted in

the vicinity of nonsynonymous sites (Figure 2D), further supporting the notion of selective

advantage for the nonsynonymous editing events that justifies the tradeoff between

transcriptome diversity and genome evolution.

Evolutionarily conserved and non-conserved editing sites

Editing at four nonsynonymous and one synonymous site is conserved between

Drosophila and honeybees (Figure 3). Three of these sites are in the *Shab* transcript (*shaker*

cognate b, two nonsynonymous and one synonymous) and the other two are is in *qvr* (*quiver*).

Editing at these sites is also observed in the bumblebee *Bombus terrestris* (Porath et al. 2019),

and editing levels are high in all of these species (Figure 3). These findings suggest a potential

functional role for these widely conserved editing sites. However, the majority of the editing

sites are not shared between *Drosophila* and honeybees. For some sites the editable adenosine

is not conserved in the honeybee genome (Figure S9), and are thus clearly uneditable, while

other adenosines are conserved but editing was not detected in RNA-Seq data (Figure S9).

Overall, of the 639 *Drosophila* editing sites in CDS for which an orthologous site was identified

in honeybees, 349 were conserved as A (55%), and 85, 127, and 78 (13%, 20%, and 12%) were

mutated to C, G and T, respectively.

Similarly, most sites are not even conserved across the two bee species. The bumblebee

study has reported 219 editing sites in CDS, 164 of which are nonsynonymous (Porath et al.

2019). Of these, we found 9 editing sites in coding regions conserved between honeybee and

bumblebee (Table S5).

One of these conserved sites is a Ser>Gly site within the *tipE* (temperature-induced

paralytic E; *GB47375*) transcript, editing levels of which correlate with task-performance in

bumblebees (nurses versus foragers) (Porath et al. 2019). Editing level at this site in honeybee

heads are ~0.7, similar to those observed in bumblebee brains. In the two other tissues this gene

is lowly expressed and poorly edited. The orthologous site in *Drosophila* genomes encodes the

post-edit Gly codon GGC (Figure S10). Thus, *tipE* recoding seems to be bee-specific, possibly

related to social behavior and task performance.

Sex-dependent and Caste-dependent editing

We further looked for potential sex-dependence and caste-dependent editing, comparing our honeybee drone head data with previously published brain RNA-seq data studying two sub-castes of female workers, foragers and reverted nurses (Herb et al. 2012) (Table S6). First, we compared the pooled editing levels between drones and workers for each well-covered (>10 reads in each pool) editing site separately. Twenty six sites exhibit differential editing (Wilcoxon rank sum test; FDR=0.05) between workers (females) and drones (males), including 4 recoding sites, 2 sites in 3'UTRs, 3 intronic sites, and 17 intergenic sites (Figure 4A and Table S7, Differentially edited sites between drones and workers, Related to Figure 4). In most cases editing in drones is higher, but this may result from looking at sites identified in drone data to begin with. The difference in *Adar* expression was not statistically significant (Figure S11).

Comparing the pooled editing levels between the two sub-castes (foragers and nurses), one finds 230 out of 407 sites with lower levels in nurses, compared to only 91 exhibiting higher levels in nurses (equal level, mostly zero, was observed in 84 sites), suggesting a globally higher editing activity (at the detected sites) in foragers (P = 1.3e-14, proportion test). The surcharge of forager-higher sites is maintained for various coverage and editing-difference cutoffs (Figure S12). Furthermore, *Adar* expression (Figure 4B) and the editing index (Figure 4C) are both higher in foragers, and the editing index correlates with *Adar* expression. However, reliable detection of specific differentially-edited sites with the available sample size (6 foragers and 6 nurses) is challenging, and not even a single site was identified with FDR=0.05 (Wilcoxon rank sum test). We have thus relaxed the statistical test, allowing for FDR=0.3, and found 61 candidate sites, most of which are differentially edited (Figure 4D and Table S8, Differentially edited sites between foragers and nurses, Related to Figure 4). These include 10 nonsynonymous, 16 3'UTR, and 35 intergenic sites. Consistently, 55 of these sites exhibit higher levels of editing in foragers. Finally, PCA analysis of the editing profile across sites results in distinct clusters of drones, nurses and foragers, a classification that is not achieved by the expression profile (Figure 4E and 4F). Taken together, these results show a global increase in editing in foragers compared to nurses.

Convergent adaptation of A-to-I editing?

One of the recoding sites conserved between the two bee species resides within *Adar*

transcript (Table S5 and Figure 5A). In bumblebee, the recoding level of the conserved site,

I482M (ATA to ATG), positively correlates with the global editing activity (Porath et al. 2019),

suggesting a possible auto-regulation mechanism. Interestingly, *Drosophila Adar* is also auto-

edited at a different position, where a Ser (AGT) to Gly (GGT) substitution leads to a less active

ADAR protein, resulting in a negative feedback loop of editing activity (Palladino et al. 2000;

Savva et al. 2012a). Intriguingly, fly-edited Ser amino-acid is conserved in honeybee and

bumblebee, but a different codon (TCA, uneditable at the first codon position) is used. Similarly,

the bee-edited Ile is conserved in flies, but the edited adenosines (at the third position of ATA)

is synonymously mutated to T, and the editable ATA bee codon is substituted by ATT codon in

flies, which could not be edited at the third position (Figure 5A). Thus, while auto-editing

S430G in flies is abolished in bees due to an uneditable Ser codon, and the auto-editing I482M

in bees is abolished in flies due to an uneditable Ile codon. However, the mechanism of auto-

editing, possibly used for global ADAR regulation, is shared by the two lineages, albeit at

different positions and possibly with different effects on the protein. This might hint at the

possibility of convergent evolution of ADAR auto-regulation strategy.

Following this example, we wondered whether there are more genes for which editing is

observed in both honeybees and *Drosophila*, even if the exact location of the editing site is not

conserved. Excluding conserved sites, there are 53 genes exhibiting CDS editing in honeybee

heads, 101 genes exhibiting CDS editing in bumblebee heads, and 312 genes edited in CDS in

male brains of *D. melanogaster*. Of these, 21 genes exhibit editing in both bee species, 14 are

edited in honeybee and *Drosophila,* 14 are edited in bumblebee and *Drosophila*, and six genes

are found to be edited in all three species (Figure 5B and Table S9). These numbers are

significantly higher than expected by random sampling, (p<1e-6, randomization test), even if

one accounts for the expression profile (Transparent Methods, Figure 5C and Figure S13). Table

S9). We have repeated the analysis using stricter criteria to characterize sites, considering as

edited only sites with an observed editing level >10% (all samples pooled), and a binomial $P <$

0.05 (excluding the null hypothesis of no editing), and considering a site to be not-edited if the

binomial $P < 0.05$ for the null hypothesis that the site is edited at 10%. Reassuringly, the results

are robust to this change of definitions (Figure S14 and S15). These results suggest an

interesting convergent evolution of A-to-I editing. Recoding sites are rarely conserved across

clades, but the same genes in different lineages tend to acquire a recoding event, apparently

independently, indicating a functional importance for recoding of this gene.

DISCUSSION

Despite over fifteen years of developments, reliable detection of CDS editing remains a

challenge. Systematic searches often lead to high false-positive rates (reflected by the fraction of

non-AG mismatches observed), especially for mammalian transcriptomes where the scope of

recoding is rather low. Important exception are *Drosophila* and cephalopods, where the recoding

signal is much more pronounced and easier to detect. Applying a novel strict alignment approach,

we were able to achieve here a high accuracy in CDS editing detection (84.3% in heads) despite the

overall modest scope of recoding sites. Another similarity between honeybees *Drosophila* and

cephalopods is related to the important question regarding the extent to which recoding is adaptive.

This is often estimated by the *N/S* ratio, where *Drosophila* and cephalopods exhibit a pattern

markedly different from the one seen in other species studied, (including human). Whereas in most

species the *N/S* ratio observed is lower or similar to the one expected under neutrality, it is

significantly higher in *Drosophila* and cephalopods. Again, we find here that honeybees show a

strikingly high *N/S* ratio, similar to the above two clades. We suggest that the two last points are

interconnected. The apparently low *N/S* ratio observed in many species (human included) may

reflect the low accuracy of the lists of CDS editing sites and not the (lack of) adaptive potential for

recoding in these species.

The importance of post-transcriptional and post-translational mechanisms in generating the

proteomic complexity of higher organisms has been emphasized in recent decades. These epigenetic

mechanisms allow for diversification of the proteome and functional heterogeneity across tissues,

developmental stages, brain regions or even among individual cells.

Recoding by A-to-I RNA editing is an epigenetic mechanism capable of diversifying the

proteome, creating a range of proteins from a single genomically encoded gene in a temporally-

regulated, tissue-specific, condition-dependent way, and providing the organism with a new means

for acclimation and adaptation. Indeed, several studies have demonstrated how recoding levels at

specific sites do change as a function of the organism's condition (Garrett and Rosenthal 2012;

Robinson et al. 2016; Gallo et al. 2017; Porath 2017; Terajima et al. 2017; Yablonovitch et al. 2017). Importantly, many studies have demonstrated altered editing of individual recoding targets in various disease states (Gallo et al. 2017). However, these interesting examples notwithstanding, the extent to which the recoding phenomenon is actually used as a means for proteome diversification is still under debate. In fact, several recent studies have raised the possibility that nonsynonymous editing may be used to compensate for otherwise deleterious G-to-A mutations, rather than allowing for the two alleles (A and G) to co-exist (Jiang and Zhang 2019; Mai and Chuang 2019; Popitsch et al. 2020). Thus, a high N/S ratio is not sufficient to prove that editing serves for adaptation through proteome diversification.

The behavioral, physiological and morphological plasticity exhibited by social insects suggests an important role for epigenetic mechanisms. Differential RNA editing was previously shown for different castes of the highly social leaf-cutting ant *Acromyrmex echinatior* (Li et al. 2014), as well as for bumblebee workers (Porath et al. 2019). Our results expand these finding to another species, further supporting the notion of RNA editing as a source of proteomic complexity, that may be recruited to provide phenotypic variation among genetically identical individuals.

ADAR enzymes, and the recoding phenomenon are widely conserved across metazoan. However, the repertoire of recoding sites varies considerably and seems to have developed almost independently in different clades. Here we point out that the same gene targets are being edited in distant species, even if the exact locations of the editing sites within the transcript vary. A particularly interesting example is that of nonsynonymous auto-editing of *Adar*, shown to affect ADAR activity in bumblebees, flies and even mammals (Rueter et al. 1999; Savva et al. 2012a; Porath et al. 2019), possibly serving as global editing autoregulation. It thus seems possible that the selective advantage in having multiple versions of the protein product is shared by the target genes, while there is more than one specific way to achieve this diversity through recoding. Clades developed along different evolutionary routes have converged on different implementations (in terms of the specific editing site) of the same diversifying solution. These ideas should be further tested in the future, using larger datasets and across additional species and clades. Moreover, as biochemical and functional understanding of the impact of the different edits are gained, it will be possible to look into a possible common effect for the different edits of the common targets.

Limitations of the Study

The editomes of the bee species analyzed here are based on a limited number of samples. They probably represent a subset of the actual repertoire of editing sites. The results should be re-tested when more data is available. In particular, the honeybee editome was built using drone samples. Thus, some individual differential editing between the nurses and foragers sub-castes may have missed worker-specific sites. However, the global increased editing in foragers, supported by a higher Adar expression, is probably robust.

Resource Availability

Lead Contact

Further information and requests for resources should be directly and will be fulfilled by the Lead Contact, Jian Lu (luj@pku.edu.cn).

Materials Availability

This study did not generate new unique materials.

Data and Code Availability

All deep-sequencing data generated in this study were deposited in the China National Genomics Data Center Genome Sequence Archive (GSA) under accession number CRA002262. Other deep-sequencing data analyzed were downloaded from SRA as follows. Brains of honeybee workers: accession numbers SRR445999 to SRR446004 (reverted nurse) and SRR446005 to SRR446010 (forager) (Herb et al. 2012). Bumblebee: SRP166322 (Porath et al. 2019). *D. mel* head: SRP067542 (Zhang et al. 2018). *D. sim* head: SRP074828 (Duan et al. 2017). *D. pse*: DRR055250 and DRR055251 (Nozawa et al. 2016). *D. wil*: SRR341127 and SRR341129 (Meisel et al. 2012). *D. moj*: SRR037508-SRR037518 (generated by the *Drosophila* modENCODE project). *D. sec*, and *D. ere* data are unpublished data generated in our own lab. The reference genomes versions used for *Drosophila* species are Dmel_r6.04 (http://flybase.org/) for *D. mel*, and versions droSim1, droSec1, droEre1, dp4, droWil1, and droMoj3 downloaded from UCSC Genome Browser (http://genome.ucsc.edu/) for the other fly species. For the *Bombus terrestris*, we used version Bter_1.0.

METHODS

All methods can be found in the accompanying Transparent Methods supplemental file.

ACKNOWLEDGMENTS

We thank the National Center for Protein Sciences at Peking University for technical assistance. Part of the analysis was performed on the High Performance Computing Platform of the Center for Life Science at Peking University. This work was supported by the joint NSFC-ISF program (NSFC grant number 3201101147 to J.L. and ISF grant number 3371/20 to E.E.), as well as grants from the Ministry of Science and Technology of the People's Republic of China (2016YFA0500800) and National Natural Science Foundation of China (91731301) to J.L. and support from the Israel Science Foundation (1945/18) to E.E.

AUTHOR CONTRIBUTIONS

J.L. designed the research; J.X.H. contributed the honeybee materials; S.Q.D. performed the research; Y.G.D., H.T.P, E.E. and J.L. performed the bioinformatics analyses; E.E. and J.L. wrote the paper.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

Alon S, Garrett SC, Levanon EY, Olson S, Graveley BR, Rosenthal JJ, Eisenberg E. 2015. The majority of transcripts in the squid nervous system are extensively recoded by A-to-I RNA editing. *Elife* **4**.

Alon S, Mor E, Vigneault F, Church GM, Locatelli F, Galeano F, Gallo A, Shomron N, Eisenberg E. 2012. Systematic identification of edited microRNAs in the human brain. *Genome Res* **22**: 1533-1540.

Athanasiadis A, Rich A, Maas S. 2004. Widespread A-to-I RNA editing of Alu-containing mRNAs in the human transcriptome. *PLoS Biol* **2**: e391.

Basilio C, Wahba AJ, Lengyel P, Speyer JF, Ochoa S. 1962. Synthetic polynucleotides and the amino acid code. V. *Proc Natl Acad Sci U S A* **48**: 613-616.

Bass BL. 2002. RNA editing by adenosine deaminases that act on RNA. *Annu Rev Biochem* **71**: 817-846.

Blow M, Futreal PA, Wooster R, Stratton MR. 2004. A survey of RNA editing in human brain. *Genome Research* **14**: 2379-2387.

Borchert GM, Gilmore BL, Spengler RM, Xing Y, Lanier W, Bhattacharya D, Davidson BL. 2009. Adenosine deamination in human transcripts generates novel microRNA binding sites. *Human Molecular Genetics* **18**: 4801-4807.

Buchumenski I, Bartok O, Ashwal-Fluss R, Pandey V, Porath HT, Levanon EY, Kadener S. 2017. Dynamic hyper-editing underlies temperature adaptation in Drosophila. *PLOS Genetics* **13**: e1006931.

Chen J-Y, Peng Z, Zhang R, Yang X-Z, Tan BC-M, Fang H, Liu C-J, Shi M, Ye Z-Q, Zhang YE et al. 2014. RNA Editome in Rhesus Macaque Shaped by Purifying Selection. *PLoS Genet* **10**: e1004274.

Danecek P, Nellaker C, McIntyre RE, Buendia-Buendia JE, Bumpstead S, Ponting CP, Flint J, Durbin R, Keane TM, Adams DJ. 2012. High levels of RNA-editing site conservation amongst 15 laboratory mouse strains. *Genome Biol* **13**: 26.

Duan Y, Dou S, Luo S, Zhang H, Lu J. 2017. Adaptation of A-to-I RNA editing in Drosophila. *PLoS Genet* **13**: e1006648.

Eisenberg E, Levanon EY. 2018. A-to-I RNA editing — immune protector and transcriptome diversifier. *Nature Reviews Genetics* **19**: 473-490.

Flomen R, Knight J, Sham P, Kerwin R, Makoff A. 2004. Evidence that RNA editing modulates splice site selection in the 5‐HT2C receptor gene. *Nucleic Acids Research* **32**: 2113-2122.

Gallo A, Vukic D, Michalik D, O'Connell MA, Keegan LP. 2017. ADAR RNA editing in human disease; more to it than meets the I. *Hum Genet* **136**: 1265-1278.

Garrett S, Rosenthal JJ. 2012. RNA editing underlies temperature adaptation in K+ channels from polar octopuses. *Sci* **335**: 848-851.

Goldstein B, Agranat-Tamir L, Light D, Ben-Naim Zgayer O, Fishman A, Lamm AT. 2017. A-to-I RNA editing promotes developmental stage–specific gene and lncRNA expression. *Genome Research* **27**: 462-470.

Gommans WM, Mullen SP, Maas S. 2009. RNA editing: a driving force for adaptive evolution? *BioEssays* **31**: 1137-1145.

Graveley BR, Brooks AN, Carlson JW, Duff MO, Landolin JM, Yang L, Artieri CG, van Baren MJ, Boley N, Booth BW et al. 2011. The developmental transcriptome of Drosophila melanogaster. *Nature* **471**: 473-479.

Herb BR, Wolschin F, Hansen KD, Aryee MJ, Langmead B, Irizarry R, Amdam GV, Feinberg AP. 2012. Reversible switching between epigenetic states in honeybee behavioral subcastes. *Nat Neurosci* **15**: 1371-1373.

Jiang D, Zhang J. 2019. The preponderance of nonsynonymous A-to-I RNA editing in coleoids is

nonadaptive. *Nat Commun* **10**: 5411.

Jin Y, Tian N, Cao J, Liang J, Yang Z, Lv J. 2007. RNA editing and alternative splicing of the insect nAChR
subunit alpha6 transcript: evolutionary conservation, divergence and regulation. *BMC*
Evolutionary Biology **7**: 1-12.

Keegan LP, Gallo A, O'Connell MA. 2001. The many roles of an RNA editor. *Nat Rev Genet* **2**: 869-878.

Keegan LP, McGurk L, Palavicini JP, Brindle J, Paro S, Li X, Rosenthal JJ, O'Connell MA. 2011. Functional
conservation in human and Drosophila of Metazoan ADAR2 involved in RNA editing: loss of
ADAR1 in insects. *Nucleic Acids Res* **39**: 7249-7262.

Kim DD, Kim TT, Walsh T, Kobayashi Y, Matise TC, Buyske S, Gabriel A. 2004. Widespread RNA editing of
embedded alu elements in the human transcriptome. *Genome Res* **14**: 1719-1725.

Klironomos FD, Berg J, Collins S. 2013. How epigenetic mutations can affect genetic evolution: Model
and mechanism. *BioEssays* **35**: 571-578.

Lev-Maor G, Sorek R, Levanon EY, Paz N, Eisenberg E, Ast G. 2007. RNA-editing-mediated exon evolution.
Genome biology **8**: R29.

Levanon EY, Eisenberg E, Yelin R, Nemzer S, Hallegger M, Shemesh R, Fligelman ZY, Shoshan A, Pollock
SR, Sztybel D et al. 2004. Systematic identification of abundant A-to-I editing sites in the human
transcriptome. *Nat Biotech* **22**: 1001-1005.

Li Q, Wang Z, Lian J, Schiott M, Jin L, Zhang P, Zhang Y, Nygaard S, Peng Z, Zhou Y et al. 2014. Caste-
specific RNA editomes in the leaf-cutting ant Acromyrmex echinatior. *Nat Commun* **5**: 4943.

Liang H, Landweber LF. 2007. Hypothesis: RNA editing of microRNA target sites in humans? *RNA* **13**:
463-467.

Licht K, Hartl M, Amman F, Anrather D, Janisiw MP, Jantsch MF. 2019. Inosine induces context-
dependent recoding and translational stalling. *Nucleic Acids Res* **47**: 3-14.

Liscovitch-Brauer N, Alon S, Porath HT, Elstein B, Unger R, Ziv T, Admon A, Levanon EY, Rosenthal JJC,
Eisenberg E. 2017. Trade-off between Transcriptome Plasticity and Genome Evolution in
Cephalopods. *Cell* **169**: 191-202 e111.

Mai TL, Chuang TJ. 2019. A-to-I RNA editing contributes to the persistence of predicted damaging
mutations in populations. *Genome Res* **29**: 1766-1776.

Mazloomian A, Meyer IM. 2015. Genome-wide identification and characterization of tissue-specific RNA
editing events in D. melanogaster and their potential role in regulating alternative splicing. *RNA*
Biology **12**: 1391-1401.

Meisel RP, Malone JH, Clark AG. 2012. Disentangling the relationship between sex-biased gene
expression and X-linkage. *Genome Res* **22**: 1255-1265.

Morse DP, Bass BL. 1999. Long RNA hairpins that contain inosine are present in Caenorhabditis elegans
poly(A)+ RNA. *Proc Natl Acad Sci U S A* **96**: 6048-6053.

Neeman Y, Levanon EY, Jantsch MF, Eisenberg E. 2006. RNA editing level in the mouse is determined by
the genomic repeat repertoire. *RNA* **12**: 1802-1809.

Nishikura K. 2010. Functions and regulation of RNA editing by ADAR deaminases. *Annu Rev Biochem* **79**:
321-349.

Nishikura K. 2016. A-to-I editing of coding and non-coding RNAs by ADARs. *Nat Rev Mol Cell Biol* **17**: 83-
96.

Nozawa M, Onizuka K, Fujimi M, Ikeo K, Gojobori T. 2016. Accelerated pseudogenization on the neo-X
chromosome in Drosophila miranda. *Nat Commun* **7**: 13659.

Page RE, Jr., Rueppell O, Amdam GV. 2012. Genetics of reproduction and regulation of honeybee (Apis
mellifera L.) social behavior. *Annu Rev Genet* **46**: 97-119.

Palladino MJ, Keegan LP, O'Connell MA, Reenan RA. 2000. dADAR, a Drosophila double-stranded RNA-
specific adenosine deaminase is highly developmentally regulated and is itself a target for RNA
editing. *RNA* **6**: 1004-1018.
Picardi E, D'Erchia AM, Lo Giudice C, Pesole G. 2017. REDIportal: a comprehensive database of A-to-I
RNA editing events in humans. *Nucleic Acids Res* **45**: D750-D757.
Popitsch N, Huber CD, Buchumenski I, Eisenberg E, Jantsch M, von Haeseler A, Gallach M. 2020. A-to-I
RNA Editing Uncovers Hidden Signals of Adaptive Genome Evolution in Animals. *Genome Biol*
Evol **12**: 345-357.
Porath HT. 2017. A-To-I RNA editing in the earliest-diverging eumetazoan phyla. *Mol Biol Evol* **34**.
Porath HT, Hazan E, Shpigler H, Cohen M, Band M, Ben-Shahar Y, Levanon EY, Eisenberg E, Bloch G. 2019.
RNA editing is abundant and correlates with task performance in a social bumblebee. *Nature*
Communications **10**.
Porath HT, Knisbacher BA, Eisenberg E, Levanon EY. 2017a. Massive A-to-I RNA editing is common across
the Metazoa and correlates with dsRNA abundance. *Genome Biol* **18**: 185.
Porath HT, Schaffer AA, Kaniewska P, Alon S, Eisenberg E, Rosenthal J, Levanon EY, Levy O. 2017b. A-to-I
RNA Editing in the Earliest-Diverging Eumetazoan Phyla. *Molecular Biology and Evolution* **34**:
1890-1901.
Ramaswami G, Li JB. 2016. Identification of human RNA editing sites: A historical perspective. *Methods*
107: 42-47.
Robinson JE, Paluch J, Dickman DK, Joiner WJ. 2016. ADAR-mediated RNA editing suppresses sleep by
acting as a brake on glutamatergic synaptic plasticity. *Nature Communications* **7**.
Rodriguez J, Menet JS, Rosbash M. 2012. Nascent-seq indicates widespread cotranscriptional RNA
editing in Drosophila. *Mol Cell* **47**: 27-37.
Rosenthal JJC. 2015. The emerging role of RNA editing in plasticity. *The Journal of Experimental Biology*
218: 1812.
Rueter SM, Dawson TR, Emeson RB. 1999. Regulation of alternative splicing by RNA editing. *Nature* **399**:
75-80.
Savva YA, Jepson JE, Sahin A, Sugden AU, Dorsky JS, Alpert L, Lawrence C, Reenan RA. 2012a. Auto-
regulatory RNA editing fine-tunes mRNA re-coding and complex behaviour in Drosophila. *Nat*
Commun **3**: 790.
Savva YA, Rieder LE, Reenan RA. 2012b. The ADAR protein family. *Genome Biol* **13**.
St Laurent G, Tackett MR, Nechkin S, Shtokalo D, Antonets D, Savva YA, Maloney R, Kapranov P, Lawrence
CE, Reenan RA. 2013. Genome-wide analysis of A-to-I RNA editing by single-molecule
sequencing in Drosophila. *Nat Struct Mol Biol* **20**: 1333-1339.
Terajima H, Yoshitane H, Ozaki H, Suzuki Y, Shimba S, Kuroda S, Iwasaki W, Fukada Y. 2017. ADARB1
catalyzes circadian A-to-I editing and regulates RNA rhythm. *Nat Genet* **49**: 146-151.
Xu G, Zhang J. 2014. Human coding RNA editing is generally nonadaptive. *Proc Natl Acad Sci U S A* **111**:
3769-3774.
Yablonovitch AL, Fu J, Li K, Mahato S, Kang L, Rashkovetsky E, Korol AB, Tang H, Michalak P, Zelhof AC et
al. 2017. Regulation of gene expression and RNA editing in Drosophila adapting to divergent
microclimates. *Nature Communications* **8**: 1570.
Yan H, Simola DF, Bonasio R, Liebig J, Berger SL, Reinberg D. 2014. Eusocial insects as emerging models
for behavioural epigenetics. *Nat Rev Genet* **15**: 677-688.
Yang X-Z, Chen J-Y, Liu C-J, Peng J, Wee YR, Han X, Wang C, Zhong X, Shen QS, Liu H et al. 2015. Selectively
Constrained RNA Editing Regulation Crosstalks with piRNA Biogenesis in Primates. *Molecular*

Biology and Evolution **32**: 3143-3157.

Yu Y, Zhou H, Kong Y, Pan B, Chen L, Wang H, Hao P, Li X. 2016. The Landscape of A-to-I RNA Editome Is
Shaped by Both Positive and Purifying Selection. *PLoS Genet* **12**: e1006191.

Zhang H, Dou S, He F, Luo J, Wei L, Lu J. 2018. Genome-wide maps of ribosomal occupancy provide
insights into adaptive evolution and regulatory roles of uORFs during Drosophila development.
PLoS Biol **16**: e2003903.

Zhang R, Deng P, Jacobson D, Li JB. 2017. Evolutionary analysis reveals regulatory and functional
landscape of coding and non-coding RNA editing. *PLoS genetics* **13**: e1006563.

Zhao HQ, Zhang P, Gao H, He X, Dou Y, Huang AY, Liu XM, Ye AY, Dong MQ, Wei L. 2015. Profiling the RNA
editomes of wild-type C. elegans and ADAR mutants. *Genome Res* **25**: 66-75.

Figure legends

Figure 1. Identification and annotation of the A-to-I RNA editing sites in honeybee. (A)

Workflow of sample collection, dissection and library construction. (B) Identification of A-to-

I RNA editing sites is facilitated by mapping to both the reference genome and the masked

genome. (C) Distribution of DNA-RNA mismatch types detected in honeybee. (D) Distribution

of the detected A-to-I editing sites over different genic regions. (E) Editing levels at the detected

as measured in three tissues of each of four individuals (D1-D4). Editing levels are higher in

the head (head versus pooled thorax and abdomen, Wilcoxon rank sum test; ***: $P < 0.001$.

Exact P values, left to right: 2.1e-35, 5.7e-20, 4.0e-29, and 3.9e-38). (F) The editing profiles

for the twelve tissues cluster according to their tissue of origin.

Figure 2. Signals of adaptation of A-to-I editing in honeybee. (A) The observed numbers of

nonsynonymous (N) and synonymous (S) editing sites in each tissue. P values are obtained from

Fisher's exact test, comparing the observed N and S counts to all adenosines in the coding

sequence of in *A. mel* (N=4,492,737 sites where A-to-G substitution is nonsynonymous, and

S=1,986,130 for synonymous). "All sites" represents the numbers of sites that appear in at least

one individual. Comparisons for sites appearing in at least two individuals are also shown. (B)

Editing levels of the nonsynonymous and synonymous sites in head. Data from four drones was

pooled to increase the statistical power. P values are calculated using Wilcoxon rank sum test.

(C) The fraction of sites in predicted stable hairpin structures. The observed editing sites were

compared to the unedited adenosines (P-values using Fisher's exact test). (D) The SNP density

in the vicinity of nonsynonymous editing sites (green), other editing sites (cyan) or unedited adenosines (grey).

Figure 3. Editing sites conserved between bees and flies. (A) One synonymous and two nonsynonymous editing sites in *Shab* transcripts are edited in heads of two bee and four *Drosophila* species. (B) Two nonsynonymous editing sites in *qvr* transcripts are edited in heads of two bee and four *Drosophila* species. The editing sites are colored red. Editing levels measured in male heads of flies (unpublished), pooled heads of honeybee drones, and brains of bumblebees (Porath et al. 2019) are shown. *D. mel*, *Drosophila melanogaster*; *D. sim*, *Drosophila simulans*; *D. sec*, *Drosophila sechellia*; *D. ere*, *Drosophila erecta; D. pse, Drosophila pseudoobscura; D. wil, Drosophila willistoni; D. moj, Drosophila mojavensis; A. mel*, *Apis mellifera*; *B. ter*, *Bombus terrestris*.

Figure 4. Sex-dependent and caste-dependent editing. (A) Comparison of editing levels between drones (males) and workers (females). Statistically significant sites (FDR < 0.05) are colored red. (B) *Adar* expression level (reads count normalized by DESeq2) in nurses and foragers. Data are presented as mean ± SEM (standard error of mean). *P* value was calculated using Wilcoxon rank sum test. (C) Spearman's correlation between *Adar* expression and editing index (sum over all G alleles in all sites divided by sum of all the coverages in all sites) for all the nurse (orange) and forager (purple) samples. (D) Comparison of editing levels between two sub-castes of workers: nurses and foragers. Statistically significant sites (FDR < 0.3; see Results) are colored red. (E) PCA analysis of the editing profile shows a different behavior for nurses and foragers, while (F) PCA of the expression profile (genes with RPKM>1) does not lead to a clear separation.

Figure 5. Convergent evolution of editing. (A) *Adar* transcripts are auto-edited in *Drosophila* and bees. The Ser>Gly site is highly conserved across *Drosophila* species. The *Drosophila*-editable Ser codon AGT is changed to an uneditable Ser codon TCA in bees. On the other hand, the Ile>Met auto-editing site is conserved between honeybee and bumblebee. The bee-editable Ile codon ATA (edited at the third position) appears as an uneditable Ile codon ATT in flies.

Editing sites are colored red. The editing levels from male heads of flies (unpublished), pooled

heads of honeybee drones and brains of bumblebee (Porath et al. 2019) are shown. *D. mel*,

Drosophila melanogaster; *D. sim*, *Drosophila simulans*; *D. sec*, *Drosophila sechellia*; *D. ere*,

Drosophila erecta; *A. mel*, *Apis mellifera*; *B. ter*, *Bombus terrestris*. (B) Venn diagram

demonstrating the overlap between orthologous genes with editing sites in coding regions. (C)

The observed number of orthologous genes with editing in coding regions, compared to the

expected distribution. Genes with RPKM > 1 in both species were chosen to calculate the

expected numbers (results are essentially the same for other expression cutoffs, Figure S13). *P*-

values calculated by randomization test. (D) The nonsynonymous to synonymous (*N*/*S*) ratios

are significantly higher than expected, for both shared and species-specific sites (except for

species-specific sites in *B. terrestris*). Here the shared and species-specific sites refer to the sites

in shared genes and species-specific genes as defined in the above Venn diagram in (B). The

expected ratio is evaluated by calculating the effect of putative editing on all adenosines in the

coding region. *P*-values by Fisher exact test.

A

	N	*S*	*N/S*	*P* value
All sites				
Head	108	8	13.5	8.5E-10
Thorax	14	5	2.8	0.81
Abdomen	16	1	16	0.032
In ≥ 2 individuals				
Head	80	8	10	2.2E-6
Thorax	11	4	2.75	1
Abdomen	13	1	13	0.078

B

P = 0.027

Editing level

Nonsyn   Syn

C

Fraction of sites in hairpin

Edited
Unedited

P = 1.3e−5

P = 0.003

All   Nonsyn

D

SNPs per Kb

Nonsyn
Other
Control

Distance (bp)

1−20   21−50   51−100   101−300   301−1K

A

Shab (shaker cognate b)

	Tyr>Cys				Val Ile>Val		
D. mel	ATC	T**A**T	CCC	... ...	CTG	GT**A**	**A**TC
D. sim	ATC	T**A**T	CCC	... ...	CTG	GT**A**	**A**TC
D. sec	ATC	T**A**T	CCC	... ...	CTG	GT**A**	**A**TC
D. ere	ATC	T**A**T	CCC	... ...	CTG	GT**A**	**A**TC
D. pse	---	---	---		---	---	---
D. wil	ATC	T**A**T	CCC	... ...	CTG	GT**A**	**A**TC
D. moj	ATC	T**A**T	CCC	... ...	CTG	GT**A**	**A**TC
B. ter	ATC	T**A**C	CCC	... ...	CTG	GT**A**	**A**TC
A. mel	ATC	T**A**C	CCC	... ...	CTG	GT**A**	**A**TC
A. ech	ATC	TAC	CCC	... ...	CTG	GTA	ATC
		site1				site2, 3	

Editing level

	Site1	Site2	Site3
D. mel	0.85	0.77	1
D. sim	0.88	0.67	0.96
D. sec	0.93	0.63	1
D. ere	0.98	0.59	0.97
	Exon Deletion		
D. wil	1	0.23	1
D. moj	0.92	0.45	0.96
B. ter	0.93	0.97	0.98
A. mel	0.31	0.79	0.81
A. ech	0	0	0

B

qvr (quiver)

	Ser>Gly						His>Arg	
D. mel	GAA	**A**GT	TCA	GGC	AAT	GGA	C**A**T	ATG
D. sim	GAA	**A**GT	TCA	GGC	AAT	GGA	C**A**T	ATG
D. sec	GAA	**A**GT	TCA	GGC	AAT	GGA	C**A**T	ATG
D. ere	GAA	**A**GT	TCA	GGC	AAT	GGA	C**A**T	ATG
D. pse	GAA	**A**GT	TCA	GGC	AAT	GGA	C**A**T	ATG
D. wil	GAA	**A**GT	TCA	GGC	AAT	GGA	C**A**T	ATG
D. moj	GAA	**A**GT	TCA	GGC	AAT	GGA	C**A**T	ATG
B. ter	GAA	**A**GT	ACC	GGC	ACT	GGT	C**A**C	ATG
A. mel	GAA	**A**GT	ACC	GGC	ACT	GGT	C**A**C	ATG
A. ech	GAA	**A**GT	ACC	GGC	ACT	GGT	C**A**C	ATG
		site1					site2	

Editing level

	Site1	Site2
D. mel	0.88	0.61
D. sim	0.96	0.60
D. sec	1	0.78
D. ere	0.96	0.82
D. pse	0.99	0.84
D. wil	0.98	0.80
D. moj	0.96	0.86
B. ter	0.88	0.79
A. mel	0.50	0.34
A. ech	0.87	0.67

	N	S	Observed N/S	Expected N/S	P value
Sites in genes shared by at least two species					
D. melanogaster	127	22	5.77	3.80	0.038
A. mellifera	37	3	12.33	2.26	4.5E-4
B. terrestris	76	21	3.62	2.29	0.036
Sites in species-specific edited genes					
D. melanogaster	539	117	4.61	3.80	0.031
A. mellifera	74	6	12.33	2.26	5.2E-7
B. terrestris	87	35	2.49	2.29	0.38

Nonsynonymous editing sites in honeybees were under positive selection

Differential editing may contribute to the phenotypic diversity between sub-castes

Target genes acquire editing in different clades, suggesting convergent evolution