Testing for functional organization of three-dimensional surface tilt encoding within visual cortex

Reuben Rideaux & Andrew E Welchman
Department of Psychology, Downing Street, University of Cambridge, CB2 3EB, UK
reuben.rideaux@gmail.com

ABSTRACT

Visual perception of three-dimensional (3D) structure is important for object recognition,

grasping, and manipulation. The 3D structure of a surface can be defined in terms of its slant

and tilt. Previous work has shown that slant and tilt are represented in the posterior and ventral

intraparietal sulcus of the human brain; however, it is unclear whether the representation of

these features is functionally organized within this region. Here we use phase-encoded

presentation of 3D planar surfaces with linear gradients defined by horizontal binocular

disparity while measuring fMRI activity to test whether the representation of 3D surface tilt is

functionally organized within visual cortex. We find functionally defined structures within V3A

and V7. Most notably, in one participant we find that the tilt preference is unilaterally organized

in a pinwheel-like structure, similar to those observed for orientation preference in V1, which

encompasses most of area V3A. These findings indicate that 3D orientation is functionally

organized within the human visual cortex, and the evidence suggesting the presence of a large

pinwheel-like structure indicates that this type of organization may be applied canonically

within the brain at multiple scales.

Acknowledgements: We would like to thank Hiroshi Ban for their help and insight on the

project. The work was supported by the Leverhulme Trust (ECF-2017-573), the Issac Newton

Trust (17.08(o)), and the Wellcome Trust (095183/Z/10/Z).

Competing interests: The authors declare no competing interests.

INTRODUCTION

Visual perception of three-dimensional (3D) structure is important for object recognition, grasping, and manipulation. Binocular disparity (i.e., the horizontal offset between the images seen by the left and right eyes) is a powerful cue from which 3D structure can be inferred. Considerable imaging work has progressed our understanding of the basic processing of binocular disparity information is processed in the human brain[1–6]; however, less is known about the regions that encode 3D orientation (parameterized here as slant and tilt).

Electrophysiological work has identified areas in monkey cortex that contain populations of neurons tuned to 3D orientation defined by binocular disparity, including the caudal bank of intraparietal sulcus[7–9] (CIP) and middle temporal area[10] (MT). These neurons encode 3D orientation; however, whereas the response of neurons in CIP to slant and tilt is co-dependent[9], these dimensions appear to be encoded separably in MT[10]. This work has revealed how 3D orientation is encoded at the single-cell level, but the nature of the technique precludes identification of population level trends such as functional organization within these areas. Further, differences between monkey and human brain organization in relation to the representation of 3D orientation restricts the application of these findings to human perception[11]. Here we define functional organization as any non-arbitrary representation of 3D orientation within one or more localized cortical regions.

Functional MRI work with humans has shown that the posterior and ventral intraparietal sulcus (PIPS/V3A & VIPS/V7, respectively) encode depth structure from binocular disparity[12–14]. These areas have also been implicated in processing 3D orientation defined by motion[11,15,16], texture[17], and shading[18]. Thus far, the study of how 3D orientation is represented within the human brain has been limited to identification of regions through contrasting patterns of fMRI activation. This technique has been successfully applied in the identification of brain areas that encode 3D orientation, but it has not revealed whether neural responses in these areas are functionally organized.

Here we use phase-encoded presentation of 3D stimuli while recording repeated in-
depth measurements of fMRI activity from four human participants to test whether the
representation of 3D orientation (i.e., tilt) is functionally organized within visual cortex. We find
functionally defined structures within V3A and V7. Most notably, in one participant we find that
tilt preference is unilaterally organized in a pinwheel-like structure, similar to those observed
for orientation preference in V1[19,20], that encompasses most of area V3A. Further, for two
participants we find an abrupt reversal in tilt preference at the (retinotopically defined) border
between V3A and V7. These findings show that it is possible to use non-invasive imaging to
discover the functional organisation of 3D orientation within the human visual cortex.
Observing pinwheel-like structures suggests that this type of organization may be applied
canonically within the brain at multiple scales.

METHODS

Participants. Four healthy participants from the University of Cambridge with normal or corrected-to-normal vision participated in the experiment. With the exception of participant 1 (author RR), participants were naïve to the aims of the experiment. The participants ages were 30, 28, 21, and 29, respectively; all were right-handed and participant 3 was female. Participants were screened for stereoacuity using a discrimination task in which they judged the (near/far) depth profile of a random-dot stereogram (RDS) depicting an annulus surrounding a disk. The difference in depth between the annulus and disk was controlled using a 2Down 1-up staircase procedure and participants were admitted into the experiment if they achieved a threshold of $<1°$ arcmin. Participants were also screened for contraindications to MRI prior to the experiment. All experiments were conducted in accordance with the ethical guidelines of the Declaration of Helsinki and were approved by the University of Cambridge STEM and all participants provided informed consent.

Apparatus and stimuli. Stimuli were programmed and presented in MATLAB (The MathWorks, Natick, MA) with Psychophysics Toolbox extensions[21,22]. Stereoscopic presentation in the scanner was achieved using a "PROPixx" DLP LED projector (VPixx Technologies) with a refresh rate of 120 Hz and resolution of 1920 × 1080, operating in RB3D mode. The left and right images were separated by a fast-switching circular polarization modulator in front of the projector lens (DepthQ; Lightspeed Design). The onset of each orthogonal polarization was synchronized with the video refresh, enabling interleaved rates of 60 Hz for each eye's image. MR-safe circular polarization filter glasses were worn by participants in the scanner to dissociate the left and right eye's view of the image. Stimuli were back-projected onto a polarization-preserving screen (Stewart Filmscreen, model 150) inside the bore of the magnet and viewed via a front-surfaced mirror attached to the head coil and angled at 45° above the participants' heads. This resulted in a viewing distance of 72 cm, from which all stimuli were visible within the binocular field of view. Stereoscopic presentation out

of the scanner was achieved using a pair of Samsung 2233RZ LCD monitors (120 Hz,

1680×1050) viewed through mirrors in a Wheatstone stereoscope configuration. The viewing

distance was 50 cm and participants' head position was stabilized using an eye mask, head

rest and chin rest. Eye movement was recorded binocularly at 1 kHz using an EyeLink 1000

(SR Research Ltd., Ontario, Canada).

Stimuli consisted of RDS presented within an annulus (inner radius, .45°; outer radius,

9°) aperture on a mid-grey background surrounded by pink noise intended to facilitate stable

vergence. The outer edge of the annulus was blurred according to a cosine profile. Dots in the

stereogram followed a black or white Gaussian luminance profile, subtending 0.17° at half

maximum. There were 12 dots/deg$_2$, resulting in ~85% coverage of the background. Dots were

allowed to overlap and where this occurred, they occluded previously positioned dots. Dots

presented to the left and right eyes were offset to generate planar surfaces with linear

gradients of horizontal disparity (**Fig. 1a**). Binocular disparity was calculated from the

cyclopean view and applied to each dot based on the specific disparity-defined slant/tilt angle.

In the centre of the annulus, we presented a fixation square (side length = 0.5°) paired with

horizontal and vertical lines.

Figure 1. Three-dimensional stimuli and travelling wave procedure. a) Example random-dot stereogram stimuli used in the experiment (slant, -60°; tilt, 45°). **b**) Illustration of varying the tilt of a planar surface. **c**) Schematic diagram of travelling wave experimental design. A single cycle is shown, 7 cycles were repeated per run.

Psychophysics procedure. We measured participants' sensitivity to change in the tilt (at 60°

slant) of the stimulus with a two-interval forced-choice experimental design in which the

reference and test stimuli were presented in randomized order. The angular difference

between test and reference stimulus was varied using the QUEST adaptive staircase

method[23]. Each stimulus was presented for 500 ms with an inter-stimulus interval of 500 ms.

Observers indicated which stimulus was tilted further clockwise using a keypress. We

measured sensitivity to change around the four cardinal and four oblique angles. For each

reference angle, we ran two randomly interleaved staircases, one that measured sensitivity to

change in the positive direction and one that measured sensitivity to change in the negative

direction; the final threshold estimate was derived from the average of the two staircases.

Each staircase consisted of between 30 to 90 trials, depending on the number of trials required

to reach a stable estimate. The order of reference angles tested for slant and tilt was

randomized for each participant.

Observers could theoretically discriminate surface angle based only on the difference

in depth at the edges (e.g., top and bottom) of a pair of stimuli. To minimize the availability of

this cue, disparity-defined position was randomized by shifting the surface relative to the

fixation plane (0° disparity) to between ±10% of the total surface depth.

Travelling wave procedure. The slant angle of the plane was held constant 60° and we

varied the tilt angle between 0-360° in 24 evenly spaced increments (**Fig. 1b,** right). A

travelling-wave paradigm was designed to generate a traveling wave of activity across/within

areas representing 3D orientation in visual cortex. This involved presenting planar surfaces

which consecutively varied in tilt in either a clockwise or counterclockwise order. Planar

surfaces at each tilt were presented for 2 s, with stimuli presented for 0.4 s separated by 0.1

s inter-stimulus-intervals consisting of only the background and fixation cross (gaps were used

to reduce adaptation). Each cycle of stimulation lasted 48 s (**Fig. 1c**). fMRI runs consisted of

7 cycles in addition to a 16 s blank interval at the beginning and end of the run, thus taking

368 s. Each participant completed the experimental scanning session, which consisted of eight

runs (four clockwise and four counterclockwise) and lasted approximately 60 min. Participant

1 (author RR) completed one additional session. This additional session was combined in the

main analysis. We also collected data for variations of slant, but an artefact in the design of

the slant travelling-wave procedure (but not the tilt procedure) meant that the data were

uninterpretable and thus were omitted.

During the stimulus presentation, participants performed an attentionally demanding

ring detection task. This served to ensured consistent attentional allocation on the stimulus

during the experiment. Participants were instructed to fixate a central crosshair fixation marker

and press a button when they detected a ring. Rings consisted of annular regions (width, 0.6°)

of the stimulus, centred on fixation, that were reduced in contrast by 40%, presented at one of

four possible eccentricities [1.8, 3.6, 5.4, 7.2]°. The shape of the ring presented to the left and

right eyes was distorted to match the 3D orientation of the current planar surface. Rings were

presented for the duration of the stimulus presentation (0.4 s) on 5% of presentations (selected

pseudorandomly).

The method used to present images stereoscopically in the scanner prohibited eye

tracking, so to assess eye movements in response to the stimuli we had participants complete

four runs (two clockwise and two counterclockwise) of the travelling wave paradigm outside

the scanner while we monitored their eye position.

Magnetic resonance imaging. Data were collected at the Wolfson Brain Imaging Center with

a 3T Siemens PRISMA system using a 32-channel head coil. Blood oxygen level-dependent

(BOLD) functional data were acquired with a gradient echo-planar imaging sequence [echo

time, 29 ms; repetition time, 2000 ms; voxel size, 1.5 × 1.5 × 2 mm; 30 slices covering occipital

cortex] for experimental and localizer scans. Slice orientation was near to coronal section but

rotated about the mediolateral axis with the ventral edge of the volume more anterior in order

to ensure coverage of lateral occipital cortex (**Fig. 2a**). A T1-weighted, 3D-MPRAGE pulse

sequence, anatomical scan (voxel size, 1 mm isotropic) was additionally acquired for each

participant. These anatomical scans were used to register the functional data across scanning

sessions, restrict the analysis of BOLD activity to grey matter voxels, and create flattened

surfaces on which cortical activity is visualized.

Data processing. Anatomical scans were aligned along the ACPC plane, grey matter and

white matter were segmented, and flattened surfaces were computationally rendered with

BrainVoyager QX (BrainInnovation, Maastricht, The Netherlands; **Fig. 2b**). Flattened surfaces

(flat maps) were rendered such that areas that were concave on the original surface (i.e., sulci)

are dark grey, and areas that were convex (i.e., gyri) are light grey. Functional data were

preprocessed with slice timing correction, head motion correction, and high-pass filtering

before being aligned to the participant's anatomical scan. Functional images from the first run

within a session were manually aligned to the anatomical, and these alignment coordinates

were used to align the images from the remaining runs. Analysis of functional imaging data

was performed in MATLAB (MathWorks, Natick, MA) using in-house scripts.

To evaluate periodic responses to 3D orientation, the data were analysed using a

Fourier transformation-based method, well documented in the standard retinotopy studies[24–

26]. As previously described by Ban et al. (2003)[27], clockwise and counterclockwise run time

series were concatenated and a discrete Fourier transformation was computed for each voxel

after converting the raw signals to percentage signal changes. The phase measurement

computed from this analysis was averaged between counterclockwise runs and reversed

before being averaged with data from the clockwise runs to remove the effect of the

hemodynamic delay in deriving the 3D orientation maps. Then, statistical testing to estimate

the significance of correlation of BOLD signal at the stimulus frequency (1/48 Hz) was

performed by comparing the squared amplitude at the stimulus frequency with the sum of

squared amplitudes at the other frequencies, which yielded an *F*-ratio. The *F*-ratio was

converted to a *P* value considering degrees of freedom of the signals (number of time points).

In mapping retinotopic responses on the cortical surface, the polar angle phase of each

significantly activated voxel was displayed using a continuous colour scale. We used a voxel-

level Fourier *F*-statistic value ($F(2,$ number of time points$) > 7.38$ and $p < 0.05$) as a criteria of

significance for all Fourier-based analysis. Additionally, coherence values were obtained by

calculating the voxel-by-voxel correlation between the concatenated time series and the

corresponding best-fitting sinusoid at the frequency of the visual stimulation; coherence values

were averaged between clockwise and counterclockwise runs. The coherence measures

signal-to-noise[24,28], and ranges from 0 to 1, where values near 1 indicate the fMRI signal

modulation at the stimulus period is large relative to the noise (at the other frequency

components) and values near 0 indicate that there is no signal modulation or that the signal is

small compared with noise.

Retinotopy. In a separate session, we performed ROI localization for each participant using

retinotopic and stimulus contrast mapping procedures; the details of which are reproduced

from Murphy, Ban, and Welchman (2013)[29]. Retinotopically organized visual areas (V1, V2,

V3v, V4, V3D, V3A, V3B/KO, and V7) were defined using polar (**Fig. 2c**) and eccentricity maps

(**Fig. 2d**), which were derived from fMRI responses to rotating wedge and expanding

concentric ring stimulus presentations, respectively[25,26]. Area V3 was separated into ventral

and dorsal quadrants in each hemisphere (V3v/d) consistent with previous delineation based

on functional and cytoarchitectonic differences[30]. V4 was defined as the ventral region of visual

cortex adjacent to V3v comprising a representation of the upper quadrant of the contralateral

visual field[31,32]. V7 was defined as the region of retinotopic activity dorsal and anterior to

V3A[3,31,32]. V3B/KO[4,33] was defined as the area containing the union of retinotopically mapped

V3B and the kinetic occipital region (KO), which was functionally localized by its preference

for motion-defined boundaries compared with transparent motion of white and black

dots[4,31,33,34]. This area contained a full hemifield representation, inferior to V7 and lateral to,

and sharing a foveal confluence with, V3A[32]. Human middle temporal complex (hMT+/V5) was

defined as the set of voxels in lateral temporal cortex that responded significantly more

strongly to transparent dot motion compared with a static array of dots[35]. Lateral occipital

complex (LOC) was defined as voxels in the lateral occipito-temporal cortex that preferred

intact, compared with scrambled, images of objects[36]. LOC subregion LO was defined based

on overlap of anatomical structures and functional activations, in line with previous work[37].

Figure 2. Retinotopic maps. a) Sagittal view of a structural MRI scan of participant 1, with coloured overlay illustrating slice orientation for functional scans. **b**) Reconstruction and flattened surface of structural scan, with coloured overlay illustrating circular regions shown in (**c-d**). (**c-d**) Flattened structural surfaces with coloured overlay illustrating phase map in response to **c**) a rotating wedge and **d**) expanding/contracting concentric rings. Black lines in (**c-d**) indicate retinotopic area demarcation.

Pinwheel analysis. Previous work in which evidence for pinwheel-like encoding structures has

been presented have relied on subjective identification of the formations. Here we developed a

novel analysis method to objectively quantify the likelihood that the pattern of encoding at a

particular location on the cortex is consistent with a pinwheel formation. To do so, we

implemented a spotlight analysis of the flat map representations of tilt angle preference. The

flat maps comprise polygons, each of which corresponds to a grey matter voxel from the

anatomical image and has a phase value that indicates the tilt preference of that voxel. The

phase values within the spotlight were compared to the position of each polygon centroid,

relative to the center of the spotlight, in polar angle. A circular-circular regression between

these values was calculated to determine their correspondence, independent of absolute

phase. Regression values varied from 0 to 1, where 0 indicates no correspondence and 1

indicates perfect correspondence. Thus, if the phase values are structured in a pinwheel

formation, their value will be predicted by the angular position from the center of the pinwheel,

regardless of the phase offset position. The spotlight analysis was performed at each centroid

location, with the coordinates of the centroid defining the center position of the spotlight. To

reduce spurious correlations, only tests that included a minimum spotlight data point density

of 1.4 (>216 data points) were included in the analysis. We present results obtained when the

spotlight radius was 7 units in the flat map coordinate space (see the white circle in **Figure 7b**

for size reference); however, to ensure that this did not bias our results, we systematically

varied the spotlight radius between 2 and 14 units, finding a similar pattern of results.

RESULTS

Following retinotopic mapping of visual areas (**Fig. 2c-d**) we measured participants'

phase-encoded responses to 3D tilt from 0-360° at 60° slant, defined by binocular disparity.

Figure 3a shows the coherence of the activity measured from visual cortex with the frequency

of the travelling wave stimulus. There was a relatively high coherence within retinoptopically

defined visual areas, especially dorsal stream regions, suggesting that the neural populations

within these regions were sensitive to aspects of the stimulus which varied as a function of tilt.

Figure 3b shows the corresponding phase maps for each participant, which were thresholded

to remove phase values with $P > .05$.

Figure 3. Cortical representation of tilt. Tilt **a**) coherence and **b**) phase maps for all subjects with retinotopic ROI borders overlaid in black. The plots on the right of the maps show the **a**) average coherence and **b**) standard deviation of phase values within each retinotopic area.

Summary analysis of voxel preference across the cortex revealed that participants 3

and 4 showed a high proportion of phase values corresponding to 0° tilt; however, this was

only weakly present in participant 1 and not present in participant 2 (**Fig. 4a**). By contrast, all

participants showed better performance for cardinal tilt angles than oblique angles (**Fig. 4b**).

Previous work has identified areas V3A, V7, and MT as regions in which 3D orientation is

encoded[10,12–14]. We reasoned that visual areas integral to encoding tilt would show selectivity

to a variety of tilt angles. Thus, to assess the heterogeneity of tilt encoding, we calculated the

variability of voxel phase preference within each visual area (**Fig. 4c**, right). In line with

previous work, we found that areas V3A and V7 had more heterogeneous tilt preference than

the other visual areas. By contrast, assessment of average coherence across visual areas

suggested similar levels of coherence (**Fig. 4d**). A possible concern is that patterns of activity

may have been influenced by eye movements during stimulus presentation. However, we

found no evidence for this; in a separate session outside the scanner in which participants

performed the same task, we found binocular eye movements were stable between different

tilt angle presentations (**Supplementary Figure 1**).

Figure 4. Summary of tilt representation. **a**) The proportion of significant voxels (P<.05) within the recorded region as a function of slant preference. **b**) Tilt sensitivity (reciprocal of the just noticeable difference) as a function of the reference slant angle. Colour labels in (**b**) match those in (**a**). (**c-d**) Average (**c**) coherence and (**d**) phase variability within retinotopic areas in the left and right hemisphere across all participants.

Another possibility is that the increased encoding variability observed in areas V3A

and V7 is not specific to the representation of tilt in these areas. That is, information may be

more heterogeneously encoded in these areas than other visual areas, regardless of the

feature. To test this possibility, we performed the same analysis on the polar angle and

eccentricity retinotopic data (**Figure 2b & c**); if all visual information is heterogeneously

represented within V3A and V7, we would expect to find higher phase variability in these areas

for polar angle and eccentricity. By contrast, the pattern of results for phase angle and

eccentricity was markedly different than that for tilt angle. For polar angle, phase variability

was highest in V1, and while it was second highest in V3A, it was relatively low in V7 (**Figure**

5a). For eccentricity, phase variability was also highest in V1, and relatively low in V3A and

V7 (**Figure 5a**). These results indicate that the heterogeneous encoding of tilt angle in V3A

and V7 was not simply due to these areas encoding all features in a varied manner.

Figure 5. Summary of polar angle and eccentricity representation. Average phase variability for (**a**) slant angle and (**b**) eccentricity within retinotopic areas in the left and right hemisphere across all participants.

266   Inspection of areas V3A and V7 revealed two types of structure (**Fig. 6a**). Participant

267   2 had pinwheel-like structure of tilt preference in V3A (**Fig. 6b**, broken white circles).

268   Participants 1 and 4 showed abrupt tilt preference reversal, between the border of V3A and

269   V7 (**Fig. 6a**, broken white rectangles). With the exception of participant 1, these structures

270   were not bilaterally present.

Figure 6. Representation of tilt in V3A/V7. a) Tilt phase maps of areas V3A and V7 for all participants. Broken black lines indicate region borders, broken white lines highlight pinwheel formation (circle) and intraregional phase transitions (rectangles). **b**) Comparison of the pinwheel like structure highlighted in the left hemisphere of participant 2 (top) and the tilt phase colour wheel (bottom).

It is possible that the pinwheel-like structure we observed in the left V3A of participant

2 was a product of chance. To test this possibility, we designed a pinwheel detection analysis

that scans a map of phase values to yield a corresponding map of ρ values that indicate how

well each location in the phase map corresponds to a pinwheel formation (see Methods for

detailed explanation of the pinwheel analysis). As a sanity check, we first used the pinwheel

detection analysis on an image of pinwheel-like structures in macaque V2 that correspond to

motion direction preference[38] (**Fig. 7a**, top). As expected, the corresponding correlation map

shows peaks at locations corresponding to the centres of the pinwheel-like structures in the

image (**Fig. 7a**, bottom). We then applied the analysis to participant 2's right hemisphere

phase map. We found that the location at which the highest correlation was found

corresponded to the centre of the pinwheel structure in V3A (ρ=.917, (cos, sin) *P*=[1.1e-5, 3.6e-

6]; **Fig. 7b,** top, white circle). A possible concern is that the pinwheel-like structure in V3A was

a result of random distribution of phase values across the cortex, rather than a representing a

functionally beneficial encoding structure. Given that the pinwheel is located in the visual area

most intrinsically involved in processing 3D orientation (V3A), this seems unlikely. However,

to test this possibility, we reran the pinwheel analysis on participant 2's phase map after

assigning phase values according to a two-dimensional pink noise (1/frequency) distribution

(**Fig. 7b**, bottom). **Figure 7c** shows the distribution of correlation values from the pink noise

simulation (pink bars) relative to the peak value found in the data (black arrow). The peak

correlation value corresponding to the pinwheel structure in V3A clearly lies outside the values

generated by pink noise, indicating that this kind of structure would be unlikely to be produced

by chance.

Figure 7. Pinwheel detection spotlight analysis. a, top) Motion direction preference map in macaque V2, shown using optical imaging[38]. **a**, bottom) Pinwheel correlation map produced by running the pinwheel detection spotlight analysis on (**a**, top). White circles indicate the size of the spotlight used in the analysis. **b**, top-left) Phase map and corresponding (**b**, top-right) pinwheel correlation map of Participant 2's left hemisphere. The black outline delineates area V3A and the position and size of the white circle indicate the point of maximum correlation and the size of the spotlight, respectively. **b**, bottom) Same as (**b**, top), but when assigning phase according to a two-dimensional pink noise distribution. **c**) Histogram of the correlation values in (**b**, bottom-right); the peak correlation value in (**b**, top-right) is indicated by the arrow.

DISCUSSION

Previous fMRI work with humans has identified areas V3A and V7 as responsive to 3D orientation[13,14]. These studies have employed experimental designs in which conditions are contrasted to reveal activity related to a particular cue. This procedure has been effective in isolating regions that support encoding of 3D orientation; however, it is blind to any putative functional organization of 3D orientation selectivity within these areas. Functional organization is likely to be the rule, not the exception, in the brain, as it has the potential to provide computational benefits that arbitrary organization cannot. There are many examples of functional organization within visual cortex at multiple scales. Classic examples include orientation/ocular dominance columns[39,40] and retinotopic organization[25]. In a more recent example, relative depth was shown to be organized in area V3A[1] in a way that persisted across scan sessions, although no identifiable functional structures were observed. Here we use phase-encoded presentation of planar surfaces with linear gradients of horizontal disparity to test whether the representation of 3D orientation is functionally organized within visual cortex. In particular, we measured fMRI activity of four participants to determine the preference of voxels within visual cortex for tilt angles between 0-360° with 60° slant angle.

We found a nonuniform distribution of tilt preference within visual cortex, there was a high proportion of voxels which preferred 0° tilt. Behaviourally, we found that observers were more sensitive to change in tilt angle around the cardinal axes, compared to the oblique. This is consistent with sensitivity to 2D orientation and motion, which is also best around cardinal axes (i.e., the oblique effect), and likely reflects an anisotropic distribution of 3D oriented surfaces in the environment. This may partially explain the anisotropic distribution of tilt preference observed. However, the large proportion of voxels that were identified as preferring 0° tilt may also reflect the anisotropic representation of near and far disparities in the upper and lower visual field. Perceptually, human observers are better at detecting targets defined by crossed (near) disparity[41] and overestimate the distance of objects in the upper visual field[42–44], while they are better at detecting targets defined by uncrossed (far) disparity and

underestimate the distance of objects in the lower visual field[45]. This bias has also been observed physiologically: clustered regions that respond to near and far stimuli are preferentially located in retinotopic regions representing the upper and lower visual fields of V2/3, respectively[46]. Although all stimuli in the tilt condition had equal absolute global disparity, the anisotropic representation of near and far disparity in the upper and lower visual field likely resulted in a larger response to the 0° tilt stimuli, thus biasing the distribution of voxel tilt preferences.

We observed two patterns of functional organization of tilt preference in V3A/V7. The first, and perhaps most striking, was a pinwheel-like structure in the left hemisphere of participant 2. This structure encompassed the majority of V3A. Pinwheel-like functional organization has been observed for orientation selectivity in V1[19,20]. However, the spatial scale of these structures is very different; orientation pinwheels in V1 are ~300 μm in radius, whereas the tilt pinwheel we observed in V3A was ~10 mm in radius. In V1, this structure is thought to bestow at least two benefits for orientation processing: i) preferences for all orientations are brought together, within reach of upper-layer dendrites, and ii) orientation preferences on opposite sides are always perpendicular[20]. These same benefits may be shared by neurons that encode tilt in V3A; however, as tilt is a vector whose singularities (i.e., the point at the center of the pinwheel) rotate through 360°, tilt angles on opposite sides of the singularity are opposite, as opposed to perpendicular. While the benefits of pinwheel-like functional organization remain an empirical question, patterns resembling this formation that we observe here might suggest that pinwheels are used to represent a different visual property (3D tilt angle) in a different region (V3A) at a different spatial scale (~30 times larger). While our results are not fully conclusive in that these were not seen across hemispheres or in all participants, they might nevertheless hint that pinwheels represents a canonical organizational structure within the brain.

We found evidence of functional structures that were common to some, but not all participants. It is possible that these structures are not universal; however, it is also possible

that we did not detect these structures across all participants due to insufficient resolution

and/or signal strength. Future work could build on the findings revealed here by using fMRI

at higher field strength[47] (e.g., 7T) to target and characterize the representation of 3D surface

orientation in areas V3A and V7. Further, although we used binocular disparity to define the

planar surfaces in the current study, V3A and V7 have been shown to respond to 3D

orientation defined by motion[11,15,16], texture[17], and shading[18]. Thus, future work could compare

the representation of 3D orientation within these areas using different depth cues in isolation

and combination.

Using phase-encoded presentation of 3D stimuli we find evidence in support of the

functional organization of 3D orientation within visual cortex. In one participant we find fMRI

response patterns suggesting that tilt preference is unilaterally organized in a pinwheel-like

structure, similar to those observed for orientation preference in V1[19,20], that encompasses

most of area V3A. Further, for two participants we find an abrupt reversal in tilt preference at

the (retinotopically defined) border between V3A and V7. These findings indicate that 3D

orientation is functionally organized within the human visual cortex, and the discovery of a

large pinwheel-like structure suggests that this type of organization may be applied canonically

within the brain at multiple scales.

REFERENCES

1. Goncalves, N. R. *et al.* 7 Tesla fMRI Reveals Systematic Functional Organization for Binocular Disparity in Dorsal Visual Cortex. *J. Neurosci.* **35**, 3056–3072 (2015).

2. Preston, T. J., Li, S., Kourtzi, Z. & Welchman, A. E. Multivoxel Pattern Selectivity for Perceptually Relevant Binocular Disparities in the Human Brain. *J. Neurosci.* **28**, 11315–11327 (2008).

3. Tsao, D. Y. *et al.* Stereopsis activates V3A and caudal intraparietal areas in macaques and humans. *Neuron* **39**, 555–568 (2003).

4. Tyler, C. W., Likova, L. T., Kontsevich, L. L. & Wade, A. R. The specificity of cortical region KO to depth structure. *Neuroimage* **30**, 228–238 (2006).

5. Neri, P., Bridge, H. & Heeger, D. J. Stereoscopic Processing of Absolute and Relative Disparity in Human Visual Cortex. *J. Neurophysiol.* **92**, 1880–1891 (2004).

6. Backus, B. T., Fleet, D. J., Parker, A. J. & Heeger, D. J. Human Cortical Activity Correlates With Stereoscopic Depth Perception. *J. Neurophysiol.* **86**, 2054–2068 (2001).

7. Taira, M., Tsutsui, K.-I., Jiang, M., Yara, K. & Sakata, H. Parietal Neurons Represent Surface Orientation From the Gradient of Binocular Disparity. *J. Neurophysiol.* **83**, 3140–3146 (2017).

8. Rosenberg, A. & Angelaki, D. E. Gravity Influences the Visual Representation of Object Tilt in Parietal Cortex. *J. Neurosci.* **34**, 14170–14180 (2014).

9. Rosenberg, A., Cowan, N. J. & Angelaki, D. E. The Visual Representation of 3D Object Orientation in Parietal Cortex. *J. Neurosci.* **33**, 19352–19361 (2013).

10. Nguyenkim, J. D. & DeAngelis, G. C. Disparity-based coding of three-dimensional

surface orientation by macaque middle temporal neurons. *J. Neurosci.* **23**, 7117–7128

(2003).

11.   Vanduffel, W. *et al.* Extracting 3D from motion: Differences in human and monkey

intraparietal cortex. *Science (80-.).* **298**, 413–415 (2002).

12.   Ban, H. & Welchman, A. E. fMRI Analysis-by-Synthesis Reveals a Dorsal Hierarchy

That Extracts Surface Slant. *J. Neurosci.* **35**, 9823–9835 (2015).

13.   Durand, J.-B., Peeters, R., Norman, J. F., Todd, J. T. & Orban, G. A. Parietal regions

processing visual 3D shape extracted from disparity. *Neuroimage* **46**, 1114–1126

(2009).

14.   Georgieva, S., Peeters, R., Kolster, H., Todd, J. T. & Orban, G. A. The Processing of

Three-Dimensional Shape from Disparity in the Human Brain. *J. Neurosci.* **29**, 727–

742 (2009).

15.   Orban, G. A., Sunaert, S., Todd, J. T., Van Hecke, P. & Marchal, G. Human Cortical

Regions Involved in Extracting Depth from Motion. *Neuron* **24**, 929–940 (1999).

16.   Murray, S. O., Olshausen, B. A. & Woods, D. L. Processing Shape, Motion and

Three-dimensional Shape-from-motion in the Human Cortex. *Cereb. Cortex* **13**, 508–

516 (2003).

17.   Shikata, E. *et al.* Surface Orientation Discrimination Activates Caudal and Anterior

Intraparietal Sulcus in Humans: An Event-Related fMRI Study. *J. Neurophysiol.* **85**,

1309–1314 (2017).

18.   Tsutsui, K.-I., Jiang, M., Yara, K., Sakata, H. & Taira, M. Integration of perspective

and disparity cues in surface-orientation-selective neurons of area CIP. *J.*

Neurophysiol. **86**, 2856–2867 (2001).

19.   Bonhoeffer, T. & Grinvald, A. Iso-orientation domains in cat visual cortex are arranged

in pinwheel-like patterns. *Nature* **353**, 429–431 (1991).

20.   Blasdel, G. G. Orientation selectivity, preference, and continuity in monkey striate

cortex. *J. Neurosci.* **12**, 3139–61 (1992).

21.   Brainard, D. H. The Psychophysics Toolbox. *Spat. Vis.* **10**, 433–436 (1997).

22.   Pelli, D. G. The VideoToolbox software for visual psychophysics: Transforming

numbers into movies. *Spatial Vision* **10**, 437–442 (1997).

23.   Watson, A. B. & Pelli, D. G. QUEST: A general multidimensional bayesian adaptive

psychometric method. *Percept. Psychophys.* **33**, 113–120 (1983).

24.   Engel, S. A. *et al.* FMRI of human visual cortex [5]. *Nature* **369**, 525 (1994).

25.   Sereno, M. I. *et al.* Borders of multiple visual areas in humans revealed by functional

magnetic resonance imaging. *Science (80-.).* **268**, 889–893 (1995).

26.   DeYoe, E. A. *et al.* Mapping striate and extrastriate visual areas in human cerebral

cortex. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 2382–6 (1996).

27.   Ban, H. *et al.* Topographic Representation of an Occluded Object and the Effects of

Spatiotemporal Context in Human Early Visual Areas. *J. Neurosci.* **33**, 16992–17007

(2013).

28.   Schluppeck, D. Topographic Organization for Delayed Saccades in Human Posterior

Parietal Cortex. *J. Neurophysiol.* **94**, 1372–1384 (2005).

29.   Murphy, A. P., Ban, H. & Welchman, A. E. Integration of texture and disparity cues to

surface slant in dorsal visual cortex. *J. Neurophysiol.* **110**, 190–203 (2013).

30.   Wilms, M. *et al.* Comparison of functional and cytoarchitectonic maps of human visual

areas V1, V2, V3d, V3v, and V4(v). *Neuroimage* **49**, 1171–1179 (2010).

31.   Tootell, R. B. H. Where is 'Dorsal V4' in Human Visual Cortex? Retinotopic,

Topographic and Functional Evidence. *Cereb. Cortex* **11**, 298–311 (2001).

32.   Tyler, C. W. *et al.* Extended Concepts of Occipital Retinotopy. *Curr. Med. Imaging*

Rev. **1**, 319–329 (2005).

33.   Zeki, S., Perry, R. J. & Bartels, A. The processing of kinetic contours in the brain.

Cereb. Cortex **13**, 189–202 (2003).

34.   Dupont, P. *et al.* The kinetic occipital region in human visual cortex. *Cereb. Cortex* **7**,

283–292 (1997).

35.   Zeki, S. *et al.* A direct demonstration of functional specialization in human visual

cortex. *J. Neurosci.* **11**, 641–649 (1991).

36.   Kourtzi, Z., Betts, L. R., Sarkheil, P. & Welchman, A. E. Distributed neural plasticity for

shape learning in the human visual cortex. *PLoS Biol.* **3**, 1317–1327 (2005).

37.   Grill-Spector, K., Kushnir, T., Hendler, T. & Malach, R. The dynamics of object-

selective activation correlate with recognition performance in humans. *Nat. Neurosci.*

3, 837–893 (2000).

38.   Lu, H. D., Chen, G., Tanigawa, H. & Roe, A. W. A Motion Direction Map in Macaque

V2. *Neuron* (2010). doi:10.1016/j.neuron.2010.11.020

39.   Hubel, D. H. & Wiesel, T. N. Sequence regularity and geometry of orientation columns

in the monkey striate cortex. *J. Comp. Neurol.* **158**, 267–293 (1974).

40.   Hubel, D. H. & Wiesel, T. N. Laminar and columnar distribution of geniculo-cortical

fibers in the macaque monkey. *J. Comp. Neurol.* **146**, 421–450 (1972).

41.   Breitmeyer, B., Julesz, B. & Kropfl, W. Dynamic random-dot stereograms reveal up-

down anisotropy and left-right isotropy between cortical hemifields. *Science (80-.).*

187, 269–270 (1975).

42.   Ooi, T. L., Wu, B. & He, Z. J. Distance determined by the angular declination below
the horizon. *Nature* **414**, 197–200 (2001).

43.   Wallach, H. & O'Leary, A. Slope of regard as a distance cue. *Percept. Psychophys.*
31, 145–148 (1982).

44.   Yang, Z. & Purves, D. A statistical explanation of visual space. *Nat. Neurosci.* **6**, 632–
640 (2003).

45.   Breitmeyer, B., Battaglia, F. & Bridge, J. Existence and Implications of a Tilted
Binocular Disparity Space. *Perception* **6**, 161–164 (1977).

46.   Nasr, S. & Tootell, R. B. H. Visual field biases for near and far stimuli in disparity
selective columns in human visual cortex. *Neuroimage* **168**, 358–365 (2018).

47.   Schluppeck, D., Sanchez-Panchuelo, R. M. & Francis, S. T. Exploring structure and
function of sensory cortex with 7 T MRI. *Neuroimage* **164**, 10–17 (2018).

470