Working paper

Wing shape and environmental energy are associated with molecular

evolutionary rates in a large avian radiation

David Alejandro Duchêne[a,1], Paola Montoya[b,c], Carlos Daniel Cadena[b]

[a]Research School of Biology, Australian National University, ACT 2601, Australia

[b]Departamento de Ciencias Biológicas, Universidad de los Andes, Apartado 4976, Bogotá,

Colombia

[c]Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Bogotá, Colombia

[1]Corresponding author

David A. Duchêne

Research School of Biology

Robertson Building, 46

Australian National University

Canberra, ACT 2601

Australia

Telephone: +61 4 12026379

Email: david.duchene@anu.edu.au

Abstract

Among the macroevolutionary drivers of molecular evolutionary rates, metabolic demands and environmental energy have been a central topic of discussion. The large number of studies examining these associations have found mixed results, and have rarely explored the interactions among various factors impacting molecular evolutionary rates. Taking the diverse avian family Furnariidae as a case study, we examined the association between several estimates of molecular evolutionary rates with a proxy of metabolic demands imposed by flight (wing morphology) and proxies of environmental energy across the geographic ranges of species (temperature and UV radiation). We found a strong positive association between molecular rates in genomic regions that can change the coded amino-acid with wing morphology, environmental temperature, and UV radiation. Strikingly, however, we did not find evidence of such associations with molecular rates at sites not impacting amino-acids. Our results suggest that the demands of flight and environmental energy primarily impact genome evolution by placing selective constraints, instead of being associated with basal mutation rates.

Keywords

Furnariidae, flight ability, hand-wing index, environmental temperature, environmental UV radiation

Introduction

The factors driving molecular evolution are a long standing matter of interest in the biological sciences. Some of the most studied factors correlating with rates of molecular evolution on a macroevolutionary scale include life-history traits (Bromham 2011), metabolic activity (e.g., Garcia-Porta *et al.* 2019) and environmental energy (Wright *et al.* 2006). These factors are also likely to interact, with the evolutionary outcome depending on the balance between the costs of sources of deleterious mutations and the requirements of the particular lifestyle of species (Bromham 2009). Although several studies have shown interactions among multiple correlates of molecular evolution (Lanfear *et al.* 2013; Bromham *et al.* 2015), there are gaps in our knowledge regarding the universality and relative contribution of different biological and environmental factors to molecular evolutionary rates.

Metabolic rate is an important biological driver of molecular evolution because metabolic activity produces mutagens (Gillooly *et al.* 2005; Allen *et al.* 2006). Specifically, metabolism produces oxygen and nitrogen free radicals (Barja 1999) which may cause novel mutations through damage to DNA (Cooke *et al.* 2003). Mass-specific metabolic rate is inversely associated with body mass (Martin & Palumbi 1993), which provides one explanation for the accelerated rates of molecular evolution in small vertebrates compared to large ones. While there is some evidence that metabolic rates are indeed associated with rates of molecular evolution (Bleiweiss 1998; Gillooly *et al.* 2005, 2007; Garcia-Porta *et al.* 2019), multiple studies have not found such an association (Mooers & Harvey 1994; Bromham *et al.* 1996; Lanfear *et al.* 2007; Galtier *et al.* 2009; Rolland *et al.* 2016).

An explanation for the mixed evidence of the impact of metabolism on molecular evolution is that there has been a heavy focus on body size and temperature as proxies of metabolic rate (Gillooly *et al.* 2001), yet multiple biological factors can impact metabolic requirements. For example, flight demand in birds is an extreme form of endurance exercise (Weber 2009), associated with high metabolic rate (Butler & Woakes 1990; Suarez 1992; Ward *et al.* 2001, 2002), high respiratory rate (Guglielmo *et al.* 2002), high rates of catabolism of lipids and proteins (Jenni-Eiermann *et al.* 2002), and oxidative stress (Jenni-Eiermann *et al.* 2014). Among the adaptations of particular avian lineages to sustain long periods of flight with high efficiency are also a high capillary density in flight muscles (Bishop *et al.* 1995; Maillet & Weber 2007), low wing loading (Norberg 1995; Alerstam *et al.* 2007), and pointed wings

(Dawideit *et al.* 2009). The high exposure to oxygen and nitrogen reactive species in birds with

long-distance flight (Jenni-Eiermann *et al.* 2014) might have a mutagenic effect that leads to

faster rates of molecular evolution beyond the effects of temperature and body mass.

Another widely studied factor that might drive molecular evolutionary rate is

environmental energy through its mutagenic effect on the genome (Martin & Palumbi 1993;

Allen *et al.* 2006). Studies in organisms including plants (Davies *et al.* 2004; Wright *et al.* 2006),

marine organisms (Wright *et al.* 2010b), and lizards (Garcia-Porta *et al.* 2019) have all found

associations between environmental energy and molecular rates. The link between energy and

molecular evolution is often inferred using environmental temperature as a proxy of available

energy (Wright *et al.* 2006). Other studies have focused on UV radiation (e.g., Davies *et al.*

2004), which is a well understood mutagen (Pawlowski *et al.* 1997). However, it is unclear how

UV might impact the germline in many organisms, like endothermic vertebrates.

Environmental energy might also affect molecular evolution by mediating the fate of

novel mutations, as opposed to affecting basal mutation rates. For example, temperature might

influence the fitness effects of mutations, with maximal fitness occurring at around global

maximal temperatures (Puurtinen *et al.* 2016). Energy might also facilitate the accumulation of

biomass and allow for larger numbers of individuals to exist in populations (Wright 1983; Currie

1991; Willig *et al.* 2003); in larger populations, selection is more efficient and novel beneficial

mutations are more likely to reach fixation rapidly (Kimura 1968; Ohta 1992; Lanfear *et al.*

2014). High energy might also lead to reduced investment in thermoregulation, allowing limited

resources to be used in other activities, such as reproduction, and ultimately allowing for novel

adaptations (Turner *et al.* 1988).

We examined the association between the rates of molecular evolution and the metabolic

demands imposed by flight and environmental energy across species of an avian family. The

avian family Furnariidae is a large radiation that includes the Neotropical ovenbirds and

woodcreepers (Sibley & Monroe 1990; Remsen Jr *et al.* 2012). They have undergone a fast and

consistent diversification over the last 30My (Derryberry *et al.* 2011) and inhabit a broad range

of habitats of south and central America (Claramunt 2010). The species richness of the family

and diversity of adaptations across broad gradients of altitude, latitude, and environmental

conditions make it an interesting group for testing the association between molecular evolution

and a range of adaptations (Fjeldså *et al.* 2005).

We focus on wing shape and environmental temperature and UV radiation as proxies of

metabolic demand and environmental energy, respectively. Using data for the majority of species

of Furnariidae (80%) we provide the first near-complete species-level examination of the

correlates of molecular evolution in a large vertebrate radiation. We tested the association that

wing shape and environmental energy have with molecular evolutionary rates in a set of nuclear

and mitochondrial genomic regions. Individual tests across codon positions were made to dissect

the independent effects on molecular signatures of selection or population size and mutation rate

(Lanfear *et al.* 2010). Our data show that wing morphology and UV radiation are associated with

molecular signatures of diversifying selection or small population sizes, suggesting an important

impact of flight demands and environmental energy on the genome whilst not necessarily

influencing the basal rate of mutation.

Methods

Data collection

We used published data on hand-wing index obtained from museum specimens for 280

species in Furnariidae (Claramunt *et al.* 2012b). The hand-wing index (Kipp 1959) is a metric of

wing-pointedness calculated as the length between the tip of the longest primary wing feather to

the tip of the first secondary feather, divided by the wing chord (Claramunt *et al.* 2012b). Pointed

wings serve as a proxy of the demands of powered flight because they are critical for long

distance bouts of flight (Pennycuick 2008; Claramunt *et al.* 2012b) and are associated with

dispersal ability (Lockwood *et al.* 1998; Burney & Brumfield 2009; Dawideit *et al.* 2009). We

thus assume that birds with more pointed wings undergo greater amounts of flight, and hence

experience greater metabolic demands associated with flight which may influence rates of

molecular evolution. Data body mass was also collected for species of Furnariidae from a

published compilation (Dunning 2008).

We collected data on environmental temperature and UV across the ranges of the species

in the family Furnariidae. For all species, we downloaded georeferenced records from GBIF

(Global Global Biodiversity Information Facility, https://www.gbif.org/) and VertNet

(http://portal.vertnet.org/), discarding duplicated records and those outside the known

distribution range for each species according to expert maps (del Hoyo *et al.* 2016). Using

145216 depurated records (mean per species=~615), we estimated the mean value for the annual

mean temperature from WorldClim data (Fick & Hijmans 2017) at 30 arc-sec resolution, and for

the annual mean UV-B from gIUV (Beckmann *et al.* 2014) with 15 arc-min resolution. Since

current geographic distributions can differ greatly from the distributions in the past,

contemporary measurements of environmental variables can introduce noise to subsequent

phylogenetic regression analyses. This noise is likely to increase the Type II error rate but it is

unlikely to cause a bias that increases Type I error rate, so the analyses of these type of data are

in fact conservative (Davies *et al.* 2004).

Genomic data were taken from a published phylogenetic study of Furnariidae that

included every species and mitochondrial and nuclear markers (Derryberry *et al.* 2011). Using

data from three nuclear and three mitochondrial loci from the source phylogenetic study of

Furnariidae, we identified species in Furnariidae for which data were available for environment,

hand-wing index, and molecular evolutionary rates ($N = 236$). The molecular data and variables

used in subsequent regression analyses are available online

(github.com/duchene/furnariidae_rates).

Estimates of rates of molecular evolution

Relative rates of substitution across branches were estimated using Bayesian inference

implemented in BEAST v2.4 (Bouckaert *et al.* 2014). We used a GTR+Γ substitution model

(Tavaré 1986) and the uncorrelated log-normal relaxed clock model of substitution rates across

branches (Drummond *et al.* 2006). We tested for potentially biased inferences from the

substitution model using the tests of model adequacy in the software PhyloMAd (Duchêne *et al.*

2018). This assesses whether empirical data adhere to the assumptions made by the model by

comparing them with data simulated under the model.

Substitution and clock model parameters were estimated independently for mitochondrial

and nuclear data, as well as independently for $1^{st}+2^{nd}$ codon positions (CP12) versus 3^{rd} codon

positions (CP3) for the coding regions available. Rates of substitutions per unit of relative time

across taxa were estimated using the phylogeny of genera extracted from the original

phylogenetic study. We fixed the mean rate at CP3 sites at 1. The rates of CP12 sites were

estimated relative to those at CP3 sites. The joint prior on the tree topology and relative node

times was generated using a birth-death process. Posterior distributions were estimated by

running analyses of 5×10^{7} Markov chain Monte Carlo steps, with samples drawn every 5×10^{3}

steps. We discarded the first 10% of samples as burn-in and assessed satisfactory convergence of
the posterior to a stationary distribution by verifying that the posterior of every sampled
parameter had an effective sample size above 200. We made an independent replicate of each
analysis to confirm that parameter estimates converged on the same results, and were not the
product of reaching local optima. The median relative rate estimates of terminal branches across
samples from the posterior distribution of each were used as rate estimates across species.
171
Regression analyses
We used phylogenetic generalized least squares (PGLS) regression models to examine
the contribution of wing shape (a proxy for metabolic rate) and variables describing
environmental energy to molecular evolutionary rates. Hypotheses were tested independently
using nuclear and mitochondrial data, and using CP12 and CP3 rates as response variables.
Estimates of CP3 rates largely represent molecular changes that do not influence the amino acid
being coded (69%, 113/163), so they primarily represent the mutation rate (Kimura 1968).
Meanwhile, CP12 rates mostly represent amino-acid substitutions (98%, 318/326), and are thus
susceptible to the mutation rate and the interaction between selection and population size (Ohta
1992).
Because rates at CP12 and CP3 are in part driven by different processes, the relationship
between the two provides additional information. The ratio between the two rates provides an
indication of rate of fixation relative to the rate of mutation, effectively cancelling out the
mutation rate and leaving the signal of the interaction between selection and population size
(Ohta 1992). For this reason, we also included models where the response variable was the ratio
between CP12 and CP3 rates. A high ratio can be a signature of historically small or fluctuating
population sizes, or it can indicate positive selection at the genomic loci examined. A low ratio
between rates can indicate the opposite historical signatures (i.e., large population sizes and/or
purifying selection).
We tested a single regression model for each variable of molecular evolution as response.
The explanatory variables in the model included the hand-wing index, environmental
temperature, and UV radiation. We also included body mass as an explanatory variable due to its
possible confounding effect of the association between the hand-wing index and molecular
evolution. In each model we also included two interaction terms that represented the possible

interactions among physiological surrogates and those among environmental variables. The

interaction terms therefore included that between body mass and hand-wing index, and the

interaction between the environmental energy and UV index.

To correct for non-independence due to relatedness between taxa, we used a species-level

phylogenetic estimate extracted from the original phylogenetic study (Derryberry *et al.* 2011).

We performed a transformation to many of the variables to adhere to least squares model

assumptions (Supplementary Table S1). In PGLS, variables are assumed to have evolved along

the phylogeny under a specified model of trait evolution. We used the lambda model of trait

evolution (Freckleton *et al.* 2002), which was the model with the best fit to our data compared

with three other common models (Brownian motion, kappa, and an Ornstein-Uhlenbeck process,

results not shown).

The parameters of PGLS regression models were optimized using the *phylolm* R package

(Ho & Ané 2014), and residuals were assessed for normality for each model. If no two-way

interactions were significant, the models were executed without these interactions. If this model

reduction did not significantly affect model fit according to a likelihood ratio test, then we also

interpreted the parameter estimates from the reduced model.

Results

We found that the hand-wing index and variables of environmental energy are positively

associated with molecular evolutionary rates, but not with the proxies of molecular evolution that

are most closely associated with mutation rates (CP3 which largely do not change amino-acids).

The overall molecular rates across all data are positively associated with hand-wing index ($P <$

0.001), environmental temperature (0.015), and UV radiation ($P < 0.001$; Table 1). Neither of the

two regression interaction terms were found to significantly explain overall molecular rates.

The evolutionary rate at the regions in the mitochondrial data that mostly change amino

acids (CP12) are associated with a negative interaction between body mass and hand-wing index

($P < 0.001$), which means that hand-wing index has a positive association with these molecular

rates primarily in small-bodied species (Table S1; Figure 1). Similarly, we found that the

evolutionary rates at nuclear regions that mostly change amino acids are positively associated

with the hand-wing index ($P = 0.038$). The proxies of molecular evolutionary rate that are

primarily impacted by selection and population size (CP12 / CP3), to the exclusion of the

mutation rate, were explained by UV radiation in both mitochondrial ($P < 0.001$) and nuclear

data ($P = 0.043$). In mitochondrial data this variable was also explained by the hand-wing index

($P < 0.001$) and environmental temperature ($P = 0.026$). Strikingly, none of the regression terms

explored explained our closest proxy to the mutation rate (CP3) in either mitochondrial or

nuclear data (Table S1).

Table 1. Overview of the PGLS regression results for the models tested for each of the variables

of molecular rates explored (rows). Shaded cells indicate terms that were not significant in

regression analyses. See Table S1 for full details on the variables and regression models tested.

Molecular rate response regression terms	Explanatory regression terms					
	Body mass	Hand-wing index	Temperature	UV radiation	Body mass × Hand-wing index	Temperature × UV radiation
Overall molecular rate		√	√	√		
Mitochondrial CP12	√	√			√	
Mitochondrial CP3						
Mitochondrial CP12 / CP3		√	√	√		
Nuclear CP12		√				
Nuclear CP3						
Nuclear CP12 / CP3				√		

Figure 1. The association between mitochondrial molecular rates at the 1st and 2nd codon

positions and hand-wing index across species of the avian family Furnariidae. Warmer colours

indicate species with greater body mass. The predicted regression line is added for the mean

body mass (orange), and each of the body masses one standard deviation from the mean (red and

yellow). Molecular rates are shown under a box-cox transformation; results with other
transformations are qualitatively identical.
246

Discussion

Our study demonstrates that the wing morphology, environmental temperature, and UV
radiation are positively associated with molecular evolutionary rates across the Neotropical avian
family Furnariidae. Strikingly, however, we did not find such association in the regions of loci
that largely reflect the mutation rate (i.e. substitution rate at the third codon position, CP3). The
effects of metabolism and environmental energy on the genome might therefore not occur via the
mutation rate, but instead through selective pressures or an effect on population size. This is
consistent with previous studies showing that flight has led to a broad range of genomic as well
as physiological adaptations (Tobalske *et al.* 2003; Wright *et al.* 2014). Our data are also
consistent with evidence that metabolic rates do not dictate the basal rate of evolution across taxa
(Lanfear *et al.* 2007), yet that environmental temperature is associated with molecular
evolutionary rates in birds (Gillman *et al.* 2012). Future genome-scale studies will be
instrumental to identify the selective role that the adaptation for flight and to high-energy
environments have in driving whole-genome evolution. Recent examples include work on the
selective impacts of skeletal structure and song on the genome (Zhang *et al.* 2014),
While theory predicts that metabolic rates can drive basal mutation rates (Allen *et al.*
2002; Gillooly *et al.* 2005), there are several reasons why this might not be the case. Most
organisms exist at an upper limit of mutations, such that even a minor increase in mutation rate
leads to an unsustainable loss in fitness (Bromham 2011). For example, large bodied mammals
have a greater opportunity to accumulate mutations than small bodied mammals. This is because
their long lifespan allows for more cell generations for producing gametes, which could interact
with the smaller population sizes to result in a faster accumulation of deleterious changes
(Bromham 2009). Nonetheless, instead of having a faster rate of accumulation of changes, large
bodied mammals have slow molecular evolutionary rates that are possibly due to the increased
selective pressures on DNA repair that avoids deleterious changes. Fast metabolic rates might
pose a similar increased selective pressure for avoiding deleterious mutations, and might explain
the several past studies that have not found an association between proxies of metabolic rate and
those of mutation rates (e.g., Lanfear *et al.* 2007; Rolland *et al.* 2016). The impact of
environmental temperature and UV radiation on mutation rates is also likely to be negligible in

species that regulate their internal energy levels and whose gametes do not undergo direct

exposure to environmental factors, such as most endotherms (but see Gillman *et al.* 2009). This

contrasts with the impact that temperature can have on the mutation rates in organisms that are

more exposed to environmental conditions, such as many marine taxa (Allen *et al.* 2006; Wright

et al. 2010b), plants (Davies *et al.* 2004; Wright *et al.* 2006; Gillman *et al.* 2010), amphibians

(Wright *et al.* 2010a), and reptiles (Garcia-Porta *et al.* 2019).

Independent sources of evidence show that metabolism has placed a set of selection

pressures on bird physiology and genomics. For instance, powered flight requires high energy

output (Suarez 1992) and efficiency (Kvist *et al.* 2001; Morris *et al.* 2010). The reduction in

genome size seen in many avian taxa is likely to be an adaptation associated with the demands of

powered flight (Wright *et al.* 2014). One explanation for our results showing increased rates in

the genomic regions that are associated with selective pressures (CP12 and CP12/CP3) is that the

demands of flight have favoured a broad range of evolutionary changes in the genome. Instead of

placing increased pressure for the genome to remain unchanged, flight has allowed for changes

in physiology and behaviour and remarkable opportunities of niche colonization (e.g., Gavrilov

2011; Claramunt *et al.* 2012a; Sol *et al.* 2012). Flight is a major driver of evolutionary change

across the genome (Zhang *et al.* 2014), and this is supported by our results of an association

between the hand-wing index and molecular evolution in both mitochondrial and nuclear

regions.

Substitution rates at sites that undergo selection are also influenced by population size,

often with a complex interaction between the two (Lanfear *et al.* 2014). For example, the

relationship between population size and the substitution rate takes drastically different shapes

depending on whether a genomic region is undergoing positive or negative selection. In cases of

genomic regions that are largely undergoing negative selection, as is common in protein-coding

loci, greater population sizes will increase the efficiency of selection and lead to a lower

substitution rate, even under a constant selection coefficient. This means that our data cannot

distinguish between an effect on selection or population size as caused by flight and

environmental energy. Therefore, our results could be caused by a negative association between

the population size and both the hand-wing index and environmental energy. However, evidence

largely supports the hand-wing index as being associated with dispersal ability and larger

effective population sizes (Claramunt *et al.* 2012b). Similarly, regions with greater

environmental energy have been associated with greater accumulation of biomass (Mittelbach *et*
al. 2007), so a negative association between our explanatory variables and population size seems
unlikely. Instead, it is likely that the impact of these variables on the genome occurs entirely via
patterns of selection.

Our results contribute to a body of work that suggests that the demands of flight and
environmental energy are not necessarily associated with basal mutation rates, but that instead
those factors have an impact on the genome through selection and adaptation. In birds, flight is
associated with a range of adaptations that lead to positive selection in large swathes of the
genome, and impact genome structure (Zhang *et al.* 2014). Similarly, adaptations to high energy
environments might include direct adaptations to thermoregulate. More complex traits to survive
in high-energy environments might be associated with the increased biotic interactions in these
environments, such as competitive and pathogenic interactions (Schemske *et al.* 2009; Díaz *et al.*
2013). Data sets of whole genomes and detailed physiology across bird taxa (e.g., Jarvis *et al.*
2014; Sheard *et al.* 2019), will bring a more complete picture of the impact of metabolism and
the environment on avian whole-genome evolution.
322

Acknowledgements
This work was supported by funding from the Australian Research Council to D.A.D. (grant
DE190100544).

References
Alerstam, T., Rosén, M., Bäckman, J., Ericson, P.G.P. & Hellgren, O. (2007). Flight speeds
among bird Species: Allometric and phylogenetic effects. *PLOS Biol.*, 5, e197.
Allen, A.P., Brown, J.H. & Gillooly, J.F. (2002). Global biodiversity, biochemical kinetics, and
the energetic-equivalence rule. *Science*, 297, 1545–1548.
Allen, A.P., Gillooly, J.F., Savage, V.M. & Brown, J.H. (2006). Kinetic effects of temperature
on rates of genetic divergence and speciation. *Proc. Natl. Acad. Sci. U. S. A.*, 103, 9130–
9135.
Barja, G. (1999). Mitochondrial oxygen radical generation and leak: Sites of production in States
4 and 3, organ specificity, and relation to aging and longevity. *J. Bioenerg. Biomembr.*, 31,
347–366.

Beckmann, M., Václavík, T., Manceur, A.M., Šprtová, L., von Wehrden, H., Welk, E., *et al.*

(2014). glUV: a global UV-B radiation data set for macroecological studies. *Methods Ecol.*

Evol., 5, 372–383.

Bishop, C.M., Butler, P.J., Egginton, S., el Haj, A.J. & Gabrielsen, G.W. (1995). Development of

metabolic enzyme activity in locomotor and cardiac muscles of the migratory barnacle

goose. *Am. J. Physiol. Integr. Comp. Physiol.*, 269, 64–72.

Bleiweiss, R. (1998). Relative-rate tests and biological causes of molecular evolution in

hummingbirds. *Mol. Biol. Evol.*, 15, 481–491.

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.-H., Xie, D., *et al.* (2014). BEAST 2:

A software platform for Bayesian evolutionary analysis. *PLOS Comput. Biol.*, 10,

e1003537.

Bromham, L. (2009). Why do species vary in their rate of molecular evolution? *Biol. Lett.*, 5,

401–404.

Bromham, L. (2011). The genome as a life-history character: why rate of molecular evolution

varies between mammal species. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, 366, 2503–13.

Bromham, L., Hua, X., Lanfear, R. & Cowman, P.F. (2015). Exploring the relationships between

mutation rates, life history, genome size, environment, and species richness in flowering

plants. *Am. Nat.*, 185, 507–524.

Bromham, L., Rambaut, A. & Harvey, P. (1996). Determinants of rate variation in mammalian

DNA sequence evolution. *J. Mol. Evol.*, 43, 610–621.

Burney, C.W. & Brumfield, R.T. (2009). Ecology predicts levels of genetic differentiation in

neotropical birds. *Am. Nat.*, 174, 358–68.

Butler, P.J. & Woakes, A.J. (1990). The physiology of bird flight. In: *Bird Migration*. Springer,

Heidelberg, pp. 300–318.

Claramunt, S. (2010). Discovering exceptional diversifications at continental scales: The case of

the endemic families of neotropical suboscine passerines. *Evolution*, 64, 2004–2019.

Claramunt, S., Derryberry, E.P., Brumfield, R.T. & Remsen, J. V. (2012a). Ecological

opportunity and diversification in a continental radiation of birds: climbing adaptations and

cladogenesis in the Furnariidae. *Am. Nat.*, 179, 649–66.

Claramunt, S., Derryberry, E.P., Remsen, J. V. & Brumfield, R.T. (2012b). High dispersal ability

inhibits speciation in a continental radiation of passerine birds. *Proc. R. Soc. B Biol. Sci.*,

279, 1567–1574.

Cooke, M.S., Evans, M.D., Dizdaroglu, M. & Lunec, J. (2003). Oxidative DNA damage:

Mechanisms, mutation, and disease. *FASEB J.*, 17, 1195–1214.

Currie, D.J. (1991). Energy and large-scale patterns of animal- and plant-species richness. *Am.*

Nat., 137, 27–49.

Davies, T.J., Savolainen, V., Chase, M.W., Moat, J. & Barraclough, T.G. (2004). Environmental

energy and evolutionary rates in flowering plants. *Proc. R. Soc. B Biol. Sci.*, 271, 2195–

2200.

Dawideit, B.A., Phillimore, A.B., Laube, I., Leisler, B. & Böhning-Gaese, K. (2009).

Ecomorphological predictors of natal dispersal distances in birds. *J. Anim. Ecol.*, 78, 388–

395.

Derryberry, E.P., Claramunt, S., Derryberry, G., Chesser, R.T., Cracraft, J., Aleixo, A., *et al.*

(2011). Lineage diversification and morphological evolution in a large-scale continental

radiation: The neotropical ovenbirds and woodcreepers (Aves: Furnariidae). *Evolution*, 65,

2973–2986.

Díaz, M., Møller, A.P., Flensted-Jensen, E., Grim, T., Ibáñez-Álamo, J.D., Jokimäki, J., *et al.*

(2013). The geography of fear: A latitudinal gradient in anti-predator escape distances of

birds across Europe. *PLOS One*, 8, e64634.

Drummond, A.J., Ho, S.Y.W., Phillips, M.J. & Rambaut, A. (2006). Relaxed phylogenetics and

dating with confidence. *PLOS Biol.*, 4, e88.

Duchêne, D.A. DA, Duchêne, S. & Ho, S.Y.W.S. (2018). PhyloMAd: Efficient assessment of

phylogenomic model adequacy. *Bioinformatics*, 34, 2300–2301.

Dunning, J.B. (2008). *CRC handbook of avian body masses 2nd Ed.* 2nd edn. Taylor & Francis

Group, Boca Raton, USA.

Fick, S.E. & Hijmans, R.J. (2017). WorldClim 2: new 1-km spatial resolution climate surfaces

for global land areas. *Int. J. Climatol.*, 37, 4302–4315.

Fjeldså, J., Irestedt, M. & Ericson, P.G. (2005). Molecular data reveal some major adaptational

shifts in the early evolution of the most diverse avian family, the Furnariidae. *J. Ornithol.*,

146, 1–13.

Freckleton, R.P., Harvey, P.H. & Pagel, M. (2002). Phylogenetic analysis and comparative data:

A test and review of evidence. *Am. Nat.*, 160, 712–726.

Galtier, N., Jobson, R.W., Nabholz, B., Glémin, S. & Blier, P.U. (2009). Mitochondrial whims:

metabolic rate, longevity and the rate of molecular evolution. *Biol. Lett.*, 5, 413–6.

Garcia-Porta, J., Irisarri, I., Kirchner, M., Rodríguez, A., Kirchhof, S., Brown, J.L., *et al.* (2019).

Environmental temperatures shape thermal physiology as well as diversification and

genome-wide substitution rates in lizards. *Nat. Commun.*, 10, 1–12.

Gavrilov, V.M. (2011). Energy expenditures for flight, aerodynamic quality, and colonization of

forest habitats by birds. *Biol. Bull.*, 38, 779–788.

Gillman, L.N., Keeling, D.J., Gardner, R.C. & Wright, S.D. (2010). Faster evolution of highly

conserved DNA in tropical plants. *J. Evol. Biol.*, 23, 1327–30.

Gillman, L.N., Keeling, D.J., Ross, H.A. & Wright, S.D. (2009). Latitude, elevation and the

tempo of molecular evolution in mammals. *Proc. Biol. Sci.*, 276, 3353–9.

Gillman, L.N., McCowan, L.S.C. & Wright, S.D. (2012). The tempo of genetic evolution in

birds: body mass and climate effects. *J. Biogeogr.*, 39, 1567–1572.

Gillooly, J.F., Allen, A.P., West, G.B. & Brown, J.H. (2005). The rate of DNA evolution: effects

of body size and temperature on the molecular clock. *Proc. Natl. Acad. Sci. U. S. A.*, 102,

140–5.

Gillooly, J.F., Brown, J.H., West, G.B., Savage, V.M. & Charnov, E.L. (2001). Effects of size

and temperature on metabolic rate. *Science*, 293, 2248–2251.

Gillooly, J.F., McCoy, M.W. & Allen, A.P. (2007). Effects of metabolic rate on protein

evolution. *Biol. Lett.*, 3, 655–659.

Guglielmo, C.G., Haunerland, N.H., Hochachka, P.W. & Williams, T.D. (2002). Seasonal

dynamics of flight muscle fatty acid binding protein and catabolic enzymes in a migratory

shorebird. *Am. J. Physiol. Integr. Comp. Physiol.*, 282, 1405–1413.

Ho, L.S.T. & Ané, C. (2014). A linear-time algorithm for gaussian and non-gaussian trait

evolution models. *Syst. Biol.*, 63, 397–408.

del Hoyo, J., Christie, D., Elliott, A., Fishpool, L.D.C., Boesman, P., Collar, N.J., *et al.* (2016).

HBW and BirdLife International illustrated checklist of the birds of the world: passerines.

Lynx Edicions, Barcelona.

Jarvis, E.D., Mirarab, S., Aberer, A.J., Li, B., Houde, P., Li, C., *et al.* (2014). Whole-genome

analyses resolve early branches in the tree of life of modern birds. *Science*, 346, 1320–1331.

Jenni-Eiermann, S., Jenni, L., Kvist, A., Lindström, Å., Piersma, T. & Visser, G.H. (2002). Fuel

use and metabolic response to endurance exercise: a wind tunnel study of a long-distance migrant shorebird. *J. Exp. Biol.*, 205, 2453–2460.

Jenni-Eiermann, S., Jenni, L., Smith, S. & Costantini, D. (2014). Oxidative stress in endurance flight: An unconsidered factor in bird migration. *PLOS One*, 9, e97650.

Kimura, M. (1968). Evolutionary rate at the molecular level. *Nature*, 217, 624–626.

Kipp, F.A. (1959). Der Handflügel-Index als flugbiologisches Maß. *Vogelwarte*, 20, 77–86.

Kvist, A., Lindström, Å., Green, M., Piersma, T. & Visser, G.H. (2001). Carrying large fuel loads during sustained bird flight is cheaper than expected. *Nature*, 413, 730–732.

Lanfear, R., Ho, S.Y.W., Jonathan Davies, T., Moles, A.T., Aarssen, L., Swenson, N.G., *et al.* (2013). Taller plants have lower rates of molecular evolution. *Nat. Commun.*, 4, 1879.

Lanfear, R., Ho, S.Y.W., Love, D. & Bromham, L. (2010). Mutation rate is linked to diversification in birds. *Proc. Natl. Acad. Sci.*, 107, 20423–20428.

Lanfear, R., Kokko, H. & Eyre-Walker, A. (2014). Population size and the rate of evolution. *Trends Ecol. Evol.*, 29, 33–41.

Lanfear, R., Thomas, J. a, Welch, J.J., Brey, T. & Bromham, L. (2007). Metabolic rate does not calibrate the molecular clock. *Proc. Natl. Acad. Sci. U. S. A.*, 104, 15388–93.

Lockwood, R., Swaddle, J.P. & Rayner, J.M. V. (1998). Avian wingtip shape reconsidered: Wingtip shape indices and morphological adaptations to migration. *J. Avian Biol.*, 29, 273.

Maillet, D. & Weber, J.M. (2007). Relationship between n-3 PUFA content and energy metabolism in the flight muscles of a migrating shorebird: Evidence for natural doping. *J. Exp. Biol.*, 210, 413–420.

Martin, A.P. & Palumbi, S.R. (1993). Body size, metabolic rate, generation time, and the molecular clock. *Proc. Natl. Acad. Sci. U. S. A.*, 90, 4087–4091.

Mittelbach, G.G., Schemske, D.W., Cornell, H. V, Allen, A.P., Brown, J.M., Bush, M.B., *et al.* (2007). Evolution and the latitudinal diversity gradient: speciation, extinction and biogeography. *Ecol. Lett.*, 10, 315–31.

Mooers, A. & Harvey, P.H. (1994). Metabolic rate, generation time, and the rate of molecular evolution in birds. *Mol. Phylogenet. Evol.*, 3, 344–350.

Morris, C.R., Nelson, F.E. & Askew, G.N. (2010). The metabolic power requirements of flight and estimations of flight muscle efficiency in the cockatiel (Nymphicus hollandicus). *J. Exp. Biol.*, 213, 2788–2796.

Norberg, U.M. (1995). How a long tail and changes in mass and wing shape affect the cost for
flight in animals. *Funct. Ecol.*, 9, 48.

Ohta, T. (1992). The nearly neutral theory of molecular evolution. *Annu. Rev. Ecol. Syst.*, 23,
263–286.

Pawlowski, J., Bolivar, I., Fahrni, J.F., de Vargas, C., Gouy, M. & Zaninetti, L. (1997). Extreme
differences in rates of molecular evolution of foraminifera revealed by comparison of
ribosomal DNA sequences and the fossil record. *Mol. Biol. Evol.*, 14, 498–505.

Pennycuick, C.J. (2008). *Modelling the flying bird. Theor. Ecol. Ser.* Academic Press,
Amsterdam.

Puurtinen, M., Elo, M., Jalasvuori, M., Kahilainen, A., Ketola, T., Kotiaho, J.S., *et al.* (2016).
Temperature-dependent mutational robustness can explain faster molecular evolution at
warm temperatures, affecting speciation rate and global patterns of species diversity.
Ecography (Cop.)., 39, 1025–1033.

Remsen Jr, J. V, Cadena, C.D., Jaramillo, A., Nores, M., Pacheco, J.F., Pérez-Emán, J., *et al.*
(2012). A classification of the bird species of South America. *Am. Ornithol. Union*.

Rolland, J., Loiseau, O., Romiguier, J. & Salamin, N. (2016). Molecular evolutionary rates are
not correlated with temperature and latitude in Squamata: an exception to the metabolic
theory of ecology? *BMC Evol. Biol.*, 16, 95.

Schemske, D.W., Mittelbach, G.G., Cornell, H. V., Sobel, J.M. & Roy, K. (2009). Is there a
latitudinal gradient in the importance of biotic interactions? *Annu. Rev. Ecol. Evol. Syst.*, 40,
245–269.

Sheard, C., Neate-Clegg, M.H.C., Alioravainen, N., Jones, S.E.I., Vincent, C., MacGregor,
H.E.A., *et al.* (2019). The latitudinal gradient in hand-wing-index: global patterns and
predictors of wing morphology in birds. *bioRxiv*, 816603.

Sibley, C.G. & Monroe, B.L. (1990). *Distribution and taxonomy of birds of the world*. Yale
University Press, New Haven, Connecticut.

Sol, D., Bartomeus, I. & Griffin, A.S. (2012). The paradox of invasion in birds: Competitive
superiority or ecological opportunism? *Oecologia*, 169, 553–564.

Suarez, R.K. (1992). Hummingbird flight: Sustaining the highest mass-specific metabolic rates
among vertebrates. *Experientia*, 48, 565–570.

Tavaré, S. (1986). Some probabilistic and statistical problems in the analysis of DNA sequences.

Lect. Math. Life Sci., 17, 57–86.

Tobalske, B.W., Hedrick, T.L., Dial, K.P. & Biewener, A.A. (2003). Comparative power curves

in bird flight. *Nature*, 421, 363–366.

Turner, J.R.G., Lennon, J.J. & Lawrenson, J.A. (1988). British bird species distributions and the

energy theory. *Nature*, 335, 539–541.

Ward, S., Bishop, C.M., Woakes, A.J. & Butler, P.J. (2002). Heart rate and the rate of oxygen

consumption of flying and walking barnacle geese (Branta leucopsis) and bar-headed geese

(Anser indicus). *J. Exp. Biol.*, 205, 3347–3356.

Ward, S., Möller, U., Rayner, J.M., Jackson, D.M., Bilo, D., Nachtigall, W., *et al.* (2001).

Metabolic power, mechanical power and efficiency during wind tunnel flight by the

European starling Sturnus vulgaris. *J. Exp. Biol.*, 204, 3311–22.

Weber, J.M. (2009). The physiology of long-distance migration: Extending the limits of

endurance metabolism. *J. Exp. Biol.*

Willig, M.R., Kaufman, D.M. & Stevens, R.D. (2003). Latitudinal gradients of biodiversity:

Pattern, process, scale, and synthesis. *Annu. Rev. Ecol. Evol. Syst.*, 34, 273–309.

Wright, D.H. (1983). Species-energy theory: An extension of species-area theory. *Oikos*, 41,

496.

Wright, N.A., Gregory, T.R. & Witt, C.C. (2014). Metabolic "engines" of flight drive genome

size reduction in birds. *Proc. R. Soc. B Biol. Sci.*, 281, 20132780.

Wright, S., Keeling, J. & Gillman, L. (2006). The road from Santa Rosalia: a faster tempo of

evolution in tropical climates. *Proc. Natl. Acad. Sci. U. S. A.*, 103, 7718–7722.

Wright, S.D., Gillman, L.N., Ross, H.A. & Keeling, D.J. (2010a). Energy and the tempo of

evolution in amphibians. *Glob. Ecol. Biogeogr.*, 19, 733–740.

Wright, S.D., Ross, H.A., Jeanette Keeling, D., McBride, P. & Gillman, L.N. (2010b). Thermal

energy and the rate of genetic evolution in marine fishes. *Evol. Ecol.*, 25, 525–530.

Zhang, G., Li, C., Li, Q., Li, B., Larkin, D.M., Lee, C., *et al.* (2014). Comparative genomics

reveals insights into avian genome evolution and adaptation. *Science*, 346, 1311–20.

Table S1. Complete results of PGLS regression models tested. Transformations used are shown

beside variables in parentheses. The full model notation used in *phylolm* is shown, with

successive model reductions indicated in letters (1a, 1b, etc.), where applicable.

1a

Overall molecular rate (log) ~ Body mass (Box-Cox) + Hand-wing index + Mean annual temperature + Mean UV radiation + Body mass (Box-Cox):Hand-wing index + Mean annual temperature:Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	-3.103	2.097	-1.480	0.140
Body mass (Box-Cox)	0.882	1.028	0.858	0.392
Hand-wing index	0.180	0.110	1.640	0.102
Mean annual temperature	0.016	0.032	0.479	0.633
Mean UV radiation	9.08×10^{-5}	7.38×10^{-5}	1.231	0.220
Body mass (Box-Cox):Hand-wing index	-0.051	0.056	-0.911	0.363
Mean annual temperature:Mean UV radiation	9.28×10^{-7}	5.72×10^{-6}	0.162	0.871

1b

Overall molecular rate (log) ~ Body mass (Box-Cox) + Hand-wing index + Mean annual temperature + Mean UV radiation + Body mass (Box-Cox):Hand-wing index

	Estimate	Standard error	t-value	p-value
(Intercept)	-3.105	2.092	-1.484	0.139
Body mass (Box-Cox)	0.847	1.003	0.845	0.399
Hand-wing index	0.176	0.107	1.649	0.101
Mean annual temperature	0.021	0.008	2.443	0.015
Mean UV radiation	1.02×10^{-4}	2.76×10^{-5}	3.688	<0.001
Body mass (Box-Cox):Hand-wing index	-0.049	0.054	-0.900	0.369

1c

Overall molecular rate (log) ~ Body mass (Box-Cox) + Hand-wing index + Mean annual temperature + Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	-1.457	1.013	-1.438	0.152
Body mass (Box-Cox)	0.017	0.394	0.043	0.966
Hand-wing index	0.081	0.014	5.866	<0.001
Mean annual temperature	0.021	0.008	2.439	0.015
Mean UV radiation	1.00×10^{-4}	2.76×10^{-5}	3.639	<0.001

2a

Molecular CP12 rate (Box-Cox) ~ Body mass (Box-Cox) + Hand-wing index + Mean annual temperature + Mean UV radiation + Body mass (Box-Cox):Hand-wing index + Mean annual temperature:Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	-8.091	2.320	-3.487	0.001
Body mass (Box-Cox)	3.489	1.138	3.066	0.002
Hand-wing index	0.496	0.122	4.075	<0.001
Mean annual temperature	0.044	0.036	1.233	0.219
Mean UV radiation	1.54×10^{-4}	8.17×10^{-5}	1.882	0.061
Body mass (Box-Cox):Hand-wing index	-0.218	0.062	-3.534	<0.001
Mean annual temperature:Mean UV radiation	-4.11×10^{-6}	6.34×10^{-6}	-0.648	0.518

3a

Molecular CP3 rate (Box-Cox) ~ Body mass (Box-Cox) + Hand-wing index + Mean annual temperature + Mean UV radiation + Body mass (Box-Cox):Hand-wing index + Mean annual temperature:Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	2.387	3.753	0.636	0.525
Body mass (Box-Cox)	1.105	1.840	0.601	0.549
Hand-wing index	0.134	0.197	0.681	0.496
Mean annual temperature	0.020	0.058	0.349	0.728
Mean UV radiation	3.84×10^{-5}	1.32×10^{-4}	0.291	0.771
Body mass (Box-Cox):Hand-wing index	-0.065	0.100	-0.653	0.515
Mean annual temperature:Mean UV radiation	-3.42×10^{-6}	1.02×10^{-5}	-0.334	0.739

3b

Molecular CP3 rate (Box-Cox) ~ Body mass (Box-Cox) + Hand-wing index + Mean annual temperature + Mean UV radiation + Body mass (Box-Cox):Hand-wing index

	Estimate	Standard error	t-value	p-value
(Intercept)	2.395	3.746	0.640	0.523
Body mass (Box-Cox)	1.234	1.796	0.687	0.493
Hand-wing index	0.149	0.191	0.776	0.438
Mean annual temperature	0.002	0.015	0.100	0.920
Mean UV radiation	-2.48×10^{-6}	4.95×10^{-5}	-0.050	0.960
Body mass (Box-Cox):Hand-wing index	-0.073	0.097	-0.759	0.449

3c

Molecular CP3 rate (Box-Cox) ~ Body mass (Box-Cox) + Hand-wing index + Mean annual temperature + Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	4.880	1.814	2.691	0.008
Body mass (Box-Cox)	-0.019	0.706	-0.027	0.978
Hand-wing index	0.005	0.025	0.187	0.852
Mean annual temperature	0.001	0.015	0.096	0.923
Mean UV radiation	-4.89×10^{-6}	4.93×10^{-5}	-0.099	0.921

4a

CP12/CP3 ratio (Box-Cox) ~ Body mass (Box-Cox) + Hand-wing index + Mean annual temperature + Mean UV radiation + Body mass (Box-Cox):Hand-wing index + Mean annual temperature:Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	-14.566	4.868	-2.992	0.003
Body mass (Box-Cox)	3.653	2.387	1.530	0.127
Hand-wing index	0.600	0.255	2.351	0.020
Mean annual temperature	0.020	0.075	0.265	0.791
Mean UV radiation	1.76×10^{-4}	1.71×10^{-4}	1.026	0.306
Body mass (Box-Cox):Hand-wing index	-0.244	0.129	-1.887	0.060
Mean annual temperature:Mean UV radiation	4.44×10^{-6}	1.33×10^{-5}	0.334	0.738

4b

CP12/CP3 ratio (Box-Cox) ~ Body mass (Box-Cox) + Hand-wing index + Mean annual temperature + Mean UV radiation + Body mass (Box-Cox):Hand-wing index

	Estimate	Standard error	t-value	p-value
(Intercept)	-14.577	4.859	-3.000	0.003
Body mass (Box-Cox)	3.486	2.330	1.496	0.136
Hand-wing index	0.581	0.248	2.339	0.020
Mean annual temperature	0.044	0.020	2.260	0.025
Mean UV radiation	2.29×10^{-4}	6.42×10^{-5}	3.566	<0.001
Body mass (Box-Cox):Hand-wing index	-0.234	0.125	-1.866	0.063

4c

CP12/CP3 ratio (Box-Cox) ~ Body mass (Box-Cox) + Hand-wing index + Mean annual temperature + Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	-6.648	2.367	-2.808	0.005

	Estimate	Standard error	t-value	p-value
Body mass (Box-Cox)	-0.511	0.921	-0.555	0.580
Hand-wing index	0.121	0.032	3.775	<0.001
Mean annual temperature	0.044	0.020	2.238	0.026
Mean UV radiation	2.21×10^{-4}	6.44×10^{-5}	3.435	0.001

5a

Molecular CP12 rate (log) ~ Body mass (log) + Hand-wing index + Mean annual temperature + Mean UV radiation + logmass:Hand-wing index + Mean annual temperature:Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	-0.384	0.671	-0.572	0.569
Body mass (log)	-0.057	0.166	-0.340	0.735
Hand-wing index	-0.003	0.032	-0.088	0.931
Mean annual temperature	0.015	0.026	0.560	0.578
Mean UV radiation	4.69×10^{-5}	6.92×10^{-5}	0.677	0.501
logmass:Hand-wing index	0.005	0.009	0.478	0.634
Mean annual temperature:Mean UV radiation	-2.66×10^{-6}	4.74×10^{-6}	-0.562	0.576

5b

Molecular CP12 rate (log) ~ Body mass (log) + Hand-wing index + Mean annual temperature + Mean UV radiation + Mean annual temperature:Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	-0.614	0.463	-1.326	0.190
Body mass (log)	0.020	0.047	0.424	0.673
Hand-wing index	0.012	0.006	2.114	0.038
Mean annual temperature	0.013	0.026	0.498	0.620
Mean UV radiation	4.5×10^{-5}	6.88×10^{-5}	0.654	0.515
Mean annual temperature:Mean UV radiation	-2.39×10^{-6}	4.68×10^{-6}	-0.512	0.610

5c

Molecular CP12 rate (log) ~ Body mass (log) + Hand-wing index + Mean annual temperature + Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	-0.429	0.288	-1.492	0.140
Body mass (log)	0.023	0.046	0.492	0.625
Hand-wing index	0.012	0.006	2.068	0.043
Mean annual temperature	-1.30×10^{-4}	0.005	-0.028	0.978
Mean UV radiation	1.14×10^{-5}	2.04×10^{-5}	0.557	0.579

6a

Molecular CP3 rate (square root) ~ Body mass (log) + Hand-wing index + Mean annual temperature + Mean UV radiation + logmass:Hand-wing index + Mean annual temperature:Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	2.021	1.499	1.348	0.183
Body mass (log)	0.270	0.372	0.726	0.471
Hand-wing index	0.059	0.071	0.835	0.407
Mean annual temperature	-0.042	0.058	-0.727	0.470
Mean UV radiation	-9.95×10^{-5}	1.55×10^{-4}	-0.642	0.523
Body mass (log):Hand-wing index	-0.014	0.021	-0.656	0.514
Mean annual temperature:Mean UV radiation	8.75×10^{-6}	1.06×10^{-5}	0.826	0.412

6b

Molecular CP3 rate (square root) ~ Body mass (log) + Hand-wing index + Mean annual temperature + Mean UV radiation + Mean annual temperature:Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	2.728	1.038	2.629	0.011
Body mass (log)	0.036	0.105	0.344	0.732
Hand-wing index	0.014	0.013	1.051	0.297
Mean annual temperature	-0.037	0.057	-0.641	0.524
Mean UV radiation	-9.37×10^{-5}	1.54×10^{-4}	-0.608	0.545
Mean annual temperature:Mean UV radiation	7.93×10^{-6}	1.05×10^{-5}	0.757	0.452

6c

Molecular CP3 rate (square root) ~ Body mass (log) + Hand-wing index + Mean annual temperature + Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	2.114	0.645	3.276	0.002
Body mass (log)	0.027	0.104	0.256	0.799
Hand-wing index	0.015	0.013	1.211	0.230
Mean annual temperature	0.006	0.010	0.565	0.574
Mean UV radiation	1.76×10^{-5}	4.59×10^{-5}	0.384	0.702

7a

CP12/CP3 ratio ~ Body mass (log) + Hand-wing index + Mean annual temperature + Mean UV radiation + logmass:Hand-wing index + Mean annual temperature:Mean UV radiation

	Estimate	Standard	t-value	p-value

	Estimate	Standard error	t-value	p-value
		error		
(Intercept)	0.365	0.496	0.737	0.464
Body mass (log)	-0.144	0.123	-1.173	0.245
Hand-wing index	-0.025	0.023	-1.084	0.283
Mean annual temperature	0.033	0.019	1.703	0.094
Mean UV radiation	5.48×10^{-5}	5.13×10^{-5}	1.070	0.289
Body mass (log):Hand-wing index	0.007	0.007	1.014	0.314
Mean annual temperature:Mean UV radiation	-6.4×10^{-5}	3.5×10^{-5}	-1.826	0.073

7b

CP12/CP3 ratio ~ Body mass (log) + Hand-wing index + Mean annual temperature + Mean UV radiation + Mean annual temperature:Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	0.004	0.345	0.012	0.991
Body mass (log)	-0.025	0.035	-0.710	0.480
Hand-wing index	-0.002	0.004	-0.476	0.635
Mean annual temperature	0.030	0.019	1.572	0.121
Mean UV radiation	5.18×10^{-5}	5.12×10^{-5}	1.013	0.315
Mean annual temperature:Mean UV radiation	-5.97×10^{-6}	3.48×10^{-6}	-1.717	0.091

7c

CP12/CP3 ratio ~ Body mass (log) + Hand-wing index + Mean annual temperature + Mean UV radiation

	Estimate	Standard error	t-value	p-value
(Intercept)	0.466	0.218	2.137	0.036
Body mass (log)	-0.018	0.035	-0.501	0.618
Hand-wing index	-0.003	0.004	-0.788	0.434
Mean annual temperature	-0.002	0.004	-0.631	0.530
Mean UV radiation	-3.2×10^{-5}	1.55×10^{-5}	-2.064	0.043