

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

Regeneration in the adult *Drosophila* brain

Kassi L. Crocker^{1,2,3}, Khailee Marischuk^{1,3}, Stacey A. Rimkus^{3#}, Hong Zhou⁴, Jerry C.P. Yin⁴ and Grace Boekhoff-Falk^{1,3*}

¹Genetics Graduate Training Program

²Science and Medicine Graduate Research Scholars Program

³Department of Cell and Regenerative Biology

⁴Department of Genetics

University of Wisconsin-Madison

School of Medicine and Public Health

1111 Highland Avenue

Madison, WI 53705

#current address: Laboratory of Genetics, University of Wisconsin-Madison 53706

*corresponding author

Tel: 608-262-1609

Fax: 608-262-7306

Email: grace.boekhoff@wisc.edu

52 **Abstract**

53 Understanding the molecular and cellular mechanisms underlying neurogenesis after injury is
54 crucial for developing tools for brain repair. We have established an adult *Drosophila*
55 *melanogaster* model for investigating regeneration after central brain injury. Within 24 hours
56 after Penetrating Traumatic Brain Injury (PTBI) to the central brain, we observe a significant
57 increase in the number of proliferating cells. Between one- and two-weeks post-injury, we detect
58 the generation of new neurons and glia and the formation of new axon tracts that target
59 appropriate brain regions, suggesting there could be functional regeneration. Consistent with
60 functional regeneration, locomotion abnormalities observed shortly after PTBI are largely
61 reversed within 2 weeks of injury. Further, we find that cells surrounding the injury site
62 upregulate neuroblast genes, such as *asense* and *deadpan*, and demonstrate that these cells
63 give rise to the new neurons and glia. Taken together, our data support the hypothesis that
64 young, adult *Drosophila* brains are capable of neuronal repair after central brain injury. We
65 anticipate that our model will facilitate the dissection of the mechanisms of neural regeneration
66 and anticipate that these processes will have relevance to humans.

67 **Introduction**

68 Different species vary in their ability to regenerate, and within a species, there is tissue to tissue
69 variation in regenerative capacity. (Birnbaum and Sanchez Alvarado, 2008, Bollini et al., 2018,
70 Michalopoulos, 2007, Stanger, 2015). One of the most arguably important organs, the human
71 brain, has become the center of intense investigation regarding the extent to which it can
72 produce new cells in adulthood. Classical studies reported that the brain stopped making new
73 neurons shortly after birth (Ramon y Cajal, 1913, Ramon y Cajal, 1914). However, in 1998, a
74 landmark study concluded that adult humans are able to create new neurons in the
75 hippocampus (Eriksson et al., 1998). Furthermore, this ability is retained throughout later stages
76 of life. These findings have been corroborated (Reif et al., 2006, Manganas et al., 2007),
77 although the exact number varies among studies (Spalding et al., 2013, Dennis et al., 2016,
78 Eriksson et al., 1998, Knoth et al., 2010). Populations of dividing progenitor cells now have been
79 observed in two major regions of the rodent brain: the subventricular zone (SVZ) of the forebrain
80 and the dentate gyrus (DG) of the hippocampus (Altman and Das, 1965) (Altman, 1969) (Kaplan
81 and Hinds, 1977). The progenitors in both brain regions generate multiple cell types, including
82 neurons, astrocytes and oligodendrocytes, over the course of the animal's lifetime (Kuhn et al.,
83 1996).

84 Identification of the cell types capable of regenerating brain structures and
85 understanding the underlying molecular mechanisms that regulate neuroregeneration is central
86 to developing novel therapeutics for neural degenerative diseases and treatment of brain
87 injuries. Analysis of SVZ and DG neural stem cells continues to provide critical insights into
88 neural regeneration. However, these cells are unlikely to provide progenitors for other brain
89 regions. Therefore, emphasis in the neuroregeneration field has been on transplanting
90 embryonic or induced pluripotent stem cells (Vishwakarma et al., 2014). However, stem cell
91 transplants often are accompanied by tumor formation (Amariglio et al., 2009) and obtaining
92 functional integration of transplanted neural cells remains a major challenge. A more recent

93 focus for the field therefore has been on coaxing resident cells in the brain to undertake
94 regeneration (Gao et al., 2016). This avenue of investigation would be facilitated by the
95 availability of a model system that mimics key aspects of mammalian neurodegeneration
96 phenotypes but is more experimentally accessible.

97 For over a century, the fruitfly *Drosophila melanogaster* has been used as a model
98 organism, furthering many fields, including developmental genetics and neuroscience. Although
99 *Drosophila* are invertebrates, there are many parallels between flies and mammals, and
100 discoveries from *Drosophila* repeatedly have proven relevant to human biology. A *Drosophila*
101 brain is composed of approximately 90,000 neurons (Chiang et al., 2011). While this is a million-
102 fold fewer than the average human brain (Meinertzhagen, 2010), both human and *Drosophila*
103 brains are complex and have analogous neural cell types (Lessing and Bonini, 2009). Common
104 neurotransmitters (GABA, glutamate, and acetylcholine) and biogenic amines (dopamine and
105 serotonin) are utilized in both humans and *Drosophila* (Bellen et al., 2010). In addition, human
106 and *Drosophila* neurons function in similar ways, with neurotransmission occurring at synapses.
107 Even the architecture of synapses is conserved between humans and *Drosophila* (Lessing and
108 Bonini, 2009). Humans and *Drosophila* also have similar physiologies, including homologous
109 sodium and calcium channels that are capable of producing action potentials and regulating
110 membrane potential (Bellen et al., 2010).

111 *Drosophila* have been utilized extensively to study neural development (Spindler and
112 Hartenstein, 2010, Sasse et al., 2015). Most of this research has focused on larval and pupal
113 stages and includes the identification of neural progenitor cells in the brain. During
114 neurogenesis in the embryonic and larval stages of *Drosophila*, neuroblasts (NBs) undergo
115 multiple rounds of asymmetric division to generate one larger and one smaller daughter cell
116 (Homem et al., 2015). At each division, cell-fate determinants are distributed from the NB to the
117 daughter cells. Numb, Prospero (Pros), and Brain tumor (Brat) are partitioned to the basal cell
118 cortex, and atypical kinase C (aPKC) to the apical cell cortex. Therefore, when mitosis is

119 complete, each daughter cell inherits a different set of determinants. As a result, the apical
120 daughter cell remains an NB and continues to proliferate while the basal daughter cell becomes
121 committed to differentiation. Central brain neurons and glia derive from two types of NBs (Boone
122 and Doe, 2008, Egger et al., 2008, Homem and Knoblich, 2012, Weng and Lee, 2011, Homem
123 et al., 2015). Type I NBs express the transcription factors Deadpan (Dpn) and Asense (Ase)
124 and undergo asymmetric cell division to self-renew and produce ganglion mother cells (GMCs)
125 (Knoblich, 2010). Each GMC then divides symmetrically to produce two neurons or glial cells.
126 Type II neuroblasts express Dpn, but not Ase and also self-renew, but, unlike type I NBs, give
127 rise to intermediate neural progenitors (INPs) (Boone and Doe, 2008, Bello et al., 2008,
128 Bowman et al., 2008). INPs undergo a maturation process, during which they do not divide but
129 sequentially initiate expression of the transcription factors Ase and Dpn. After this initial stage,
130 each mature INP undergoes three to five additional rounds of asymmetric division, with each
131 round generating another INP and a GMC that divides terminally into two neurons or glial cells.
132 Other transcription factors required during neurogenesis in the Type I and/or Type II lineages
133 include Prospero (Pros), Inscuteable (Insc), and Earmuff (Erm) which are required for
134 asymmetric cell division (Chia et al., 2008).

135 While neurogenesis is very well studied during development, there remain gaps in our
136 knowledge about homeostasis and response to damage, particularly in the adult brain. This can
137 be partly attributed to previous studies indicating that the adult *Drosophila* brain has limited
138 mitotic activity (~1 dividing cell/brain) (von Trotha et al., 2009) and the fact that known neural
139 progenitors undergo terminal differentiation or apoptosis during metamorphosis (Siegrist et al.,
140 2010). Both of these characteristics are very similar to mammalian brains (Ming and Song,
141 2011, Stiles and Jernigan, 2010). The combination of rare cell proliferation and NB death or
142 terminal differentiation has made the adult *Drosophila* brain an unlikely candidate model in
143 which to investigate neuroregeneration. Nonetheless, other studies have demonstrated that the
144 adult *Drosophila* is capable of neurogenesis after injury (Fernandez-Hernandez et al., 2013,

145 Moreno et al., 2015). In fact, slowly cycling neural progenitor cells were discovered in the
146 medulla cortex of the optic lobes. These progenitors are activated upon injury (Fernandez-
147 Hernandez et al., 2013) and create new neurons in response to damage. These data support
148 the idea that if there are slowly cycling cells in the central brain, they also may be able to
149 proliferate post-injury and function in regeneration.

150 The limited information regarding not only the regeneration potential of the adult
151 *Drosophila* brain, led us to examine the regenerative potential of the adult *Drosophila* central
152 brain after injury. We have developed a novel model of brain injury that we call Penetrating
153 Traumatic Brain Injury (PTBI). To inflict PTBI, we use a sterilized metal needle to penetrate the
154 adult head cuticle and puncture the brain. Our goal is to fully exploit the adult *Drosophila* brain
155 to characterize its neurogenic potential post-injury and discover mechanisms that control
156 neuroregeneration. This will further our knowledge of development and homeostasis of the
157 nervous system and may lead to novel neurodegenerative disease therapies.

158

159 **Results**

160 PTBI as a new model for neuroregeneration

161 To investigate the neuroregenerative capacity of the adult *Drosophila melanogaster* central
162 brain, we first needed to develop a reproducible injury method. The model we devised is called
163 penetrating traumatic brain injury (PTBI). To induce PTBI, we employed a method of injury
164 termed the penetrating small targeted area of the brain (STAB) injury. This was accomplished
165 using a thin metal insect needle (~12.5µm diameter tip, 100 µm diameter rod) to penetrate the
166 head cuticle and the brain, specifically near the mushroom body (MB; **Fig. 1A, B**). We focused
167 on the mushroom body (MB), a region of the central brain critical for learning and memory (Aso
168 et al., 2014), because MB neuroblasts are the last to stop proliferating during development
169 (Siegrist et al., 2010, Ito and Hotta, 1992, Ito et al., 1997). We therefore reasoned that the MB
170 may have the most mitotic potential. Located dorsally in the central brain, the MB is made up of

171 complex dendrite and axon arbors in large and highly stereotyped arrays. This architecture
172 makes it useful for assaying the regeneration of neurites (Aso et al., 2014). Another factor that
173 we took advantage of is that the cell bodies of the MB lie at the surface of the brain and can be
174 visualized through the head cuticle using the expression of green fluorescent protein (GFP),
175 facilitating for reproducible damage to the MB.

176 Surprisingly, when we injured 0-6-hour post-eclosion *OK107/yw* adult males and allowed
177 them to recover for 24 hours, we saw no significant decrease in survival of injured males
178 compared to uninjured age-matched controls, with each having an approximate 99% survival
179 rate (**Fig. 1G**). Yet, when adult *OK107/yw* males were aged to 14 days post-eclosion, and then
180 injured, there was a significant increase in mortality to ~13% following PTBI (p-value<0.0001)
181 (**Fig. 1G**). A similar trend was seen when flies were aged to 28 days then injured, with mortality
182 increasing further to ~25% (p-value<0.006). Thus, as flies grow older, their ability to survive a
183 PTBI decreases. This trend showed that resilience in response to injury is specific to younger
184 flies. In subsequent experiments, we therefore focused on injuring flies that were 0-6 hours old.

185 To quantify the neurodegeneration caused by PTBI, we used a standardized index (Cao
186 et al., 2013) to analyze histological samples from brains 25 days after injury. We observed that
187 there was some increase neurodegeneration following PTBI compared to age-matched
188 uninjured controls (**Fig. 1C-E**). While younger flies were able to survive PTBI for at least 24
189 hours, we wanted to see how this would impact their lifespan. We therefore examined the
190 lifespan of *OK107/yw* injured males and observed a substantially reduced lifespan compared to
191 uninjured controls, with only 50% survival at around 48 days and no survivors at approximately
192 74 days post-injury (**Fig. 1F**).

193

194 PTBI stimulates cell proliferation

195 Because one of the first signs of regeneration is cell proliferation, we assayed cell division post-
196 injury using a mitotic marker, phospho-histone H3 (PH3). Histone H3 is transiently

197 phosphorylated during M phase of the cell cycle (Hans and Dimitrov, 2001). Brains from
198 *OK107/yw* adult males that were assayed 24 hours post-PTBI using anti-PH3 had a significant
199 increase of PH3-positive cells, around 11 cells, compared to uninjured controls of the same sex,
200 age, and genotype, in which we observed about 2 cells per brain (p -value<0.0001)(**Fig. 2A-D,**
201 **I**). These PH3-positive cells primarily were located near the area of injury. Similar numbers of
202 proliferating cells were seen at 24 hours using the mitotic marker 5-ethynyl-2'-deoxyuridine
203 (EdU) (**Fig. 3-S1**). To quantify this, we mapped each PH3-positive cell to a hemisphere within
204 the central brain: left or right, with the right hemisphere the location of injury (**Fig. 3-S2A,B**). 24
205 hours post-injury, there is an increased number of PH3-positive cells in the right hemisphere,
206 where the penetrating STAB injury occurred (**Fig. 3-S2C**). Although not statistically significant,
207 we also detect a slight increase in cell proliferation in the left hemisphere at 24 hours post-PTBI
208 (**Fig. 3-S2C**). These data demonstrate that a penetrating STAB injury stimulates cell division in
209 young flies and that the mitotically active cells primarily are located near the area of injury. The
210 new cells could be created to replace the cells damaged by the injury and/or to remove debris
211 caused by the injury.

212 We next wanted to determine whether these newly created cells were maintained or
213 were eliminated soon after injury. Because PH3 only transiently labels dividing cells, we used a
214 different assay for these experiments. Specifically, we utilized EdU which is incorporated into
215 newly synthesized DNA of dividing cells and therefore permanently labels dividing cells and
216 their progeny. We used both pulse-chase and continuous feeding experimental designs and
217 assayed for EdU incorporation using fluorescent 'click chemistry' (Invitrogen®). The pulse-
218 chase experiments allowed us to ask whether the cells incorporating EdU were maintained. If a
219 cell died after synthesizing DNA, the incorporated EdU would either be undetectable or punctate
220 instead of being uniformly allocated within the nuclei following the chase. We saw similar
221 numbers of evenly labeled nuclei in both pulse-chase labeled brains and in brains from animals
222 continuously fed EdU (**Fig. 3-S3**). This indicates that the EdU labeled cells were indeed

223 proliferating and not dying. Consistent with the PH3 labeling at 24 hours, we observed more
224 EdU-positive cells 7- and 14-days post-injury in PTBI samples compared to uninjured controls
225 (p -value <0.0001 and p -value <0.0002) (**Fig. 3-2E-I**). This indicates that cell proliferation
226 continues between 7 and 14 days post-PTBI. Interestingly, at the later timepoints we saw a
227 statistically significant increase in the number of EdU-positive cells not only in the right
228 hemisphere (region near the injury), but also in the left hemisphere (**Fig. 3-S2D**). This suggests
229 that the penetrating STAB injury may induce both local and widespread proliferation. However,
230 we have not ruled out the possibility that some of the EdU+ cells distant to the injury migrated
231 from areas closer to the wound. Similar effects at a distance from an injury have been observed
232 in rodent models of TBI (Urrea et al., 2007, Ngwenya and Danzer, 2018).

233 Introducing even a mild injury may also induce cell death, and our histology experiments
234 suggested that PTBI did induce some cell death. To assay cell death, we therefore used
235 terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) (**Figs. 3-S4, 3-S5**).
236 TUNEL marks the terminal stages of death, both apoptotic and necrotic, where nuclear DNA
237 has been cleaved and degraded by DNases (Grasl-Kraupp et al., 1995). 0-6-hour old *OK107/yw*
238 adult males were injured and allowed to recover for 4 hours before dissecting the brains and
239 TUNEL staining. We classified the brains into groups based on the number of TUNEL-positive
240 cells/brain, with Group 1 having 0 to 9, Group 2 having 10-19, and Group 3 having 20-29
241 TUNEL-positive cells, respectively. The uninjured controls fell into Group 1, indicating that there
242 is not much cell death occurring in uninjured young adult brains (**Fig. 3-S4A-A',C**). Following
243 PTBI, the brains were classified as either Group 2 or Group 3 (**Fig. 3-S4B-B',C**). This indicates
244 that our standard injury method causes cell death in the brain within 4 hours-post-injury. Next,
245 we wanted to examine the temporal dynamics of the cell death observed after injury. When we
246 analyzed cell death later, at 24 hours post-injury, there were more TUNEL-positive cells in
247 injured brains compared to controls, with controls falling into Group 2 and injured brains falling
248 in the Group 3 category (**Fig 3-S5**). However, when we examined cell death at 10 days post-

249 injury, there was no significant difference in the numbers of TUNEL-positive cells, with both
250 control and injured brains falling into Group 3 (**Fig 3-S5**). These data indicate two things: 1)
251 PTBI-induced cell death is transient and peaks at earlier timepoints after injury; and 2) that cell
252 death in uninjured adult brains gradually increases following eclosion.

253
254 Proliferative ability is age-dependent

255 Because PTBI in older adult flies results in higher mortality than PTBI in young adult flies (**Fig.**
256 **3-1G**), we hypothesized that age might be an important factor in the proliferative ability post-
257 injury. To test this, we aged *OK107/yw* adult males out to 7, 14, and 28 days, then injured them
258 using our standard penetrating STAB injury, and assayed for cell proliferation 24 hours later. For
259 this experiment, we used anti-PH3 to identify cells that were actively dividing. *OK107/yw* adult
260 male brains injured at 7 days had significantly fewer PH3-positive cells 24 hours after injury
261 compared to flies that had been injured 0-6 hours post-eclosion (p -value<0.004) (**Fig. 3-2J**).
262 Although flies injured at 7 days had more PH3-positive cells after injury than the age-matched
263 uninjured controls, this difference was not statistically significant (**Fig. 3-2J**). However, flies
264 injured at either 14- or 28-days exhibited little cell proliferation post-injury, and this was not
265 significantly different compared to age-matched uninjured controls (**Fig. 3-2J**). Taken together,
266 these data support the idea that there is a specific window of time when adult *Drosophila* brains
267 possess the ability for cells to initiate division, but this timeframe is limited to very early
268 adulthood, from eclosion until ~7 days of age. At 7 days of age and further, we see a decrease
269 in the ability of adult *Drosophila*'s ability to respond to a central brain injury. Nonetheless,
270 uninjured flies at 24 hours and 7 days do not have significantly different baseline cell
271 proliferation, despite the fact that response to injury is significantly different (**Fig. 3-2J**). This
272 suggests that the presence of PH3-positive cells does not reflect the ability of the fly brain to
273 regenerate after injury. At 14 and 28 days of age, uninjured flies exhibit a noticeable decrease in
274 baseline cell division compared to 24 hour and 7-day uninjured controls, and injured flies no

275 longer mount a proliferative response to injury (**Fig. 3-2J**). Taken together, these data indicate
276 that the the climate/microenvironment of young brains may be more permissive of/conducive to
277 cell proliferation.

278 Importantly, while age is an important factor in proliferation rates post-PTBI, once
279 proliferation is initiated, it can persist for more than 7 days. This is supported by the EdU data
280 (**Fig. 3-2I**), in which there is a clear increase in the total number of EdU+ cells between 7 and 14
281 days. This raised the question of whether EdU+ cells observed at 14 days are continuing to
282 proliferate. The EdU labeling is cumulative, and we needed a real-time assay to address this
283 question. We therefore used anti-PH3 staining to label actively dividing cells. Newly eclosed
284 *OK107/yw* adult males were subjected to PTBI and aged to 24 hours, 7 days, or 14 days post-
285 injury prior to dissecting and fixing their brains and staining them with anti-PH3. At 7 days post-
286 injury, the numbers of PH3+ cells were similar to those at 24 hours post-injury (**Fig. 3-S6**).
287 However, when flies were allowed to recover for 14 days, we observed fewer PH3-positive cells
288 post-injury compared to injured flies aged to 24 hours and 7 days. Additionally, while the
289 number of PH3+ cells in 14-day post-PTBI brains appears to be slightly higher than controls,
290 this difference is not statistically significant (**Fig. 3-S6**). This is consistent with the EdU data in
291 **Fig. 3-2I**, and supports the idea that there is a specific window of time between eclosion and 7
292 days of age when adult fly brains have a relatively high capacity for (to initiate?) cell division.

293

294 Characterizing the cell populations of dividing cells post-injury

295 Next, we set out to determine the identities of the mitotically active cells. This was accomplished
296 by feeding *OK107/yw* adult males with EdU, then and assaying their brains for EdU, the glial
297 protein Reversed polarity (Repo) and the neuronal protein Embryonic lethal, abnormal vision
298 (Elav). At 7 days post-injury, we observed 4 classes of cells (**Fig. 3-3A-D**). Class I consists of
299 cells that were EdU+ but did not express Elav or Repo and therefore had an unknown identity
300 (arrowheads in **Fig. 3-3A-A**). Out of 172 EdU+ cells counted, approximately 44% were Class I

301 (Fig. 3-3E). Cells that were EdU+ and Repo+ were defined as Class II (arrowheads in Fig. 3-
302 3B-B’’). At 7 days post-injury, approximately 38% of EdU+ cells expressed a glial identity (Fig.
303 3-3E). Class III cells were EdU+ and expressed Elav (arrowheads in Fig. 3-3C-C’’).
304 Approximately 13% of cells that had divided by 7 days were EdU+ and Elav+ (Fig. 3-3E). These
305 data show that cells actively divide after PTBI, and that the dividing cells either are or become
306 glia and neurons. While most of the new cells that expressed mature cell type markers
307 generated after injury are glia, there were some new neurons. Interestingly, we also observed a
308 fourth class, cells that were EdU+, Repo+, and Elav+ (arrowheads in Fig. 3-3D-D’’).
309 Approximately 4% of EdU+ cells were classified as Class IV (Fig. 3-3E). These cells that
310 expressed both glial and neuronal identity were not detected in uninjured adult brains. At seven
311 days post-eclosion, approximately 46% of EdU+ cells in control brains are Class I and 53% of of
312 EdU+ cells are Class II (Fig. 3-3F). These numbers are similar to what has been seen in
313 previous studies of wildtype brains using a different nucleotide analog, bromodeoxyuridine
314 (BrDU) (von Trotha et al., 2009). In control brains, not only were there many fewer EdU+ cells,
315 we also did not observe either Class III or Class IV cells (Fig. 3-3F). This supports the idea that
316 injury stimulates neurogenesis.

317 Because age plays an important role in the proliferative ability of brain cells, we asked
318 whether the amount of time after injury impacts the types of cells observed. Using the same four
319 classes described in the 7-day post-PTBI experiment, we also looked at the cell types
320 generated at 24 hours and 14 days post-injury (Fig. 3-3E). The proliferation marker we used in
321 the 24-hour assay was anti-PH3 and the marker used in the 14-day assay was EdU
322 incorporation. At 24 hours post-injury, we observed the following distribution of classes: Class I,
323 56%; Class II, 38%; Class III, 0%; and Class IV, 5% (Fig. 3-3E). While there is a slightly lower
324 proportion of Class I (EdU+) cells at 7 days than at 24 hours, the proportions of Class II
325 (EdU+/Repo+) and IV (EdU+/Repo+/Elav+) cells are approximately the same. However, at 24
326 hours post-PTBI, there are no Class III (EdU+/Elav+) cells (Fig. 3-3F). Control uninjured flies at

327 24 hours had 39% Class I cells and 60% Class II cells (**Fig. 3-3F**). In comparing 7-day to 24-
328 hour control flies, there appears to be a slight trend of increasing Class I cell types and
329 decreasing Class II cells (**Fig. 3-3F**). Together these data demonstrate that new neurons are
330 created later than new glial cells and support the hypothesis that some of the Class I cells
331 (EdU+) are giving rise to Class III cells (EdU+/Elav+).

332 If we look later, 14 days post-PTBI, there is another striking change in cell type
333 distribution, with only 25% of cells falling into Class I, 45% in Class II, and 28% of cells in Class
334 III (**Fig. 3-3E**). Intriguingly, we did not observe any Class IV cell types 14 days post-PTBI. Class
335 IV (EdU+/Repo+/Elav+) cells express a hybrid glial and neuronal identity that is not observed in
336 control uninjured brains. We speculated that this could represent an abnormal transitional status
337 enroute to differentiation of new glia and neurons. By 14 days post-PTBI, these cells are no
338 longer detected, suggesting that they have committed to either neuronal or glial fate. Control
339 uninjured flies at 14 days had similar patterns as earlier timepoints, with approximately 66% of
340 cells in Class I, 33% of cells in Class II, and no cells in Classes III or IV (**Fig. 3-3F**).

341

342 Presence of neuroblast-like cells post-PTBI

343 To test whether the generation of new neurons post-PTBI follows a normal developmental
344 progression from NBs, we probed PTBI and control brains from *ase-Gal4, UAS-Stinger; UAS-*
345 *Gal80^{ts}* adult males with anti-Dpn and anti-PH3 antibodies to test whether NB genes were
346 upregulated post-injury. These flies were reared at 18°C until eclosion. At 18°C, the Gal80^{ts} is
347 functional and prevents expression of the fluorescent Stinger protein in *ase*-expressing cells. 0-
348 6 hours after eclosion, adult male flies were subjected to PTBI and placed at 30°C for 24 hours
349 prior to dissection and immunostaining. At 30°C, the Gal80^{ts} is not functional and fluorescent
350 Stinger protein is expressed in *ase*-expressing cells. 24 hours post-injury, we observe both
351 GFP+ cells and cells that are Dpn+/PH3+ in PTBI, but not control, brains (**Fig. 3-4A-D**). This
352 indicates that proliferating cells have key features of neuroblast identity, including *ase* and *dpn*

353 expression. Interestingly, we did not observe cells that were GFP+/Dpn+/PH3+. However, cells
354 that were GFP+ and cells that were Dpn+/PH3+ were clustered together and found adjacent to
355 each other, suggesting the two types of cells might share a lineage.

356 Upon cell division, Pros is asymmetrically segregated to the non-NB daughter cell where
357 it migrates from the cytoplasm into the nucleus. In the nucleus, Pros assumes the role of
358 transcription factor, which establishes the neural fate of the daughter cell. Indeed, using *ase-*
359 *Gal4*, *UAS-mCD8-GFP* flies, we observe large GFP+ cells near the wound at 24 hours post-
360 injury (arrowheads and asterisks in **Fig. 3-4E-F'**). A subset of the GFP+ cells also express
361 cytoplasmic Pros (asterisks in **Fig. 3-4F-F'**). These GFP+/cytoplasmic Pros+ cells are
362 consistent with the presence of NB-like cells post-PTBI. The GFP+/cytoplasmic Pros+ cells are
363 not seen in uninjured controls, which indicates that PTBI is somehow stimulating the generation
364 of these NB-like cell types.

365 To quantify levels of neural progenitor gene expression, we collected and injured
366 *OK107/yw* adult males at 0-6 hours post-eclosion. We then extracted total mRNA at 5 different
367 time points (4 hours, 24 hours, 3 days, 7 days, and 14 days) from post-PTBI and age-matched,
368 uninjured control heads. Relative transcript levels of neural progenitor genes were measured
369 using quantitative real-time PCR (qRT-PCR). At 4 and 24 hours of age, *ase* expression is
370 increased more than 2-fold in injured flies compared to controls (**Fig. 3-4G**). This difference
371 decreases by 3 days and is not significant at later timepoints (**Fig. 3-4G**). Although we observed
372 Dpn+ cells using immunohistochemistry near the area of injury at 24 hours post-PTBI, *dpn*
373 mRNA levels were not detectably increased at any timepoints (**Fig. 3-4H**). This could be
374 because we isolated RNA from whole heads and relatively few cells activate *dpn* following PTBI.

375 Consistent with the idea that NB-like cells are generated following PTBI, transcript levels
376 of *insc* were significantly increased at 4 hours, 24 hours, and 3 days, while *erm* transcript levels
377 also were increased at 4 hours (**Fig. 3-4I,J**). Together these data support the hypothesis that
378 there are NB-like precursor cells in the adult brain that are either generated following PTBI or

379 activated by PTBI. These cells are the potential precursor cells of adult-born neurons and glia
380 and represent a previously unknown cell type within the adult *Drosophila* brain.

381

382 Lineage-tracing the origins of newly created cells

383 As mentioned previously, in the uninjured wildtype adult central *Drosophila* brain, there is no
384 evidence of NBs. However, post-PTBI, we observe NB-like cells and increases in NB-related
385 transcript levels. The presence of these NB-like cells could be explained in two ways: 1)
386 differentiated cells such as glia or neurons are giving rise to new glia and neurons after injury,
387 via dedifferentiation to NB-like fates; or 2) a quiescent population of adult neural stem cells
388 exists and is activated by damage, similar to what is described in the optic lobe (Fernandez-
389 Hernandez et al., 2013). To test whether glial cells can give rise to neuronal stem cells, we
390 carried out lineage-tracing of glial-derived cells and asked whether they could become neurons.
391 To do this, we used *repo-Gal4* in conjunction with a *flipout-GFP* construct to permanently mark
392 glial lineages. F1 males that were $w[*]; repo-Gal4/P\{w[+mC]=Ubi-$
393 $p63E(FRT.STOP)Stinger\}15F2$ were injured at 0-6 hours post-eclosion and aged for 14 days
394 prior to dissection and immunostaining. We observed no GFP+ cells that were also Elav+. This
395 indicates that glia do not generate neurons through either trans-or dedifferentiation. We note
396 that ~50% of PH3+ and EdU+ cells post-PTBI are glia (**Fig. 3-3E**). Thus, although we found no
397 evidence that glia give rise to neurons, glia are nonetheless proliferating, especially very early
398 post-PTBI. To test whether neurons can give rise to glia, we used a similar lineage-labeling
399 technique with an *Nsyb-Gal4* driver in a system called Gal4 technique for real-time and clonal
400 expression (G-TRACE) (Evans et al., 2009). F1 males that were $w[*]; Nsyb-$
401 $Gal4/P\{w[+mC]=UAS-RedStinger\}6, P\{w[+mC]=UAS-FLP.Exel\}3, P\{w[+mC]=Ubi-$
402 $p63E(FRT.STOP)Stinger\}15F$ were injured at 0-6 hours post-eclosion and aged for 14 days
403 prior to dissection and immunostaining. We observed no GFP+/Repo+ cells. This indicates that
404 neurons do not give rise to new glia post-injury.

405 To address the second possibility, that new cells are created by a quiescent NB-like
406 population, we used a similar lineage labeling technique, this time in combination with a
407 neuroblast driver, *dpn-Gal4*. Although we did not detect GFP+/Dpn+ cells using Dpn antibody in
408 the *ase-Gal4* lineage experiments, we did observe GFP+ (i.e. *ase*-expressing) cells near Dpn+
409 cells, suggesting there might be a clonal relationship. Indeed, in Type II NB lineages during
410 normal development, *dpn* expression in NBs is followed by *ase* expression in the INPs to which
411 those NBs give rise. Also, *dpn* is expressed in both Type I and Type II NBs. We therefore
412 decided to use *dpn-Gal4* to search for quiescent NBs and other NB-like cells. To ensure that
413 neuroblast cells were not labeled during development, we added a temperature sensitive Gal80
414 and reared the crosses at 18°C. Under these conditions, the Gal80 prevents transcriptional
415 activation by Gal4, thus keeping the lineage tracing system off. F1 males that were *w[*]; dpn-*
416 *Gal4/P{w[+mC]=tubP-GAL80[ts]}20; P{w[+mC]=UAS-RedStinger}6, P{w[+mC]=UAS-*
417 *FLP.Exel}3, P{w[+mC]=Ubi-p63E(FRT.STOP)Stinger}15F* were collected or injured at 0-6 hours
418 post-eclosion and aged for 14 days at 30°C prior to dissection and immunostaining. At 30°C, the
419 temperature sensitive Gal80 protein is inactivated and Gal4 can activate transcription. Indeed,
420 14 days post-injury, we observed GFP+/Elav+ cells in injured brains (arrowheads in **Fig. 3-5A-**
421 **A''**), but not in uninjured age-matched controls. These results are consistent with the existence
422 of a quiescent stem cell-like population in the adult *Drosophila* brain that is activated by injury to
423 create new neurons and new glia. Several lines of evidence strongly support this view, including
424 the presence of Dpn+ and Ase+ cells near the area of injury 24 hours post-PTBI (**Figs. 3-4A-**
425 **D''** and **3-5A-A''**) and the elevated expression levels *ase*, *erm*, and *insc* post-PTBI. However,
426 because no Dpn+ cells are observed in our control uninjured central brains, these putative NB-
427 like cells differ from the NBs present during development because they lack detectable *dpn*
428 expression until stimulated by PTBI.
429
430

431 Structural and Functional Regeneration post-PTBI

432 To test the extent new neurons were generated following PTBI, we utilized a derivative of the
433 mosaic analysis with a repressible cell marker (MARCM) method (Lee and Luo, 1999). This
434 technique, the Perma-twin system, permanently labels dividing cells and their progeny with
435 either green fluorescent protein (GFP) or red fluorescent protein (RFP). We used adult F1 male
436 flies of the genotype: *w; FRT40A, UAS-CD8-GFP, UAS- CD2-Mir/ FRT40A, UAS-CD2-RFP,*
437 *UAS-GFP-Mir; act-Gal4 UAS-flp/tub-Gal80^{ts}* that had been reared at 17°C during development
438 to keep the system off. These flies were subjected to PTBI within 24 hours of eclosion, and
439 allowed to recover at 30°C for either 2, 7, or 14 days post-injury before dissecting and analyzing
440 their brains. As expected, based on our prior finding that PTBI stimulates cell proliferation, we
441 observed more clones in injured samples than controls at all timepoints (**Fig. 3-6A-E**). We also
442 found that injured flies had significantly more clones at later timepoints compared to earlier
443 ones, suggesting that regeneration and proliferation were progressive and did not only occur
444 immediately following the initial injury (**Fig. 3-6A-G, N**). Interestingly, we also observed large
445 clones at later timepoints that produced new MB neurons (**Fig. 3-6A-G, N**). These new neurons
446 project dendrites correctly to the MB calyx and axons that correctly project to the MB lobes. This
447 robust regeneration is fairly common in PTBI flies at 14 days post-injury and was observed in
448 approximately 50% of the assayed brains to varying degrees and with varying clone sizes (**Fig.**
449 **3N**). Other areas of the brain were also found to regenerate using the Perma-twin system.
450 These include the antennal lobes (AL), the ellipsoid body (EB), and the lateral horn (LH) (**Fig. 3-**
451 **6H-M**). We observed large clones in each of these regions approximately 26%, 26%, and 20%
452 of the time, respectively (**Fig. 3-6N**). These data suggest that there is structural repair of the
453 damaged MB region.

454 In order to assay for functional recovery post-PTBI, we assayed locomotor function. Due
455 to their relatively simple nervous system and compact behavioral repertoire, we can ask
456 whether *Drosophila* locomotor circuit function is damaged by PTBI and, if so, whether function is

457 restored at later timepoints. 2-day control uninjured flies displayed stereotypic locomotory
458 patterns throughout a 24-hour period (**Fig. 3-6O**). However, 2-day post-PTBI flies, exhibited
459 significantly different locomotor profiles (p -value <0.001) (**Fig. 3-6O**). Nonetheless, by 14 days,
460 by which time we observe increased cell proliferation and generation of new neurons,
461 PTBI flies display comparable locomotor profiles to age-matched controls (**Fig. 3-6P**). These
462 data indicate that PTBI significantly impacts motor function, and that this damage is largely
463 repaired by 14 days.

464

465 **Discussion**

466 Using a novel penetrating STAB injury, we have shown that the adult *Drosophila* central brain
467 has regenerative potential (**Fig. 3-7**). We demonstrate that PTBI stimulates cell proliferation
468 within 24 hours post-injury. However, the robust proliferative response is primarily in younger
469 flies and as flies age, this ability decreases. These data indicate that age plays an important role
470 in the adult *Drosophila*'s ability to survive a traumatic injury and warrants further investigation to
471 understand what cellular and molecular events transpire that allow for survival. While there is
472 some cell death post-PTBI, this is limited to early timepoints after injury; by 10 days post-PTBI,
473 the amount of cell death is not significantly different from non-injured age-matched controls. At
474 early timepoints post-PTBI, but not in control brains, we observe dividing neuroblast-like cells
475 that are Ase⁺ or Dpn⁺. Other neural progenitor genes such as *insc*, and *erm* exhibit elevated
476 transcript levels at early timepoints post-injury. By 7 days post-PTBI, new glia and new neurons
477 have been created. Using cell lineage-tracing techniques, we found that new neurons are
478 generated by cells that had once expressed *dpn*. These *dpn*-expressing cells were not found in
479 uninjured controls. Taken together, our data support the idea that there is a quiescent NB-like
480 population of cells that is activated upon injury, specifically in young adult flies. The newly
481 created cells are able to contribute to the overall regeneration of damaged brain tissue,
482 particularly near the mushroom body.

483 Although glia do not give rise to new neurons, and neurons do not give rise to new glia,
484 we have not ruled out the possibility that a process of dedifferentiation is occurring to give rise to
485 neural progenitors. For instance, it could be that mature cell types, such as neurons which are
486 not known to have proliferative abilities, are able to adopt more plastic fates upon injury and
487 create more neurons. Although we detect new neurons and new glia at 7 days post-injury, these
488 cell types are not generated in equal proportions. This trend continues even out to 14 days post-
489 PTBI. There appears to be an initial wave of gliogenesis, followed by a delayed wave of
490 neurogenesis. We originally thought that this was because glia were the precursors of both new
491 glia and new neurons, and that glial cells dedifferentiated to a more primitive state before giving
492 rise to new neurons. However, our lineage studies do not support this hypothesis. While glia
493 proliferate following PTBI, they do not give rise to new neurons.

494 Taking the 24 hour, 7 day, and 14 day timepoints together, we can identify noticeable
495 trends. At the earliest timepoint post-injury, we see the greatest proportion of Class I cells which
496 have incorporated EdU but are neither neurons nor glia. As time progresses post-PTBI, the
497 proportion of Class I cells decreases. This is consistent with progressive differentiation of the
498 newly born cells. Also consistent is that at earlier timepoints, there are no Class III cells
499 (EdU+/Elav+, i.e. new neurons). However, the further we assay after injury, the higher the
500 proportion of Class III cells. At 14 days post-injury the proportion of Class III cells is more than
501 double what was observed at 7 days (**Fig. 3-3E**).

502 Not only is cell proliferation quite low in control brains, but the identities of the new cells
503 exhibit different trends. Specifically, there are no Class III (proliferation+/Elav+) or IV
504 (proliferation+/Elav+/Repo+) cells at any of the three timepoints. There also is a slight increase
505 in the proportion of Class I (proliferation+/) cells over time in the controls, in direct proportion to
506 the decrease in the proportion of Class II (proliferation+/Repo+) cells (**Fig. 3-3F**). This suggests
507 that the cells that are dividing in control brains are slowly losing their glial identity over time. This
508 also corresponds with the decrease in the adult *Drosophila* brain cells ability to proliferate in

509 response to injury as they age and could be evidence of a progenitor cell type, specifically found
510 in younger adult flies, that is lost with age. Most important however, is that Class III
511 (proliferation+/Elav+) cells are not observed in controls at any timepoint, indicating that PTBI
512 brains make new neurons, but control brains do not (**Fig. 3-3F**).

513 Nonetheless, our studies do not rule out the possibility that glia undergo a process of
514 partial deregulation before making more glial cells. This is supported by the observation of Class
515 IV cells (proliferation+/Repo+/Elav+/) cells seen at 24 hours and 7 days post-PTBI.
516 Repo+/Elav+ cells are not found in uninjured control adult brains, but have been reported during
517 larval development (Berger et al., 2007) and in certain brain tumors (Beaucher et al., 2007).
518 Elav also is known to be transiently expressed in some NB-like cells during development
519 (Beaucher et al., 2007). Thus, the presence of dividing cells that are Repo+/Elav+ is consistent
520 with a less differentiated state. Further investigation is needed to conclusively identify the origins
521 of the glia created post-injury.

522 Previous studies have indicated a higher occurrence of certain brain tumors, such as
523 glioblastomas, in people that have previously experienced TBI (Tyagi et al., 2016). In our model
524 of PTBI, we do not find evidence of unregulated growth. Although there is an increase in new
525 cells over time post-injury as assayed EdU labeling, this is a cumulative view of all the cells that
526 proliferated and does not represent how many cells are still mitotically active. To measure the
527 number of cells that are actively proliferating at later timepoints, we injured flies at 0-6 hours
528 post-eclosion, allowed them to recover for 14 days and then assayed brains using anti-PH3,
529 which specifically marks cells in mitosis. At 14 days post-PTBI, although injured flies have
530 slightly more PH3+ cells than age-matched uninjured controls, the number is significantly lower
531 than at 7 days post-PTBI (**Fig. 3-S6**). Additionally, we have observed that the proportion of
532 Class I (prolif+ only) cells decreases over time, with 56% of cell in Class I at 24 hours, 44% of
533 cells in Class I at 7 days, and 26% of cells in Class I at 14 days post-PTBI. We also observed a
534 unique cell type, that was dividing and expressed Repo and Elav (Class IV cell). The timing of

535 these shifts in proportions suggest that a process of deregulation is occurring, and that a mature
536 cell type may be in the process of adopting a more plastic fate. While there are ~5% Class IV
537 cells at both 24 hours and 7 days post-PTBI, we did not observe any Class IV cells in 14-day
538 injured brains. Together, these data support the idea that PTBI does not stimulate uncontrolled
539 cell division. The mechanisms behind this need to be further analyzed to understand how
540 *Drosophila* are able to give a measured proliferative cell response to regenerate damaged
541 tissue.

542 In addition to proliferating glial cells, we also have identified a novel population of cells
543 that upregulate the expression of *dpn* post-PTBI and give rise to new neurons. Using Dpn and
544 PH3 antibodies, we found that uninjured adult central brains lack Dpn and have few mitotic
545 cells, while injured brains have Dpn+ cells that are mitotically active. Our lineage analysis is
546 consistent with this and indicates that *dpn*-expressing cells can generate neurons. We propose
547 that these cells represent a previously unknown population of NB-like cells that are quiescent in
548 uninjured brains. This is significant for several reasons. First, uninjured brains have cells that
549 cannot be identified with standard neural progenitor markers, but that nonetheless have
550 regenerative potential. If such cells exist in *Drosophila*, perhaps they also exist in humans.
551 Second, regeneration can be stimulated by mild injury. This provides us with an avenue for
552 identifying these cells for future study. It also provides us with a novel model system for
553 screening pharmacologic agents for those that activate the regenerative program. This could
554 lead to novel therapeutic approaches for both neurodegenerative diseases and brain injuries.

555 Our results in the central brain are somewhat reminiscent of optic lobe regeneration after
556 injury. However, while we have identified 2 distinct populations of new cells: glia and neurons,
557 the Fernandez-Hernandez *et al.* study identified one, neurons. Another intriguing difference is
558 that the proliferating optic lobe cells express the neural stem cell gene *deadpan (dpn)*, but not
559 the proneural gene *asense (ase)*. In the central brain after PTBI, there is *ase* upregulation as
560 assayed by both qRT-PCR and immunohistochemistry. Interestingly, we did not observe cells

561 expressing Ase that were Dpn+/PH3+. Instead, the *ase*→GFP cells were located adjacent to
562 Dpn+/PH3+ cells. This could indicate a lineage relationship between the Dpn+ and Ase+ cells.
563 For example, Dpn+/PH3+ cells could indicate a type II NB cell type while cells expressing Ase
564 might be progenitors, similar to INPs or GMCs. Two different populations of progenitor cells
565 could also explain the differences seen in the creation of glia versus neurons. Perhaps glia are
566 able to adopt more plastic like fates, akin to GMCs, and create more glia post-injury. It could
567 also be the case that the proliferating cells seen in uninjured control brains that express Repo
568 represent an unknown glial progenitor cell type that respond by creating more glia post-PTBI.
569 Further investigation is needed to understand how these cells contribute to proliferation post-
570 PTBI and what mechanisms allow this to occur. Together, these results indicate that
571 neurogenesis in adult *Drosophila* is different in the central brain than it is in the optic lobe and
572 that the neural progenitors in these brain regions have distinct developmental potentials.

573 Using our novel injury paradigm, we have been able to establish that young adult
574 *Drosophila* are capable of robust regeneration, with the creation of new neurons and glia and
575 functional recovery from locomotor defects by 14 days post-PTBI. We have demonstrated that
576 regeneration is accomplished via proliferation and by a neuroblast-like population that is
577 activated by injury. Further questions remain about the origin and properties of the NB-like cells.
578 Additionally, the molecular mechanisms that trigger regeneration and the activation of the
579 neuroblast-like population(s) have yet to be identified. In order to translate promise into
580 progress, we began at a basic level to elucidate the complex process of regeneration. By
581 utilizing *Drosophila melanogaster*, we can reap the benefits of a low-cost model and powerful
582 genetic and molecular tools. Once the myriad of components involved in neuroregeneration are
583 dissected and teased apart, we can work our way towards converting our knowledge of
584 *Drosophila* neurogenesis to humans.

585

586

587 **Materials & Methods**

588 Fly Stocks and Rearing

589 Unless otherwise specified, flies were reared at 25°C on a standard cornmeal-sugar medium.

590 The following stocks were obtained from the Bloomington Drosophila Stock Center (BDSC):

591 **#854** (*w*^{*}; *P*{*w*⁺*mW.hs*]=*GawB*}*OK107 ey*[*OK107*]/*ln*(4)*ci*[*D*], *ci*[*D*] *pan*[*ciD*] *sv*[*spa-pol*]; **#1495**

592 (*y*[1] *w*[1]); **#4539** *y*[1] *w*^{*}; *P*{*w*⁺*mC*]=*UAS-FLP.D*}*JD1*; **#5130** (*y*[1] *w*[1]; *Pin*[*Yt*]/*CyO*;

593 *P*{*w*⁺*mC*]=*UAS-mCD8::GFP.L*}*LL6*); **#7018** (*w*^{*}; *sna*[*Sco*]/*CyO*;*P*{*w*⁺*mC*]=*tubP-*

594 *GAL80*[*ts*] *ncd*[*GAL80ts-7*]); **#7019** (*w*^{*}; *P*{*w*⁺*mC*]=*tubP-GAL80*[*ts*]20; *TM2/TM6B*, *Tb*[1];

595 **#7415** (*w*¹¹¹⁸; *P*{*w*⁺*m*^{*}]=*GAL4*}*repo/TM3*, *Sb*¹); **#28281** (*w*^{*}; *P*{*w*⁺*mC*]=*UAS-*

596 *RedStinger*}6, *P*{*w*⁺*mC*]=*UAS-FLP.Exel*}3, *P*{*w*⁺*mC*]=*Ubi-p63E(FRT.STOP)Stinger*}15F);

597 **#32251** (*w*^{*}; *P*{*w*⁺*mC*]=*Ubi-p63E(FRT.STOP)Stinger*}15F2); **#47859** *w*[1118]; *P*{*y*⁺*t7.7*}

598 *w*⁺*mC*]=*GMR13CO2-GAL4*}*attP2*; **#51635** (*y*¹ *w*^{*}; *P*{*w*⁺*m*^{*}]=*nSyb-GAL4S*}3); **#65408**

599 (*P*{*w*⁺*mC*]=*UAS-Stinger*}2, *P*{*w*⁺*mC*]=*UAS-hid.Z*}2/*CyO*). Other lines used were *ase-*

600 *Gal4/CyO*; *Dr/TM6B* (a gift of Dr. Cheng-Yu Lee); *w*; *FRT40A*, *UAS-CD8-GFP*, *UAS- CD2-Mir*;

601 *act-Gal4 UAS-flp/TM6B*; and *w*; *FRT40A*, *UAS-CD2-RFP*, *UAS-GFP-Mir*; *tub-Gal80ts/TM6B*

602 (both gifts of Dr. Eduardo Moreno).

603

604 Standard cross

605 To minimize variations due to genetic background and sex, we routinely analyzed F1 males

606 from the following cross: *w*^{*}; *UAS-mCD8-GFP*; *OK107-Gal4* X *y*[1] *w*[1]. This cross was

607 maintained at 25°C. PTBI flies were kept at 25°C until their brains were dissected and fixed for

608 analysis. For simplicity, this genotype is referred to as “*OK107/yw*” hereafter.

609

610 Perma-twin flies

611 Perma-twin flies were generated by crossing *w*; *FRT40A*, *UAS-CD2-RFP*, *UAS-GFP-Mir*; *tub-*

612 *Gal80^{ts}/TM6B* virgin females to *w*; *FRT40A*, *UAS-CD8-GFP*, *UAS- CD2-Mir*; *act-Gal4 UAS-*

613 *flp/TM6B* males (Fernandez-Hernandez et al., 2013). These crosses were maintained at 17°C.
614 F1 progeny of the genotype: *w; FRT40A, UAS-CD8-GFP, UAS- CD2-Mir/ FRT40A, UAS-CD2-*
615 *RFP, UAS-GFP-Mir; act-Gal4 UAS-flp/tub-Gal80^{ts}* were collected at eclosion, subjected to PTBI
616 or kept as uninjured controls and maintained at 30°C until their brains were dissected and fixed
617 for analysis.

618

619 G-TRACE crosses

620 Lineage-labeling was accomplished using a G-TRACE line (**#28281** (*w[*];P{w[+mC]=UAS-*
621 *RedStinger}6,P{w[+mC]=UAS-FLP.Exel}3,P{w[+mC]=Ubi-p63E(FRT.STOP)Stinger}15F*)
622 crossed to various Gal4 driver strains listed above. These crosses were maintained at 17°C. F1
623 progeny of the desired genotyped were selected at eclosion, subjected to PTBI or kept as
624 uninjured controls and maintained at 30°C for 14 days when their brains were dissected and
625 fixed for analysis.

626

627 Penetrating Traumatic Brain Injury

628 To induce PTBI, we used thin metal needles (~12.5 µm diameter tip, 100 µm diameter rod; Fine
629 Science Tools) sterilized in 70% ethanol to penetrate the head capsule of CO₂-anesthetized
630 adult flies. Injured flies were transferred back to our standard sugar food for recovery and aging.

631

632 Immunohistochemistry

633 Brains were dissected in PBS (phosphate-buffered saline; 100 mM K₂HPO₄, 140 mM NaCl pH
634 7.0) and fixed in a 3.7% formaldehyde in a PEM (100 mM PIPES, 2 mM EGTA, 1 mM MgSO₄)
635 solution for 20 minutes at 25°C. Fixed brain samples were washed in PT (PBS and 0.1% Triton
636 X-100), blocked with 2% BSA in PT solution (PBT), and then incubated with primary antibodies
637 overnight at 4°C in PBT. Following primary antibody incubation, the samples were washed with
638 PT (5 times over the course of an hour) and incubated overnight in secondary antibody at 4°C.

639 The next day, samples were washed in PT, stained with DAPI (1:10,000, ThermoFisher) for 8
640 minutes, and mounted in Vectashield anti-fade mountant (Vector Labs) and imaged using a
641 Nikon A1RS system and analyzed using the Nikon NIS Elements software. Cell counting was
642 done both manually and using the Nikon NIS-Elements software to analyze regions of interest
643 (ROIs) with a threshold of over 1000 and an area of at least 10 μ m.

644 The primary antibodies used in this study were: rabbit anti-PH3 (1:500, Santa Cruz
645 Biotechnology, Inc); mouse anti-FasII (1:20, Developmental Studies Hybridoma Bank; DSHB);
646 mouse anti-Repo (1:20, DSHB); rat anti-Elav (1:20, DSHB); mouse anti-Pros (1:20, DSHB); and
647 rat anti-Dpn (1:50, AbCam). Secondary antibodies used were: anti-rabbit Alexa Fluor 568
648 (1:200, ThermoFisher); anti-rabbit Cy5 (1:400, Jackson ImmunoResearch, Inc.); anti-mouse
649 Cy5 (1:100, Jackson ImmunoResearch, Inc.); anti-rat Alexa Fluor 488 (1:400, ThermoFisher);
650 anti-rat Alexa Fluor 568 (1:400, ThermoFisher); and anti-rat Cy5 (1:200, Jackson
651 ImmunoResearch, Inc.).

652

653 EdU labeling

654 The standard injury method was used on flies for 5-ethynyl-2'-deoxyuridine (EdU) labeling,
655 except flies were fed 50 mM EdU in 10% sucrose solution on a size 3 Whatman filter for six
656 hours prior to PTBI and allowed to recover on the same solution for the desired amount of time.
657 The EdU solution was replaced every 24 hours. Brains were dissected, processed, and antibody
658 stained as described above with the exception of using buffers without azide. To detect EdU
659 incorporation, Click-IT[®] reagents from InVitrogen were used according to the manufacturer's
660 instructions. The brains then were antibody stained mounted and imaged as described above.

661

662 Quantitative Real-Time PCR

663 Transcript levels of target genes were measured by quantitative real-time PCR (qRT-PCR)
664 using methods described in (Ihry et al., 2012). RNA was isolated from appropriately staged

665 animals using TRIzol Reagent used according to the manufacturer's instructions (Thermo Fisher
666 Scientific). cDNA was synthesized from 40 to 400 ng of total RNA using the SuperScript III First-
667 Strand Synthesis System (Invitrogen). qPCR was performed on a Roche 480 LightCycler using
668 the LightCycler 480 DNA SYBR Green I Master kit (Roche). In all cases, samples were run
669 simultaneously with three independent biological replicates for each target gene, and *rp49* was
670 used as the reference gene. To calculate changes in relative expression, the Relative
671 Expression Software Tool was used (Pfaffl et al., 2002). We used the following primers to detect
672 transcript levels: *ase* Forward: 5'-CAGTGATCTCCTGCCTAGTTTG-3' & Reverse: 5'-
673 GTGTTGGTTCCTGGTATTCTGATG-3' (gift from Stanislava Chtarbanova); *dpn* Forward: 5'-
674 CGCTATGTAAGCCAAATGGATGG-3' & Reverse: 5'-CTATTGGCACACTGGTTAAGATGG-3'
675 (gift from Stanislava Chtarbanova); *elav* Forward: 5'-CGCAGCCCAATACGAATGG-3' &
676 Reverse: 5'-CATTGTTTGC GGCAA GTAGTTG-3' (Fly Primer Bank); *erm* Forward: 5'-
677 GTCCCCTAAAGTTTTTCGATAGCC-3' & Reverse: 5'- GAGTCATAGTTGACAGTGGATGG-3'
678 (Fly Primer Bank); *insc* Forward: 5'- CCCTGGGCAATCTGTCCTG-3' & Reverse: 5'-
679 GAGAAGCCCGAATCCTGACT-3' (Fly Primer Bank); *myc* Forward: 5'-
680 AGCCAGAGATCCGCAACATC-3' & Reverse: 5'- CGCGCTGTAGAGATTCGTAGAG-3' (Fly
681 Primer Bank); *repo* Forward: 5'-TCGCCCAACTATGTGACCAAG-3' & Reverse: 5'-
682 CGGCGCACTAATGTA TCG-3' (Fly Primer Bank); *Rp49* Forward: 5'-
683 CCAGTCGGATCGATATGCTAA-3' & Reverse: 5'-ACGTTGTGCACCAGGAACTT-3' (Denton et
684 al., 2009).

685

686 Locomotor assays

687 0-6-hour post-eclosion *OK107/yw* males were collected, subjected to PTBI, and aged to 2 days
688 and 14 days, respectively. The 2 and 14 day injured and age-matched uninjured controls were
689 placed in the *Drosophila* Activity Monitor (DAM) system (TriKinetics, Waltham, MA) to record

690 locomotory behavior. The circadian locomotor activity of flies was assayed and analyzed as
691 previously described (Hamblen et al. 1986; Sehgal et al, 1992).

692

693 Statistical analysis

694 For all cell/clone counting and locomotor assays, counts were expressed as means \pm standard
695 deviations. Two-tailed t-tests were performed using Graphpad Prism Version 8.3.0 for Mac
696 (GraphPad Software, La Jolla California USA, www.graphpad.com). An alpha value of 0.05 was
697 considered significant. The following symbols represent significance; * significant at $p \leq 0.05$; **
698 significant at $p \leq 0.01$; *** significant at $p \leq 0.001$; **** significant at $p \leq 0.0001$.

699

700 **Acknowledgements**

701 We are grateful to: Becky Katzenberger and Sarah Neuman for technical assistance; Eduardo
702 Moreno for sharing the Perma-twin stocks; and Stanislava Chtarbanova for the *ase* and *dpr*
703 primers. We also would like to thank Barry Ganetzky and David Wassarman for lively
704 discussions that undoubtedly improved the science and Kent Mok, Cayla Guerra, and Bailey
705 Spiegelberg for their contributions to the lab. The FasII, Elav and Repo monoclonal antibodies
706 developed by Corey Goodman (FasII and Repo) and Gerry Rubin (Elav) were obtained from the
707 Developmental Studies Hybridoma Bank, created by the NICHD of the NIH and maintained at
708 The University of Iowa, Department of Biology, Iowa City, IA 52242. Most of the *Drosophila*
709 strains used in this study were obtained from the Bloomington *Drosophila* Stock Center (BDSC;
710 NIH P40OD018537). This work was supported by NIH T32 GM007133 (KLC and KM); NIH
711 NS090190 (GBF); NIH NS102698 (GBF; the University of Wisconsin Graduate School (GBF);
712 and the Women in Science and Engineering Leadership Institute (WISELI) (GBF).

713 **References cited**

- 714 ALTMAN, J. 1969. Autoradiographic and histological studies of postnatal neurogenesis.
715 IV. Cell proliferation and migration in the anterior forebrain, with special reference
716 to persisting neurogenesis in the olfactory bulb. *J Comp Neurol*, 137, 433-57.
- 717 ALTMAN, J. & DAS, G. D. 1965. Autoradiographic and histological evidence of
718 postnatal hippocampal neurogenesis in rats. *J Comp Neurol*, 124, 319-35.
- 719 AMARIGLIO, N., HIRSHBERG, A., SCHEITHAUER, B. W., COHEN, Y.,
720 LOEWENTHAL, R., TRAKHTENBROT, L., PAZ, N., KOREN-MICHOWITZ, M.,
721 WALDMAN, D., LEIDER-TREJO, L., TOREN, A., CONSTANTINI, S. &
722 RECHAVI, G. 2009. Donor-derived brain tumor following neural stem cell
723 transplantation in an ataxia telangiectasia patient. *PLoS Med*, 6, e1000029.
- 724 ASO, Y., HATTORI, D., YU, Y., JOHNSTON, R. M., IYER, N. A., NGO, T. T., DIONNE,
725 H., ABBOTT, L. F., AXEL, R., TANIMOTO, H. & RUBIN, G. M. 2014. The
726 neuronal architecture of the mushroom body provides a logic for associative
727 learning. *Elife*, 3, e04577.
- 728 BEAUCHER, M., GOODLIFFE, J., HERSPERGER, E., TRUNOVA, S., FRYDMAN, H. &
729 SHEARN, A. 2007. Drosophila brain tumor metastases express both neuronal
730 and glial cell type markers. *Dev Biol*, 301, 287-97.
- 731 BELLEN, H. J., TONG, C. & TSUDA, H. 2010. 100 years of Drosophila research and its
732 impact on vertebrate neuroscience: a history lesson for the future. *Nat Rev*
733 *Neurosci*, 11, 514-22.
- 734 BELLO, B. C., IZERGINA, N., CAUSSINUS, E. & REICHERT, H. 2008. Amplification of
735 neural stem cell proliferation by intermediate progenitor cells in Drosophila brain
736 development. *Neural Dev*, 3, 5.
- 737 BERGER, C., RENNER, S., LUER, K. & TECHNAU, G. M. 2007. The commonly used
738 marker ELAV is transiently expressed in neuroblasts and glial cells in the
739 Drosophila embryonic CNS. *Dev Dyn*, 236, 3562-8.
- 740 BIRNBAUM, K. D. & SANCHEZ ALVARADO, A. 2008. Slicing across kingdoms:
741 regeneration in plants and animals. *Cell*, 132, 697-710.
- 742 BOLLINI, S., SMITS, A. M., BALBI, C., LAZZARINI, E. & AMERI, P. 2018. Triggering
743 Endogenous Cardiac Repair and Regeneration via Extracellular Vesicle-
744 Mediated Communication. *Front Physiol*, 9, 1497.
- 745 BOONE, J. Q. & DOE, C. Q. 2008. Identification of Drosophila type II neuroblast
746 lineages containing transit amplifying ganglion mother cells. *Dev Neurobiol*, 68,
747 1185-95.
- 748 BOWMAN, S. K., ROLLAND, V., BETSCHINGER, J., KINSEY, K. A., EMERY, G. &
749 KNOBLICH, J. A. 2008. The tumor suppressors Brat and Numb regulate transit-
750 amplifying neuroblast lineages in Drosophila. *Dev Cell*, 14, 535-46.
- 751 CAO, Y., CHTARBANOVA, S., PETERSEN, A. J. & GANETZKY, B. 2013. Dnr1
752 mutations cause neurodegeneration in Drosophila by activating the innate
753 immune response in the brain. *Proc Natl Acad Sci U S A*, 110, E1752-60.
- 754 CHIA, W., SOMERS, W. G. & WANG, H. 2008. Drosophila neuroblast asymmetric
755 divisions: cell cycle regulators, asymmetric protein localization, and
756 tumorigenesis. *J Cell Biol*, 180, 267-72.
- 757 CHIANG, A. S., LIN, C. Y., CHUANG, C. C., CHANG, H. M., HSIEH, C. H., YEH, C. W.,
758 SHIH, C. T., WU, J. J., WANG, G. T., CHEN, Y. C., WU, C. C., CHEN, G. Y.,

- 759 CHING, Y. T., LEE, P. C., LIN, C. Y., LIN, H. H., WU, C. C., HSU, H. W.,
760 HUANG, Y. A., CHEN, J. Y., CHIANG, H. J., LU, C. F., NI, R. F., YEH, C. Y. &
761 HWANG, J. K. 2011. Three-dimensional reconstruction of brain-wide wiring
762 networks in *Drosophila* at single-cell resolution. *Curr Biol*, 21, 1-11.
- 763 DENNIS, C. V., SUH, L. S., RODRIGUEZ, M. L., KRIL, J. J. & SUTHERLAND, G. T.
764 2016. Human adult neurogenesis across the ages: An immunohistochemical
765 study. *Neuropathol Appl Neurobiol*, 42, 621-638.
- 766 DENTON, D., SHRAVAGE, B., SIMIN, R., MILLS, K., BERRY, D. L., BAEHRECKE, E.
767 H. & KUMAR, S. 2009. Autophagy, not apoptosis, is essential for midgut cell
768 death in *Drosophila*. *Curr Biol*, 19, 1741-6.
- 769 EGGER, B., CHELL, J. M. & BRAND, A. H. 2008. Insights into neural stem cell biology
770 from flies. *Philos Trans R Soc Lond B Biol Sci*, 363, 39-56.
- 771 ERIKSSON, P. S., PERFILIEVA, E., BJORK-ERIKSSON, T., ALBORN, A. M.,
772 NORDBORG, C., PETERSON, D. A. & GAGE, F. H. 1998. Neurogenesis in the
773 adult human hippocampus. *Nat Med*, 4, 1313-7.
- 774 EVANS, C. J., OLSON, J. M., NGO, K. T., KIM, E., LEE, N. E., KUOY, E.,
775 PATANANAN, A. N., SITZ, D., TRAN, P., DO, M. T., YACKLE, K., CESPEDES,
776 A., HARTENSTEIN, V., CALL, G. B. & BANERJEE, U. 2009. G-TRACE: rapid
777 Gal4-based cell lineage analysis in *Drosophila*. *Nat Methods*, 6, 603-5.
- 778 FERNANDEZ-HERNANDEZ, I., RHINER, C. & MORENO, E. 2013. Adult neurogenesis
779 in *Drosophila*. *Cell Rep*, 3, 1857-65.
- 780 GAO, X., WANG, X., XIONG, W. & CHEN, J. 2016. In vivo reprogramming reactive glia
781 into iPSCs to produce new neurons in the cortex following traumatic brain injury.
782 *Sci Rep*, 6, 22490.
- 783 GRASL-KRAUPP, B., RUTTKAY-NEDECKY, B., KOUDELKA, H., BUKOWSKA, K.,
784 BURSCH, W. & SCHULTE-HERMANN, R. 1995. In situ detection of fragmented
785 DNA (TUNEL assay) fails to discriminate among apoptosis, necrosis, and
786 autolytic cell death: a cautionary note. *Hepatology*, 21, 1465-8.
- 787 HANS, F. & DIMITROV, S. 2001. Histone H3 phosphorylation and cell division.
788 *Oncogene*, 20, 3021-7.
- 789 HOMEM, C. C. & KNOBLICH, J. A. 2012. *Drosophila* neuroblasts: a model for stem cell
790 biology. *Development*, 139, 4297-310.
- 791 HOMEM, C. C., REPIC, M. & KNOBLICH, J. A. 2015. Proliferation control in neural
792 stem and progenitor cells. *Nat Rev Neurosci*, 16, 647-59.
- 793 IHRY, R. J., SAPIRO, A. L. & BASHIRULLAH, A. 2012. Translational control by the
794 DEAD Box RNA helicase belle regulates ecdysone-triggered transcriptional
795 cascades. *PLoS Genet*, 8, e1003085.
- 796 ITO, K., AWANO, W., SUZUKI, K., HIROMI, Y. & YAMAMOTO, D. 1997. The
797 *Drosophila* mushroom body is a quadruple structure of clonal units each of which
798 contains a virtually identical set of neurones and glial cells. *Development*, 124,
799 761-71.
- 800 ITO, K. & HOTTA, Y. 1992. Proliferation pattern of postembryonic neuroblasts in the
801 brain of *Drosophila melanogaster*. *Dev Biol*, 149, 134-48.
- 802 KAPLAN, M. S. & HINDS, J. W. 1977. Neurogenesis in the adult rat: electron
803 microscopic analysis of light radioautographs. *Science*, 197, 1092-4.

- 804 KNOBLICH, J. A. 2010. Asymmetric cell division: recent developments and their
805 implications for tumour biology. *Nat Rev Mol Cell Biol*, 11, 849-60.
- 806 KNOTH, R., SINGEC, I., DITTER, M., PANTAZIS, G., CAPETIAN, P., MEYER, R. P.,
807 HORVAT, V., VOLK, B. & KEMPERMANN, G. 2010. Murine features of
808 neurogenesis in the human hippocampus across the lifespan from 0 to 100
809 years. *PLoS One*, 5, e8809.
- 810 KUHN, H. G., DICKINSON-ANSON, H. & GAGE, F. H. 1996. Neurogenesis in the
811 dentate gyrus of the adult rat: age-related decrease of neuronal progenitor
812 proliferation. *J Neurosci*, 16, 2027-33.
- 813 LEE, T. & LUO, L. 1999. Mosaic analysis with a repressible cell marker for studies of
814 gene function in neuronal morphogenesis. *Neuron*, 22, 451-61.
- 815 LESSING, D. & BONINI, N. M. 2009. Maintaining the brain: insight into human
816 neurodegeneration from *Drosophila melanogaster* mutants. *Nat Rev Genet*, 10,
817 359-70.
- 818 MANGANAS, L. N., ZHANG, X., LI, Y., HAZEL, R. D., SMITH, S. D., WAGSHUL, M. E.,
819 HENN, F., BENVENISTE, H., DJURIC, P. M., ENIKOLOPOV, G. & MALETIC-
820 SAVATIC, M. 2007. Magnetic resonance spectroscopy identifies neural
821 progenitor cells in the live human brain. *Science*, 318, 980-5.
- 822 MEINERTZHAGEN, I. A. 2010. The organization of the invertebrate brains: cells,
823 synapses and circuits. *Acta Zoologica*, 91, 64-71.
- 824 MICHALOPOULOS, G. K. 2007. Liver regeneration. *J Cell Physiol*, 213, 286-300.
- 825 MING, G. L. & SONG, H. 2011. Adult neurogenesis in the mammalian brain: significant
826 answers and significant questions. *Neuron*, 70, 687-702.
- 827 MORENO, E., FERNANDEZ-MARRERO, Y., MEYER, P. & RHINER, C. 2015. Brain
828 regeneration in *Drosophila* involves comparison of neuronal fitness. *Curr Biol*, 25,
829 955-63.
- 830 NGWENYA, L. B. & DANZER, S. C. 2018. Impact of Traumatic Brain Injury on
831 Neurogenesis. *Front Neurosci*, 12, 1014.
- 832 PFAFFL, M. W., HORGAN, G. W. & DEMPFFLE, L. 2002. Relative expression software
833 tool (REST) for group-wise comparison and statistical analysis of relative
834 expression results in real-time PCR. *Nucleic Acids Res*, 30, e36.
- 835 RAMON Y CAJAL, S. 1913. *Estudios sobre la degeneración y regeneración*
836 *del sistema nervioso. Tomo I, Degeneración y regeneración de los*
837 *nervios.*, Madrid, Imprenta de Hijos de Nicola`s Moya.
- 838 RAMON Y CAJAL, S. 1914. *Estudios sobre la degeneración y regeneración*
839 *del sistema nervioso Tomo II, Degeneración y regeneración de los*
840 *centros nerviosos.*, Madrid, Imprenta de Hijos de Nicola`s Moya.
- 841 REIF, A., FRITZEN, S., FINGER, M., STROBEL, A., LAUER, M., SCHMITT, A. &
842 LESCH, K. P. 2006. Neural stem cell proliferation is decreased in schizophrenia,
843 but not in depression. *Mol Psychiatry*, 11, 514-22.
- 844 SASSE, S., NEUERT, H. & KLAMBT, C. 2015. Differentiation of *Drosophila* glial cells.
845 *Wiley Interdiscip Rev Dev Biol*, 4, 623-36.
- 846 SIEGRIST, S. E., HAQUE, N. S., CHEN, C. H., HAY, B. A. & HARIHARAN, I. K. 2010.
847 Inactivation of both Foxo and reaper promotes long-term adult neurogenesis in
848 *Drosophila*. *Curr Biol*, 20, 643-8.

- 849 SPALDING, K. L., BERGMANN, O., ALKASS, K., BERNARD, S., SALEHPOUR, M.,
850 HUTTNER, H. B., BOSTROM, E., WESTERLUND, I., VIAL, C., BUCHHOLZ, B.
851 A., POSSNERT, G., MASH, D. C., DRUID, H. & FRISEN, J. 2013. Dynamics of
852 hippocampal neurogenesis in adult humans. *Cell*, 153, 1219-1227.
- 853 SPINDLER, S. R. & HARTENSTEIN, V. 2010. The *Drosophila* neural lineages: a model
854 system to study brain development and circuitry. *Dev Genes Evol*, 220, 1-10.
- 855 STANGER, B. Z. 2015. Cellular homeostasis and repair in the mammalian liver. *Annu*
856 *Rev Physiol*, 77, 179-200.
- 857 STILES, J. & JERNIGAN, T. L. 2010. The basics of brain development. *Neuropsychol*
858 *Rev*, 20, 327-48.
- 859 TYAGI, V., THEOBALD, J., BARGER, J., BUSTOROS, M., BAYIN, N. S., MODREK, A.
860 S., KADER, M., ANDERER, E. G., DONAHUE, B., FATTERPEKAR, G. &
861 PLACANTONAKIS, D. G. 2016. Traumatic brain injury and subsequent
862 glioblastoma development: Review of the literature and case reports. *Surg*
863 *Neurol Int*, 7, 78.
- 864 URREA, C., CASTELLANOS, D. A., SAGEN, J., TSOULFAS, P., BRAMLETT, H. M. &
865 DIETRICH, W. D. 2007. Widespread cellular proliferation and focal neurogenesis
866 after traumatic brain injury in the rat. *Restor Neurol Neurosci*, 25, 65-76.
- 867 VISHWAKARMA, S. K., BARDIA, A., TIWARI, S. K., PASPALA, S. A. & KHAN, A. A.
868 2014. Current concept in neural regeneration research: NSCs isolation,
869 characterization and transplantation in various neurodegenerative diseases and
870 stroke: A review. *J Adv Res*, 5, 277-94.
- 871 VON TROTHA, J. W., EGGER, B. & BRAND, A. H. 2009. Cell proliferation in the
872 *Drosophila* adult brain revealed by clonal analysis and bromodeoxyuridine
873 labelling. *Neural Dev*, 4, 9.
- 874 WENG, M. & LEE, C. Y. 2011. Keeping neural progenitor cells on a short leash during
875 *Drosophila* neurogenesis. *Curr Opin Neurobiol*, 21, 36-42.
- 876




Figure 3-1A,B. Penetrating Traumatic Brain Injury (PTBI) as a new model to study neuroregeneration. In our standard injury protocol, we anesthetize young flies within 0-6 hours of eclosion and unilaterally injure the right central brain. Specifically, we target the right mushroom body (MB) region using a sterilized steel minuten pin (~12.5 μ m diameter tip, 100 μ m rod diameter; Fine Science Tools) to penetrate the head cuticle and the brain. **A.** This schematic shows an external frontal view of the *Drosophila* head and the location of injury with the needle trajectory indicated in blue). In this diagram, the right side of the brain is on the left because it is a frontal view. **B.** As described above, our penetrating STAB injury (trajectory indicated in grey) damages multiple cell types, including the mushroom body (MB, green), parts of the brain (white), such as glia and neurons, and cells outside the brain, such as the fat body (blue) and hemocytes (red). CB = central brain region. OL= optic lobe region. In this diagram, the right side of the brain is to the right.




Figure 3-1C-E. PTBI increases neurodegeneration. To understand the long-term consequences of PTBI, we evaluated control and injured brains by histology at 25 days post-eclosion. Controls exhibited little neurodegeneration at 25 days (C). In PTBI flies, we observed an increase in the number of lesions (D). We used the neurodegeneration index described in Cao *et al.*, 2013 to quantify this neurodegeneration. Controls had an average neurodegeneration index score of 1.7 ± 0.2 , while PTBI flies had an average neurodegeneration index score of 3.0 ± 0.4 (E). This represented a statistically significant difference (p -value=0.0089). Error bars reflect the standard deviation (SD).




Figure 3-1F,G. PTBI decreases lifespan. **F.** To assess the impact of PTBI on viability, we performed a lifespan assay with control and PTBI flies. Within the first 12 days, there was no significant difference in survival between control and injured flies. However, beyond 12 days, we saw a significant drop in survival in injured flies. Control flies reached 50% survival at 70 days of age, while injured flies reached 50% survival at 48 days. The maximum lifespan was 84 days for uninjured flies and 74 days for injured flies. This indicates that PTBI does have long-term consequences, similar to what is observed with other forms of TBI. Yet, in the window of 0-12 days, there is no significant difference in survival, suggesting that something is occurring to combat or delay the negative effects of injury. **G.** Because we observed little death within the first 12 days post-injury, we suspected that age might play an important role in this outcome. To test this, we collected, aged, and injured flies that were 0-6 hours post-eclosion, 14 days old, and 28 days old. Survival was then assayed 24 hours post-PTBI. Young flies (0-6 hours post-eclosion), exhibited no significant difference in survival compared to control flies (p -value=0.69), while older flies at 14 days (p -value<0.0001) and 28 days (p <0.0001) of age exhibited decreased survival compared to age-matched uninjured controls. Error bars reflect the standard deviation (SD).




Figure 3-2A-H. PTBI stimulates cell proliferation. **A**, uninjured, control and **B**, **PTBI** brain schematics. The blue boxes in the upper right corner of these brains indicate the area near the MB imaged in **C-H**. **C** and **D**. To assay for cell proliferation, we used an anti-PH3 antibody (red) and stained brains 24 hours after injury. In control brains (**C**) there are few PH3+ cells, and none near the MB. However, in PTBI brains (**D**), there is an increase in the number of PH3+ cells specifically near the MB. **E** and **F**. To test whether newly created cells are surviving or being eliminated, we conducted a pulse-chase EdU experiment. We fed flies EdU (red) for 4 days post-injury (a pulse) and followed this with a 3-day period without EdU (the chase). In the 7-day control brain (**E**), there is little EdU incorporation. In 7-day PTBI brains (**F**), there are more EdU+ cells and many of these cells are near the MB. **G**. In 14-day control brains, there are few EdU+ cells. **H**. However, in 14-day injured brains, there is an increase in EdU+ cells near the MB. All brains are from males of our standard genotype (*OK107/yw*).




Figure 3-2I. The number of proliferating cells increases over time post-PTBI. We quantified proliferating cells using anti-PH3 at 24 hours and EdU at 7 and 14 days. EdU permanently marks dividing cells and would therefore allow us to measure the cumulative number of cells that had divided since the PTBI. At 24 hours, uninjured control brains had an average of 2.5 PH3+ cells per brain (n=11), while 24-hour post-PTBI brains had an average of approximately 11 PH3+ cells per brain (n=17). This is significantly different (p-value>0.0001). At 7 days, uninjured controls have very few EdU+ cells, with an average of approximately 1 EdU+ cell per brain (n=14), while 7-day post-PTBI brains had approximately 12 EdU+ cells per brain (n=22; p-value<0.0001). At a later timepoint, 14 days post-PTBI, uninjured controls have very few EdU+ cells, with an average of approximately 1 EdU+ per brain (n=8). In 14-day post-PTBI brains, there were an average of 29 EdU+ cells per brain (n=14; p-value=0.0002). Interestingly, although there was no significant difference between the number of proliferating cells between 24 hours and 7 days, by 14 days post-PTBI there was an increase in the number of dividing cells (p-value=0.0002). This indicates that cell proliferation continues between 7 and 14 days. Error bars reflect the standard deviation (SD).




Figure 3-2J. Robust proliferative responses decrease with age. To explore whether age impacts the amount of cell proliferation that occurs post-injury, we aged adult flies to 24 hours, 7 days, 14 days, and 28 days prior to PTBI and used anti-PH3 to assay cell proliferation 24 hours after injury. Flies injured at 24 hours had an average of 11 PH3+ cells/brain compared to an average of 2.5 PH3+ cells/brain in age-matched controls. (p-value<0.0001). Flies that were aged to 7 days, then subjected to PTBI and allowed to recover for 24 hours had an average of 6 PH3+ cells per brain (n=11). While this is a significant increase from uninjured control 7-day brains that had an average number of 2 PH3+ cells per brain (n=5; p-value=0.04), this was significantly fewer PH3+ cells than when flies were injured at 24 hours post-eclosion (p-value=0.004). When flies were aged to 14 days, subjected to PTBI, and assayed 24 hours later, there was an average of 1 PH3+ cell per brain. In 14-day controls, there were also very few PH3+ cells, with an average of 0.5 PH3+ per brain. This was not significantly different from the average number of PH3+ cells in injured brains. In 28-day controls, there were an average of 0.25 PH3+ cells in control brains and an average of 0.33 PH3+ cells in injured brains.




Figure 3-3A-D'''. **Analysis of new cell identities 7 days post-PTBI.** To determine what types of new cells had been generated in the first 7 days post-PTBI, we used pulse-chase experiments with EdU in combination with the glial marker anti-Repo and the neuronal marker anti-Elav. At 7 days post-PTBI, we found four classes of cells resulting from proliferation: Class I cells were cells that were EdU+ and did not express either Repo or Elav (**A-A'''**); Class II cells were Repo+/EdU+ (**B-B'''**); Class III were Elav+/EdU+ (**C-C'''**); and Class IV cells were Repo+/Elav+/EdU+ (**D-D'''**). Arrowheads indicate representative cells in each class. The nuclear dye DAPI is in blue.




Figure 3-3E. New cells induced by PTBI express different cell markers at different times post injury. We used cell proliferation markers, such as anti-PH3 at 24 hours and EdU at 7 and 14 days in combination with the glial marker anti-Repo and the neuronal marker anti-Elav to quantify the cells in each class. EdU permanently marks dividing cells and would therefore allow us to see the cumulative number of cells that had been generated since the injury. At 24 hours post-injury, we observed the following approximate distribution of cell classes: 56% Class I (proliferation+ only), 39% Class II (Repo+/proliferation+), 0% Class III (Elav+/proliferation+), and 5% Class IV (Repo+/Elav+/ proliferation+). For 24-hour PTBI brains, n=17 and 181 PH3+ cells were counted. At 7 days post-injury, we saw the following distribution of cell classes: 44% Class I (proliferation+ only), 38% Class II (Repo+/proliferation+), 13% Class III (Elav+/proliferation+), and 5% Class IV (Repo+/Elav+/proliferation +). For the 7-day PTBI brains, n=15 and 172 EdU+ cells were counted. At 14 days post-injury, we saw the following distribution of cell classes: 26% Class I (proliferation+ only), 45% Class II (Repo+/proliferation+), 29% Class III (Elav+/proliferation+), and 0% Class IV (Repo+/Elav+/proliferation +). For the 14-day PTBI brain, n=8 and 278 EdU+ cells were counted. These data demonstrate that cell classes change over time. There are several trends that appear. Class I cells (proliferation+ only) are most frequent at 24 hours post-PTBI at 55%. As time goes on, the proportion of Class I cells decreases, with 44% at 7 days and 26% at 14 days. Class II cells (Repo+/proliferation+) are found in approximately the same percentage at 24 hours and 7 days post-injury, 39% and 38%, respectively. However, at 14 days, there are proportionately more Class II cells, with about 45% of cells in this class. Interestingly, at 24 hours post-PTBI, we see no Class III cells (Elav+/proliferation+). However, at 7 days PTBI, there are 13% of Class III cells, and this percentage more than doubles at 14 days PTBI to 29%. The most unusual cell class, Class IV (Repo+/Elav+/proliferation+) composed 5% of proliferating cells at 24 hours post-injury and was similar at 7 days post-PTBI. However, at 14 days PTBI, we observed 0% of proliferating cells Class IV, suggesting that these cells had transitioned into other classes, such as Class II or III.




Figure 3-3F. Proliferating cells in uninjured brains express different cell markers at different ages. To quantify the proportions of cell classes in control brains, we used the same cell proliferation markers, i.e. anti-PH3 at 24 hours and EdU at 7 and 14 days. At 24 hours, we saw the following distribution of cell classes: 39% Class I (proliferation+ only) and 61% Class II (Repo+/proliferation+). For the 24-hour control brains, n=11 and 28 PH3+ cells were counted. At 7 days of age, we saw the following distribution of cell classes: 47% Class I (proliferation+ only) and 53% Class II (Repo+/proliferation+). For the 7-day control brains, n=9 and 15 EdU+ cells were counted. At 14 days post-injury, we saw the following distribution of cell classes: 67% Class I (proliferation+ only) and 33% Class II (Repo+/proliferation+). For the 14-day PTBI brain, n=2 and 3 EdU+ cells were counted. These data demonstrate that cell classes change over time in control brains and there are several trends that appear. Class I cells (proliferation+ only) appear to increase in proportion over time, from 39% at 24 hours, 47% at 7 days, and 67% at 14 days. This is the opposite trend in Class II cells (Repo+/proliferation+). Class II cells decrease in proportion over time, with 61% at 24 hours, 53% at 7 days, and 33% at 14 days. Notably, neither Class III (Elav+/proliferation+) or Class IV (Repo+/Elav+/proliferation+) were observed in the control brains.




Figure 3-4A-D'''. Neuroblast gene expression is activated by PTBI. (A) and (C) are low magnification views of the control and PTBI brains shown at higher resolution in B-B''' and D-D''', respectively. A-B'''. Images from an uninjured *ase-Gal4, UAS-Stinger; UAS-Gal80^{ts}* probed with anti-PH3 (red) and anti-Dpn (cyan). Stinger expressed under control of the *ase* regulatory sequence is in green. The nuclear dye DAPI is in blue. Arrowheads in (A) and (C) indicate the regions where higher magnification images were collected. At 24 hours in injured (C-D'''), but not in control (A-B''') brains, we observe Stinger+ cells, indicating the expression of *ase*, which is a neuroblast and neural progenitor gene. (D-D'). We observed cells that were Dpn+ (cyan) and PH3+ (red) in injured brains (D''-D'''), but not in controls (B''-B'''). *dpn* also is a neuroblast and neural progenitor gene. We did not observe cells that were GFP+/Dpn+/PH3+ in either injured or control brains. However, the Dpn+/PH3+ cells were often in close proximity to Stinger+ cells (D).




Figure 3-4E-F'. Cytoplasmic Pros seen in cells expressing *ase* at 24 hours post-PTBI. E-F'. An *ase-Gal4>>mCD8-GFP* brain 24 hours post-PTBI immunostained with anti-Pros (red). In this brain, GFP (green) is expressed in the cells that express *ase*. **E.** Low magnification view. Arrowhead indicates region that is seen in higher magnification in **F** and **F'**. **F** and **F'**. High magnification view of area near arrowhead in **E**. There are several cells in which Pros is cytoplasmic, one of which also expresses *ase* (asterisk) and another which lacks *ase* (arrow). Cytoplasmic Pros is characteristic of neuroblasts and neural progenitors and is not observed in control uninjured brains, while nuclear Pros is a hallmark of differentiated neurons and glia.




Figure 3-4G-J. qRT-PCR reveals increases in neural progenitor gene expression following PTBI. The mRNA levels of four different neural progenitor genes were assayed at 4 hours, 24 hours, 3 days, 7 days, and 14 days. **G.** The level of *ase* mRNA is increased more than 5-fold by 4 hours and remains elevated at 24 hours. However, at 3, 7, and 14 days, *ase* mRNA levels are no longer higher than in controls. **H.** The level of *dpn* mRNA was not detectably increased at any timepoints. **I.** mRNA levels of *erm* are increased almost 3-fold at 4 hours post-injury. However, by 24 hours, 3 days, and 7 days, *erm* mRNA levels have returned to baseline. **J.** *insc* mRNA levels are increased 6-fold at 4 hours, 24 hours, and 3 days post-injury. At later timepoints, 7 and 14 days, *insc* mRNA levels return to near baseline. The qPCR results reflect triplicate biological samples, represented relative to the levels of *rp49*, and then normalized to the corresponding levels in time-matched controls. Error bars calculated by Relative Expression Software Tool analysis and reflect the standard error of the mean (SEM). Note that scales on Y axes differ among the graphs.




Figure 3-5A-A''. Lineage-tracing shows that neurons are created through Dpn expressing cells 14 days post-PTBI. Using *dpn-Gal4*, the G-TRACE lineage-tracing system, and a temperature sensitive Gal80, we observed cells that were GFP+/Elav+ at 14 days post-PTBI near the mushroom body, indicated with arrowheads. Other GFP+ cells that did not stain with Elav were also observed, indicated with asterisk. No GFP+/Elav+ cells were observed in age-matched uninjured control brains. DAPI is shown in blue and Elav is shown in Cyan.




Figure 3-6A-D. Perma-twin lineage tracing demonstrates brain regeneration and appropriate targeting of axons following PTBI. To analyze neurogenesis after PTBI, we utilized a lineage-tracing system, called the Perma-twin system, which permanently labels dividing cells and their progeny with either green fluorescent protein (GFP) or red fluorescent protein (RFP). Flies were reared at 17°C to keep the system off during development. Upon eclosion, F1 males carrying transgenes of the Perma-twin system were collected and injured using our standard protocol. Flies were then placed at 30°C to recover for either 2 days or 14 days. **A.** In 2-day uninjured controls, there are some GFP+ cells, and these cells are scattered throughout the brain. **B.** At 14 days, there are relatively few GFP+ cells present in the control central brain. **C.** In comparison, 2-day injured brains have significantly more GFP+ cells, and these cells appear in a cluster near the injury, indicated with arrowhead. **D.** At 14 days post-injury, there are large clones near the site of injury. These clones also have axons and dendrites that correctly project to target areas that were damaged during PTBI. Only the GFP channel is shown here for simplicity; there were similar RFP+ clones in the PTBI samples.




Figure 3-6E. The number of clones increases over time post-PTBI. Control uninjured brains (n=13) have approximately 10 clones at 2 days whereas 2-day PTBI brains (n=20) have a much higher mean of 23 clones ($p < 0.00002$). At 7 days, control brains had an average of 9 clones per brain (n=18), while 7-day PTBI brain had substantially more clones, with an average of 39 clones per brain (n=16) ($p\text{-value} < 0.00000002$). This is significantly more compared to the number of clones seen at 2 days post-injury ($p\text{-value} > 0.0009$). In 14-day uninjured brains, there are ~10 clones per brain, which is not significantly different from the 2-day and 7-day controls. However, at 14 days post-PTBI, there are 66 GFP+ clones, which is significantly more than in age-matched controls ($p < 0.0000003$). The number of clones seen at 14-days post-PTBI is significantly increased compared to 2-day post-PTBI brains ($p\text{-value} < 0.0001$). Error bars reflect SD.




Figure 3-6F-M. PTBI stimulates clone formation in multiple regions in the brain. Panels on the left side show schematics of regions where large clones were found 14 days post-PTBI (**A**, **H**, **J**, **L**). Panels on the right show high magnification of representative images (**G**, **I**, **K**, **M**). Many 14-day brains exhibited clones that had striking large clones that seem to project to correct target areas. There were several structural regions that seem to be able to regenerate post-injury, including the mushroom body (MB) (**F**, **G**), the ellipsoid body (EB) (**H**, **I**), the antennal lobe (AL) (**J**, **K**), and the lateral horn (LH) (**L**, **M**).




Figure 3-6N. Brain regeneration increases over time post-PTBI. The proportion of anatomical structures that regenerated were calculated at 2, 7, and 14 days in controls and injured brains. At 2 days, approximately 8% of control brains (n=13) showed AL regeneration, while in 2-day injured brains (n=20), there were no areas of regeneration. 7-day control brains also exhibited some regions of regeneration (n=18), with 6% showing AL and 6% showing EB clones. At 7 days post-injury, there are many more areas with large clones (n=16), with 6% of brains showing AL, 6% showing EB, and 19% of brain showing large MB clones. At 14 days, control brains (n=9) did not exhibit any specific areas that had regenerated, but 47% of PTBI brains (n=15) had MB clones, 20% had AL clones, 27% had EB clones, and 27% had LH clones.




Figure 3-6O,P. Locomotor defects observed at 2 days post-PTBI have recovered by 14 days post-PTBI. In order to assay for functional recovery post-PTBI, we examined locomotor function. The 2 and 14 day injured, and age-matched uninjured controls were placed in the *Drosophila* Activity Monitor (DAM) system (TriKinetics, Waltham, MA) to record locomotory behavior. **O.** 2-day control uninjured flies displayed stereotypic locomotory patterns throughout a 24-hour period. However, 2-day post-PTBI flies, exhibited significantly different locomotor profiles (p -value <0.001). **P.** Nonetheless, by 14 days, PTBI flies display comparable locomotor profiles to age-matched controls.




Figure 3-7. Summary model for regeneration following penetrating traumatic brain injury (PTBI). We propose that in young adult *Drosophila* there are quiescent NB-like cells within the central brain that lack expression of canonical NB genes. By 24 hours post-PTBI, the quiescent NB-like cells are activated, express NB genes, and have begun to proliferate. At both 4 hours and 24 hours post-PTBI, there is a wave of cell death as assayed using TUNEL. At 7 days, the proliferation rate is still high, and many of the new cells have adopted mature cell identities, becoming neurons or glia. At 10 days post-PTBI, there is no longer a difference in TUNEL+ cells between uninjured brains and injured brains, indicating that the wave of cell death has ended. Because the peaks of both cell death and proliferation occur at the same time post-injury, this could explain why there is not a significant increase in the number of EdU+ cells seen at 7 days compared to the number of PH3+ cells seen at 24 hours. However, by 10 days, cell death is back to control levels while proliferation has decreased but is still slightly above baseline levels. This could explain why there is an increase in the number of EdU+ by 14 days. At 14 days post-PTBI, there are large clones of new neurons with axons and dendrites correctly projecting to their respective target areas. Locomotor defects are also restored by 14 days, suggesting that adult *Drosophila* are able to regenerate functionally as well as structurally.




Figure 3-S1. There is no significant difference in the number of PH3+ and EdU+ cells observed 24 hours post-PTBI. We use two methods, anti-PH3 immunocytochemistry to label mitotic cells and EdU labeling of newly synthesized DNA to assay cell division. To assess the extent to which anti-PH3 and EdU labeling are comparable, we evaluated both control and PTBI brains with both methods at 24 hours. In control brains, there were an average of 2.5 PH3+ (n=11) cells and 2.2 EdU+ cells (n=6). In PTBI brains, there were an average of 10.7 PH3+ cells (n=16) and 10.8 EdU+ cells (n=6). Thus, while control and injured brains displayed significant differences in cell proliferation with both assays (PH3: p-value>0.0001, EdU: p-value>0.0005), the number of proliferating cells detected with the two methods was not significantly different. Error bars reflect the standard deviation (SD).




Figure 3-S2A-D. Cell proliferation is concentrated in the damaged right hemisphere. In order to determine where cell proliferation occurs post-PTBI, we counted PH3+ or EdU+ cells in the left and right hemispheres of the central brain. The schematics in **(A)** and **(B)** illustrate the brain regions assayed. **C.** At 24 hours, control brains (n=11) have ~2 PH3+ cells and in PTBI brains (n=17) have ~3 PH3+ cells in the left hemisphere. In the right hemispheres, control brains have ~1 PH3+ cell while PTBI brains have ~7 PH3+ cells (p-value<0.00006). **D.** At 7 days, in the left hemispheres, control brains (n=8) have an average of 0.6 EdU+ cells while PTBI brains have an average of 4 EdU+ cells (p-value=0.04). In the right hemisphere, control brains (n=8) have an average of 1 EdU+ cell, while PTBI brains (n=14) have an average of 7 EdU+ cells (p-value=0.002). Error bars reflect the standard deviation (SD).




Figure 3-S3. Pulse-chase and continuously fed EdU animals exhibit no difference in the number of EdU+ cells at 7 days post-PTBI. To assess whether mitotically active cells survive post-PTBI, we used two methods of feeding EdU, pulse-chase (flies are fed EdU for 4 days, then placed on standard sugar food for 3 days) and continuously fed (flies are fed EdU every day before being assayed). At 7 days post-PTBI, we find that pulse-chase PTBI brains have an average of 13.5 EdU+ cells (n=8) while continuously fed PTBI brains have an average of 9.1 EdU+ cells (n=7), this is not significantly different. Error bars reflect the standard deviation (SD).




Figure 3-S4A-C. Cell death following PTBI is increased after injury. We assayed cell death using TUNEL staining (in green) in control and PTBI brains. At 24 hours, there is little cell death following our standard penetrating STAB injury (compare **B** and **B'** to **A** and **A'**). Quantification of neurodegeneration is provided in **C**. Groups 1, 2, 3 and 4 have 0-9, 10-19, 20-29, and 30+ TUNEL+ cells/brain respectively. Error bars reflect the standard deviation (SD).




Figure 3-S5. Increase in cell death is inversely correlated with time after injury. In order to understand the temporal dynamics of cell death after PTBI, we used the TUNEL assay to determine how much cell death was present at three time points. We used 4 hours after injury to detect primary cell death resulting from the mechanical injury and 24 hours and 10 days after injury to detect cell death resulting from secondary injury. We found that the both the 4 hour and 24 hour injured samples exhibited significantly higher levels of cell death than the time-matched controls. However, there is no difference between amount of cell death seen in 10-day control and injured brains. Average number of TUNEL+ cells are 13.4 ± 2.5 for 4 hour control and 36.6 ± 5.0 for 4 hour injured samples (p -value=0.0033); 10.4 ± 1.6 for 24 hour controls and 21.4 ± 3.4 for 24 hour injured samples (p -value=0.0203); and 24.2 ± 4.7 for 10 day control and 26.4 ± 3.1 for 10-day injured samples (p -value=0.7053). Error bars reflect the standard deviation (SD).




Figure 3-S6. Mitotic activity highest between 24 hours and 7 days. To assay how many cells were actively dividing as time progressed, we used anti-PH3 at three timepoints: 24 hours, 7 days, and 14 days post-injury. At 24 hours, uninjured control brains had an average of 2.5 PH3+ cells per brain (n=11), while 24-hour post-PTBI brains had an average of approximately 11 PH3+ cells per brain (n=17), which is significantly different (p-value<0.0001). At 7 days, control brains had an average of 1.8 PH3+ cells per brain, while 7 days post-PTBI brains had an average of 11.7 PH3+ cells per brain, which is significantly different (p-value=0.0005). However, 24-hour PTBI and 7-day PTBI brains did not have significantly different numbers of PH3+ cells. At 14 days, control brains had an average of 1.5 PH3+ cells and 14-day post-PTBI brains had an average of 4 PH3+. While this is slightly more than controls, it is not quite statistically significant. Error bars reflect the standard deviation (SD).