A decade of arbovirus emergence in the temperate

southern cone of South America: dengue, *Aedes*

aegypti and climate dynamics in Córdoba, Argentina

Elizabet L. Estallo[1*¶], Rachel Sippy[2,3¶], Anna M. Stewart-Ibarra[2,4], Marta G. Grech[5],

Elisabet M. Benitez[1], Francisco F. Ludueña-Almeida[1,6], Mariela Ainete[7], María Frias-

Cespedes[7], Michael Robert[8], Moory M. Romero[2,9] and Walter R. Almirón[1]

[1] Instituto de Investigaciones Biológicas y Tecnológicas (IIBYT) CONICET-

Universidad Nacional de Córdoba. Centro de Investigaciones Entomológicas de Córdoba,

Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba.

Ciudad Universitaria, Córdoba Capital, Córdoba, Argentina

[2] Institute for Global Health & Translational Sciences, SUNY Upstate Medical

University, Syracuse, NY, USA

[3] Department of Geography, University of Florida, Gainesville, FL, USA

[4] InterAmerican Institute for Global Change Research (IAI), Montevideo, Department of

Montevideo, Uruguay

[5] Centro de Investigación Esquel de Montaña y Estepa Patagónica (CIEMEP), CONICET

and Universidad Nacional de la Patagonia San Juan Bosco. Facultad de Ciencias

Naturales y Ciencias de la Salud, Sede Esquel. Esquel, Chubut, Argentina

[6] Cátedra de Matemática (Cs. Biológicas), Facultad de Ciencias Exactas, Físicas y

Naturales, Universidad Nacional de Córdoba, Ciudad Universitaria, Córdoba Capital,

Córdoba, Argentina

[7] Ministerio de Salud de la Provincia de Córdoba- Dirección de Epidemiología, Hospital

San Roque Viejo, Córdoba Capital, Córdoba, Argentina

[8] Department of Mathematics, Statistics, and Physics, University of the Sciences,

Philadelphia, PA, USA

[9] Department of Environmental Studies, State University of New York College of

Environmental Science and Forestry (SUNY ESF), Syracuse, NY, USA

*Corresponding author:

Email: eelizabet@gmail.com (ELE)

¶These authors contributed equally to this work.

Abstract

Background: Argentina is located at the southern temperate range of arboviral transmission by the mosquito *Aedes aegypti* and has experienced a rapid increase in disease transmission in recent years. Here we present findings from an entomological surveillance study that began in Córdoba, Argentina, following the emergence of dengue in 2009.

Methods: From 2009 to 2017, larval surveys were conducted monthly, from November to May, in 600 randomly selected households distributed across the city. From 2009 to 2013, ovitraps (n=177) were sampled weekly to monitor the oviposition activity of *Ae. aegypti*. We explored seasonal and interannual dynamics of entomological variables and dengue transmission. Cross correlation analysis was used to identify significant lag periods.

Results: *Aedes aegypti* were detected over the entire study period, and abundance peaked during the summer months (January to March). We identified a considerable increase in the proportion of homes with juvenile *Ae. aegypti* over the study period (from 5.7% of homes in 2009-10 to 15.4% of homes in 2016-17). *Aedes aegypti* eggs per ovitrap and larval abundance were positively associated with temperature in the same month. Autochthonous dengue transmission peaked in April, following a peak in imported dengue cases in March; autochthonous dengue was not positively associated with vector or climate variables.

Conclusions: This longitudinal study provides insights into the complex dynamics of arbovirus transmission and vector populations in a temperate region of arbovirus emergence. Our findings suggest that Córdoba is well suited for arbovirus disease

transmission, given the stable and abundant vector populations. Further studies are

needed to better understand the role of regional human movement.

Author summary

There is an increasing risk of arbovirus transmission in temperate regions. Argentina is

located at the southern range of dengue virus transmission by the *Aedes*

aegypti mosquito. In the last decade, epidemics of dengue fever have emerged for the

first time in the city of Córdoba, Argentina. We present the study design and findings

from an entomological surveillance study in Córdoba, which began following the

emergence of dengue in 2009. We found that *Ae. aegypti* were most abundant from

January to March, followed by a peak in local dengue transmission in April. Over the

study period, we noted a considerable increase in the proportion of homes with *Ae.*

aegypti. Vector indices were positively associated with warmer temperatures, which have

been increasing in this region. However, the timing of local dengue transmission appears

to be driven by the appearance of imported dengue cases associated with human

movement. These results highlight the important role of long term surveillance studies in

areas of disease emergence.

Keywords*: Aedes aegypti*; Argentina; dengue; ovitrap; larval surveys; surveillance;

climate.

1. Introduction

Over the last two decades, Argentina has experienced the emergence of epidemics of arboviral diseases vectored by *Aedes* mosquitoes (Aviles et al., 1999; Vezzani and Carbajo, 2008). Cases of dengue fever, chikungunya, and Zika have been reported from northern and central provinces (PAHO, 2019). Argentina is located at the southern limit of arboviral transmission by *Aedes aegypti*, the primary disease vector, thus the increasing burden of disease suggests a southern expansion of the vector and the arboviruses they carry (Masuh, 2008; Robert et al., 2019).

Córdoba, the second largest city in Argentina, experienced its first dengue outbreak in 2009 followed by subsequent outbreaks in 2013, 2015, and 2016 (Robert et al., 2019). In the midst of the COVID-19 global pandemic, Córdoba is currently experiencing the largest dengue outbreak to date (caused by DENV serotypes 1 and 4), which began in early 2020 (Ministerio de Salud de La Nación, 2020). Córdoba is among the southernmost cities in the Western Hemisphere to report autochthonous transmission of *Aedes*-borne arboviruses, highlighting its importance in the investigation of arbovirus emergence in the Americas and globally.

Aedes aegypti was considered eradicated from all of Argentina between 1963-1986 (Curto et al., 2002) and first re-appeared in Córdoba in 1996 (Almirón and Almeida, 1998). Since then, the vector has become established in the city. It is well adapted to the urban environment, living in and around human dwellings and feeding preferentially on human hosts (WHO, 2009).

Climate plays an important role in the *Ae. aegypti* life cycle as well as in its ability to transmit viruses. *Aedes aegypti* juvenile habitat consist of water-bearing

containers such as man-made cisterns, discarded buckets, and tires (Manrique-Saide et

al., 2011). Rainfall and drought can both potentially increase the availability of larval

habitat, depending on local water storage practices and housing characteristics (Lowe et

al., 2018; Stewart Ibarra et al., 2013). Prior studies in the subtropical northwest region of

Argentina revealed a year-round vector population in areas with an annual mean

temperature around 20ºC. A low number of eggs was recorded during the winter season,

and peak in oviposition was detected during summer months (December to March in the

southern hemisphere) (Estallo et al., 2016, 2015, 2013; Micieli and Campos, 2003).

Studies from the temperate central region, including Buenos Aires (Campos and Maciá,

1996; Carbajo et al., 2006, 2004; Micieli et al., 2006; Vezzani et al., 2004) and Córdoba

provinces (Avilés et al., 1997; Domínguez et al., 2000; M. G. Grech, 2013), revealed that

oviposition activity discontinued during the cool winter months due to low ambient

temperatures (below 17ºC).

Vector surveillance studies provide local empirical data on the distribution and

density of arthropod species of public health concern. This information can be used to

trigger public health alert systems, to guide decisions about interventions, and to evaluate

control programs (Focks, 2003). These data can also be used to inform our understanding

of the linkages between vector abundance, disease risk, and local climate conditions. In

response to the first dengue outbreak in Córdoba in 2009, the province created the

"Master Plan to Fight Dengue" and the Ministry of Health (MoH) implemented an

ambitious *Ae. aegypti* surveillance program in cooperation with local academic partners.

Here we report on the study design, methods, and empirical data from this

longitudinal surveillance study for the first time, following the first dengue outbreak in

the temperate city of Córdoba. We describe the seasonal and annual variation in Ae.

aegypti and dengue cases and explore the lagged associations with local climate

variables. We explored the hypotheses that larval abundance is positively associated with

warmer temperatures and greater rainfall in the months prior, and that dengue incidence

is positively associated with greater vector abundance in the months prior. Few studies of

this nature have been conducted in southern temperate zones of arbovirus emergence.

The data generated by this study lays the groundwork for further investigation

regarding the dynamics of arbovirus emergence.

2. Methods

2.1 Ethics

Entomological data were collected from households by the MoH of Córdoba as

part of the routine surveillance program, thus no ethical review or informed consent was

required. Dengue case data at the city-level were extracted from National MoH bulletins

and aggregated to weekly and monthly case counts; no identifying information was

provided.

2.2 Study site

This study was conducted in the city of Córdoba (31.4°S, 64.2°W, population 1.3

million), which is located in the central region of Argentina (elevation: 360-480 m above

sea level) (INDEC, 2010). The city has grown outward from the center to the periphery;

agricultural fields with patches of forest surround the urban core (Maristany et al., 2008)

(Fig. 1). The local climate of Córdoba is warm temperate with hot summers and four

seasons (Cwa under the Köppen-Geiger classification system) (Peel et al., 2007). On

average, the city center is several degrees warmer than the urban periphery, due to dense

construction and location in a topographic depression (Asar et al., 2019). Summer is a

warm, rainy season from November to March (monthly mean temp: 22.7°C, max temp:

37.6°C, min temp: 11.8°C, monthly total rainfall: 123.2 mm), and winter is a cool, dry

season from June to September (monthly mean temp: 14.8°C, max temp: 30.9°C, min

temp: 2.4°C, monthly total rainfall: 36.1 mm). *Aedes aegypti* is the only known vector of

dengue in Córdoba. The vector does not reproduce (oviposit) during the winter,

presumably due to cool temperatures (M. Grech, 2013); it is thought that the vectors

persist through the cold season as egg. Vector control by the MoH is mostly focal around

homes with dengue cases, and includes indoor and outdoor fumigation, larviciding with

Bacillus thuringiensis israelensis (BTI), and eliminating standing water (Ministerio de

Salud de la Nación, 2009).

Fig 1. **Location of Córdoba in South America.** The location of Córdoba Province (dark

purple) shown within the country of Argentina (light purple), with an inset map showing

the city of Córdoba, the location of this study, with roads in grey and waterways in blue.

Note that Córdoba is located in the temperate southern cone of South America.

Field studies were conducted under the *Ae. aegypti* surveillance program of the

Department of Epidemiology of the Córdoba Province Ministry of Health in cooperation

with the Córdoba Entomological Research Centre (CIEC) of Córdoba National

University (UNC) and the Biological and Technological Research Institute (IIByT) of

Córdoba National University (UNC) and the National Research Council (CONICET). No

formal ethical review or consent procedure was required for this study, as the field

operations were conducted as part of routine surveillance activities by the MoH.

Entomological samples were collected from households that verbally agreed to

participate in the study and were taken to the CIEC laboratory to be processed.

2.3 Larval sampling

From 2009 to 2017, the distribution and abundance of *Ae. aegypti* were monitored

through monthly larval surveys conducted across the city. Sample periods each year were

chosen by the CIEC laboratory team to span the period before, during and after mosquito

activity peak in the region (Domínguez et al., 2000). Each month, 600 different homes

were randomly selected for sampling. The city was divided into 215 quadrants (1.2 km

per side), which were distributed across five areas of approximately the same land area:

Central, Northeast, Northwest, Southeast, and Southwest (Fig 1). In each quadrant, we

randomly selected 6 neighborhoods, and the field technicians randomly selected 20

homes to be inspected per neighborhood.

At each home, an inspector noted all containers inside and outside the home with

standing water. Inspectors noted the presence of juvenile mosquitoes and collected

specimens for species identification. Whenever possible, all juvenile mosquitoes were

collected; for large containers where it was not possible to collect all specimens, an

inspector collected three samples using a white dipper (62 ml volume). Specimens were

transported to the CIEC laboratory. Pupae were reared to adults, and 3^{rd} and 4^{th} instar

larvae were preserved in 80% ethanol and were identified using taxonomic keys (Darsie

Jr, 1985). First and 2^{nd} instar larvae were reared until reaching the 3^{rd} instar in plastic

trays with 500 ml of water from the natural larval habitat or dichlorine water. Each day

larvae were fed 0.25 mg of liver powder per larva, and we cleaned the surface of the

water using absorbent paper to avoid contamination by fungi and/or bacteria. The

juvenile mosquitoes were counted and identified to species. For the purposes of this

study, data were aggregated to the city level and we calculated the proportion of

households and neighborhoods with juvenile *Ae. aegypti* present during each sampling

period.

198 2.4 Ovitraps

The weekly oviposition activity of *Ae. aegypti* was observed from November to

May over four years (2009 – 2013) using ovitraps, a sensitive method of detecting the

presence of gravid female of *Ae. aegypti* (Fay & Eliason, 1966). Ovitraps were placed in

randomly selected households (N=177) that were distributed evenly across the city. To

assess oviposition during the winter months, a subset of evenly distributed ovitraps

(n=40) were selected to continue sampling during June (late autumn-early austral winter

seasons) to September 2010 (late winter-early austral spring seasons).

Ovitraps consisted of plastic bottles (350 ml volume, 8 cm diameter x 13 cm

height) with filter paper as an oviposition substrate. An attractive infusion (250 ml) was

prepared by fermenting dry cut grass with tap water (Reiter et al., 1991). Each week the

traps were inspected and replaced in the same dwelling during the sampling years. Traps

were transferred to the laboratory and the number of *Ae. aegypti* eggs per trap were

counted using a stereomicroscope. For the purposes of this study, ovitrap data were

aggregated to the city level. We calculated the proportion of traps that were

positive/negative for *Ae. aegypti* eggs and the mean number of eggs per trap during each

sampling period.

215 2.5 Climate information

The Argentinian National Meteorological Service provided daily weather data for

the study period (rainfall, min/mean/max temperature, min/mean/max relative humidity)

from the Observatory station (31.42°S; 64.20°W). This is one of the oldest weather

stations in Latin America (in operation since 1871) and the most reliable in Córdoba (de

la Casa and Nasello, 2010). We calculated summary monthly values (mean, standard

deviation) over the study period, summary values by epidemiologic week, and summary

values corresponding with the timing of ovitrap and larval survey collection periods.

2.6 Dengue case data

Weekly dengue case reports (January 2009 to December 2017) were extracted

from the weekly epidemiological bulletins of the Argentina Health Secretary (Ministerio

de Salud de Nación (MSN)., 2017). Dengue diagnostic procedures, case definitions and

data extraction procedures have been previously described (Robert et al., 2019). Cases

include suspected and laboratory confirmed cases aggregated at the city-level and annual

incidence was calculated using the total city population for the corresponding year.

Autochthonous and imported cases are reported separately.

2.7 Statistical analysis

Statistical analyses were conducted using time series of vector variables, dengue

incidence (Supplemental Fig 1), and local climate (Supplemental Fig 2) at the city-level

using R version 3.3.3 (*R Studio*, 2015) using the packages splines, TSA, geepack, MASS,

lubridate, TSimpute (Bates et al., 2011; Chan and Ripley, 2012; Grolemund and

Wickham, 2011; Halekoh et al., 2006; Moritz and Bartz-Beielstein, 2017; Team, 2018;

Venables and Ripley, 2013). Missing values in the ovitrap data (n=10) were imputed via

seasonally decomposed structural modeling with Kalman smoothing. Missing values in

the larval data (n=20) were imputed via seasonally decomposed random selection. For

each climate, mosquito abundance, and dengue incidence variable, we performed a

spectral analysis and subsequently tested for the presence of inter-annual variability with

a restricted cubic spline using model fit to determine the periodic frequency and the

number of knots. These descriptive analyses formally test for the presence of periodicity

in a time series and can be used to determine frequency and timing of peaks in the time

series. Assessing inter-annual variability can detect the presence of trends in data

covering multiple years.   Models were fit using a generalized linear model with

appropriate distributions for each variable using generalized estimating equations (auto-

regressive correlation); best fit was determined using quasi-likelihood information

criteria (QIC) as compared to a null model. Residual plots and model assumptions were

examined to inform final model selection. Data were analyzed in relation to the collection

periods of each respective variable; means and frequencies for these collection periods

were calculated for reference.

We assumed a unidirectional temporal relationship between the variables as

follows: climate affecting all other variables, *Ae. aegypti* eggs affecting *Ae. aegypti*

larvae and dengue incidence, and *Ae. aegypti* larvae variables affecting dengue incidence.

Both autochthonous and imported cases were included in the analysis. To examine the

correlations between these variables, cross-correlation functions were calculated between

differenced monthly summary data for each variable with lags up to two months. We

selected this period as it is a biologically plausible period of time that includes the

combined time for *Ae. aegypti* egg hatching and larval development to adult mosquitoes.

3. Results

3.1 *Aedes aegypti* detection

Aedes aegypti were detected over the entire study period, with the exception of

the winter months (June to September), where winter was sampled in 2010 only. There

was a notable increase in *Ae. aegypti* larval abundance over the study period

(Supplemental Fig. 1). On average, *Ae. aegypti* abundance peaked once annually (from

January to March), followed by a peak in autochthonous dengue transmission in April

(Supplemental Fig 1). Our findings support the hypothesis that larval abundance was

associated with warmer temperatures in months prior; however, autochthonous dengue

transmission was not positively associated with larval abundance or climate variables.

Results are described in the following and summary statistics for all variables are shown

in S1 Table.

Aedes aegypti egg abundance was highest in January (Fig. 2), with a median of 33

eggs per trap and 61% of traps positive for eggs during this month. *Aedes aegypti* egg

abundance was highest in the 2009-10 and 2011-12 seasons (Fig. 2), with a mean of 21

eggs per ovitrap per month for both seasons, and 36.4% and 41.6% of traps positive for

Ae. aegypti eggs in 2009-10 and 2011-12, respectively. Note that ovitraps were used only

until 2013 (4 collection seasons).

Fig 2. Seasonal and annual *Aedes aegypti* egg counts and positivity from ovitraps

(2009-2013). Box and whisker plots show the monthly (left) and annual (right) median

and quartiles of *Ae. aegypti* eggs counts and positivity of ovitraps distributed across the

city of Córdoba from 2009 to 2013. Note that ovitraps were sampled during the winter

months (June to September) only in 2010. Annual averages calculated using data

collected during the sampling season (November—May) each year, from 2009 to 2013.

Top: Number of *Ae. aegypti* eggs collected per ovitrap (left) or per ovitrap per annual

sampling season (right). Bottom: The percent of ovitraps with *Ae. aegypti* eggs (left) or

per annual sampling season (right).

Aedes aegypti larval abundance was highest in March (at the residences level) and

February (at the neighborhood level) in each year (Fig. 3). During these months, 14% of

residences and 85% of neighborhoods had containers positive for larvae. Larval

abundance increased over the study period, with the highest infestation levels detected in

2015-16 and 2016-17 (Fig. 3). Fourteen percent and 15.8% of residences, and 83.1 and

71.3% of neighborhoods, had containers positive for *Ae. aegypti* larvae in 2015-16 and

2016-17, respectively.

Fig 3. Seasonal and annual *Aedes aegypti* larval abundance from household larval

surveys (2009-2017). Box and whisker plots show the monthly (left) and annual (right)

median and quartiles of *Ae. aegypti* larval abundance from household larval surveys

conducted in households in Córdoba from 2009 to 2017. Annual averages were

calculated using data collected during the sampling season each year (November—May).

Top: Percent of homes with water-bearing containers with juvenile *Ae. aegypti*. Bottom:

Percent of neighborhoods with water-bearing containers with juvenile *Ae. aegypti*.

Dengue cases were highest during April across all years, with a local transmission

season from March to May (mean 65 autochthonous cases and mean 9 imported cases,

Fig. 4). The number of dengue cases was highest in 2016 (Fig. 4), with 689 total

autochthonous and 139 total imported cases.

Fig 4. Seasonal and annual imported and autochthonous dengue cases in the city of

Córdoba (2009-2017). Box and whisker plots (left) show the monthly median and

quartiles of dengue cases reported in Córdoba from 2009 to 2017. Annual (right)

imported and autochthonous dengue cases in Córdoba as reported by the Ministry of

Health from 2009 to 2017. Top: Total annual reported autochthonous dengue cases (no

travel history). Bottom: Total annual imported dengue cases.

3.2 Periodicity

The results of the seasonal analysis for climate, mosquito abundance, and dengue

incidence variables are presented in Supplemental Table 2. All climate variables were

found to have a 12-month periodicity with nonlinear interannual variability, except for

minimum relative humidity, which had no interannual variability, and maximum relative

humidity, which had no periodicity nor interannual variability (*i.e.* temporal trend). The

detection of temporal trends can be limited with shorter sets of time series data.

Aedes aegypti eggs were collected from ovitraps every 7 days on average; an

average of 26 times within each collection season. Both *Ae. aegypti* egg abundance

variables (trap positivity and mean number of eggs) exhibited a peak about once each

year on average, and the mean number of eggs had an additional peak occurring every 2

years on average. Trap positivity had no interannual variability while the mean number of

eggs had nonlinear interannual variability.

Over the study period, on average, *Ae. aegypti* larvae were collected every 39

days, six times per collection season. *Aedes aegypti* larvae abundance variables

(proportion of positive homes, proportion of positive neighborhoods) were found to have

different seasonal patterns: there was no periodicity nor interannual variability for the

proportion of positive homes, but the proportion of positive neighborhoods peaked once

per year, as well as every 10 years.

Dengue surveillance reports of autochthonous and imported cases occurred every

8 days on average; an average of 39 times per year. Autochthonous dengue incidence

peaked approximately twice per year and annually. Imported dengue incidence peaked

approximately every 1.5 and 3 years. There was no interannual variability in

autochthonous or imported dengue incidence.

3.3 Cross correlation analysis

The results of the cross-correlation between *Ae. aegypti* eggs and larval

abundance, climate, and dengue incidence variables are shown in Table 1. Cross-

correlation function plots for all tested relationships are available in Supplemental Fig. 3

Ae. aegypti egg abundance (mean number of eggs per ovitrap) was positively correlated

with mean temperature in the same month. *Aedes aegypti* larval abundance (percent of

neighborhoods with positive larval traps) was positively correlated with minimum

temperature in the same month, while the percentage of residences with positive larval

traps was not correlated with any climate variable. *Aedes aegypti* egg abundance was

positively correlated with larval abundance (percent of neighborhoods with positive

larval traps) at lag 1.

Autochthonous dengue incidence was negatively correlated with mean and

maximum temperature at lag 2, and negatively correlated with total precipitation in the

same month, with no correlation to any relative humidity variables. Autochthonous

dengue incidence was not correlated with *Ae. aegypti* egg abundance. *Aedes aegypti*

larvae abundance (percent of neighborhoods with larval presence) was negatively

correlated with autochthonous dengue incidence at lag 1.

Table 1. Cross-Correlation Between Climate, Mosquito Abundance, and Dengue Cases for 2009—2017. Using the cross-correlation function, we examined the correlation between pairs of variables, with variable A hypothesized to occur 0 to 2 months before variable B.

Pairing	Variable A	Variable B	Correlated Lags	Direction of Correlation
Climate & Egg Abundance	Min Temp	Mean Number of Eggs	none	
	Mean Temp		0 months	positive
	Max Temp		none	
	Min Rel Humidity		none	
	Mean Rel Humidity		none	
	Max Rel Humidity		none	
	Total Precipitation		none	
Climate & Larvae Abundance	Min Temp	% Positive Homes	none	
	Mean Temp		none	

	Max Temp		none	
	Min Rel Humidity		none	
	Mean Rel Humidity		none	
	Max Rel Humidity		none	
	Total Precipitation		none	
	Min Temp	% Positive Neighborhoods	0 months	positive
	Mean Temp		none	
	Max Temp		none	
	Min Rel Humidity		none	
	Mean Rel Humidity		none	
	Max Rel Humidity		none	
	Total Precipitation		none	
Climate & Dengue Incidence	Min Temp	Autochthonous Dengue Incidence	none	
	Mean Temp		2 months	negative
	Max Temp		2 months	negative
	Min Rel Humidity		none	
	Mean Rel Humidity		none	
	Max Rel Humidity		none	
	Total Precipitation		0 months	negative
Mosquito Egg & Dengue Incidence	Mean Eggs/ovitrap	Autochthonous Dengue Incidence	none	
Mosquito Egg & Larvae Abundance	Mean Eggs/ovitrap	% Positive Homes	none	
		% Positive Neighborhoods	1 months	positive
Mosquito Larvae Abundance & Dengue Incidence	Positive Homes	Autochthonous Dengue Incidence	none	
	Positive Neighborhoods		1 months	negative

4. Discussion

Herein we present the findings from a vector surveillance study in the temperate city of Córdoba, Argentina, during the decade after which the first major epidemic of dengue occurred in the city. This study is among few such studies investigating arbovirus emergence in a temperate climate as it is actively occurring, providing a unique dataset with insights into dengue emergence and the drivers thereof. Among the key findings in this work, we identified a notable increase in the proportion of homes with juvenile *Ae. aegypti* (from 5.7% of homes in 2009-10 to 15.4% of homes in 2016-17). We also found that vector indices increased with warmer temperatures. In prior research, we noted an increase in ambient temperature anomalies over the last 30 years in Córdoba, suggesting a warming climate that could increase suitability for the mosquito vector (Robert et al., 2020). We did not identify positive associations between dengue disease transmission and lagged climate or vector variables, suggesting that disease dynamics in this zone of emergence may be driven more by intrinsic disease dynamics (e.g., herd immunity, importation of new serotypes) than by external environmental factors.

Prior research has shown that temperature is a critical driver of *Ae. aegypti* dynamics due largely to temperature effects on mosquito physiology (Mordecai et al., 2019). In our investigation, we found that the temperatures of Córdoba were warm enough to support the survival and breeding of *Ae. aegypti* populations from October to May. *Aedes aegypti* abundance peaked once annually (from January to March), followed by a peak in autochthonous dengue cases in April. Prior studies from Córdoba found that *Ae. aegypti* egg-to-adult survival began at mean minimum ambient temperatures greater

than 13°C (Domínguez et al., 2000), suggesting that *Ae. aegypti* populations in Córdoba

are not likely to survive when temperatures drop below this threshold. Taken together,

these findings indicate that the temporal window for a stable *Ae. aegypti* population in

Córdoba is rather short, which limits the period in which potential autochthonous dengue

can occur. Indeed, Estallo *et al*. (2014) showed that low autumnal temperatures were an

important factor limiting the spread of dengue during the first dengue outbreak in 2009.

Of note, we found that imported cases had an annual seasonal peak in March,

followed by a peak in autochthonous transmission in April. Local dengue transmission

was not positively associated with lagged climate or vector variables. We suspect that the

negative associations with lagged climate variables may be spurious, and likely due to the

timing of seasonal human movement to neighboring endemic regions within and outside

the country (Estallo et al., 2014). For example, during 2009 dengue outbreak in the city,

the introduction of DENV was associated with outbreaks in the neighboring countries of

Bolivia, Brazil and Paraguay at the end of 2008, and also with dengue transmission in the

Northern provinces of Argentina, where 92% of dengue cases in the country occurred that

year (Estallo et al., 2014). It is possible that human movement into Córdoba from

neighboring areas corresponds with summer breaks in academic schedules in Argentina

(December-February) before university classes begin each year in March. An estimated

60% of the approximately 132,000 university students in the National University of

Córdoba originate from other provinces or countries (Telediario Digital, 2014;

Universidad Nacional de Cordoba, 2017), which could lead to importation of cases by

students who travel between Córdoba and other dengue-endemic regions during the

summer months. A prior study in Brazil also found that dengue transmission was most

closely related to the movement of viremic humans rather than vector densities (Honório

et al., 2009).

This study contributes to a growing body of knowledge on the linkages among

vector abundance, disease cases, and climate. Local studies such as this are necessary, as

these linkages may vary in space and time when comparing regions with distinct eco-

epidemiological contexts (e.g., tropical endemic region versus temperate emergence

region) and when different entomological surveillance methodologies are used (e.g.,

ovitraps, larval surveys, adult trapping). The lack of association between dengue and

juvenile *Ae. aegypti* abundance was confirmed in prior studies in Córdoba (Estallo et al.,

2014). Studies in Venezuela and Malaysia similarly found no relation (Barrera et al.,

2002; Sulaiman et al., 1996), whereas studies in Puerto Rico, Vietnam, and Trinidad

found that vector density measurements were associated with dengue transmission

(Barrera et al., 2011; Chadee, 2009; Pham et al., 2011). It is possible that surveillance of

juvenile vectors did not capture the entomological risk presented by adults of *Ae. aegypti*

and that spatial variation in entomological risk leads to localized hotspots of transmission

risk that are not captured in the aggregated city-level data. We also hypothesize that in a

zone of emergence, the timing of transmission depends on the timing of the introduction

of the virus (as a high proportion of the population is immunologically naïve to dengue)

and transmission may be sustained with low vector densities.

As indicated by the appearance and continued transmission of dengue in Córdoba,

this temperate region is actively experiencing the emergence of dengue (Masuh, 2008;

Robert et al., 2019). The ongoing dengue outbreak is the largest to date, with over 4,000

cases reported thus far (Ministerio de Salud de La Nación, 2020). Emergence of dengue

in temperate regions has been occurring with more frequency in the past two decades

(Lourenço & Recker, 2014; Messina *et al*., 2014; Radke *et al*., 2012). As the climate

becomes increasingly favorable for *Ae. aegypti* in Argentina, we anticipate that the

southern range of the vector will continue to increase, with prolonged periods of activity

during the year. Empirical studies such as this, from the lower thermal range of arbovirus

transmission, can be used to improve models that investigate arbovirus dynamics in

temperate zones of emergence, predict future outbreaks, and explore potential control

measures.

It is important to note that we do not expect that the reported correlations between

climate, mosquito abundance, dengue incidence to be causal in any way. Cross-

correlation analyses do not adjust for potential confounding and we expect that the

correlation found is due to the coincidental timing of Argentina vacation periods and

peak larval activity. One limitation of our work is irregular sampling in the surveillance

data, creating gaps in some periods, though the sampling design was strong and carefully

designed by the team of researchers at the CIEC. It is also worth noting that the monthly

summaries used in this study do not allow for an analysis of finer scale climate effects on

mosquito vectors (e.g., daily diurnal temperature fluctuations) (Lambrechts et al., 2011).

In addition, the shorter time series of ovitrap data does not allow for a more robust

comparison to dengue transmission data.

5. Conclusions

This longitudinal study provides insights into the complex dynamics of arbovirus

transmission and vector populations in a temperate region of arbovirus emergence. Our

findings suggest that Córdoba is well suited for arbovirus disease transmission, given the

stable and abundant vector populations. Further studies are needed to better understand

the role of regional human movement, including origins, destinations, and timing,

throughout South America to understand local patterns of disease transmission.

Conflict of interest

The authors report no conflicts of interest.

Acknowledgements

The authors wish to acknowledge the United States Embassy in Argentina and Fulbright

Commission as well as the Department of Epidemiology of the Córdoba Province

Ministry of Health. AMSI and MAR were support in Córdoba Argentina by the USA

Zika program of the United States Embassy in Argentina administered by the Fulbright

commission. ELE, MGG, and WRA are members of the Consejo de Investigaciones

Cientificas y Tecnológicas (CONICET) from Argentina, EMB is a PhD Student with

scholarship support from CONICET.

References

Almirón, W.R., Almeida, F.F., 1998. Aedes aegypti (Diptera: Culicidae) en Córdoba,

Argentina. Rev. Soc. Entomológica Argent. 57.

Asar, M.L., Estallo, E.L., Benítez, E.M., Di Benedetto, H.M., Ludueña-Almeida, F.F.,

2019. Un estudio sobre el efecto Islas de Calor Urbano en la ciudad de Córdoba,

Argentina.

Avilés, G., Cecchini, R., Harrington, M.E., Cichero, J., Asis, R., Rios, C., 1997. Aedes

aegypti in Córdoba province, Argentina. J. Am. Mosq. Control Assoc.-Mosq.

News 13, 255–258.

Aviles, G., Rangeón, G., Vorndam, V., Briones, A., Baroni, P., Enria, D., Sabattini, M.S.,

1999. Dengue reemergence in Argentina. Emerg. Infect. Dis. 5, 575.

Barrera, R., Amador, M., MacKay, A.J., 2011. Population Dynamics of Aedes aegypti

and Dengue as Influenced by Weather and Human Behavior in San Juan, Puerto

Rico. PLoS Negl Trop Dis 5, e1378. https://doi.org/10.1371/journal.pntd.0001378

Barrera, R., Delgado, N., Jiménez, M., Valero, S., 2002. Eco-epidemiological factors

associated with hyperendemic dengue haemorrhagic fever in Maracay City,

Venezuela. Dengue Bull. 26, 84–95.

Bates, M.D., Venables, B., Team, M.R.C., 2011. Package 'splines.' R Version 2.

Campos, R.E., Maciá, A., 1996. Observaciones biológicas de una población natural de

Aedes aegypti (Diptera: Culicidae) en la provincia de Buenos Aires, Argentina.

Rev Soc Entomol Argent 55, 67–72.

Carbajo, A.E., Curto, S.I., Schweigmann, N.J., 2006. Spatial distribution pattern of

oviposition in the mosquito Aedes aegypti in relation to urbanization in Buenos

Aires: southern fringe bionomics of an introduced vector. Med. Vet. Entomol. 20,

209–218.

Carbajo, A.E., Gomez, S.M., Curto, S.I., Schweigmann, N.J., 2004. Variación espacio-

temporal del riesgo de transmisión de dengue en la Ciudad de Buenos Aires. Med.

B. Aires 64, 231–234.

Chadee, D.D., 2009. Dengue cases and< i> Aedes aegypti</i> indices in Trinidad, West

Indies. Acta Trop. 112, 174–180.

Chan, K.-S., Ripley, B., 2012. TSA: time series analysis. R package version 1.01. URL

HttpCRAN R-Proj. Orgpackage TSA.

Curto, S., Boffi, R., Carbajo, A.E., Plastina, R., 2002. Reinfestación del territorio

Argentino por Aedes aegypti. Distribución geográfica (1994-1999), in:

Actualizaciones En Artropodología Sanitaria Argentina. Fundación Mundo Sano,

Buenos Aires, pp. 127–137.

Darsie Jr, R.F., 1985. Mosquitoes of Argentina. I. Keys for identification of adult females

and fourth stage larvae in English and Spanish (Diptera, Culicidae). Mosq. Syst.

USA.

de la Casa, A., Nasello, O., 2010. Breakpoints in annual rainfall trends in Córdoba,

Argentina. Atmospheric Res. 95, 419–427.

Domínguez, M.C., Ludueña Almeida, F.F., Almirón, W.R., 2000. Dinámica poblacional

de Aedes aegypti (Diptera: Culicidae) en Córdoba capital. Rev Soc Entomol

Argent 59, 41–50.

Estallo, E.L., Benitez, E.M., Lanfri, M.A., Scavuzzo, C.M., Almirón, W.R., 2016.

MODIS environmental data to assess Chikungunya, Dengue, and Zika diseases

through Aedes (Stegomia) aegypti oviposition activity estimation. IEEE J. Sel.

Top. Appl. Earth Obs. Remote Sens. 9, 5461–5466.

Estallo, E.L., Carbajo, A.E., Grech, M.G., Frías-Céspedes, M., López, L., Lanfri, M.A.,

Ludueña-Almeida, F.F., Almiron, W.R., 2014. Spatio-temporal dynamics of

dengue 2009 outbreak in Córdoba City, Argentina. Acta Trop. 136, 129–136.

Estallo, E.L., Ludueña-Almeida, F.F., Introini, M.V., Zaidenberg, M., Almirón, W.R.,

2015. Weather variability associated with Aedes (Stegomyia) Aegypti (Dengue

vector) Oviposition dynamics in northwestern Argentina. PLoS One 10,

e0127820.

Estallo, E.L., Más, G., Vergara-Cid, C., Lanfri, M.A., Ludueña-Almeida, F., Scavuzzo,

C.M., Introini, M.V., Zaidenberg, M., Almirón, W.R., 2013. Spatial patterns of

high Aedes aegypti oviposition activity in northwestern Argentina. PloS One 8,

e54167.

Focks, D.A., 2003. A review of entomological sampling methods and indicators for

dengue vectors. Geneva WHO.

Grech, M., 2013. Bioecologia de culícidos (Diptera) urbanos de importancia sanitaria de

la ciudad de Córdoba, Córdoba (Argentina). Facultad de Ciencias Exactas Físicas

y Naturales. Universidad Nacional de Córdoba, Córdoba, Argentina.

Grech, M.G., 2013. Bioecología de culícidos (Diptera) urbanos de importancia sanitaria

de la ciudad de Córdoba, Córdoba (Argentina) (Doctorado en Ciencias

Biológicas). National University of Cordoba, Cordoba, Cordoba Province,

Argentina.

Grolemund, G., Wickham, H., 2011. Dates and times made easy with lubridate. J. Stat.

Softw. 40, 1–25.

Halekoh, U., Højsgaard, S., Yan, J., 2006. The R package geepack for generalized

estimating equations. J. Stat. Softw. 15, 1–11.

Honório, N., Codeço, C., Alves, F., Magalhães, M., Louren\cco-de-Oliveira, R., 2009.

Temporal distribution of Aedes aegypti in different districts of Rio de Janeiro,

Brazil, measured by two types of traps. J. Med. Entomol. 46, 1001–1014.

INDEC, 2010. Censo Nacional de Población, Hogares y Viviendas 2010.

La UNC en cifras – Centenario de la Reforma Universitaria, n.d. URL

https://centenariodelareforma.unc.edu.ar/la-unc-en-cifras/ (accessed 10.8.19).

Lambrechts, L., Paaijmans, K.P., Fansiri, T., Carrington, L.B., Kramer, L.D., Thomas,

M.B., Scott, T.W., 2011. Impact of daily temperature fluctuations on dengue virus

transmission by Aedes aegypti. Proc. Natl. Acad. Sci. 108, 7460–7465.

https://doi.org/10.1073/pnas.1101377108

Lowe, R., Gasparrini, A., Van Meerbeeck, C.J., Lippi, C.A., Mahon, R., Trotman, A.R.,

Rollock, L., Hinds, A.Q., Ryan, S.J., Stewart-Ibarra, A.M., 2018. Nonlinear and

delayed impacts of climate on dengue risk in Barbados: A modelling study. PLoS

Med. 15, e1002613.

Manrique-Saide, P., Che-Mendoza, A., Rizzo, N., Arana, B., Pilger, D., Lenhart, A.,

Kroeger, A., 2011. Operational guide for assessing the productivity of Aedes

aegypti breeding sites. Geneva Switz. WHO-TDR 1–30.

Maristany, A., Abadía, L., Angiolini, S., Pacharoni, A., Pardina, M., 2008. Estudio del

fenómeno de la isla de calor en la ciudad de Córdoba-Resultados preliminares.

Av. Energ Renov Medio Ambient 12, 11–69.

Masuh, H., 2008. Re-emergence of dengue in Argentina: Historical development and

future challenges.

Micieli, M.V., Campos, R.E., 2003. Oviposition activity and seasonal pattern of a

population of Aedes (Stegomyia) aegypti (L.) (Diptera: Culicidae) in subtropical

Argentina. Mem. Inst. Oswaldo Cruz 98, 659–663. https://doi.org/10.1590/S0074-

02762003000500013

Micieli, M.V., García, J.J., Achinelly, M.F., Martí, G.A., 2006. Dinámica poblacional de

los estadios inmaduros del vector del dengue Aedes aegypti (Diptera: Culicidae):

un estudio longitudinal (1996-2000). Rev. Biol. Trop. 54, 979–983.

Ministerio de Salud de La Nación, 2020. Boletín Integrado de Vigilancia. 2020 15, 42.

Ministerio de Salud de la Nación, 2009. Plan nacional para la prevención y el control del

dengue y la fiebre amarilla.

Ministerio de Salud de Nación (MSN)., 2017. Boletines Epidemiologicos 2009-2017.

Mordecai, E.A., Caldwell, J.M., Grossman, M.K., Lippi, C.A., Johnson, L.R., Neira, M.,

Rohr, J.R., Ryan, S.J., Savage, V., Shocket, M.S., Sippy, R., Ibarra, A.M.S.,

Thomas, M.B., Villena, O., 2019. Thermal biology of mosquito-borne disease.

Ecol. Lett. 22, 1690–1708. https://doi.org/10.1111/ele.13335

Moritz, S., Bartz-Beielstein, T., 2017. imputeTS: time series missing value imputation in

R. R J. 9, 207–218.

PAHO, 2019. PLISA Health Information Platform for the Americas [WWW Document].

Pan Am. Health Organ. World Health Organ. URL

http://www.paho.org/data/index.php/en/?option=com_content&view=article&id=

524&Itemid= (accessed 10.8.19).

Peel, M.C., Finlayson, B.L., McMahon, T.A., 2007. Updated world map of the Köppen-

Geiger climate classification. Hydrol. Earth Syst. Sci. Discuss. 4, 439–473.

Pham, H., Doan, H., Phan, T., Minh, N.T., 2011. Ecological factors associated with

dengue fever in a central highlands Province, Vietnam. BMC Infect. Dis. 11, 172.

Población estudiantil: bajo la lupa de la UNC | Telediario Digital [WWW Document],

n.d. URL https://www.telediariodigital.net/2014/11/poblacion-estudiantil-bajo-la-

lupa-de-la-unc/ (accessed 10.8.19).

R Studio, 2015. . RStudio Team. Integrated Development for R Inc., Boston, MA, USA.

Reiter, P., Amador, M.A., Colon, N., others, 1991. Enhancement of the CDC ovitrap with

hay infusions for daily monitoring of Aedes aegypti populations. J. Am. Mosq.

Control Assoc. 7, 52.

Robert, M., Stewart Ibarra, A.M., Estallo, E.L., 2020. Climate change and viral

emergence: Evidence from Aedes-borne arboviruses. Curr. Opin. Virol. in press.

Robert, M.A., Tinunin, D.T., Benitez, E.M., Ludueña-Almeida, F., Romero, M., Stewart-

Ibarra, A.M., Estallo, E.L., 2019. Arbovirus emergence in temperate climates: the

case of Córdoba, Argentina, 2009-2018. bioRxiv 602003.

https://doi.org/10.1101/602003

Stewart Ibarra, A.M., Ryan, S.J., Beltrán, E., Mejía, R., Silva, M., Muñoz, Á., 2013.

Dengue Vector Dynamics (Aedes aegypti) Influenced by Climate and Social

Factors in Ecuador: Implications for Targeted Control. PLOS ONE 8, e78263.

Sulaiman, S., Pawanchee, Z.A., Arifin, Z., Wahab, A., 1996. Relationship between

Breteau and House indices and cases of dengue/dengue hemorrhagic fever in

Kuala Lumpur, Malaysia. J. Am. Mosq. Control Assoc. 12, 494.

Team, R.C., 2018. R: A language and environment for statistical computing; 2015.

Telediario Digital, 2014. Población estudiantil: bajo la lupa de la UNC [WWW

Document]. URL https://www.telediariodigital.net/2014/11/poblacion-estudiantil-

bajo-la-lupa-de-la-unc/ (accessed 4.29.20).

Universidad Nacional de Cordoba, 2017. La UNC en cifras – Centenario de la Reforma

Universitaria,.

Venables, W.N., Ripley, B.D., 2013. Modern applied statistics with S-PLUS. Springer

Science & Business Media.

Vezzani, D., Carbajo, A.E., 2008. Aedes aegypti, Aedes albopictus, and dengue in

Argentina: current knowledge and future directions. Mem. Inst. Oswaldo Cruz

103, 66–74.

Vezzani, D., Velázquez, S.M., Schweigmann, N., 2004. Seasonal pattern of abundance of

Aedes aegypti (Diptera: Culicidae) in Buenos Aires city, Argentina. Mem. Inst.

Oswaldo Cruz 99, 351–356.

WHO, 2009. Dengue: guidelines for diagnosis, treatment, prevention and control. World

Health Organization.

Supplemental Figure 1. Time series of monthly dengue, *Aedes aegypti* **larval and** *Ae.*

aegypti **eggs abundance.** Top: Dengue cases. Center: proportion of homes with *Ae.*

aegypti larvas. Bottom: proportion of ovitraps (N=177) positive for *Ae. aegypti* and the

mean number of eggs per ovitrap.

Supplemental Figure 2. Times series of monthly climate variables. Mean monthly

temperatures (°Celsius) are in the top panel (red) (minimum and maximum in dashed

lines), total monthly precipitation (mm) is in the central panel (blue) and mean monthly

relative humidity (percentage) is in the bottom panel (green) (minimum and maximum in

dashed lines).

Supplemental Figure 3. Cross-correlation function plots for climate variables, *Aedes*

aegypti **abundance, and dengue cases.** Cross-correlation function (CCF) plots are given

for tested correlations between climate variables and egg abundance (a), climate variables

and larval abundance (b), climate variables and dengue incidence (c), egg abundance and

dengue incidence (d), egg abundance and larval abundance (e), and larval abundance and

dengue incidence (f). We assumed a unidirectional temporal relationship between the

variables as follows: climate affecting all other variables, *Aedes aegypti* eggs affecting

Aedes aegypti larvae and dengue incidence, and *Aedes aegypti* larvae variables affecting

dengue incidence. Lags of 0-2 months were tested.

Supplemental Table 1. Summary Statistics of Climate, *Aedes aegypti***, and Dengue**

Measures. Mean and range by year are given for climate, *Aedes aegypti*, and dengue

measures used in this study.

Supplemental Table 2. Seasonality and Interannual Variability of Data.

Equator

Tropic of Capricorn

1000   0   1000   2000   3000   Kilometers

A

N
W   E
S

2.5   0   2.5   5   7.5   Kilometers

B

a. Climate Variables & Egg Abundance

b) Climate Variables & Larval Abundance

c) Climate Variables & Dengue Incidence

d) Egg Abundance & Dengue Incidence

e) Egg Abundance & Larval Abundance

f) Larval Abundance & Dengue Incidence