Title

The relationship between longevity and diet is genotype dependent and sensitive to desiccation

Running title

Dietary reaction norms are genotype dependent

Authors

Andrew W McCracken, Eleanor Buckle, Mirre J. P. Simons

Affiliations

Department of Animal and Plant Sciences & Bateson Centre, The University of Sheffield, Sheffield S10 2TN, UK.

Corresponding Author

Mirre J. P. Simons

Alfred Denny Building, Department of Animal and Plant Sciences, The University of Sheffield, Sheffield S10 2TN, UK.

+44 (0)114 222 0111

m.simons@sheffield.ac.uk

Additional email addresses

awmccracken1@sheffield.ac.uk

eleanor.buckle96@gmail.com

Keywords

Dietary restriction, Drosophila melanogaster, reaction norm, overfeeding, starvation, desiccation

Summary

Dietary restriction (DR) is a key focus in ageing research. Specific conditions and genotypes

were recently found to negate lifespan extension by DR, questioning its universal relevance.

However, the conceptual framework of dietary reaction norms explains why DR's effects might

not be apparent in some situations. We tested comprehensively the importance of dietary

reaction norms by measuring longevity and fecundity on five diets in five genotypes, with and

without water supplementation in the fly (N>25,000). We found substantial genetic variation in

the reaction norm between diet and lifespan. Environments supplemented with water rescued

putative desiccation stress but only at the richest diets. Fecundity declined at these richest diets,

but was unaffected by water and is thus most likely caused by nutritional toxicity. Our results

demonstrate empirically that any conclusion on the absence of DR is only justified when a range

of diets is considered in a reaction norm framework.

Introduction

Dietary restriction (DR), the limitation of food intake but avoiding malnutrition, extends lifespan. The

generality of the DR response has been questioned, however, by reports that DR does not extend

lifespan under certain experimental conditions (Austad, 2012; Dick et al., 2011; Ja et al., 2009 [but

see Piper et al., 2010]) or in a considerable proportion of the genotypes tested (Dick et al., 2011; Liao

et al., 2010; Mitchell et al., 2016; Rikke et al., 2010; Swindell, 2012; Wilson et al., 2018 preprint).

These conclusions are routinely based upon experiments using two diets (dietary dyad) alone, whereas

it is recognised that a change in the continuous relationship between diet and lifespan (reaction norm)

can obscure lifespan extension by DR (Flatt, 2014; Tatar, 2011). The bell-shaped nature of the dietary

reaction norm dictates that one particular diet concentration, in one genotype or environment, will

result in the longest lifespan; lower or higher diet concentrations will induce a shortened lifespan due

to malnutrition or overfeeding, respectively. Where a particular dietary dyad falls on this reaction

norm will determine the magnitude of the DR effect and can even lead to the erroneous conclusion

that DR shortens lifespan (Fig.1).

Few studies have examined dietary reaction norms in more detail by titrating the supply of protein or

calories across multiple genotypes or environments, and none have tested both genetic and

environmental effects on dietary reaction norms simultaneously. Of these studies, a proportion

employed transgenic or lab strains (Clancy et al., 2002; Grandison et al., 2009; Min et al., 2008;

Skorupa et al., 2008; Tatar, 2011; Wang et al., 2009) and demonstrated varying degrees of genetic

variance in the plastic response to diet. Across these studies, shifts in dietary reaction norms on the x-

or y-plane are more apparent than changes in the overall shape of the relationship between diet and

longevity (Flatt, 2014; Tatar, 2011). Whether genetic variation in transgenic and lab strain

experiments is representative of standing genetic variation of natural populations is, however, unclear.
A naturalistic appreciation of the genetic variation of the DR response becomes particularly important
when null responses are interpreted to question the universal properties of DR important in translating
its benefits to our own species. One previous study did measure detailed reaction norms using wild-
derived outbred populations and found a degree of genetic variance for the relationship between diet
and lifespan (Metaxakis and Partridge, 2013). However, the estimate of genetic variance of a
population level trait, such as lifespan, when estimated from between outbred stains (Whitlock and
Fowler, 1999) will be affected by mortality heterogeneity (Chen et al., 2013), which can bias the
estimated level of genetic variance upwards or downwards.
46
When specific environmental effects interact or interfere with the DR reaction norm, the use of dietary
dyads - or the neglect of environmental confounds - could similarly lead to misleading conclusions.
For flies specifically, water supplementation has been suggested to diminish the effect of DR on
lifespan (Dick et al., 2011; Ja et al., 2009). The conclusion that water completely explains DR has
been discredited (Piper et al., 2010), but flies nonetheless consider water a nutrient and consume 1-2μl
per day, with higher consumption at higher dietary yeast (Fanson et al., 2012) and sugar
concentrations (van Dam et al., 2020). Hence, erroneous conclusions could be drawn from diet
responses if desiccation presents a genotype- or diet-specific hazard.
55
56 Here, we present DR reaction norms for fecundity and longevity across five genotypes in female flies
57 (*Drosophila melanogaster*) with and without water supplementation using high sample sizes. We
58 show empirically across five wild-derived, inbred lines that there are strong genetic and
59 environmental elements to dietary reaction norms, and therefore the thorough appreciation of reaction
60 norms is critical when interpreting diet effects across genotypes and environments.
61
62
63 **Materials and Methods**
64
Fly husbandry, experimental protocol and dietary regimes
For lifespan experiments adults were provided with either 0.5%, 2%, 5%, 8% or 14% autolysed yeast
media. All other media components (13% table sugar, 6% cornmeal, 1% agar and 0.225% [w/v]
nipagin)l remained the same, given the dietary protein axis is the main lifespan determinant in flies
(Jensen et al., 2015; Lee et al., 2008). Note, cornmeal concentration was halved in 14% yeast media to
allow dispensing of the media. Purpose-built demography cages included two openings, one for the
supplementation of food, and one for water-agar (2% agar) or empty vial. Cages contained between
70-125 females each (mode of ~ 100 females), with 5 cages per treatment, per genotype (N = 50 cages

per genotype). For one genotype, DGRP-195, sample size was even higher: an additional two cages of

water-supplemented, and control cages at 2% media. All experimental flies were mated on 8% media

for 48 hours, and kept in cages on 8% media until age 3-4 days, when experimental dietary treatments

started.

To establish dietary reaction responses, flies were exposed to continuous diets with the addition, or

absence of water-agar supplementation. To test the effect of water supplementation on longevity, we

provided an additional vial of water-agar ('water supplementation'), or an empty vial ('control'), to

each cage. Separation of food and water sources allowed flies to choose their source of nourishment,

and eliminated the need for hydration to be coupled with caloric intake. Dietary treatments were

balanced for age, and date of eclosion. All flies presented were grown within one batch. The

experiment was carried out on a small collection of DGRP lines (Mackay et al., 2012); DGRP-195;

217; 239; 362; 853).

Fecundity

Feeding vials were imaged and analysed using image analysis software QuantiFly (Waithe et al.,

2015) to determine the relative quantity of egg laying. Vials were removed, during normal scoring

periods, from all cages containing eggs from flies aged 11 or 12.

Data analysis

Mixed Cox-proportional hazard models were used that included 'cage' as random term to correct for

uncertainty of pseudo-replicated effects within demography cages (Ripatti and Palmgren, 2000;

Therneau et al., 2003). Additional specific tests of coefficients are provided that combine the single

and interaction term (in a z-test, using the maximum s.e. of the factors compared) to test how survival

was changing in water-treated flies, compared to respective control treatments. Note, formal tests for

proportionality of hazards are not available for mixed effects Cox regressions. For survival data

comparisons, we report the full model, and models fitted within each genotype separately (see

supplement). By splitting the analysis between genotypes, bias introduced by deviations in

proportionality of hazards between genotypes is avoided. Qualitative conclusions remain similar,

irrespective of how these models are fitted. Coefficients are reported as logged hazards with

significance based on z-tests. Right-censoring was included, and dietary treatments were considered

categorical factors.

Egg laying was analysed using linear models of log-transformed fecundity count data. Flies only

differed by one day in age, and age was equally distributed across treatment and measured in a

balanced design. BIC with backward elimination of terms was used for model comparisons and
selection, and resulted in a model that only contained the interaction between genotype and diet.
Water was added to our models to directly test for any effect on fecundity, but this proved negligible
(Table S13,14).
113
114 For hazard ratio figures, ratios are plotted as coefficients derived from within-line Cox mixed-effect
115 models, with error bars representing 95% confidence intervals.
116
117

<u>Results and Discussion</u>
The genotypes tested showed a classical bell-shaped response to diet. Longer lifespans were observed
at intermediate dietary yeast concentrations, consistent with DR (Fig.2A,S4; Table S1-6; $P < 0.001$).
All genotypes also exhibited a reduction in survival at very lowest yeast concentrations (starvation),
and at the very highest (maximal performance or nutritional toxicity). We detected considerable
genetic variation in the response to diet ($\chi2=162$, df=16, P< 0.001) with the diet of maximum
longevity differing between genotypes.
125

To test the effect of desiccation, we compared longevity under control conditions to water-
supplemented. Supplemental water reduced mortality particularly at higher yeast concentrations, and
we found genetic variance for this environmental effect ($\chi2=160$, df=16, P<0.001; Fig.2B; Table S2-
6). At the highest yeast concentrations, this amounted to a one and a half- to fifty-fold reduction in
hazard rate. Given this, particular caution should be afforded when considering the effect of
desiccation, especially in organisms without ad libitum access to water and when fed a concentrated
diet. To assess statistically whether water supplementation abolished DR-induced life extension (Ja et
al., 2009; Piper et al., 2010) we ran our statistical models within the water treatment only, but found
no evidence for this suggestion (Fig.S1, Table S7-12). Given that water supplementation ameliorated,
but did not eliminate, elevated mortality under the highest yeast concentrations, we conclude
desiccation can play an experimentally confounding role in DR, but is not causal.
137

DR is known to reduce reproductive output and is interpreted as a response to decreased energy
availability (Moatt et al., 2016). The effect of overfeeding on reproduction, although appreciated in
humans (Broughton and Moley, 2017), has received little attention (McCracken et al., 2020). These
two responses were evident in egg laying: an increase with yeast concentration, and a stabilisation, or
decline at the highest yeast concentrations (Fig.S2,3; Table S13,14). As with mortality, genetic lines
also differed in the reproductive response to diet (F=6.3, df=16, P< 0.001). Reduced egg laying
together with a reduction in survival, lowered predicted lifetime reproductive at the richest diet

(Fig.S3). Egg laying was not affected by water supplementation (Fig.S2; Table S13,14; P>0.15.

Notably, even when water rescued mortality caused by desiccation at the high yeast concentrations,

egg laying was unaffected (Fig.2,S2). Given this, we infer the decline in reproductive output at the

highest yeast concentration was not due to desiccation stress, but nutritional toxicity. By contrast, the

rescue of mortality at high yeast concentrations by water supplementation is therefore likely to be

separate and driven by desiccation.

In conclusion, we observe significant genetic, and environmentally induced, variation in the lifespan

and fecundity responses to diet. These data now directly demonstrate that specific care is needed when

interpreting effects of DR across genotypes, experimental conditions or environments. We

acknowledge that carrying out full reaction norms in all DR experiments would be laborious.

Therefore, we suggest selecting dietary dyads that differ only minimally when genetic variance in DR

is the object of study. Such a strategy reduces the chance that tested diets diverge considerably from

maximal lifespans, leading to starvation or nutritional toxicity (Fig.1). Furthermore, we suggest when

environmental conditions, such as water (Ja et al., 2009), sex (Regan et al., 2016) and microbiome

(Wong et al., 2014) are presumed to negate the DR response, that a post-hoc reaction norm is

performed. Similar considerations hold for mechanistic research. Should, for example, a genetic

manipulation remove the DR response, only a full dietary reaction norm can demonstrate how such an

effect arises: by either a shift in, or compression of, the reaction norm (Flatt, 2014; Tatar, 2011). The

importance of reaction norms when studying DR has been stressed before, but this is the first high

sample size data across multiple wild-type inbred genotypes and diets, including an environmental

confound, that demonstrates this empirically.

Acknowledgements

We thank the Deplancke laboratory for supplying DGRP lines and members of Simons' lab for their

valuable assistance.

Competing interests

The authors declare no competing interests.

Author Contributions

AWM and MJPS designed and interpreted the experiments. AWM drafted the first version of the

manuscript. AWM and MJPS revised the manuscript. AWM led the data acquisition with help from

EB. MJPS supervised the project.

Funding

AWM is supported by the NERC ACCE DTP. EB was funded by the SURE scheme and the 301

academic skills centre. MJPS was supported by a Sir Henry Wellcome, a Sheffield Vice Chancellor's

Fellowship, the Natural Environment Research Council (M005941 & N013832) and currently by a Sir

Henry Dale Fellowship (Wellcome & Royal Society) and an Academy of Medical Sciences

Springboard Award (the Wellcome Trust, the Government Department of Business, Energy and

Industrial Strategy (BEIS), the British Heart Foundation and Diabetes UK).

References

Austad, S. N. (2012). Mixed results for dieting monkeys. *Nature* **489**, 210–211.

Broughton, D. E. and Moley, K. H. (2017). Obesity and female infertility: potential
mediators of obesity's impact. *Fertil. Steril.* **107**, 840–847.

Chen, H.-Y., Zajitschek, F. and Maklakov, A. A. (2013). Why ageing stops: heterogeneity
explains late-life mortality deceleration in nematodes. *Biol. Lett.* **9**, 20130217.

Clancy, D. J., Gems, D., Hafen, E., Leevers, S. J. and Partridge, L. (2002). Dietary
restriction in long-lived dwarf flies. *Science* **296**, 319.

Dick, K. B., Ross, C. R. and Yampolsky, L. Y. (2011). Genetic variation of dietary
restriction and the effects of nutrient-free water and amino acid supplements on lifespan and
fecundity of Drosophila. *Genet. Res.* **93**, 265–273.

Fanson, B. G., Yap, S. and Taylor, P. W. (2012). Geometry of compensatory feeding and
water consumption in Drosophila melanogaster. *J. Exp. Biol.* **215**, 766–773.

Flatt, T. (2014). Plasticity of lifespan: a reaction norm perspective. *Proc. Nutr. Soc.* **73**, 532–
542.

Grandison, R. C., Wong, R., Bass, T. M., Partridge, L. and Piper, M. D. W. (2009). Effect
of a standardised dietary restriction protocol on multiple laboratory strains of Drosophila
melanogaster. *PLoS One* **4**, e4067.

Ja, W. W., Carvalho, G. B., Zid, B. M., Mak, E. M., Brummel, T. and Benzer, S. (2009).
Water- and nutrient-dependent effects of dietary restriction on Drosophila lifespan. *PNAS*
USA. **106**, 18633–18637.

Jensen, K., McClure, C., Priest, N. K. and Hunt, J. (2015). Sex-specific effects of protein
and carbohydrate intake on reproduction but not lifespan in Drosophila melanogaster. *Aging*
Cell **14**, 605–615.

Lee, K. P., Simpson, S. J., Clissold, F. J., Brooks, R., Ballard, J. W. O., Taylor, P. W.,
Soran, N. and Raubenheimer, D. (2008). Lifespan and reproduction in Drosophila: New
insights from nutritional geometry. *PNAS USA.* **105**, 2498–2503.

Liao, C.-Y., Rikke, B. A., Johnson, T. E., Diaz, V. and Nelson, J. F. (2010). Genetic
variation in the murine lifespan response to dietary restriction: from life extension to life
shortening. *Aging Cell* **9**, 92–95.

Mackay, T., Richards, S., Stone, E. A., Barbadilla, A., Ayroles, D. Z., Zhu, D., Casillas,
S., Han, Y., Magwire, M. M., Cridland, J. M., et al. (2012). The Drosophila melanogaster
Genetic Reference Panel. *Nature* **482**, 173–178.

McCracken, A. W., Adams, G., Hartshorne, L., Tatar, M. and Simons, M. J. P. (2020).
The hidden costs of dietary restriction: Implications for its evolutionary and mechanistic
origins. *Sci. Adv.* **6**, eaay3047.

Metaxakis, A. and Partridge, L. (2013). Dietary restriction extends lifespan in wild-derived
populations of Drosophila melanogaster. *PLoS One* **8**, e74681.

Min, K.-J., Yamamoto, R., Buch, S., Pankratz, M. and Tatar, M. (2008). Drosophila
lifespan control by dietary restriction independent of insulin-like signaling. *Aging Cell* **7**, 199–
206.

Mitchell, S. J., Madrigal-Matute, J., Scheibye-Knudsen, M., Fang, E., Aon, M., González-
Reyes, J. A., Cortassa, S., Kaushik, S., Gonzalez-Freire, M., Patel, B., et al. (2016).
Effects of Sex, Strain, and Energy Intake on Hallmarks of Aging in Mice. *Cell Metab.* **23**,
1093–1112.

Moatt, J. P., Nakagawa, S., Lagisz, M. and Walling, C. A. (2016). The effect of dietary
restriction on reproduction: a meta-analytic perspective. *BMC Evol. Biol.* **16**,.

Piper, M. D. W., Wong, R., Grandison, R. C., Bass, T. M., Martinez, P. M. and Partridge,
L. (2010). Water-independent effects of dietary restriction in Drosophila. *PNAS USA.* **107**,
E54–6; author reply E57.

Regan, J. C., Khericha, M., Dobson, A. J., Bolukbasi, E., Rattanavirotkul, N. and
Partridge, L. (2016). Sex difference in pathology of the ageing gut mediates the greater
response of female lifespan to dietary restriction. *eLife* **5**, e10956.

Rikke, B. A., Liao, C.-Y., McQueen, M. B., Nelson, J. F. and Johnson, T. E. (2010).
Genetic dissection of dietary restriction in mice supports the metabolic efficiency model of
life extension. *Exp. Gerontol.* **45**, 691–701.

Ripatti, S. and Palmgren, J. (2000). Estimation of multivariate frailty models using
penalized partial likelihood. *Biometrics* **56**, 1016–1022.

Skorupa, D. A., Dervisefendic, A., Zwiener, J. and Pletcher, S. D. (2008). Dietary
composition specifies consumption, obesity, and lifespan in Drosophila melanogaster. *Aging*
Cell **7**, 478–490.

Swindell, W. R. (2012). Dietary restriction in rats and mice: a meta-analysis and review of
the evidence for genotype-dependent effects on lifespan. *Ageing Res. Rev.* **11**, 254–270.

Tatar, M. (2011). The plate half-full: status of research on the mechanisms of dietary
restriction in Drosophila melanogaster. *Exp. Gerontol.* **46**, 363–368.

Therneau, T. M., Grambsch, P. M. and Shane Pankratz, V. (2003). Penalized Survival
Models and Frailty. *J. Comput. Graph. Stat.* **12**, 156–175.

van Dam, E., van Leeuwen, L. A. G., Dos Santos, E., James, J., Best, L., Lennicke, C.,
Vincent, A. J., Marinos, G., Foley, A., Buricova, M., et al. (2020). Sugar-Induced Obesity
and Insulin Resistance Are Uncoupled from Shortened Survival in Drosophila. *Cell Metab.*
31, 710–725.

Waithe, D., Rennert, P., Brostow, G. and Piper, M. D. W. (2015). QuantiFly: Robust
Trainable Software for Automated Drosophila Egg Counting. *PLoS One* **10**, e0127659.

Wang, P.-Y., Neretti, N., Whitaker, R., Hosier, S., Chang, C., Lu, D., Rogina, B. and
Helfand, S. L. (2009). Long-lived Indy and calorie restriction interact to extend life span.
PNAS USA. **106**, 9262–9267.

Whitlock, M. C. and Fowler, K. (1999). The changes in genetic and environmental variance
with inbreeding in Drosophila melanogaster. *Genetics* **152**, 345–353.

Wilson, K. A., Nelson, C. S., Beck, J. N., Brem, R. B. and Kapahi, P. (2018). Genome-
wide analysis reveals distinct genetic mechanisms of diet-dependent lifespan and
healthspan in D. melanogaster. *bioRxiv* doi: https://doi.org/10.1101/153791

Wong, A. C.-N., Dobson, A. J. and Douglas, A. E. (2014). Gut microbiota dictates the
metabolic response of Drosophila to diet. *J. Exp. Biol* **217**, 1894–1901.

Figures

Fig.1 Schematic of multiple thresholds in the lifespan reaction norm to diet. Diet concentration
has a bell-shaped relationship with lifespan, ranging from malnutrition (A), DR (B), maximal
performance - or highest Darwinian fitness - at a relatively rich diet (C), to overfeeding, leading to
nutritional toxicity (D). As a detailed reaction norm is rarely known, a dietary dyad although often
used can lead to misleading conclusions. A dietary dyad (A & C) can show no response at all due to
the symmetry in the shape of the reaction norm. Furthermore, genetic or environmental effects can
alter the shape or shift the reaction norm (dashed line), or lead to effects at only specific parts of the
reaction norm (solid gray, e.g. desiccation). For example, diets B & C result in a DR response on the
focal curve, but malnutrition on the dashed curve.

Fig.2. Log hazard ratios of diet and water supplementation in a panel of DGRP genotypes. **A -** Dietary reaction norms vary in a genotype-specific manner. **B -** Water-supplementation, relative to control treatment, rescues desiccation in a diet and genotype-dependent manner. Hazard ratios represent risk to die, therefore higher values indicate shorter lifespans and are relative. Ratios are plotted as coefficients derived from within-line Cox mixed-effect models, with error bars representing 95% confidence intervals. For **A**, 8% yeast treatment was treated as a reference and as such, no CIs are available. Rates here are relative to 8% yeast diet, and lines represent this standard. N = 25,519 females total; 4,800-5,282 per genotype. For **B**, hazard rates are relative to the corresponding control for each diet. Horizontal lines represent a water effect size of 0. N = 12,737 females total; 2,396-2629 per genotype.

Fig. S1. Log hazard ratios of diet within water-treated cages in a panel of DGRP genotypes.

Reaction norms to diet still differ in water-treated circumstances. Hazard ratios represent the inverse

of typical survival reaction norms to diet. 8% yeast treatment was treated as a reference and as such,

no CIs are available. Rates here are relative to 8% yeast diet, and lines represent this standard. N =

12,737 females total; 2,396-2629 per genotype. Hazard ratios have the benefit over median lifespan in

that they are directly related to the appropriate statistics used for time-to-event data. In addition, they

are directly comparable in a quantitative fashion across genotypes of different lifespans, as they

express a relative risk.

Fig. S2. Fecundity analysis of panel under all conditions. Fecundity has a positive relationship with

dietary yeast concentration, except at the highest yeast concentration assayed (14%) for most

genotypes. **A** - raw egg counts. **B** - mortality-corrected counts. Counts generated using QuantiFly

software. Counts are relative, but directly comparable. Flies assayed at age 11-12 days, with boxplots

aggregating totals (median, with the box depicting a quartile each way, and whiskers showing the

range; outliers plotted as dots). Each cage was assayed on 1 scoring day at this age. Mortality

corrected counts (**B**) generated by dividing raw counts, by N flies remaining in cage at the time of

assaying. N = 25,519 females total; 4,800-5,282 per genotype. Note that DGRP-362 experienced

significant mortality at this age under 14% yeast dietary treatment. This is the cause of the

discrepancy in significance between raw, and age-adjusted fecundity counts. Note, egg-laying was not

assessed throughout life and in natural circumstances lifespan of the fly is truncated by extrinsic

factors, e.g. predation. We nonetheless, tentatively conclude that the enhanced mortality and reduced

egg laying on very rich diets is caused by nutritional toxicity.

Fig. S3. Lifetime reproductive fitness estimates of panel under all conditions. Lifetime fitness has

a positive relationship with dietary yeast concentration, except at the highest yeast concentration

assayed (14%) for most genotypes. Mortality-adjusted egg counts from Fig. S2 were multiplied by the

area under the relevant survival curve (restricted mean) to generate lifetime estimates. N = 25,519

females total; 4,800-5,282 per genotype.

Fig. S4. Survival curves of panel in response to diet. Dietary reaction norms vary in a genotype-

specific manner. Survival curves are separated by genotype, and water-supplementation status. N =

25,519 females total; 4,800-5,282 per genotype.

Tables

Table S1. Effect of diet and water supplementation on mortality across 5 DGRP lines (DGRP-195 is reference).

| | Full Model | | | |
coefficient	estimate	exp	s.e.	p
water	-0.390	0.677	0.228	0.086
0.5% yeast	-0.796	0.451	0.228	<0.001
2% yeast	-1.856	0.156	0.206	<0.001
5% yeast	-1.296	0.274	0.225	<0.001
14% yeast	1.043	2.837	0.226	<0.001
217	-0.444	0.641	0.226	0.05
239	0.763	2.145	0.227	0.001
362	0.104	1.110	0.227	0.645
853	-1.090	0.336	0.223	<0.001
0.5% yeast * water	0.470	1.601	0.322	0.144
2% yeast * water	0.030	1.031	0.300	0.92
5% yeast * water	0.277	1.319	0.322	0.39
14% yeast * water	-0.156	0.856	0.322	0.629
217 * 0.5% yeast	1.195	3.302	0.320	<0.001
217 * 2% yeast	0.632	1.881	0.310	0.042
217 * 5% yeast	0.786	2.195	0.321	0.014
217 * 14% yeast	0.643	1.903	0.322	0.046
239 * 0.5% yeast	1.793	6.008	0.321	<0.001
239 * 2% yeast	1.887	6.599	0.308	<0.001
239 * 5% yeast	1.429	4.174	0.321	<0.001
239 * 14% yeast	-0.382	0.683	0.324	0.238
362 * 0.5% yeast	1.754	5.775	0.319	<0.001
362 * 2% yeast	0.262	1.300	0.311	0.399
362 * 5% yeast	-1.322	0.267	0.315	<0.001
362 * 14% yeast	1.880	6.551	0.316	<0.001
853 * 0.5% yeast	3.087	21.901	0.318	<0.001
853 * 2% yeast	2.029	7.604	0.307	<0.001
853 * 5% yeast	0.287	1.332	0.320	0.37
853 * 14% yeast	2.100	8.168	0.320	<0.001
217 * water	0.551	1.736	0.322	0.087
239 * water	-0.101	0.904	0.322	0.754
362 * water	-2.015	0.133	0.319	<0.001
853 * water	-0.101	0.904	0.323	0.753
217 * 0.5% yeast * water	-0.568	0.567	0.455	0.212
217 * 2% yeast * water	-0.114	0.893	0.441	0.797
217 * 5% yeast * water	-0.702	0.496	0.456	0.123
217 * 14% yeast * water	-0.448	0.639	0.456	0.327
239 * 0.5% yeast * water	-0.145	0.865	0.457	0.751
239 * 2% yeast * water	0.126	1.134	0.440	0.775
239 * 5% yeast * water	-0.427	0.652	0.455	0.348
239 * 14% yeast * water	-0.383	0.682	0.457	0.402
362 * 0.5% yeast * water	1.990	7.319	0.456	<0.001
362 * 2% yeast * water	2.068	7.912	0.439	<0.001
362 * 5% yeast * water	2.007	7.444	0.457	<0.001
362 * 14% yeast * water	-1.942	0.143	0.451	<0.001
853 * 0.5% yeast * water	-0.027	0.974	0.458	0.954
853 * 2% yeast * water	0.351	1.420	0.441	0.427
853 * 5% yeast * water	0.433	1.541	0.457	0.344
853 * 14% yeast * water	-2.056	0.128	0.455	<0.001

Table S2. Effect of diet and water supplementation on mortality within DGRP-195.

coefficient	Estimates from individual model				Effect of water, compared to no water		
	estimate	exp	s.e.	p	estimate	exp	p
water supplementation	-0.481	0.618	0.113	<0.001			
0.5% yeast	-0.954	0.385	0.112	<0.001			
2% yeast	-2.139	0.118	0.103	<0.001			
5% yeast	-1.620	0.198	0.112	<0.001			
14% yeast	1.612	5.012	0.112	<0.001			
0.5% yeast * water	0.475	1.608	0.158	0.003	-0.006	0.994	0.971
2% yeast * water	0.156	1.169	0.147	0.289	-0.324	0.723	0.028
5% yeast * water	0.369	1.446	0.160	0.021	-0.112	0.894	0.483
14% yeast * water	-0.301	0.740	0.159	0.059	-0.781	0.458	<0.001

Table S3. Effect of diet and water supplementation on mortality within DGRP-217.

coefficient	Estimates from individual model				Effect of water, compared to no water		
	estimate	exp	s.e.	p	estimate	exp	p
water supplementation	0.190	1.209	0.213	0.373			
0.5% yeast	0.454	1.574	0.211	0.032			
2% yeast	-1.395	0.248	0.208	<0.001			
5% yeast	-0.549	0.578	0.212	0.01			
14% yeast	1.905	6.716	0.200	<0.001			
0.5% yeast * water	-0.108	0.898	0.306	0.724	0.082	1.08	0.789
2% yeast * water	-0.105	0.901	0.306	0.732	0.085	1.08	0.781
5% yeast * water	-0.470	0.625	0.300	0.117	-0.280	0.75	0.350
14% yeast * water	-0.668	0.513	0.306	0.029	-0.479	0.62	0.118

Table S4. Effect of diet and water supplementation on mortality within DGRP-239.

coefficient	Estimates from individual model				Effect of water, compared to no water		
	estimate	exp	s.e.	p	estimate	exp	p
water supplementation	-0.343	0.710	0.105	0.001			
0.5% yeast	0.946	2.575	0.104	<0.001			
2% yeast	0.070	1.072	0.104	0.502			
5% yeast	0.094	1.098	0.103	0.363			
14% yeast	0.572	1.771	0.103	<0.001			
0.5% yeast * water	0.173	1.188	0.148	0.243	-0.170	0.843	0.249
2% yeast * water	0.079	1.082	0.148	0.593	-0.264	0.768	0.075
5% yeast * water	-0.108	0.898	0.147	0.464	-0.451	0.637	0.002
14% yeast * water	-0.476	0.621	0.147	0.001	-0.819	0.441	<0.001

Table S5. Effect of diet and water supplementation on mortality within DGRP-362.

coefficient	Estimates from individual model				Effect of water, compared to no water		
	estimate	exp	s.e.	p	estimate	exp	p
water supplementation	-2.330	0.097	0.363	<0.001			
0.5% yeast	0.763	2.144	0.356	0.032			
2% yeast	-1.520	0.219	0.383	<0.001			
5% yeast	-2.638	0.072	0.351	<0.001			
14% yeast	2.393	10.950	0.339	<0.001			
0.5% yeast * water	2.358	10.571	0.545	<0.001	0.028	1.029	0.958
2% yeast * water	2.016	7.509	0.482	<0.001	-0.314	0.731	0.515
5% yeast * water	2.213	9.145	0.597	<0.001	-0.116	0.890	0.845
14% yeast * water	-1.564	0.209	0.549	0.004	-3.894	0.020	<0.001

Table S6. Effect of diet and water supplementation on mortality within DGRP-853.

	Estimates from individual model				Effect of water, compared to no water		
coefficient	estimate	exp	s.e.	p	estimate	exp	p
water supplementation	-0.473	0.623	0.259	0.067			
0.5% yeast	1.987	7.293	0.239	<0.001			
2% yeast	0.150	1.162	0.254	0.554			
5% yeast	-0.967	0.380	0.248	<0.001			
14% yeast	2.955	19.204	0.223	<0.001			
0.5% yeast * water	0.436	1.547	0.399	0.274	-0.037	0.964	0.926
2% yeast * water	0.379	1.461	0.365	0.299	-0.094	0.910	0.797
5% yeast * water	0.680	1.973	0.373	0.068	0.206	1.229	0.58
14% yeast * water	-2.083	0.125	0.339	<0.001	-2.556	0.078	<0.001

Table S7. Effect of diet on mortality across 5 water-supplemented DGRP lines (DGRP-195 is reference).

	Full Model			
coefficient	estimate	exp	s.e.	p
0.5% yeast	-0.345	0.709	0.228	0.131
2% yeast	-1.921	0.146	0.202	<0.001
5% yeast	-1.060	0.347	0.224	<0.001
14% yeast	0.930	2.535	0.225	<0.001
217	0.107	1.113	0.228	0.638
239	0.683	1.980	0.228	0.003
362	-1.991	0.137	0.218	<0.001
853	-1.242	0.289	0.221	<0.001
217 * 0.5% yeast	0.658	1.931	0.320	0.04
217 * 2% yeast	0.567	1.764	0.311	0.068
217 * 5% yeast	0.096	1.101	0.321	0.764
217 * 14% yeast	0.206	1.229	0.324	0.524
239 * 0.5% yeast	1.771	5.877	0.318	<0.001
239 * 2% yeast	2.119	8.326	0.305	<0.001
239 * 5% yeast	1.042	2.834	0.321	0.001
239 * 14% yeast	-0.799	0.450	0.322	0.013
362 * 0.5% yeast	3.937	51.246	0.315	<0.001
362 * 2% yeast	2.434	11.404	0.307	<0.001
362 * 5% yeast	0.691	1.995	0.318	0.03
362 * 14% yeast	-0.071	0.932	0.323	0.826
853 * 0.5% yeast	3.228	25.234	0.312	<0.001
853 * 2% yeast	2.482	11.960	0.303	<0.001
853 * 5% yeast	0.739	2.094	0.322	0.022
853 * 14% yeast	0.033	1.033	0.319	0.918

Table S8. Effect of diet on mortality within water-supplemented DGRP-195.

	Estimates from individual model			
coefficient	estimate	exp	s.e.	p
0.5% yeast	-0.512	0.599	0.098	<0.001
2% yeast	-2.124	0.120	0.100	<0.001
5% yeast	-1.322	0.267	0.102	<0.001
14% yeast	1.410	4.095	0.100	<0.001

Table S9. Effect of diet on mortality within water-supplemented DGRP-217.

| | Estimates from individual model | | | |
coefficient	estimate	exp	s.e.	p
0.5% yeast	0.357	1.429	0.214	0.095
2% yeast	-1.487	0.226	0.215	<0.001
5% yeast	-1.011	0.364	0.216	<0.001
14% yeast	1.260	3.527	0.215	<0.001

Table S10. Effect of diet on mortality within water-supplemented DGRP-239.

| | Estimates from individual model | | | |
coefficient	estimate	exp	s.e.	p
0.5% yeast	1.114	3.046	0.119	<0.001
2% yeast	0.150	1.162	0.118	0.204
5% yeast	-0.012	0.988	0.117	0.919
14% yeast	0.096	1.101	0.118	0.413

Table S11. Effect of diet on mortality within water-supplemented DGRP-362.

| | Estimates from individual model | | | |
coefficient	estimate	exp	s.e.	p
0.5% yeast	3.631	37.764	0.386	<0.001
2% yeast	0.518	1.678	0.379	0.172
5% yeast	-0.487	0.614	0.380	0.2
14% yeast	0.920	2.509	0.379	0.015

Table S12. Effect of diet on mortality within water-supplemented DGRP-853.

| | Estimates from individual model | | | |
coefficient	estimate	exp	s.e.	p
0.5% yeast	2.811	16.623	0.222	<0.001
2% yeast	0.564	1.759	0.216	0.009
5% yeast	-0.284	0.753	0.216	0.19
14% yeast	0.923	2.518	0.216	<0.001

Table S13. Effect of diet and water supplementation on fecundity across 5 DGRP lines, derived from linear model estimates oflog-transformed raw fecundity counts (DGRP-195 is reference). Counts generated using QuantiFly software. Counts are relative, but directly comparable.

coefficient	Full Model			Compared to 8%	
	estimate	s.e.	p	estimate	p
intercept	2.731	0.067	<0.001		
water	0.039	0.027	0.15		
0.5% yeast	-1.374	0.092	<0.001		
2% yeast	-0.595	0.092	<0.001		
5% yeast	-0.367	0.092	<0.001		
14% yeast	0.157	0.092	0.09	2.888	<0.001
217	-0.044	0.092	0.634		
239	-0.331	0.095	0.001		
362	-0.294	0.092	0.002		
853	-0.221	0.095	0.021		
217 * 0.5% yeast	0.113	0.130	0.387		
217 * 2% yeast	-0.268	0.134	0.048		
217 * 5% yeast	-0.164	0.132	0.216		
217 * 14% yeast	-0.224	0.132	0.092	-0.110	0.404
239 * 0.5% yeast	0.509	0.139	<0.001		
239 * 2% yeast	0.310	0.139	0.027		
239 * 5% yeast	-0.119	0.134	0.373		
239 * 14% yeast	-0.262	0.136	0.056	-0.436	0.001
362 * 0.5% yeast	0.524	0.130	<0.001		
362 * 2% yeast	0.212	0.130	0.106		
362 * 5% yeast	0.180	0.130	0.168		
362 * 14% yeast	-0.202	0.130	0.122	-0.339	0.009
853 * 0.5% yeast	0.249	0.134	0.065		
853 * 2% yeast	-0.233	0.132	0.079		
853 * 5% yeast	-0.137	0.132	0.3		
853 * 14% yeast	-0.657	0.134	<0.001	-0.721	<0.001

Table S14. Effect of diet and water supplementation on fecundity across 5 DGRP lines, derived from linear model estimates oflog-transformed mortality-adjusted fecundity counts (DGRP-195 is reference). Counts generated using QuantiFly software. Counts are relative, but directly comparable.

coefficient	Full Model			Compared to 8%	
	estimate	s.e.	p	estimate	p
intercept	0.776	0.067	<0.001		
water	-0.004	0.027	0.873		
0.5% yeast	-1.379	0.093	<0.001		
2% yeast	-0.602	0.093	<0.001		
5% yeast	-0.384	0.093	<0.001		
14% yeast	0.148	0.093	0.114	0.925	<0.001
217	-0.050	0.093	0.594		
239	-0.350	0.096	<0.001		
362	-0.239	0.093	0.011		
853	-0.241	0.096	0.013		
217 * 0.5% yeast	0.101	0.132	0.444		
217 * 2% yeast	-0.301	0.136	0.028		
217 * 5% yeast	-0.113	0.134	0.398		
217 * 14% yeast	-0.165	0.134	0.218	-0.067	0.617
239 * 0.5% yeast	0.542	0.141	<0.001		
239 * 2% yeast	0.311	0.141	0.028		
239 * 5% yeast	-0.120	0.136	0.378		
239 * 14% yeast	-0.237	0.138	0.087	-0.439	0.001
362 * 0.5% yeast	0.485	0.132	<0.001		
362 * 2% yeast	0.177	0.132	0.182		
362 * 5% yeast	0.186	0.132	0.16		
362 * 14% yeast	0.166	0.132	0.209	0.075	0.568
853 * 0.5% yeast	0.284	0.136	0.038		
853 * 2% yeast	-0.227	0.134	0.091		
853 * 5% yeast	-0.111	0.134	0.406		
853 * 14% yeast	-0.550	0.136	<0.001	-0.643	<0.001