1

Title: Expected impacts of climate change on tree ferns distribution and diversity patterns in subtropical Atlantic Forest

Running title: Tree ferns distribution and diversity in future scenarios

André Luís de Gasper[a,b,c], Guilherme Salgado Grittz[b], Carlos Henrique Russi[a], Carlos Eduardo Schwartz[c], Arthur Vinicius Rodrigues[d]

[a] Regional University of Blumenau, Rua Antônio da Veiga, 140 - Itoupava Seca, 89030-903 - Blumenau - SC – Brasil. +55 47 98446-5810. algasper@furb.br.
[b] Postgraduate program in Biodiversity, Regional University of Blumenau, Rua Antônio da Veiga, 140 - Itoupava Seca, 89030-000 - Blumenau - SC – Brasil.
[c] Postgraduate program in Forestry Engineering, Regional University of Blumenau, Rua São Paulo, 3360 - Itoupava Seca, 89030-903 - Blumenau - SC – Brasil.
[d] Postgraduate program in Ecology, Federal University of Rio Grande do Sul, Av. Bento Gonçalves, 9500 - Agronomia - Porto Alegre - RS - Brasil.

Corresponding author: André Luís de Gasper (algasper@furb.br)

Acknowledgments: The authors are thankful to Fundação de Amparo à Pesquisa e Inovação de Santa Catarina (FAPESC) for supporting IFFSC and for Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001, for postgraduate research grants. We are also grateful for FUMDES (SC) and PIPE (SC, FURB) for funding research at the undergraduate level. Finally, we thank Raphael A. Saavedra for reviewing the manuscript. The authors declare that they have no conflict of interest.

ABSTRACT
Tree ferns are common elements in subtropical Atlantic Forest, sometimes reaching more than half of dominance at forest sites. Climate change could impact the distribution and diversity of tree ferns, hence impacting the ecological processes provided by them. We investigate the impacts of future climate changes in the potential distribution of tree fern species, as well as in the α- and β-diversity. Our first hypothesis (H_1) is that warmer climate in the future will increase the distribution of Cyatheaceae species and decrease the distribution of Dicksoniaceae species. Due to a larger number of Cyatheaceae than Dicksoniaceae, the richness will tend to increase. However, the increase in richness might lead to homogenization, consequently decreasing β-diversity (H_2). To test these hypotheses, we used distribution models to predict the potential species distribution in the present and future. Then, we calculated the changes in potential distribution areas, α-, and β-diversity components between scenarios. Finally, we assessed the impact of these changes within Protected Areas. We found that Dicksoniaceae distribution tends to shrink in the future, while half of Cyatheaceae tend to lose distribution. Species richness tends to decrease in the future, as well as β-diversity. At least 43% of the sites tend to have their species richness reduced, while only 26% of sites tend to gain species. Our results suggest that species associated with cold environments will lose suitable areas and are more threatened. Richness tends to decrease mainly in sites with high precipitation seasonality. Furthermore, the tree ferns assemblage tends to homogenize.
Keywords: Beta-diversity, climate change, Cyatheaceae, Dicksoniaceae, species distribution modelling, tree ferns.

1. INTRODUCTION

Tree ferns are expressive elements in (sub)tropical forest formations (Tryon & Tryon, 1982), sometimes establishing monodominant forests (Gasper et al., 2011). For this reason, tree ferns act in the dynamics of the ecosystem and may affect the regeneration of wood species and nutrient cycling (Brock, Perry, Lee & Burns, 2016). In addition, they contribute to the process of ecological succession (Arens & Baracaldo, 1998), to biomass stock in tropical forests (Sarmiento, Pinillos, & Garay, 2005), and provide microhabitat for several epiphytic plants, many of them occurring exclusively on tree ferns caudices (Wagner, Mendieta-Leiva, & Zotz, 2015).

Tree ferns have suffered intense exploitation in the tropical forest due to ornamental use of their caudices (Eleutério & Pérez-Salicrup, 2006; Hoshizaki, & Moran, 2001), causing populational exhaustion of many species (Santiago, Mynssen, Maurenza, Penedo, & Sfair, 2013). Combined with this, the actual high forest fragmentation in the Atlantic Forest (Ribeiro, Metzger, Martensen, Ponzoni, & Hirota, 2009) plus climate change scenarios (IPCC, 2014; Lima, Ribeiro, Grelle, & Pinto, 2019) are other potential sources of threat to tree ferns. These threats might change the density and distribution of these species as well as the locations of the suitable area to grow and reproduce.

Moreover, despite tree ferns being an important group in forest structure, these plants are historically neglected in floristic and ecological studies in Brazil (Weigand & Lehnert, 2016). There are two main families of tree ferns in subtropical Atlantic Forest: Dicksoniaceae and Cyatheaceae. The former is represented by *Dicksonia sellowiana*, a species that inhabits high and cold environments (Gasper et al., 2011), and *Lophosoria*

quadripinnata, that grows in ravines, doing best on moist, well-drained soil and in full
sun until 2000 m in eastern Brazil (Lehnert & Kessler, 2018). The later family (about 14
species) exhibits a preference for warm, humid and low seasonal climates (Bystriakova,
Schneider, & Coomes, 2011) and may benefit from warmer climate regimes.
The Intergovernmental Panel on Climate Change (IPCC) predicts shifts in
rainfall and temperature in the subtropical Atlantic Forest, and since water availability is
correlated to fern species richness (Kessler, Kluge, Hemp, & Ohlemüller, 2011), rainfall
regimes modifications could impact in ferns distribution. Therefore, a reduction of cold
environments and increasing of warm and humid environments may impact *D.*
sellowiana, an already endangered species, through the restriction of its occurrence area.
In this regard, our study sought to predict the impacts of future climate changes
in α- and β-diversity of tree ferns in the subtropical Atlantic Forest, as well as to predict
the impact in the potential distribution of each species. Our first hypothesis (H_1) is that
species from both families will change their potential distribution areas. We expect
Dicksoniaceae species will have their potential distribution area reduced (specially
Dicksonia sellowiana) because of their association with cold habitats — and
Cyatheaceae species will increase their potential distribution areas since they generally
occur along hot and humid regions. Our second hypothesis (H_2) is that the changes in
species distribution will affect α- and β-diversity. Since we expect the Cyatheaceae
species will increase their distribution range, hence increasing the overlap in species
areas, we expect an increase in α-diversity and a decrease in β-diversity, i.e., less
variation in species composition among sites, leading to a homogenization of our study
region.
105
106
107 **2. MATERIAL AND METHODS**
108
2.1 Study area
The study area is delimited by the subtropical Atlantic Forest, floristically distinct from
the tropical Atlantic Forest (Eisenlohr & Oliveira-Filho, 2014). The subtropical Atlantic
Forest occurs in southern Brazil as well as in parts of Argentina and Paraguay (C.
Galindo-Leal & Câmara, 2005). The predominant climate type is Cfa (temperate humid
with hot summer), with some areas fluctuating to Cfb (temperate humid with temperate
summer). The relief ranges from sea level to altitudes near 1200 m, including peaks that
reach almost 1900 m. Distinct forest types can be found, which includes Restinga, on
the coastal areas; Rainforests, in low altitudes at the coastal region (< 800–900 m);
Mixed Forests (*Araucaria* forest), generally in areas with altitudes over 800 m; and
Semideciduous Forests in low areas with high seasonality in precipitation (Figure 1).
120
2.2 Data gathering
Among the species that occur in subtropical Atlantic Forest (following Brazilian
Flora 2020 in construction 2019; Fernandes 1997; Lehnert & Weigand, 2016), we
identified 1167 records and 15 species from the literature (Fernandes, 1997; Vibrans,
Sevegnani, Lingner, Gasper, & Sabbagh, 2010) and herbaria (using SpeciesLink –
http://splink.cria.org and GBIF Global Biodiversity Information Facility –
https://www.gbif.org). Uncertain records were removed, as well as *Cyathea uleana* due
to its low number of registers.
The data available for the study area was under the effect of sampling biases.
This issue arose because, in the state of Santa Catarina, the Floristic and Forest

Inventory of Santa Catarina (IFFSC; Vibrans et al., 2010) had strongly sampled the
state's territory — an effort that did not take place in other locations of the study area.
To minimize sample biases, we performed spatial filtering on the occurrence data by
defining a buffer of $\cong$ 30 km as a minimum distance between occurrence records in all
study areas (Fourcade, Engler, Rödder, & Secondi, 2014).
Climate data were obtained for the present and future climate change scenarios
from WorldClim v2.0 (Fick & Hijmans, 2017) and WorldClim v1.4 (Hijmans,
Cameron, Parra, Jones, & Jarvis, 2005), respectively. Data for the future scenario refers
to the year 2050 and was based on the climate projections made by IPCC. We used the
mean of four Earth System Models (ESM) for future modeling: NorESM1-M (Bentsen
et al., 2013), MIROC-ESM (Watanabe et al., 2011), HadGEM2-ES (Jones et al., 2011),
and CNRM-CM5 (Voldoire et al., 2013). For these models, two future scenarios were
selected: the most optimistic (RCP 2.6) and the most pessimistic (RCP 8.5), which were
elaborated and named accordingly to the concentration of carbon dioxide in the
atmosphere (W/m^2) between the pre-industrial period and the year 2100 (IPCC, 2014).
In all tested ESMs we chose 19 environmental variables (bioclim) with 2.5' resolution.
We used the Variance Inflation Factor (VIF) via the *vifstep* function on *usdm* R-package
(Naimi, Hamm, Groen, Skidmore, & Toxopeus, 2014; R Core Team 2015) to remove
collinearity. This function generates 5,000 random points across the climate layers and
calculates the VIF for each variable using a step-by-step procedure. At each step, the
layer with the highest VIF is removed from the set. The process is repeated until only
variables with a VIF value below a defined threshold remain. We used a restrictive VIF
value (VIF = 5) that reduced the 19 climate variables to six: Annual Mean Temperature
(BIO01), Isothermality (BIO03), Temperature Annual Range (BIO07), Mean
Temperature of Driest Quarter (BIO09), Precipitation of Wettest Month (BIO13) and
Precipitation of Coldest Quarter (BIO19).
157

158 **2.3 Species Distribution Models (SDMs)**
159
We used SDMs to predict current and future distributions of each species. Our
SDMs were formulated using the *sdm* R-package (Naimi & Araújo, 2016) through five
different algorithms: MaxEnt (Phillips, Dudík, & Schapire, 2004), Boosted Regression
Trees (BRT; Friedman, 2001), Generalized Linear Models (GLM; McCullagh &
Nelder, 1989), Multivariate Adaptive Regression Spline (MARS; Friedman, 1991), and
Support Vector Machine (SVM; Vapnik, 2013), all with default settings. Moreover, as
proposed by Barbet-Massin, Jiguet, Albert, & Thuiller (2012), we applied 10 000
pseudo-absences for each species randomly created across the study area.
We evaluated the performance of the models generated by each algorithm
through a cross-validation procedure with 10 repetitions, where data were randomly
divided into training, amounting to 70% of the data, and testing, comprising the
remaining 30%. Two statistical methods were responsible for evaluating the robustness
of the models: Area Under the Curve (AUC) of the receiver operating characteristic
curve (ROC; Fielding & Bell, 1997) — a threshold-independent metric — and True
Skill Statistic (TSS; Allouche, Tsoar, & Kadmon, 2006) — a threshold-dependent
metric. All generated models were above the previously established cutoff values (AUC
> 0.70 and TSS > 0.40).
To avoid biases created by choosing a single statistical algorithm, as pointed out
by several studies (e.g., Araújo & New, 2007; Grenouillet, Buisson, Casajus, & Lek,

2011), we built a single final model through an ensemble approach. In this sense, the
models generated by statistical algorithms were projected using current and future
climate, and the ensemble model for each scenario was considered as the mean
weighted by the AUC value of the projections of each algorithm. Binary maps
(presence-absence) were made based on the threshold that maximizes the sensitivity
plus specificity of the models (maxSSS; Liu, Newell, & White, 2016), calculated with
the R-package *SDMtools* (VanDerWal, Falconi, Januchowski, Shoo, & Storlie, 2014).
The threshold was calculated for each species based on the final ensemble map of the
current scenario. Likewise, the resulting threshold value for the current scenario was
also used to build binary maps for future scenarios.

2.4 Species distribution and diversity patterns
191
To understand how climate change will affect species distribution and diversity
patterns, the study area was divided into hexagonal cells of $\cong$ 200 km². Based on SDM
predictions for tree ferns, we measured for each scenario the changes of suitable cells
for all species and the α- and β-diversity indexes. We also evaluate the same metrics
exclusively in grid cells that overlap Protected Areas (PAs) in the study region. We
obtained the boundaries of PAs from the World Database on Protected Areas (UNEP-
WCMC, 2019). Therefore, we can compare species distribution and diversity indexes
between protected and non-protected areas for any specific scenario.
We measured α-diversity as the number of species present in each cell, i.e., the
predicted species richness. We calculated the β-diversity as the total variance of the
community (Total β-diversity, or BD_{TOTAL}; *sensu* Legendre & De Cáceres, 2013) and it
was subsequently decomposed in Local Contributions to β-Diversity (LCBD). LCBD
represents the ecological uniqueness of each hexagonal cell regarding species
composition. We calculated BD_{TOTAL} based on the Jaccard dissimilarity coefficient (1–
similarity). Then, the LCBD was determined based on the partition of BD_{TOTAL} between
the cells (Legendre & De Cáceres, 2013). The significance of the LCBD values for each
cell was obtained through 999 permutations, where the species are distributed randomly
and independently along the grid and the LCBD values are calculated for each random
distribution. Significant LCBD values were those with p-value < 0.05.
Finally, we followed Legendre (2014) by partitioning the BD_{TOTAL} in species
replacement and richness difference components, using the replacement ($Repl_J$) and
richness difference ($RichDiff_J$) indexes of the Podani family (*sensu* Podani & Schmera,
2011; Legendre, 2014). All β-diversity metrics were calculated with the *adespatial* R-
package (Dray et al., 2019) using the functions *beta.div* and *beta.div.comp*. We used
QGIS 3.8 (QGIS Development Team 2019) to generate the maps. Lastly, to test
differences in α-diversity between current and future scenarios, we conducted the
Wilcoxon Rank-Sum Test, a non-parametric unpaired test, for the current scenario ×
RCP 2.6 scenario and current scenario × RCP 8.5 scenario.

220
221   **3. Results**
222
All models had a good predictive power (AUC: mean = 0.825, sd = 0.05; TSS:
mean = 0.623, sd = 0.14 — for individual distribution maps and statistical performance

of each SDM see the supporting information S1, and the interactive maps in
https://avrodrigues.shinyapps.io/tferns/). Overall, the predicted impacts of climate
change on species were similar in both RCP 2.6 and RCP 8.5 (Table 1), where
Dicksoniaceae species tend to lose area of distribution and half of Cyatheaceae species
tend to increase its area of occurrence — while the other half tends to decrease. Also,
currently suitable areas inside PAs also suffer a significant reduction for many tree
ferns.

232
233 **3.1 α-diversity**

234
The highest α-diversity values (current and future scenarios, Figure 2) were
found in the Rainforest areas. In the west, where α-diversity is low in the current
scenario, many sites tend to become unsuitable for all the species in the future.
Furthermore, PAs harbor a higher average of α-diversity when compared to the
subtropical Atlantic Forest. At the same time, PAs will lose fewer species than non-PAs
and gain more species than non-PAs regions (Table 2).

241
242 **3.2 β-diversity**

243
The BD_{TOTAL} is predicted to decrease with climate change (Table 3), indicating
community homogenization along the study area. The LCBD values (Figure 3) ranged
from 3.2×10^{-4} to 5.5×10^{-4} in the current scenario, 2.7×10^{-4} to 5.8×10^{-4} in the RCP
2.6 scenario and 2.5×10^{-4} to 6.0×10^{-4} in the RCP 8.5 scenario, indicating an increase
in LCBD at some locations in the future and a decrease in others. Considering only PAs,
community homogenization is like the rest of the study area (Table 3).

250
251
252 **4. DISCUSSION**

253
254 **4.1 Changes in species distribution**

255
Our first hypothesis was that the species potential distribution would change in
future climate scenarios — where Dicksoniaceae species would have its distribution
reduced and Cyatheaceae species would have its distribution increased. We partially
confirmed our expectation since Dicksoniaceae species indeed had their potential
distribution reduced in both future scenarios, but only half of Cyatheaceae species had
their potential distribution increased. *Cyathea phalerata*, a common species in southern
Brazil, had its potential area reduced in both scenarios, and *Alsophila setosa* — one of
the most common tree ferns in forest communities in subtropical Atlantic Forest
(Lingner et al., 2015) — increased its area of occurrence in scenario RCP 2.6 but
decreased in RCP 8.5. *Cyathea feeana*, recently segregated from *C. corcovadensis*
(Lehnert & Weigand, 2013), had the most reduced potential occurrence area. All
Cyatheaceae species that were predicted to lose potential area in the future by our model
showed climatic suitability for the coldest regions of the study area at the current
scenario (Figures S1-S14 and interactive maps in
https://avrodrigues.shinyapps.io/tferns/).
A global analysis of Cyatheaceae distribution patterns showed a clear preference
of the group for hotter and wetter locations with low seasonality, although some species
show a capacity to occupy relatively cold regions, rarely where minimum temperatures

drop below freezing (Bystriakova et al., 2011). Half of the Cyatheaceae species
analyzed here were able to occupy the coldest areas of the current Mixed Forest.
However, few Cyatheaceae species were able to occupy areas of Semideciduous Forest,
where there is higher seasonality in precipitation than areas of Rainforest and eastern
Mixed Forest (Oliveira-Filho, Budke, Jarenkow, Eisenlohr, & Neves, 2015; Oliveira-
Filho & Fontes, 2000). The effects of high temperatures, reducing the cold intensity
(Wilson, Walters, Mayle, Lingner, & Vibrans, 2019), set up favorable conditions for
some of Cyatheaceae, since they expanded to it.
Dicksonia sellowiana, a species able to withstand colder environments (Gasper et
al., 2011), was restricted to higher and colder locations, wherein the Mixed Forest
predominates. In the same way, *L. quadripinnata* seems able to inhabit the colder
regions of the Mixed Forest. In addition, the warmer areas of the Rainforest also
showed high climatic suitability for this species. In fact, *L. quadripinnata* has been
recorded in both temperate (Ricci, 1996) and tropical (Bernabe et al., 1999) Rainforests.
In this sense, precipitation rather than temperature appears to be the main determinant of
L. quadripinnata occurrence.
290
291 **4.2 Changes in diversity patterns**
292
The second hypothesis was that the different responses of Dicksoniaceae and
Cyatheaceae species should cause significant changes in diversity patterns. Our
expectation was that species richness (α-diversity) would increase, following the
increase in overlapping distributions of Cyateaceae species, which in turn would lead to
a decrease of β-diversity. Our expectation about α-diversity was not confirmed since we
found evidence for less species richness per cell in the future scenarios. Nevertheless,
we found evidence for future homogenization of cells in the subtropical Atlantic Forest,
since both BD_{TOTAL} and LCBD values tend to decrease in future scenarios.
Our estimates indicate that at least 40% of the sites (cells) will lose richness in
the subtropical Atlantic Forest. This is due to a reduction in the area of distributions for
Cyatheaceae species, against our expectations. This loss can affect the conservation of
other biological groups that depend on tree ferns caudices, such as epiphytes — some of
these growths exclusively in tree ferns (Mehltreter, 2008). These changes may be
associated with variations in precipitation since water availability seems to be an
important species richness predictor for ferns, as pointed out by several authors
(Aldasoro, Cabezas, & Aedo, 2004; Gasper, Eisenlohr, & Salino, 2015; Kessler et al.,
2011). The current low α-diversity of tree ferns observed in the midwest and far west, as
well as the reduction of α-diversity in the future scenarios, may be caused by the rainfall
regime and high climatic seasonality (Bystriakova et al., 2011; Cabré, Solman, &
Núñez, 2016).
In the study region, the BD_{TOTAL} is more influenced by *RichDiff$_J$* among the
sites than by *Repl$_J$* (Figure 3 and Table 3). Together, these results indicate that local
extinctions are important drivers of homogenization in species composition of tree
ferns. The high influence of *RichDiff$_J$* in β-diversity could be explained by the
environmental filtering process (Atmar & Patterson, 1993). This process is evidenced
by the loss of species in the west region, with high precipitation seasonality, and by the
loss of more suitable areas by species from a colder climate. Working with ferns
communities in the northern Atlantic Forest, da Costa, Arnan, de Paiva Farias, & Barros
(2019) found a higher contribution of species replacement than of species differences in
ferns on a regional and local scale in tropical Atlantic Forest. The authors listed high

dispersal ability, niche preferences, and the heterogeneous structure of the Atlantic forest as the three main factors that contribute to species replacement. Two differences between the studies could explain the different results: i) our study used local assemblage in coarse scale, provided by the stack of SDMs, which is not able to capture differences in microhabitats as in da Costa et al. (2019), and ii) we focused on the tree ferns niches rather than all ferns species in the subtropical Atlantic Forest.

4.3 Implications for biodiversity conservation agenda

Our results provide relevant insights to the conservation of tree ferns by predicting which species will lose larger proportions of adequate habitat in the future (Table 1). We note that this phenomenon will mainly affect species that occupy higher areas such as *A. capensis, C. feeana, C. villosa, S. gardneri, L. quadripinnata*, and *D. sellowiana*. Although extinctions did not occur in our study area, historically endangered species such as *D. sellowiana* had their potential distribution area greatly diminished (–42%). Also, the climate changes can alter the threatened status from endangered (Santiago et al. 2013) to critically endangered (considering IUCN 2012 criteria of population size reduction: A1cB1b). *C. feeana* and *S. gardneri* (currently present in 424 and 386 cells, respectively), both endemic to eastern Brazil, seem to be the most threatened species in the future considering the reduction in cell numbers (to 208 in RCP 2.6 and 95 in RCP 8.5 for *C. feeana;* and to 217 in RCP 2.6 and 134 cells in RCP 8.5 for *S. gardneri*). *Cyathea feeana* occurs from 500 to 1750 m and was recently segregated from *C. corcovadensis* (Lehnert & Weigand, 2013), which may bias our results since not all herbaria (despite our best efforts) review these records. *Sphaeropteris gardneri* occurs between 450–600 m and few individuals were found by the IFFSC project (Gasper et al., 2018).

More than half of the studied species are predicted to lose suitable areas, even inside PAs (Table 2). So, in order to safeguard these species, new PAs will be needed as they play an essential role in protecting species *in situ* (Chape, Harrison, Spalding, & Lysenko, 2005). However, these new PAs may too have reduced long-term effectiveness in species conservation since they are also going to be impacted by climate change. Then, to optimize cost-benefit analysis of implementing new PAs, lawmakers and specialists should always consider species conservation now and in the future (Araújo, Alagador, Cabeza, Nogués-Bravo, & Thuiller, 2011). Also, LCBD values should be interpreted with caution when looking at alternative indexes for conservation use since they are not always positively correlated to species richness (Legendre & De Cáceres, 2013; Silva & Hernández, 2014).

Unfortunately, some PAs in higher altitudes, such as Parque Nacional de São Joaquim — where *D. sellowiana*, as well as other threatened species (such as *Araucaria angustifolia*; Wilson et al., 2019) may find suitable areas in the future — are threatened to be downsized (see the workgroup created to study the protected area boundaries; ICMBio 2019).

At last, we expected that some tree fern species could expand their distributions as a result of climate change. However, we must not extrapolate, without deep analysis, this expansion to other groups. It is well known that animals (Radchuk et al., 2019) and plants (Corlett & Westcott, 2013) will not be able to keep up, in general, with the changes affecting the planet.

REFERENCES

Aldasoro, J. J., Cabezas, F., & Aedo, C. (2004). Diversity and distribution of ferns in sub-Saharan Africa, Madagascar and some islands of the South Atlantic. *Journal of Biogeography*, *31*(10), 1579–1604. https://doi.org/10.1111/j.1365-2699.2004.01106.x

Allouche, O., Tsoar, A., & Kadmon, R. (2006). Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). *Journal of Applied Ecology*, *43*(6), 1223–1232. https://doi.org/10.1111/j.1365-2664.2006.01214.x.

Araújo, M. B., Alagador, D., Cabeza, M., Nogués-Bravo, D., & Thuiller, W. (2011). Climate change threatens European conservation areas. *Ecology Letters*, *14*(5), 484–492. https://doi.org/10.1111/j.1461-0248.2011.01610.x

Arens, N. C., & Baracaldo, P. S. (1998). Distribution of Tree Ferns (Cyatheaceae) across the Successional Mosaic in an Andean Cloud Forest, Nariño, Colombia. *American Fern Journal*, *88*, 60–71. https://doi.org/10.2307/1547225

Barbet-Massin, M., Jiguet, F., Albert, C. H., & Thuiller, W. (2012). Selecting pseudo-absences for species distribution models: How, where and how many? *Methods in Ecology and Evolution*, *3*(2), 327–338. https://doi.org/10.1111/j.2041-210X.2011.00172.x

Bentsen, M., Bethke, I., Debernard, J. B., Iversen, T., Kirkevåg, A., Seland, Ø., … Kristjánsson, J. E. (2013). The Norwegian Earth System Model, NorESM1-M – Part 1: Description and basic evaluation of the physical climate. *Geoscientific Model Development*, *6*(3), 687–720. https://doi.org/10.5194/gmd-6-687-2013

Bernabe, N., Williams-Linera, G., & Palacios-Rios, M. (1999). Tree Ferns in the Interior and at the Edge of a Mexican Cloud Forest Remnant: Spore Germination and Sporophyte Survival and Establishment. *Biotropica*, *31*(1), 83. https://doi.org/10.2307/2663961

Brazilian Flora 2020 in construction. (2019). Retrieved July 16, 2019, from http://floradobrasil.jbrj.gov.br

Brock, J. M. R., Perry, G. L. W., Lee, W. G., & Burns, B. R. (2016). Tree fern ecology in New Zealand: A model for southern temperate rainforests. *Forest Ecology and Management*, *375*, 112–126. https://doi.org/10.1016/j.foreco.2016.05.030

Bystriakova, N., Schneider, H., & Coomes, D. (2011). Evolution of the climatic niche in scaly tree ferns (Cyatheaceae, Polypodiopsida). *Botanical Journal of the Linnean Society*, *165*(1), 1–19. https://doi.org/10.1111/j.1095-8339.2010.01092.x

Cabré, M. F., Solman, S., & Núñez, M. (2016). Regional climate change scenarios over southern South America for future climate (2080-2099) using the MM5 Model. Mean, interannual variability and uncertainties. *Atmósfera*, *29*(1), 35–60. https://doi.org/10.20937/ATM.2016.29.01.04

Chape, S., Harrison, J., Spalding, M., & Lysenko, I. (2005). Measuring the extent and effectiveness of protected areas as an indicator for meeting global biodiversity targets. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *360*(1454), 443–455. https://doi.org/10.1098/rstb.2004.1592

Corlett, R. T., & Westcott, D. A. (2013). Will plant movements keep up with climate change? *Trends in Ecology and Evolution*, *28*(8), 482–488. https://doi.org/10.1016/j.tree.2013.04.003

Costa, L. E. N., Arnan, X., Farias, R. P. et al. (2019). Community responses to fine-scale environmental conditions: Ferns alpha and beta diversity along Brazilian Atlantic forest remnants. Acta Oecologica, 101: 103475.

421   https://doi.org/10.1016/j.actao.2019.103475

422 Dray, S., Bauman, D., Blanchet, G., Borcard, D., Clappe, S., Guénard, G., … Wagner,
423   H. H. (2019). adespatial: Multivariate multiscale spatial analysis. R package
424   version 0.3–7. Retrieved from https://cran.r-project.org/package=adespatial

425 Eisenlohr, P. V., & Oliveira-Filho, A. T. (2014). Tree species composition in areas of
426   Atlantic Forest in southeastern Brazil is consistent with a new system for
427   classifying the vegetation of South America. *Acta Botanica Brasilica*, *28*(2), 227–
428   233. https://doi.org/10.1590/S0102-33062014000200009

429 Eleutério, A. A., & Pérez-Salicrup, D. (2006). Management of tree ferns (Cyathea spp.)
430   for handicraft production in Cuetzalan, Mexico. *Economic Botany*, *60*(2), 182–186.
431   Retrieved from https://www.jstor.org/stable/4257089

432 Fernandes, I. (1997). *Taxonomia e fitogeografia de Cyatheaceae e Dicksoniaceae nas*
433   *regiões Sul e Sudeste do Brasil*. Universidade de São Paulo.

434 Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate
435   surfaces for global land areas. *International Journal of Climatology*, *37*(12).
436   https://doi.org/10.1002/joc.5086

437 Fourcade, Y., Engler, J. O., Rödder, D., & Secondi, J. (2014). Mapping species
438   distributions with MAXENT using a geographically biased sample of presence
439   data: A performance assessment of methods for correcting sampling bias. *PLoS*
440   *ONE*, *9*(5), 1–13. https://doi.org/10.1371/journal.pone.0097122

441 Friedman, J. H. (1991). Multivariate adaptive regression splines. *The Annals of*
442   *Statistics*, *19*(1), 1–67. Retrieved from https://www.jstor.org/stable/2241837

443 Friedman, J. H. (2001). Greedy function approximation: a gradient boosting machine.
444   *The Annals of Statistics*, *29*(5), 1189–1232.

445 Galindo-Leal, C., & Câmara, I. G. (2005). *Mata Atlântica⬚: biodiversidade, ameaças e*
446   *perspectivas*. (Carlos Galindo-Leal & I. de Gusmão Câmara, Eds.). Fundação SOS
447   Mata Atlântica — Belo Horizonte⬚: Conservação Internacional.

448 Gasper, A. L. de, Eisenlohr, P. V., & Salino, A. (2015). Climate-related variables and
449   geographic distance affect fern species composition across a vegetation gradient in
450   a shrinking hotspot. *Plant Ecology & Diversity*, *8*(1), 25–35.
451   https://doi.org/10.1080/17550874.2013.843604

452 Gasper, A. L. de, Oliveira, L. Z., Lingner, D. V., & Vibrans, A. C. (2018). *Inventário*
453   *Florístico Florestal de Santa Catarina, Volume VII, Espécies arbóreas raras de*
454   *Santa Catarina*. (A. L. de Gasper, L. Z. Oliveira, D. V. Lingner, & A. C. Vibrans,
455   Eds.) (Vol. VII). Blumenau: Edifurb.

456 Gasper, A. L. de, Sevegnani, L., Vibrans, A. C., Uhlmann, A., Lingner, D. V., Verdi,
457   M., … Klemz, G. (2011). Inventário de Dicksonia sellowiana Hook. em Santa
458   Catarina. *Acta Botanica Brasilica*, *25*(4), 776–784. https://doi.org/10.1590/S0102-
459   33062011000400005

460 Grenouillet, G., Buisson, L., Casajus, N., Lek, S. (2011). Ensemble modelling of
461   species distribution: the effects of geographical and environmental ranges.
462   *Ecography*, *34*(1), 9–17. https://doi.org/10.1111/j.1600-0587.2010.06152.x

463 Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high
464   resolution interpolated climate surfaces for global land areas. *International Journal*
465   *of Climatology*, *25*(15), 1965–1978. https://doi.org/10.1002/joc.1276

466 Hoshizaki, B. J., & Moran, R. C. (2001). *Fern grower's manual*. Portland, Oregon:
467   Timber Press.

468 ICMBio – Instituto Chico Mendes de Conservação da Biodiversidade. (2019). Portaria
469   N^o 116, de 22 de março de 2019. Brasília: Diáro Oficial da União. Retrieved from

http://www.in.gov.br/materia/-/asset_publisher/Kujrw0TZC2Mb/content/id/68744392

IPCC. (2014). *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. (Core Writing Team, R. K. Pachaur, & L. A. Meyer, Eds.). Geneva, Switzerland: IPCC.

IUCN. (2012). *IUCN red List Categories and Criteria: Version 3.1*. Gland, Switzerland and Cambridge: IUCN. Retrieved from www.iucn.org/publications

Jones, C. D., Hughes, J. K., Bellouin, N., Hardiman, S. C., Jones, G. S., Knight, J., … Zerroukat, M. (2011). The HadGEM2-ES implementation of CMIP5 centennial simulations. *Geosci. Model Dev.*, *4*(3), 543–570. https://doi.org/10.5194/gmd-4-543-2011

Kessler, M., Kluge, J., Hemp, A., & Ohlemüller, R. (2011). A global comparative analysis of elevational species richness patterns of ferns. *Global Ecology and Biogeography*, *20*(6), 868–880. https://doi.org/10.1111/j.1466-8238.2011.00653.x

Legendre, P. (2014). Interpreting the replacement and richness difference components of beta diversity. *Global Ecology and Biogeography*, *23*(11), 1324–1334. https://doi.org/10.1111/geb.12207

Legendre, P., & De Cáceres, M. (2013). Beta diversity as the variance of community data: dissimilarity coefficients and partitioning. *Ecology Letters*, *16*(8), 951–963. https://doi.org/10.1111/ele.12141

Lehnert, M., & Kessler, M. (2018). Prodromus of a fern flora for Bolivia. XXI. Dicksoniaceae. *Phytotaxa*, *344*(1), 69. https://doi.org/10.11646/phytotaxa.344.1.9

Lehnert, M., & Weigand, A. (2013). A proposal to distinguish several taxa in the Brazilian tree fern Cyathea corcovadensis (Cyatheaceae). *Phytotaxa*, *155*(April), 35–49. https://doi.org/10.11646/phytotaxa.155.1.3

Lehnert, M., & Weigand, A. (2016). A synopsis of the Neotropical species of Cyathea (Cyatheaceae; Polypodiopsida) with bipinnate fronds. *Brittonia*, 1–20. https://doi.org/10.1007/s12228-016-9445-1

Lima, A. A. de, Ribeiro, M. C., Grelle, C. E. de V., & Pinto, M. P. (2019). Impacts of climate changes on spatio-temporal diversity patterns of Atlantic Forest primates. *Perspectives in Ecology and Conservation 17*(2), 50–56. https://doi:10.1016/j.pecon.2019.04.004

Lingner, D. V., Schorn, L. A., Sevegnani, L., de Gasper, A. L., Meyer, L., & Vibrans, A. C. (2015). Floresta Ombrófila Densa de Santa Catarina-Brasil: agrupamento e ordenação baseados em amostragem sistemática. *Ciência Florestal*, *25*(4), 933–946. https://doi.org/10.5902/1980509820595

Liu, C., Newell, G., & White, M. (2016). On the selection of thresholds for predicting species occurrence with presence-only data. *Ecology and Evolution*, *6*(1), 337–348. https://doi.org/10.1002/ece3.1878

McCullagh, P., & Nelder, J. A. (1989). *Generalized Linear Models. Monograph on Statistics and Applied Probability*. Londo: Chapman & Hall. https://doi.org/10.1007/978-1-4899-3242-6

Mehltreter, K. (2008). Phenology and habitat specificity of tropical ferns. In T. A. Ranker & C. H. Haufler (Eds.), *Biology and evolution of ferns and lycophytes*. Cambridge: Cambridge University Press.

Naimi, B., & Araújo, M. B. (2016). sdm: a reproducible and extensible R platform for species distribution modelling. *Ecography*, *39*(4), 368–375. https://doi.org/10.1111/ecog.01881

Naimi, B., Hamm, N. A. S., Groen, T. A., Skidmore, A. K., & Toxopeus, A. G. (2014).
Where is positional uncertainty a problem for species distribution modelling?
Ecography, *37*(2), 191–203. https://doi.org/10.1111/j.1600-0587.2013.00205.x
Oliveira-Filho, A. T., Budke, J. C., Jarenkow, J. A., Eisenlohr, P. V., & Neves, D. R. M.
(2015). Delving into the variations in tree species composition and richness across
South American subtropical Atlantic and Pampean forests. *Journal of Plant*
Ecology, *8*(3), 242–260. https://doi.org/10.1093/jpe/rtt058
Oliveira-Filho, A. T., & Fontes, M. A. L. (2000). Patterns of floristic differentiation
among Atlantic forests in southeastern Brazil and the influence of climate.
Biotropica, *32*(4B), 793–810. https://doi.org/10.1111/j.1744-7429.2000.tb00619.x
Phillips, S. J., Dudík, M., & Schapire, R. (2004). A maximum entropy approach to
species distribution modeling. *Twenty-First International Conference on Machine*
Learning - ICML '04, 83. https://doi.org/10.1145/1015330.1015412
QGIS Development Team. (2019). QGIS Geographic Information System. Open Source
Geospatial Foundation Project. Retrieved from http://www.qgis.org
R Core Team. (2015). R: a language and environment for statistical computing. Viena.
Radchuk, V., Reed, T., Teplitsky, C., van de Pol, M., Charmantier, A., Hassall, C., …
Kramer-Schadt, S. (2019). Adaptive responses of animals to climate change are
most likely insufficient. *Nature Communications*, *10*(1), 1–14.
https://doi.org/10.1038/s41467-019-10924-4
Ribeiro, M. C., Metzger, J. P., Martensen, A. C., Ponzoni, F. J., & Hirota, M. M.
(2009). The Brazilian Atlantic Forest: How much is left, and how is the remaining
forest distributed? Implications for conservation. *Biological Conservation*, *142*,
1141–1153. https://doi.org/10.1016/j.biocon.2009.02.021
Ricci, M. (1996). Variation in distribution and abundance of the endemic flora of Juan
Fernandez Islands, Chile Pteridophyta. *Biodiversity and Conservation*, *5*(12),
1521–1532. https://doi.org/10.1007/BF00052113
Salino, A., Mota, N. F. O., & Almeida, T. E. (2013). Lycophytes and monilophytes in
Rio Preto State Park, Minas Gerais, Brazil. *Acta Botanica Brasilica*, *27*(2), 252–
263. https://doi.org/10.1590/S0102-33062013000200002
Santiago, A. C. P., Mynssen, C. M., Maurenza, D., Penedo, T. S. A., & Sfair, J. C.
(2013). Dicksoniaceae. In G. Martinelli & M. A. Moraes (Eds.), *Livro vermelho da*
flora do Brasil (pp. 475–476). Rio de Janeiro: Andrea Jakobsson: Instituto de
Pesquisas Jardim Botânico do Rio de Janeiro.
Sarmiento, G., Pinillos, M., & Garay, I. (2005). Biomass variability in tropical american
lowland rainforests. *Ecotropicos*, *18*(1), 1–20.
Silva, P. G., & Hernández, M. I. M. (2014). Local and regional effects on community
structure of dung beetles in a Mainland-Island scenario. *PLOS ONE*, *9*(10),
e111883. Retrieved from https://doi.org/10.1371/journal.pone.0111883
Tryon, R. M., & Tryon, A. F. (1982). *Ferns and allied plants: with special reference to*
tropical America. New York: Springer New York. https://doi.org/10.1007/978-1-
4613-8162-4
UNEP-WCMC. (2019). Protected Area Profile for Latin America & Caribbean from the
World Database of Protected Areas. Retrieved December 5, 2019, from
www.protectedplanet.net
VanDerWal, J., Falconi, L., Januchowski, S., Shoo, L., & Storlie, C. (2014).
SDMTools: Species Distribution Modelling Tools: Tools for processing data
associated with species distribution modelling exercises. R package version
1.1□221. Retrieved from https://cran.r-project.org/package=SDMTools

Vapnik, V. (2013). *The nature of statistical learning theory*. Springer science &
business media.
Vibrans, A. C., Sevegnani, L., Lingner, D. V., Gasper, A. L. de, & Sabbagh, S. (2010).
Inventário florístico florestal de Santa Catarina (IFFSC): aspectos metodológicos e
operacionais. *Pesquisa Florestal Brasileira*, *30*(64), 291–302.
https://doi.org/10.4336/2010.pfb.64.291
Voldoire, A., Sanchez-Gomez, E., Salas y Mélia, D., Decharme, B., Cassou, C., Sénési,
S., … Chauvin, F. (2013). The CNRM-CM5.1 global climate model: description
and basic evaluation. *Climate Dynamics*, *40*(9), 2091–2121.
https://doi.org/10.1007/s00382-011-1259-y
Wagner, K., Mendieta-Leiva, G., & Zotz, G. (2015). Host specificity in vascular
epiphytes: A review of methodology, empirical evidence and potential
mechanisms. *AoB PLANTS*, *7*(1), 1–25. https://doi.org/10.1093/aobpla/plu092
Watanabe, S., Hajima, T., Sudo, K., Nagashima, T., Takemura, T., Okajima, H., …
Kawamiya, M. (2011). MIROC-ESM 2010: model description and basic results of
CMIP5-20c3m experiments. *Geoscientific Model Development*, *4*(4), 845–872.
https://doi.org/10.5194/gmd-4-845-2011
Weigand, A., & Lehnert, M. (2016). The scaly tree ferns (Cyatheaceae-Polypodiopsida)
of Brazil. *Acta Botanica Brasilica*, *30*(3), 336–350. https://doi.org/10.1590/0102-
33062016abb0065
Wilson, O. J., Walters, R. J., Mayle, F. E., Lingner, D. V., & Vibrans, A. C. (2019).
Cold spot microrefugia hold the key to survival for Brazil's Critically Endangered
Araucaria tree. *Global Change Biology*, *25*(12), 4339–4351.
https://doi.org/10.1111/gcb.14755

Table 1. Comparisons between potential areas occupied by the tree ferns in all
scenarios (current, RCP 2.6, and RCP 8.5). The percentages are based on the subtropical
area of the Atlantic Forest ($\cong$ 731.788 km²) and Protected Areas inserted into the study
region. Negative values (**bold**) indicate a reduction of the potential area, while positive
values indicate an increase in the potential area. The "Current" column indicates the
predicted number of suitable cells at the present.

Protected Areas			Species (record number)	Subtropical Atlantic Forest		
Current	RCP 2.6	RCP 8.5		Current	RCP 2.6	RCP 8.5
			CYATHEACEAE			
340	**-27.06**	**-43.82**	*Alsophila capensis* (20)	815	**-40.61**	**-55.71**
477	**-9.43**	**-24.32**	*Alsophila setosa* (144)	1239	4.68	**-17.68**
219	41.10	12.33	*Alsopahila sternbergii* (19)	297	86.87	21.89
481	**-9.15**	**-10.81**	*Cyathea atrovirens* (204)	1084	4.89	3.97
349	**-0.57**	2.87	*Cyathea corcovadensis* (142)	615	7.15	17.40
385	1.56	17.92	*Cyathea delgadii* (95)	849	14.02	39.58
188	**-40.43**	**-69.68**	*Cyathea feeana* (29)	424	**-50.94**	**-77.59**
301	3.32	**-2.33**	*Cyathea hirsuta* (29)	470	28.30	20.21
248	24.60	41.53	*Cyathea leucofolis* (20)	312	85.58	145.19
371	**-6.47**	**-9.16**	*Cyathea phalerata* (205)	777	**-5.15**	**-6.18**
-	-	-	*Cyathea uleana*[1] (6)	-	-	-
354	**-25.99**	**-49.72**	*Cyathea vilosa* (21)	801	**-41.32**	**-68.16**
205	**-35.61**	**-57.56**	*Sphaeropteris gardneri* (21)	386	**-43.78**	**-65.28**
			DICKSONIACEAE			
304	**-22.70**	**-39.80**	*Dicksonia sellowiana* (146)	923	**-25.79**	**-42.15**
336	12.50	0	*Lophosoria quadripinnata* (66)	1255	**-22.23**	**-29.48**

622 [1]Species not considered in the analysis due to the low number of records.
623
624
625

Table 2. Mean species per cell and changes in α-diversity in the future scenarios (RCP 2.6, and RCP 8.5) for the whole study area and PAs only. "Gain", "equal" and "loss" indicates the proportion of cells that will respectively gain, keep the same number, or lose species in each future scenario.

Protected Areas			α-diversity	Subtropical Atlantic Forest		
Current	RCP 2.6	RCP 8.5		Current	RCP 2.6	RCP 8.5
3.6	3.4	3.1	Mean per cell[2]	5.53	5.15	4.68
-	26.56	21.83	Gain (%)	-	23.52	18.06
-	30.32	25.97	Equal (%)	-	28.00	20.36
-	43,12	52.20	Loss (%)	-	48.48	61.58

[2]Mean differences between current and future scenarios were supported by the Wilcoxon Rank-Sum Test (p-value < 0.001).

Table 3. BD$_{TOTAL}$ and its components (*RichDiff$_J$* and turnover) between the current and future climate scenarios for the whole study area and PAs only.

Protected Areas			β-diversity	Subtropical Atlantic Forest		
Current	RCP 2.6	RCP 8.5		Current	RCP 2.6	RCP 8.5
0.316	0.303	0.3	BD$_{TOTAL}$	0.320	0.305	0.294
83.3	91.2	91.3	*RichDiff$_J$* (%)	80.8	88.3	90.4
16.7	8.8	8.7	Turnover (%)	19.2	11.7	9.6

709
710 **Figure 1.** Study area and forest types of subtropical Atlantic Forest. The occurrence
711 data (black dots) was grouped into $\cong$ 30km² clusters to simplify the visualization. The
712 respective numbers on the dots represent the number of records found in each cluster.

726
727 **Figure 2.** α-diversity of tree ferns in the subtropical Atlantic Forest based on current
728 and future (RCP 2.6 and RCP 8.5) climate scenarios. The bar chart demonstrates the
729 predicted proportion of cells that will lose, keep and gain α-diversity in the two future
730 scenarios.

745
746 **Figure 3.** Current and future (RCP 2.6 and RCP 8.5) distribution of LCDB (left
747 column) and LCBD p-values (right column) of tree ferns in the subtropical Atlantic
748 Forest. Cells with p-value < 0.05 show significantly higher than average LCBD values.
749

Changes in α-diversity

Changes in β-diversity

LCBD

p-value

CURRENT

LCBD
High
Low

CURRENT

p-value
< 0.05
> 0.05

RCP 2.6

LCBD
High
Low

RCP 2.6

p-value
< 0.05
> 0.05

RCP 8.5

LCBD
High
Low

RCP 8.5

p-value
< 0.05
> 0.05