Parent of origin gene expression in the bumblebee, *Bombus terrestris*, supports Haig's kinship theory for the evolution of genomic imprinting.

Hollie Marshall[1], Jelle S. van Zweden[2], Anneleen Van Geystelen[2], Kristof Benaets[2], Felix Wäckers[3,4], Eamonn B. Mallon[1] and Tom Wenseleers[2]

[1]Department of Genetics and Genome Biology, The University of Leicester, Leicester, UK.
[2]Laboratory of Socioecology and Social Evolution, Department of Biology, KU Leuven, Leuven, Belgium.
[3]Biobest Belgium N.V., Westerlo, Belgium.
[4]The Lancaster Environmental Centre, University of Lancaster, Lancaster, UK.

Submitted: February 28[th], 2020

1 **Corresponding authors:** Eamonn Mallon
2 **Tel:** +44(0)1162523488
3
4 **Emails:** hollie_marshall@hotmail.co.uk, js.vanzweden@cbs.nl, anneleen.vangeystelen@gmail.com,
5 kristof_benaets@hotmail.com, felix.wackers@biobest.be, ebm3@le.ac.uk and tom.wenseleers@kuleuven.be.
6
7 **Abstract work count:** 150.

Abstract

Genomic imprinting is the differential expression of alleles in diploid individuals, with the expression being dependent upon the sex of the parent from which it was inherited. Haig's kinship theory hypothesizes that genomic imprinting is due to an evolutionary conflict of interest between alleles from the mother and father. In social insects, it has been suggested that genomic imprinting should be widespread. One recent study identified parent-of-origin expression in honeybees and found evidence supporting the kinship theory. However, little is known about genomic imprinting in insects and multiple theoretical predictions must be tested to avoid single-study confirmation bias. We, therefore, tested for parent-of-origin expression in a primitively eusocial bee. We found equal numbers of maternally and paternally biased expressed alleles. The most highly biased alleles were maternally expressed, offering support for the kinship theory. We also found low conservation of potentially imprinted genes with the honeybee, suggesting rapid evolution of genomic imprinting in Hymenoptera.

Impact summary

Genomic imprinting is the differential expression of alleles in diploid individuals, with the expression being dependent upon the sex of the parent from which it was inherited. Genomic imprinting is an evolutionary paradox. Natural selection is expected to favour expression of both alleles in order to protect against recessive mutations that render a gene ineffective. What then is the benefit of silencing one copy of a gene, making the organism functionally haploid at that locus? Several explanations for the evolution of genomic imprinting have been proposed. Haig's kinship theory is the most developed and best supported.

Haig's theory is based on the fact that maternally (matrigene) and paternally (patrigene) inherited genes in the same organism can have different interests. For example, in a species with multiple paternity, a patrigene has a lower probability of being present in siblings that are progeny of the same mother than does a matrigene. As a result, a patrigene will be selected to value the survival of the organism it is in more highly, compared to the survival of siblings. This is not the case for a matrigene.

Kinship theory is central to our evolutionary understanding of imprinting effects in human health and plant breeding. Despite this, it still lacks a robust, independent test. Colonies of social bees consist of diploid females (queens and workers) and haploid males created from unfertilised eggs. This along with their social structures allows for novel predictions of Haig's theory.

In this paper, we find parent of origin allele specific expression in the important pollinator, the buff-tailed bumblebee. We also find, as predicted by Haig's theory, a balanced number of genes showing matrigenic or patrigenic bias with the most extreme bias been found in matrigenically biased genes.

Keywords— reciprocal cross, bumblebee, genomic imprinting, Hymenoptera, kinship theory, RNA-seq

Introduction

Genomic imprinting is the differential expression of alleles in diploid individuals, with the expression being dependent upon the sex of the parent from which it was inherited [1]. Multiple evolutionary theories attempt to explain its existence [reviewed in 2]. The most widely accepted explanation is the kinship theory developed by Haig [3]. This theory predicts genomic imprinting arose due to natural selection acting differently on the matrigenes (maternal alleles) and the patrigenes (paternal alleles) of an individual for given processes. For example, in a polyandrous mating system with maternal care (e.g. mammals), patrigenes are predicted to be subject to selection pressures which increase resource allocation from the mother at the expense of siblings. Whereas matrigenes in this scenario are predicted to be selected for equal resource distribution amongst offspring.

The majority of support for this theory comes from studies based on mammals and flowering plant systems [2]. However, it has been suggested haplodiploid social insects can provide an ideal system to independently test Haig's kinship theory [4]. Colonies of social bees consist of diploid females (queens and workers) and haploid males created from unfertilised eggs. This along with their social structures allows for novel predictions of Haig's theory.

Research exploring parent-of-origin effects in social insects has focused on the behavioural and physiological outputs of genetic crosses. In the Argentine ant (*Linepithema humile*) paternal effects were observed in care-giving associated behaviours and in sex allocation of offspring [5, 6]. Paternal effects on dominance and stinging behaviour have also been observed in crosses of European and Africanized honeybees [7]. Additionally, Oldroyd *et al.* [8] found a parent-of-origin effect of increased ovary size in honeybees but could not definitively determine which parent this effect was driven by.

More recently, reciprocal crosses and next generation sequencing technologies have been used to identify genes with parent-of-origin allele specific expression patterns in honeybees [9, 10]. Both groups used RNA-Seq to study parent-of-origin gene expression in hybrid crosses of honeybee

sub-species. The logic of their test was that as honeybee queens are multiply mated, matrigenes can

occur in half sisters and therefore should be selected to moderate worker reproduction in queenless

colonies. Patrigenes, on the other hand will not be in half sisters and will be selected to reproduce at

any cost [10]. Therefore, the prediction is that parent of origin allele specific expression should exist

in honeybees and patrigenic expression will dominate [10]. Surprisingly, Kocher *et al*. [9] found

a matrigenic bias in gene expression however, it was later shown that sub-species incompatibility

effects influenced the results obtained [11]. Showing support for the kinship theory, Galbraith *et al*.

[10] found greater patrigenic expression in reproductive workers compared to sterile workers, with

increased patrigenic expression in the reproductive tissues.

The lack of agreement between these studies weakens their support for Haig's theory. Another

weakness of the evidence is that it is limited to only one species and tests only one prediction from the

many predictions made [4] for Haig's theory and genomic imprinting's role in social insect biology.

To test the robustness of Haig's kinship theory we present gene expression data (RNA-seq) of

reproductive and sterile workers from reciprocal crosses of sub-species of the primitively eusocial

bumblebee, *Bombus terrestris*. This species is naturally singly-mated. As such, the predictions for

the presence of matrigenic/patrigenic expression bias are different of those predicted for the naturally

multiply-mated honeybee [4]. As in Galbraith *et al*. [10] we test the effect of queenless conditions.

The same patrigene is present in all sisters (Fig.1), therefore patrigenes should be selected to moderate

worker reproduction to reduce the cost to nephews. This is the opposite of the honeybee prediction.

Matrigenes, similar to honeybees, have a non-zero probability of being in any given sister (Fig.1),

therefore matrigenes should also be selected to moderate worker reproduction, although possibly at

a lower level that patrigenes. Therefore we make three predictions; 1) that parent-of-origin allele

specific expression exists in bumblebees, 2) that parent-of-origin allele specific expression should

be balanced between genes biased patrigenically and matrigenically, with perhaps a slight surplus

of matrigenically expressed genes due to the decreased probability of a matrigene occurring in a

nephew as opposed to a son and 3) that genes showing parent-of-origin expression will be enriched

96   for reproductive related processes.

Figure 1: **Parental genome inheritance probabilities under queenless conditions in bumblebees**. Schematic adapted from Drewell *et al*. [12]. As the bumblebee is singly mated, there is only one patrigene (blue) shared between all workers. This means there is an equal probability of the patrigene ending up in the worker's son or nephew. The worker of interest has inherited the purple matrigene. Her sister has a 50% change of inheriting the purple or the green matrigene. For the purple matrigene, there is a 50% chance of ending up in the son but a 25% chance of ending up in the nephew.

Results

Differential expression between reproductive castes

Ten bases were trimmed from the 3' end of each worker RNA-seq read due to base bias generated by the Illumina protocol [13]. The mean number of uniquely mapped reads to the maternal genomes was 91.31% ± 0.57% (mean ± standard deviation) for the head samples and 91.53% ± 1.86% for the abdomen samples. This equated to a mean of 13,602,926 ± 2,774,040 and 14,181,003 ± 1,303,486 uniquely mapped reads respectively (supplementary 1.0.1). The mean number of uniquely mapped reads to the paternal genomes was 91.50% ± 0.68% (mean ± standard deviation) for the head samples and 91.70% ± 1.83% for the abdomen samples. This equated to a mean of 13,630,611 ± 2,776,779 and 14,208,273 ± 1,307,054 uniquely mapped reads respectively (supplementary 1.0.1).

Tissue type explains the majority of variation within all of the RNA-seq samples, followed by reproductive status; either reproductive or sterile (supplementary 2.0: Fig.S1). Following differential expression analysis a total of 3,505 genes were up-regulated in the abdomen of reproductive workers compared to sterile workers and 4,069 genes were down-regulated (q<0.01) (supplementary 1.0.2).

The enriched GO terms for the differentially expressed genes between reproductive and sterile castes in the abdomen included mostly regulatory processes but also "*reproduction*" (GO:0000003) (supplementary 1.0.3). Enriched GO terms associated specifically with up-regulated genes in reproductive workers in the abdomen also included: "*reproduction*" (GO:0000003) and "*DNA methylation*" (GO:0006306) (supplementary 1.0.4), these terms were not found in the enriched GO terms for genes up-regulated in sterile workers (supplementary 1.0.5).

Considerably fewer genes were differentially expressed in the head samples; 86 up-regulated genes in reproductive compared to sterile workers and 41 down-regulated genes (q<0.01) (supplementary 1.0.6). The majority of the GO terms associated with these differentially expressed genes involved biosynthetic processes (supplementary 1.0.7). Up-regulated genes in the head tissue of

121 reproductive workers also included "*reproduction*" (GO:0000003) whereas the up-regulated genes in

122 the head tissue of sterile workers consisted of mostly metabolic processes (supplementary 1.0.8 and

123 1.0.9).

Parent-of-origin gene expression

A total of 10,211 genes had a minimum of two SNPs with at least a coverage of five reads each. Of

those, 7,508 genes occurred in every cross, worker-type and tissue type. 700 genes had significant

maternal/paternal expression bias in both cross-directions for reproductive workers (q <0.05), and

747 were significant for sterile workers. The expression bias was averaged across; tissue type, worker

type, family and direction of cross to obtain an extremely conservative expression proportion. The

significant genes were then filtered to also have an average maternal expression proportion of >0.6 or

<0.4 to give a final confident list of genes showing parent-of-origin expression.

Reproductive workers have 163 genes showing significant parent-of-origin expression (Fig.2)

(supplementary 1.1.0). Sterile workers have 170 genes showing significant parent-of-origin expression

(Fig.2) (supplementary 1.1.1). There is no significant difference between the number of genes

showing maternal expression bias compared to paternal expression bias based on reproductive status

(chi-squared test of independence, χ-squared = 0, df = 1, p-value = 1). There is also no difference

in the number of genes showing paternal expression bias compared to maternal expression bias in

reproductive and sterile workers, assessed independently (chi-squared goodness of fit, reproductive:

χ-squared = 1.3804, df = 1, p-value = 0.24, sterile: χ-squared = 1.5059, df = 1, p-value = 0.2198).

The most extreme expression bias is seen in the maternally expressed alleles in both castes, with 17

genes showing a maternal expression proportion of >0.9 in both reproductive and sterile workers

(Fig.2). There were no genes showing >0.9 paternal expression bias. Additionally we did not find

any genes with significant sub-species expression bias.

Figure 2: **Maternal expression proportion of all genes by worker caste.** Each point represents a gene. Blue points are genes with significant parental expression bias (q<0.05 and expression proportion >0.6). Red points are genes with significant parental expression bias (q<0.05) with the proportion of expression >0.9. The top left quadrant of each plot represents genes with a *B. terrestris audax* expression bis, the bottom right quadrant represents gene with a *B. terrestris dalmatinus* expression bias. The top right represents genes with a maternal expression bias and the bottom left represents genes with a paternal expression bias.

Reproductive and sterile workers share a significant number of genes showing parent-of-origin

expression with the same parental bias (Fig.3, maternal expression bias: hypergeometric test, p =

9.20×10^{-108}, paternal expression bias: hypergeometric test, p = 7.66×10^{-90}). There were no

genes with maternal/paternal bias in one caste which also had the opposite bias in the other caste.

Additionally the majority of genes identified as showing parental expression bias show the same

bias in both abdomen and head tissue as well as across behaviourally defined castes (dominant and

subordinate reproductives and sterile foragers and nurses), see supplementary 2.0: Fig.S2-S7.

Figure 3: **Overlapping genes showing parent-of-origin expression in reproductive and sterile workers.** The set size indicates the number of genes in each list. The intersection size shows how many genes the corresponding lists have in common. A single dot refers to the number of genes unique to each list.

Overall genes showing parent-of-origin expression have enriched GO terms for multiple

biological processes (supplementary 1.1.2), specifically the GO terms "*negative regulation of*

reproductive processes" (GO:2000242) and "*female germ-line sex determination*" (GO:0019099) are

enriched. Genes with maternal and paternal bias in both reproductive and sterile workers also have

enriched GO terms for multiple biological processes (supplementary 1.1.3 and 1.1.4). Specifically,

paternally expressed genes in both reproductive and sterile workers are enriched for the GO term;

"*behaviour*" (GO:0007610).

GO terms for genes showing parent-of-origin expression in only reproductive or sterile

workers were also enriched for various biological processes (supplementary 1.1.5). Including

"*histone ubiquitination*" (GO:0016574) and "*histone H2A monoubiquitination*" (GO:0035518) in

reproductive paternally expressed genes.

Genes showing maternal parent-of-origin expression are enriched for genes which are also

differentially expressed in head tissue (Supplementary 2.0: Fig.S8, hypergeometric test, p = 0.004).

Specifically *Serine Protease Inhibitor 3/4* (LOC100652301) shows maternal expression bias in both

reproductive and sterile workers and is up-regulated in the head tissue of reproductive workers.

Genes with paternal parent-of-origin expression do not significantly overlap with differentially

expressed genes (supplementary 2.0: Fig.S8, hypergeometric test, p = 0.148). There is also no

significant overlap between genes showing parent-of-origin expression bias and differential expression

in the abdomen (supplementary 2.0: Fig.S9, hypergeometric test, p = 0.996).

Honeybee homology

A custom database of putative orthologs was made between *A. mellifera* and *B. terrestris*, as in [14],

containing 6,539 genes. 68% of differentially expressed abdominal genes were identified within the

database, 43% of differentially expressed head genes and 46% of genes showing parent-of-origin

expression bias (supplementary 1.1.6). Gene lists were obtained from Galbraith *et al*. [10], the

ortholog database contained 38% of the differentially expressed genes identified between honeybee

reproductive worker castes and 53% of the genes found to show parent-of-origin expression bias

(supplementary 1.1.6).

There was no significant overlap between the genes identified as showing parent-of-origin

expression between both studies (hypergeometric p = 0.64), with only two genes overlapping

(supplementary 2.0 Fig.S10). One of these is uncharacterised in both species (honeybee id:

LOC552195, bumblebee id: LOC100648162) and the second is a serine protease inhibitor (honeybee

id: LOC411889, bumblebee id: LOC100644680). The serine protease inhibitor shows paternal

expression bias in honeybees and maternal bias in both bumblebee reproductive and sterile castes.

It is not differentially expressed in the honeybee but it shows up-regulation in the abdomen tissue

of reproductive bumblebee workers compared to sterile workers. Additionally, [10] identified

numerous genes of interest which are involved in reproductive behaviour (*vitellogenin, yolkless,*

ecdysone-receptor and ecdysone-induced protein) which show paternal expression bias in honeybees,

none of which show significant parent-of-origin expression in the bumblebee (supplementary 2.0

Fig.S11).

There was also no significant overlap between differentially expressed genes identified in

head tissue and abdomen tissue between *B. terrestris* reproductive castes with those identified as

differentially expressed between reproductive castes of *A. mellifera* from [15] (supplementary 2.0

Fig.S12, head: hypergeometric p = 1, abdomen hypergeometric p = 1).

Discussion

Using parental genome sequencing and offspring RNA-seq we have identified genes showing parent-of-origin allele specific expression in a primitively eusocial bumblebee species. There was no difference in the number of genes showing maternal or paternal expression bias in either reproductive or sterile workers. The genes showing the highest proportion of expression bias were all maternally expressed in both worker castes. Additionally, reproductive related GO terms were enriched in both maternally and paternally biased genes.

Reproductive and sterile workers were chosen in order to provide a robust test for the kinship theory. Imprinted genes should maintain their expression bias regardless of the current reproductive state of the individual, i.e. higher matrigenic expression compared to patrigenic expression should be present in queenless workers regardless of whether they have become reproductive or remained sterile. Additionally the use of sterile and reproductive workers allowed us to assess differential expression between castes. This meant we could investigate if there was a relationship between differentially expressed genes between reproductive castes and genes showing parent-of-origin expression. A high overlap of differentially expressed genes between reproductive castes with genes showing parent-of-origin expression would suggest a role for imprinted genes in reproductive caste determination.

As in Galbraith *et al.* [10] we also found no significant overlap of paternally expressed genes with differentially expressed genes between reproductive and sterile workers. However, we did find a significant overlap with maternally expressed genes and genes differentially expressed in head tissue between reproductive and sterile castes. This significant overlap should be interpreted cautiously however as only seven of the 103 unique maternally biased genes were differentially expressed in head tissue between reproductive castes. The lack of overlap of differentially expressed genes with paternally expressed genes and the small overlap with maternally expressed genes suggests parent-of-origin expression may not directly influence reproductive status in bumblebee workers.

One of the overlapping genes found to be differentially expressed in head tissue between

reproductive castes which also shows maternal expression bias is a serine protease inhibitor. One of

the two homologous genes identified as showing parent-of-origin expression in this study and in

honeybees was also a serine protease inhibitor. Serine proteases (also know as serpins) have been

shown to be involved in insect immunity in various species including; the silkworm *Bombyx mori*

[16], another species of silk producing moth *Antheraea pernyi* [17] and the mosquito *Anopheles*

gambiae [18]. Most recently a kazal-type serine protease inhibitor has been directly linked to

oocyte development in the desert locust *Schistocerca gregaria* [19]. Future work identifying the

function of serine protease inhibitors in social insects is needed to better understand the function of

parent-of-origin expression of these genes in both *B. terrestris* and *A. mellifera*.

We found genes involved in histone modifications to be paternally expressed in reproductive

workers. Histone modifications have been identified as an imprinting mark in plants [20] and thought

to be involved in imprinting maintenance in mammals [21]. Histone modifications can alter gene

expression by affecting gene accessibility via chromatin [22]. Chromatin modifications have been

associated with parent-of-origin expression in the fruit fly *Drosophilia melanogaster* [23]. Galbraith

et al. [10] also found genes involved in histone modifications showing parent-of-origin expression in

the honeybee.

Only two genes were found in common between those found in Galbraith *et al*. [10] as

showing parent-of-origin gene expression in the honeybee *A. mellifera* and those identified here in

B. terrestris. Galbraith *et al* [10] used ovaries and fat bodies as their tissues samples whereas we

selected to test whole head and whole abdomen samples for strong signals of expression bias. Some

imprinted genes in mammals are known to be tissue specific, such as *GBR10* which has been found

to be maternally expressed in brain and muscle tissue but not in growth plate cartilage [24]. Tissue

specificity could account for the lack of concordance in parentally expressed genes found between *B.*

terrestris and *A. mellifera*. Additionally 51% of bumblebee genes and 52% of honeybee genes were

not present in the homology database created.

Imprinted genes in mammals show much more consistency across species, with mice and

humans reportedly sharing around 50 imprinted genes out of around 150 and 100 characterised in

each respectively [25]. Additionally, domestic cattle and pigs have been shown to share 14 imprinted

genes out of 26 and 18 respectively [26]. However, imprinted genes in plants generally show less

conservation, with one study reporting 14% of maternally expressed genes and 29% of paternally

expressed genes in *Capsella rubella* show the same imprinting status in *Arabidopsis thaliana*, even

though both species belong to the Brassicaceae family [27]. Hatorangan *et al*. [27] suggest the lack

of consistency between species could be the result of a historical shift in mating-systems. Given the

differences in mating-systems between honeybees and bumblebees, and that variable predictions

from the kinship theory apply to each species, rapid evolution of imprinted genes in Hymenoptera is

a feasible explanation for the lack of consistency in potentially imprinted genes identified here and in

honeybees.

The GO terms associated with both maternally and paternally expressed genes are diverse.

It has been suggested imprinted genes can function as a mechanism for plasticity, allowing gene

regulation to change depending on environmental conditions by activating the silenced allele and

increasing dosage of that gene [28]. Social insects display, sometimes extreme, phenotypic plasticity,

where multiple discrete phenotypes (castes) can arise from a single genome within a colony. In some

species this is genetically determined [29], however there is growing evidence epigenetic factors

may play a role in caste determination in some species [14, 30, 31]. Matsuura *et al*. [32] modelled a

genomic imprinting mediated caste determination system in the termite *Reticulitermes speratus* and

found this better explained the influence of parental phenotype on offspring than a purely genetic

model. Given the diversity of genes found here showing both maternal and paternal expression

bias we believe, along with Matsuura [33], that further experimental investigation into the role of

genomic imprinting in caste determination in social insects is needed.

The identification of genes showing parent-of-origin expression in this study lays the ground

work for future research to identify potential epigenetic mechanisms of allele specific expression in

social insects. Genes showing allele specific expression and DNA methylation have been previously

identified in *B. terrestris* [34], and genes involved in the reproductive process have been shown to be

differentially methylated between reproductive castes [14, 35]. DNA methylation is the mechanism by

which some genes are imprinted in mammals and plants [36] and so investigation of parent-of-origin

methylation in *B. terrestris* may be fruitful.

These results provide support for Haig's kinship theory. We found as predicted a balanced

number of genes showing matrigenic or patrigenic bias. Also as predicted, the most extreme bias

was found in matrigenically biased genes. This is novel, independent support for this important

evolutionary theory. The results of this study create a base for many future avenues of research

including; gene function analysis of serine protease inhibitors in Hymenoptera, epigenetic mechanisms

of imprinting in insects and imprinted genes as a mechanism for plasticity, caste determination and

social evolution.

Methods

Sample collection

Reciprocal crosses of *B. terrestris dalmatinus* (native to southern Europe) and *B. terrestris audax* (native to the UK) were carried out by Biobest, Leuven. Reciprocal crosses allow subspecies-of-origin effects (i.e. the effect of genotype) to be disentangled from parent-of-origin effects [9, 10]. In order to obtain enough successful colonies multiple males and females for each cross (Fig. 4) were released into cages to mate. Once mating had occurred the male and female were removed. The male was immediately frozen at -80°C and the female was placed in cold conditions for eight weeks to induce diapause. Ten matings were carried out for each cross.

Four successful colonies (one of each cross direction) from two 'families' (Fig.4) were housed at the University of Leuven and kept in 21°C with red light conditions, they were fed *ad libitum* with pollen and a sugar syrup. Callow workers were labelled with numbered disks in order to determine age and allow behaviour to be recorded. Once each colony contained approximately 30 workers the queen was removed. The colonies were then filmed under queenless conditions for 30mins per day for 14 days in order to score individual behaviour. The following behaviours were used to classify workers: incubating, feeding larvae, inspecting brood cells, building egg cups, ventilation, biting, pushing, egg-laying, egg-eating, foraging, feeding and grooming. Workers were classified based on the frequency of each of the above behaviours as either: sterile foragers, sterile nurses, dominant reproductives or subordinate reproductives (supplementary 1.0.0).

Figure 4: **Graphic display of the family-wise reciprocal crosses carried out.** Each colour refers to related individuals, i.e. the queen from colony 02 is the sister of the male used in colony 12. This design reduces genetic variability between the initial and reciprocal crosses.

Worker reproductive status was confirmed by ovary dissection, ovaries were scored on a

0-4 scale as in Duchateau and Velthuis [37], entire bodies were then stored at -80°C along with

the original queen mothers and male fathers. Workers were selected for sequencing based on their

behavioural classification and ovary status. Two of each behavioural type per colony were selected

(i.e. two dominant reproductives, two subordinate reproductives, two nurses and two foragers),

with the exception of colony 22 (Fig.4) which contained three subordinate reproductives and one

dominant reproductive. All sterile and reproductive samples were age matched. This gave a total of

32 samples, 8 per colony, four of each reproductive status, reproductive or sterile. See supplementary

1.0.0 for behavioural and ovary scoring per sample.

DNA and RNA extraction and sequencing

DNA was extracted from the mother and father of each colony using the Qiagen DNeasy® Blood & Tissue Kit. Thirty-two workers were selected for RNA sequencing. The head and abdomen were dissected and RNA extracted separately for each, using the Qiagen RNeasy® Lipid Tissue Kit, giving 64 total RNA samples. The quality of the DNA and RNA extraction were measured by Nanodrop and Qubit® fluorometer. Whole genome parental DNA was sequenced using 91bp paired-end reads and worker RNA was sequenced using 90bp paired-end reads on an Illumina HiSeq 2000 by BGI, China. Lane effects were minimised for the RNA samples by spreading colony and tissue samples across five lanes.

Generation of alternative reference genomes

Whole genome sequencing data was checked using Fastqc (v.0.11.05) (www.bioinformatics.ba-braham.ac.uk/projects/fastqc/) and adapters and low quality reads were trimmed using Cutadapt v.1.11 [38]. Reads were aligned to the bumblebee reference genome (Bter_1.0, Refseq accession no. GCF_000214255.1 [39]) using BWA-mem v.0.7.15 [40] with standard parameters. SNPs were then called using freebayes v.1.1.0 [41] which can account for the difference in ploidy between males and females, a minimum of five reads per SNP were required. Queen SNPs were then filtered so only the homozygous alternative SNPs remained. The subtractbed command from BEDtools v.2.25.0 [42] was then used to create files containing SNPs unique to either the mother/father of each colony. The individual parental SNP files were then used to create alternate reference genomes for each parent using the 'fasta alternate reference maker' command in GATK v.3.6 [43].

Identification of parent-of-origin expression

RNA-Seq data was quality checked and trimmed as above. STAR v.2.5.2b [44] was used to align worker RNA-seq reads to each of that colony's specific parental genomes with zero mismatches

allowed. This ensures any reads containing a SNP will only be matched to the parent that allele was inherited from. Alignment files were then filtered using BEDtools v.2.25.0 [42] so only alignments which contain an informative SNP (a unique SNP from either the mother or father) were kept. Reads were counted for the maternal/paternal alignments, also using BEDtools v.2.25.0 [42] and SNP positions were annotated with a gene ID taken from the Bter_1.0 annotation file (Refseq accession no. GCF_000214255.1) using a custom R script. SNPs which had zero maternal reads in at least one sample were removed completely from the analysis to avoid possible inflation of paternal counts. This would occur if the queen position was mis-called as homozygous with the missing allele matching that of the male [10].

Genes showing parent-of-origin expression were determined using a logistic regression model in R v.3.4.0 (https://cran.r-project.org). Only genes occurring in both cross directions and in both family combinations, with a minimum of two SNPs per gene were analysed, this left a total of 7508 genes. If any gene showed zero reads for paternal counts this was changed to 1 to avoid complete separation. A quasibionimal distribution was also used to account for overdispersion within the data. Fixed factors included the direction of the cross, family and reproductive status (reproductive or sterile). Correction for multiple testing was carried out using the Benjamini–Hochberg method [45]. Genes were determined as showing parent-of-origin expression if the allelic ratio (maternal/paternal) corrected p-value was <0.05 and the parental expression proportion was >0.6.

Differential expression

All RNA-seq samples were aligned to the reference genome (Bter_1.0, Refseq accession no. GCF_000214255.1 [39]) using STAR v.2.5.2b [44] with standard parameters. HTseq v.0.8.0 [46] was then used to count the number of reads per gene for each sample. Differential gene expression between reproductive and sterile workers for head and abdomen samples was assessed using the DESeq2 package v.1.16.1 [47] in R. DESeq2 allows the incorporation of a general linear model (GLM) to identify differential expression; family, age, weight, direction of the cross, tissue type

359 and reproductive status were all factors. P-values were corrected for multiple testing using the the

360 Benjamini–Hochberg method [45].

Gene ontology enrichment

GO enrichment analysis was carried out using the hypergeometric test with Benjamini-Hochberg

[45] multiple-testing correction in a custom R script, (q <0.05). This script utilised a previously

made GO database for the *B. terrestris* genome, Bter_1.0 [48]. GO terms for differentially expressed

genes were tested for enrichment against GO terms associated with all genes identified in either the

the RNA-seq data from the abdomen or head. Up-regulated genes in either reproductive or sterile

workers were tested for GO term enrichment against all differentially expressed genes from the

respective tissue type as a background set.

Genes showing parent-of-origin expression were tested for enrichment against GO terms

associated with all genes identified in both abdomen and head RNA-Seq data sets. Genes maternally

or paternally biased were checked for GO term enrichment against all genes showing parent-of-origin

expression as a background set. REVIGO [49] was used to obtain the GO descriptions from the GO

identification numbers.

Comparative analyses

375 A hypergeometric test was applied to gene lists from the differential expression analysis and the

376 parent-of-origin expression analysis to identify potential enrichment. *B. terrestis* and *A. mellifera*

377 orthologous genes were determined as in Marshall *et al.* [14] and a custom R script was then

378 used to check for overlap between genes identified as showing parent-of-origin expression here and

379 orthologous *A. mellifera* genes identified in Galbraith *et al.* [10].

Acknowledgements

We thank Biobest N.V. for providing us with the forty bumblebee colonies of the desired reciprocal crosses. Hollie Marshall had a Ph.D-scholarship from the Central England NERC Training Alliance (NERC, UK). Kristof Benaets had a Ph.D.-scholarship of the Flemish Agency for Innovation by Science and Technology (IWT-Vlaanderen). This study was funded in the context of the KU Leuven BOF Centre of Excellence Financing on 'Eco- and socio-evolutionary dynamics' (Project number PF/2010/07) and a grant from the Research Foundation-Flanders (FWO-Vlaanderen, grant G.0463.12). E.B.M. was funded by NERC grant NE/N010019/1. This research used the ALICE2 High Performance Computing Facility at the University of Leicester.

Author Contributions

T.W. and E.B.M. conceived the study. The reciprocal crosses were carried out by Biobest (Westerlo, Belgium) under supervision of F.W. J.S.Z. and K.B. carried out the behavioural observations and lab work. H.M. and T.W. carried out the analyses with input from A.V.G. H.M. and E.B.M. wrote the initial manuscript. All authors contributed to and reviewed the manuscript.

Data Accessibility

All RNA-seq and whole genome sequencing data is available under NCBI BioProject number PRJNA329487. Custom scripts are available at the following DOI: `https://zenodo.org/record/3235636`.

Conflict of Interest

All authors declare no conflict of interests.

References

[1] Pegoraro, M., Marshall, H., Lonsdale, Z.N. and Mallon, E.B. Do social insects support Haig's kin theory for the evolution of genomic imprinting? *Epigenetics*, 12(9):725–742, (2017).

[2] Patten, M.M., Ross, L., Curley, J.P., Queller, D.C., Bonduriansky, R. and Wolf, J.B. The evolution of genomic imprinting: theories, predictions and empirical tests. *Heredity*, 113(2): 119–128, (2014).

[3] Haig, D. The Kinship Theory of Genomic Imprinting. *Annual Review of Ecology, Evolution, and Systematics*, (31):9–32, (2000).

[4] Queller, D.C. Theory of genomic imprinting conflict in social insects. *BMC Evolutionary Biology*, 3(15), (2003).

[5] Libbrecht, R. and Keller, L. Genetic compatibility affects division of labor in ants. *Evolution*, 67(2):517–524, (2012).

[6] Libbrecht, R., Schwander, T. and Keller, L. Genetic Components to Caste Allocation in a Multiple-Queen Ant Species. *Evolution*, 65(10):2907–2915, (2011).

[7] Guzman-Novoa, E., Hunt, G.J., Page, R.E., Uribe-Rubio, J.L., Prieto-Merlos, D. and Becerra-Guzman, F. Paternal effects on the defensive behavior of honeybees. *Journal of Heredity*, 96 (4):376–380, (2005).

[8] Oldroyd, B.P., Allsopp, M.H., Roth, K.M., Remnant, E.J., Drewell, R.A. and Beekman, M. A parent-of-origin effect on honeybee worker ovary size. *Proceedings of the Royal Society B: Biological Sciences*, 281(1775):20132388, (2014).

[9] Kocher, S.D. et al. A Search for Parent-of-Origin Effects on Honey Bee Gene Expression. *G3*, 5(8):1657–1662, (2015).

[10] Galbraith, D.A. et al. Testing the kinship theory of intragenomic conflict in honey bees (*Apis mellifera*). *Proceedings of the National Academy of Sciences*, 113(4):1020–1025, (2016).

[11] Gibson, J.D., Arechavaleta-Velasco, M.E., Tsuruda, J.M. and Hunt, G.J. Biased allele expression and aggression in hybrid honeybees may be influenced by inappropriate nuclear-cytoplasmic signaling. *Frontiers in Genetics*, 6(343), (2015).

[12] Drewell, R., Lo, N., Oxley, P.R. and Oldroyd, B.P. Kin conflict in insect societies: A new epigenetic perspective. *Trends in Ecology and Evolution*, 27(7):367–373, (2012).

[13] Krueger, F., Andrews, S.R. and Osborne, C.S. Large scale loss of data in low-diversity illumina sequencing libraries can be recovered by deferred cluster calling. *PLoS ONE*, 6(1):4–10, (2011).

[14] Marshall, H., Lonsdale, Z.N. and Mallon, E.B. Methylation and gene expression differences between reproductive and sterile bumblebee workers. *Evolution Letters*, 3(5):485–499, (8 2019).

[15] Galbraith, D.A., Yi, S.V. and Grozinger, C.M. Evaluation of possible proximate mechanisms underlying the kinship theory of intragenomic conflict in social insects. *Integrative and Comparative Biology*, 56(6):1206–1214, (2016).

[16] Zou, Z., Picheng, Z., Weng, H., Mita, K. and Jiang, H. A comparative analysis of serpin genes in the silkworm genome. *Genomics*, 93(4):367–375, (2009).

[17] Yu, H.M. et al. Characterization and functional analysis of serpin-1 like gene from oak silkworm *Antheraea pernyi*. *Bulletin of Entomological Research*, (2017):1–7, (2017).

[18] Gorman, M.J. and Paskewitz, S.M. Serine proteases as mediators of mosquito immune responses. *Insect Biochemistry and Molecular Biology*, 31(3):257–262, (2001).

[19] Guo, W., Wu, Z., Yang, L., Cai, Z., Zhao, L. and Zhou, S. Juvenile hormone-dependent Kazal-type serine protease inhibitor Greglin safeguards insect vitellogenesis and egg production. *FASEB Journal*, 33(1):917–927, (2019).

[20] Rodrigues, J.A. and Zilberman, D. Evolution and function of genomic imprinting in plants. *Genes and Development*, 29(24):2517–2531, (2015).

[21] Delaval, K. and Feil, R. Epigenetic regulation of mammalian genomic imprinting. *Current Opinion in Genetics & Development*, 14:188–195, (2004).

[22] Bannister, A.J. and Kouzarides, T. Regulation of chromatin by histone modifications. *Cell Research*, 21(3):381–395, (2011).

[23] Joanis, V. and Lloyd, V. Genomic imprinting in Drosophila is maintained by the products of Suppressor of variegation and trithorax group, but not Polycomb group, genes. *Molecular Genetics and Genomics*, 268:103–112, (2002).

[24] McCann, J.A., Zheng, H., Islam, A., Goodyer, C.G. and Polychronakos, C. Evidence against GRB10 as the gene responsible for Silver-Russell syndrome. *Biochemical and Biophysical Research Communications*, 286(5):943–948, (2001).

[25] Babak, T. et al. Genetic conflict reflected in tissue-specific maps of genomic imprinting in human and mouse. *Nature Genetics*, 47(5):544–549, (2015).

[26] Tian, X.C. Genomic Imprinting in Farm Animals. *Annual Review of Animal Biosciences*, 2(1): 23–40, (2013).

[27] Hatorangan, M.R., Laenen, B., Steige, K.A., Slotte, T. and Köhler, C. Rapid Evolution of Genomic Imprinting in Two Species of the Brassicaceae. *The Plant Cell*, 28(8):1815–1827, (2016).

[28] Radford, E.J., Ferrón, S.R. and Ferguson-Smith, A.C. Genomic imprinting as an adaptative model of developmental plasticity. *FEBS Letters*, 585(13):2059–2066, (2011).

[29] Mott, B.M., Gadau, J. and Anderson, K.E. Phylogeography of Pogonomyrmex barbatus and *P. rugosus* harvester ants with genetic and environmental caste determination. *Ecology and Evolution*, 5(14):2798–2826, (2015).

[30] Lyko, F., Foret, S., Kucharski, R., Wolf, S., Falckenhayn, C. and Maleszka, R. The honey bee epigenomes: Differential methylation of brain DNA in queens and workers. *PLoS Biology*, 8 (11), (2010).

[31] Bonasio, R. et al. Genome-wide and caste-specific DNA methylomes of the ants *Camponotus floridanus* and *Harpegnathos saltator*. *Current Biology*, 22(19):1755–1764, (2012).

[32] Matsuura, K. et al. A Genomic Imprinting Model of Termite Caste Determination: Not Genetic but Epigenetic Inheritance Influences Offspring Caste Fate. *The American Naturalist*, 191(6), (2018).

[33] Matsuura, K. Genomic imprinting and evolution of insect societies. *Population Ecology*, pages 1–15, (2019).

[34] Lonsdale, Z. et al. Allele specific expression and methylation in the bumblebee, *Bombus terrestris*. *PeerJ*, 5(e3798), (2017).

[35] Amarasinghe, H.E., Clayton, C.I. and Mallon, E.B. Methylation and worker reproduction in the bumble-bee (*Bombus terrestris*). *Proceedings of the Royal Society B: Biological Sciences*, 281(20132502), (2014).

[36] Scott, R.J. and Spielman, M. Genomic imprinting in plants and mammals: How life history constrains convergence. *Cytogenetic and Genome Research*, 113(1-4):53–67, (2006).

[37] Duchateau, M.J. and Velthuis, H.H.W. Development and reproductive strategies in *Bombus terrestris* colonies. *Behaviour*, 107(3):186–207, (1988).

[38] Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal*, 17(1):10, (2011).

[39] Sadd, B.M. et al. The genomes of two key bumblebee species with primitive eusocial organization. *Genome Biology*, 16(1):76, (2015).

[40] Li, H. and Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*, 25(14):1754–1760, (2009).

[41] Garrison, E. and Marth, G. Haplotype-based variant detection from short-read sequencing. *arXiv*, 1207.3907, (2012).

[42] Quinlan, A.R. and Hall, I.M. BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics*, 26(6):841–842, (2010).

[43] McKenna, A. et al. The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, 20:1297–1303, (2010).

[44] Dobin, A., Gingeras, T.R. and Spring, C. Mapping RNA-seq Reads with STAR Alexander. *Current Protocols in Bioinformatics*, (51):1–11, (2016).

[45] Benjamini, Y. and Hochberg, Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society*, 57(1):289–300, (1995).

[46] Anders, S. et al. Count-based differential expression analysis of RNA sequencing data using R and Bioconductor. *Nature Protocols*, 8(9):1765–1786, (2013).

[47] Love, M.I., Huber, W. and Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome biology*, 15(12):550, (2014).

[48] Bebane, P.S.A. et al. The effects of the neonicotinoid imidacloprid on gene expression and DNA methylation in the buff-tailed bumblebee *Bombus terrestris*. *Proc. R. Soc. B*, 286:20190718, (2019).

[49] Supek, F., Bošnjak, M., Škunca, N. and Šmuc, T. Revigo summarizes and visualizes long lists of gene ontology terms. *PLoS ONE*, 6(7), (2011).