1

Transcription factors involved in stem cell maintenance are

downstream of Slug/Snail2 and repressed by TGF-β in bronchial

epithelial progenitors from Chronic Obstructive Pulmonary Disease

Pierre de la Grange[3], Ariane Jolly[3], Charlotte Courageux[1,2], Chamseddine Ben Brahim[1,2],

Pascale Leroy[1,2*]

[1]Inserm UMR1152, Physiopathology and Epidemiology of Respiratory Diseases, Paris, France

[2]Faculty of Medicine, Paris Diderot University, Bichat Campus, Paris, France.

[3]GenoSplice, Paris, France

*Correspondence: pascale.leroy@inserm.fr

Abstract

Objectives: Patients with Chronic Obstructive Pulmonary Disease (COPD) have a bronchial

epithelium with many anomalies and basal/progenitor cells showing a decrease of self-renewal

and differentiation potential. The objective of this study was to identify deregulations in the

genetic program of COPD bronchial progenitors that could account for their exhaustion. The

transcription factor Slug/Snail2 is highly expressed in bronchial progenitors and we aimed at

identifying genes downstream of Slug whose expression is deregulated in COPD progenitors.

Results: We knocked down Slug in primary basal cells from COPD subjects and, since COPD

subjects have higher levels of Transforming Growth Factor (TGF)-β and Slug is regulated by

TGF-β, we selected genes downstream of Slug involved in differentiation that respond to TGF-β.

We identified transcription factors involved in stem cell maintenance downstream of Slug and

repressed by TGF-β in COPD but not normal progenitors. We found that the effect of TGF-β on

the expression of these genes is correlated to Slug knockdown effect. We also found a correlation

between the mRNA levels of Slug and these genes only in presence of TGF-β. These results

reveal that stem cell maintenance genes are deregulated in COPD bronchial progenitors, Slug and

TGF-β being involved in that deregulation.

Key words: COPD, Bronchial epithelium, Primary progenitors, TGF-β, Slug/Snail2,

Somatic stem cell maintenance genes, Transcription factors, Microarray analysis

Introduction

Subjects with chronic obstructive pulmonary disease (COPD), a respiratory disease mainly

caused by cigarette smoke and characterized by a progressive and irreversible loss of respiratory

capacity, have a remodeling of their airways with many anomalies of the epithelium that increase

with disease progression (1, 2). These anomalies are found all along the airway epithelium as

cells keep the memory of the exposure to cigarette smoke establishing a "field of injury", and

would result from an imbalance of the fate of basal cells, the airway epithelium adult

stem/progenitor cells that can self-renew and/or differentiate to repair the epithelium after injury

(3, 4). A decrease in the count and functional ability of basal progenitors has been reported in

smokers with a higher decrease in COPD smokers, leading to a statistically significant difference

between non-smokers and smokers-COPD that correlates with the loss of respiratory capacity (5,

6).

Slug/Snail2 belongs to the Snail family of transcription factor and is involved in

epithelial-mesenchymal transition (EMT). In contrast to other EMT-inducing transcription

factors, Slug is highly expressed in adult basal/progenitor cells in both mouse and human normal

airway epitheliums (7, 8). We hypothesized that Slug is involved in the fate of basal/progenitor

cells, and, as such, deregulation of its function could lead to the decrease in count and functional

ability of progenitors observed in COPD. To test this hypothesis, we exploited the Gene

Expression Omnibus (GEO) dataset GSE123129 that we generated from a microarray analysis of

Slug knockdown in basal/progenitor cells of COPD and normal subjects (9). Transforming

Growth Factor (TGF)-β is found at higher levels in COPD lung tissues and, as it has been shown

to regulate Slug expression and to play a role in stem/progenitor cell fate, we also used the GEO

dataset GSE122957 to select genes downstream of Slug in COPD that respond to TGF-β (9-12).

Main text

Material and methods

Study subjects and cell isolation

Human lung tissues, smokers (n=6) and COPD-smokers (n=6) (see additional file 2) were obtained from patients undergoing lung lobectomy for peripheral lung carcinoma for removal of a primary lung tumor. COPD patients were diagnosed according to the GOLD (Global initiative for chronic Obstructive Lung Disease) guidelines. Lung tissues used in this study were dissected as far away as possible from the tumor. Primary human bronchial epithelial cells were isolated from a piece of large bronchus according to standard protocol.

Cell culture

Primary human bronchial epithelial cells (Passage 1 or 2) were expanded on flasks coated with collagen I (BD Biocoat) in bronchial epithelial growth medium (BEGM), composed of bronchial epithelial basal medium (BEBM) supplemented with the SingleQuots kit (Lonza) and incubated at 37°C in 5% de CO_2. TGF-β1 (Peprotech) was added at a concentration of 1ng/ml for 2 days.

RNA extraction

Cells were rinsed with phosphate buffer saline (PBS) before to be homogenized in RNA lysis buffer (NucleoSpin-RNA Kit, Macherey-Nagel), supplemented with 1% β-Mercaptoethanol. Lysates were vortexed and proceeded immediately or snapped Frozen and stored at -80°C. Total RNA was extracted on column using the NucleoSpin-RNA Kit (Macherey-Nagel). DNaseI treatment was done directly on the column. RNA concentration was determined using a NanoDrop.

cDNA synthesis

For each sample, 0.3μg of total RNA was reverse transcribed as previously described (13) with

the following modifications: briefly, total RNA was annealed with 0.1mg/ml oligo(dT)15 primer

(Promega) and cDNA synthesis was performed with M-MLV Reverse Transcriptase (Promega)

for 1h at 42˚C. A control without reverse transcriptase was done for each series of cDNAs.

Quantitative Real-Time Polymerase Chain Reaction (qPCR)

Quantitative PCR was done on the QuantStudio6 Flex with the QuantStudio Real-Time PCR

software using SYBR green PCR master mix (Applied BioSystems). Primers (see additional file

3) were designed to be used with an annealing temperature of 60°C and an elongation time of

1min. For a given gene target, cDNA volume used was chosen to be in a Ct range from 20 to 30,

using 0.125μM each forward and reverse primer. Glyceraldehyde-3-Phosphate Dehydrogenase

(GAPDH) was used to normalize cDNA amounts between samples.

Gene expression profiling

Microarray analysis was performed on biological triplicate samples. Total RNA were amplified

and labeled before hybridization onto Affymetrix human Gene 2.1ST GeneChip according to the

manufacturer, by the Genomics platform at Curie Institute, Paris (14). Array datasets were

controlled using Expression console (Affymetrix) and further analyses and visualization were

made using EASANA (GenoSplice, www.genosplice.com), which is based on the FAST DB

annotations (15, 16). Gene Array data were normalized using quantile normalization. Background

corrections were made with antigenomic probes and probes were selected as described previously

(17). Only probes targeting exons annotated from FAST DB transcripts were selected to focus on

well-annotated genes whose mRNA sequences are in public databases (16). Bad-quality selected

probes (e.g. probes labeled by Affymetrix as 'cross-hybridizing') and probes whose intensity

signal was too low compared to antigenomic background probes with the same GC content were

removed from the analysis. Only probes with a DABG P-value ≤ 0.05 in at least half of the arrays

were considered for statistical analysis (17). Only genes expressed in at least one compared

condition were analyzed. To be considered expressed, the DABG P-value had to be ≤ 0.05 for at

least half of the gene probes. We performed an unpaired Student's t-test to compare gene

intensities in the different biological replicates. Genes were considered significantly regulated

when fold-change was ≥1.2 and P-value ≤ 0.05. Significant KEGG and REACTOME pathways

and GO terms were retrieved using DAVID (18-20) from union of results of all, up- and down-

regulated genes separately. Data set GEO ID numbers are GSE122957 and GSE123129.

Statistical Analysis

Biological replicates were n ≥3 for the knockdown experiments and n ≥5 for expression analysis

with data generated by at least 2 independent experiments. For fold-change, mean is ±SEM and

statistical analysis was carried out by a one sample two-sided t-test. Correlations were computed

as Pearson correlation coefficients and P-value determined by two-sided test. Significance was

accepted when P-value < 0.05.

Results

Transcription factor genes involved in somatic stem cell maintenance are

repressed downstream of Slug in COPD progenitors

When setting the minimum fold-change threshold at 1.2 with P-value ≤5.00E-0, 808 genes were

found upregulated in COPD cells knocked down for Slug, i.e. repressed downstream of Slug.

Among these, 307 genes are responding to both differentiation and TGF-β and we classified them

in 4 groups according to their combined type of response. Fig. 1a shows histograms representing

the number of genes for each group and small bars indicating the mean fold-increase induced by

Slug knockdown for these genes in COPD (red) and normal (blue) cells. The large majority of the

genes are upregulated during differentiation and repressed by TGF-β, and for these the mean fold-

increase induced by Slug knockdown is much higher in COPD cells. A search for enriched gene

pathways using KEGG, REACTOME and Gene Ontology (GO) databases (18-20) identified 2

pathways related to stem cells, Somatic stem cell population maintenance and transcriptional

regulation of pluripotent stem cell, comprising 5 genes coding all for transcription factors. We

first confirmed by RT-qPCR the fold-change induced by Slug knockdown in COPD and normal

cells and Fig. 1b shows the correlation with the values obtained in the microarray. The good

correlation (R^2 values of 0.7309 and 0.5519, respectively for normal and COPD), validates the

microarray analysis and confirms that all the genes are repressed downstream of Slug in COPD

cells while they are not downstream of Slug in normal cells, except for ELF5 that is also

repressed in normal cells but less than in COPD.

Transcription factor genes involved in stem cell maintenance are more

repressed by TGF-β in COPD progenitors

The transcription factor genes involved in stem cell maintenance, which we identified as

repressed downstream of Slug in COPD progenitors, were also found by microarray analysis

repressed by TGF-β. We confirmed that all 5 genes are repressed by TGF-β in COPD, with the

decrease of expression statistically significant except for STAT3. In contrast, different effects,

mostly mild, are observed in normal progenitors, none being statistically significant **(**Fig. 2a**).** By

plotting Slug knockdown effect with that of TGF-β on the expression of these genes, we find a

good correlation with a R^2 value of 0.6813 (Fig. 2b). Moreover, for all the genes, except STAT3,

a strong negative correlation coefficient is found between their expression levels and that of Slug

but only in presence of TGF-β. The correlation is statistically significant for SOX2 and ELF5

while it is not for EPAS1 and PBX1; however, for these later genes, the correlation strength

increases significantly in presence of TGF-β (Table1, additional file 1).

Taken together, these results show that in COPD progenitors, when compared to normal, genes

coding for transcription factors involved in stem cell maintenance that became direct or indirect

repressed targets of Slug, have their response to TGF-β deregulated and a link exists between the

repressive effect of TGF-β and that of Slug.

164 **Discussion**

We identified 5 genes coding for transcription factors involved in somatic stem cell maintenance

that are repressed downstream of Slug and by TGF-β in COPD bronchial basal/progenitor cells;

in contrast, in normal cells, these genes are less or not repressed by TGF-β, and 4 of them are not

downstream of Slug. In COPD, we find a correlation between the effect of Slug knockdown and

that of TGF-β on their expression, as well as between their mRNA levels and that of Slug when in

presence of TGF-β.

Previous works have shown at the cellular level that bronchial basal/progenitor cells from

COPD are deregulated when compared to normal, with a decrease in self-renewal and a loss in

differentiation potential leading to progenitor exhaustion (5, 6). Our work reveals that key

transcription factors involved in stem cell maintenance, direct or indirect repressed targets of Slug,

have their expression deregulated in COPD, being in particular repressed when in presence of

TGF-β. A high level of TGF-β is found in COPD airways and our results are in line with the

epithelial progenitor exhaustion described for this disease.

TGF-β represses these transcription factors with different efficiency, and it correlates with

the effect of Slug knockdown, showing a link in COPD bronchial progenitors between Slug

expression and the response of these genes to TGF-β. This is supported by the high negative

correlation coefficient between the mRNA levels of Slug and that of these transcription factors,

with the exception of STAT3, found in presence of TGF-β. For EPAS1 and PBX1, despite strong

correlation coefficients, P-values are above the significance threshold, showing only a tendency

for these genes. However, the small number of patients studied is likely to limit the statistical

significance. No such correlation is found for STAT3 but Slug knockdown has the least effect on

this gene and it is repressed by TGF-β only slightly and with no statistical significance. Each gene

have a different profile of regulation by Slug and TGF-β and it is likely that this reflects the

existence of other regulations, specific for each gene. TGF-β may coregulate Slug and these

genes or, as in other cells, Slug may act as a mediator of TGF-β to regulate these transcription

factors (21, 22).

Our work reveals that the genetic program of COPD bronchial progenitors is deregulated;

this deregulation involves a change in Slug downstream genes and in the response to TGF-β, with

consequences on self-renewal and differentiation potential. There is a higher level of TGF-β in

COPD and it increases with disease progression (23-25) and we can speculate that progenitor

deregulation worsen with the level of TGF-β, participating to COPD progression.

Limitations

In the present work, the analysis of the stem cell maintenance genes was limited to the

respective effect of Slug knockdown and TGF-β on their expression, and the study of Slug and

TGF-β link was limited to expression correlation. Further research on functional analysis is

warranted to define the role of these stem cell maintenance transcription factors in bronchial

progenitors and the mechanisms involved in their regulation by Slug and TGF-β.

List of abbreviations

205   COPD Chronic obstructive pulmonary disease, TGF-β Transforming growth factor beta, EMT

206   epithelial-mesenchymal transition, GAPDH Glyceraldehyde-3-Phosphate Dehydrogenase

Declarations

Authors' contributions

210   PL conceptualized project, conceived and designed the study, conducted experiments, interpreted

211   the data, obtained financial support. PdlG designed the study, analyzed and interpreted the data.

212   AJ analyzed the data. CC and CBB performed the experiments. PL and PdlG wrote the

213   manuscript. All authors read and approved the final manuscript.

Author details

215   [1]Inserm UMR1152, Physiopathology and Epidemiology of Respiratory Diseases, Paris, France.

216   [2]Faculty of Medicine, Paris Diderot University, Bichat Campus, Paris, France. [3]GenoSplice, Paris,

217   France.

Acknowledgements

219   We thank the Pulmonary Department, the Pathology Department and the Thoracic Surgery

220   Department at Bichat-Claude Bernard University Hospital (Paris, France) and INSERM UMR

221 1152 for providing lung tissues and isolating the cells. We thank Audrey Rapinat and David

222 Gentien at the Genomics platform, Curie Institute (Paris, France) for Affymetrix GeneChip

223 hybridization.

Competing interests

225 The authors declare that they have no competing interests.

Availability of data and material

227 The data analyzed in this publication have been deposited in the NCBI's Gene

228 Expression Omnibus (GEO) and will be accessible through GEO Series Accession

229 Number: GSE122957 and GSE123129

Consent for publication

231 Not applicable.

Ethics approval and consent to participate

233 Human lung tissues were obtained from patients undergoing lung lobectomy for peripheral lung

234 carcinoma after receiving written informed consent. The study was approved by the ethics

235 committee of Paris Nord, IRB 00006477 Paris 7 University, France.

Funding

237 P.L. is supported by the French National Center for Scientific Research (CNRS). This work was

238 supported by a donation from Association Science et Technologie (Groupe Servier) to P.L. and

239 by funding from French National Institute for Medical Research (INSERM). The funders had no

240 role in study design, data collection and analysis, interpretation of data and preparation of the

241 manuscript.

References

1.   Rigden HM, Alias A, Havelock T, O'Donnell R, Djukanovic R, Davies DE, et al.
      Squamous Metaplasia Is Increased in the Bronchial Epithelium of Smokers
      with Chronic Obstructive Pulmonary Disease. PLoS One
      2016;11(5):e0156009.

2.   Sohal SS, Walters EH. Role of epithelial mesenchymal transition (EMT) in
      chronic obstructive pulmonary disease (COPD). Respir Res 2013;14:120.

3.   Rock JR, Randell SH, Hogan BL. Airway basal stem cells: a perspective on
      their roles in epithelial homeostasis and remodeling. Dis Model Mech
      2010;3(9-10):545-56.

4.   Steiling K, van den Berge M, Hijazi K, Florido R, Campbell J, Liu G, et al. A
      dynamic bronchial airway gene expression signature of chronic obstructive
      pulmonary disease and lung function impairment. Am J Respir Crit Care Med
      2013;187(9):933-42.

5.   Ghosh M, Miller YE, Nakachi I, Kwon JB, Baron AE, Brantley AE, et al.
      Exhaustion of Airway Basal Progenitor Cells in Early and Established Chronic
      Obstructive Pulmonary Disease. Am J Respir Crit Care Med 2018;197(7):885-
      896.

6.   Staudt MR, Buro-Auriemma LJ, Walters MS, Salit J, Vincent T, Shaykhiev R, et
      al. Airway Basal stem/progenitor cells have diminished capacity to
      regenerate airway epithelium in chronic obstructive pulmonary disease. Am J
      Respir Crit Care Med 2014;190(8):955-8.

265   7.   Hackett NR, Shaykhiev R, Walters MS, Wang R, Zwick RK, Ferris B, et al. The

266   human airway epithelial basal cell transcriptome. PLoS One

267   2011;6(5):e18378.

268   8.   Rock JR, Onaitis MW, Rawlins EL, Lu Y, Clark CP, Xue Y, et al. Basal cells as

269   stem cells of the mouse trachea and human airway epithelium. Proc Natl

270   Acad Sci U S A 2009;106(31):12771-5.

271   9.   Ben Brahim C, Courageux C, Jolly A, Ouine B, Cartier A, de la Grange P, et al.

272   Slug/Snail2 is involved in the repression of proliferation genes by TGF-• in

273   bronchial epithelial progenitor cells and is deregulated in abnormal

274   epithelium. BioRxiv 2019.

275   10.   Mahmood MQ, Reid D, Ward C, Muller HK, Knight DA, Sohal SS, et al.

276   Transforming growth factor (TGF) beta1 and Smad signalling pathways: A

277   likely key to EMT-associated COPD pathogenesis. Respirology

278   2017;22(1):133-140.

279   11.   Slabakova E, Pernicova Z, Slavickova E, Starsichova A, Kozubik A, Soucek K.

280   TGF-beta1-induced EMT of non-transformed prostate hyperplasia cells is

281   characterized by early induction of SNAI2/Slug. Prostate 2011;71(12):1332-

282   43.

283   12.   Yumoto K, Thomas PS, Lane J, Matsuzaki K, Inagaki M, Ninomiya-Tsuji J, et al.

284   TGF-beta-activated kinase 1 (Tak1) mediates agonist-induced Smad

285   activation and linker region phosphorylation in embryonic craniofacial

286   neural crest-derived cells. J Biol Chem 2013;288(19):13467-80.

287   13.   Leroy P, Mostov KE. Slug is required for cell survival during partial epithelial-

288   mesenchymal transition of HGF-induced tubulogenesis. Mol Biol Cell

289   2007;18(5):1943-52.

290   14.   Maubant S, Tahtouh T, Brisson A, Maire V, Nemati F, Tesson B, et al. LRP5

291   regulates the expression of STK40, a new potential target in triple-negative

292   breast cancers. Oncotarget 2018;9(32):22586-22604.

293   15.   de la Grange P, Dutertre M, Correa M, Auboeuf D. A new advance in

294   alternative splicing databases: from catalogue to detailed analysis of

295   regulation of expression and function of human alternative splicing variants.

296   BMC Bioinformatics 2007;8:180.

297   16.   de la Grange P, Dutertre M, Martin N, Auboeuf D. FAST DB: a website

298   resource for the study of the expression regulation of human gene products.

299   Nucleic Acids Res 2005;33(13):4276-84.

300   17.   de la Grange P, Gratadou L, Delord M, Dutertre M, Auboeuf D. Splicing factor

301   and exon profiling across human tissues. Nucleic Acids Res 2010;38(9):2825-

302   38.

303   18.   Haw R, Stein L. Using the reactome database. Curr Protoc Bioinformatics

304   2012;Chapter 8(Unit8 7.).

305   19.   Huang da W, Sherman BT, Lempicki RA. Systematic and integrative analysis

306   of large gene lists using DAVID bioinformatics resources. Nat Protoc

307   2009;4(1):44-57.

308   20.   Kanehisa M, Goto S, Sato Y, Furumichi M, Tanabe M. KEGG for integration and

309   interpretation of large-scale molecular data sets. Nucleic Acids Res

310   2012;40(Database issue):D109-14.

311   21.   Geismann C, Arlt A, Bauer I, Pfeifer M, Schirmer U, Altevogt P, et al. Binding of

312   the transcription factor Slug to the L1CAM promoter is essential for

313   transforming growth factor-beta1 (TGF-beta)-induced L1CAM expression in

314   human pancreatic ductal adenocarcinoma cells. Int J Oncol 2011;38(1):257-

315   66.

316   22.   Joseph MJ, Dangi-Garimella S, Shields MA, Diamond ME, Sun L, Koblinski JE,

317   et al. Slug is a downstream mediator of transforming growth factor-beta1-

318   induced matrix metalloproteinase-9 expression and invasion of oral cancer

319   cells. J Cell Biochem 2009;108(3):726-36.

320   23.   Godinas L, Corhay JL, Henket M, Guiot J, Louis R, Moermans C. Increased

321   production of TGF-beta1 from sputum cells of COPD: Relationship with

322   airway obstruction. Cytokine 2017;99:1-8.

323   24.   Stoll P, Wuertemberger U, Bratke K, Zingler C, Virchow JC, Lommatzsch M.

324   Stage-dependent association of BDNF and TGF-beta1 with lung function in

325   stable COPD. Respir Res 2012;13:116.

326   25.   Zhang S, Xie JG, Su BT, Li JL, Hu N, Chen J, et al. MFG-E8, a clearance

327   glycoprotein of apoptotic cells, as a new marker of disease severity in chronic

328   obstructive pulmonary disease. Braz J Med Biol Res 2015;48(11):1032-8.

Figures, table and additional Files

Fig. 1 Slug knockdown in COPD bronchial progenitors identifies repressed genes

involved in somatic stem cell maintenance. Primary bronchial epithelial basal cells from COPD

and normal subjects were knocked down for Slug, and RNA were analyzed by microarray on an

Affymetrix chip. Data are for n ≥ 3 each, normal and COPD. **a** Genes significantly upregulated by

Slug knockdown in COPD cells and responding to both differentiation and TGF-β were classified

in 4 groups according to their type of response. Histograms represent the number of genes in each

group with genes upregulated by TGF-β in red and genes downregulated by TGF-β in green, and

genes upregulated during differentiation as checkboard and genes downregulated during

differentiation as oblique lines. Horizontal bars represent the mean fold-increase for each group

of genes, in red for COPD cells and in blue for normal cells. **b** Validation of microarray by RT-

qPCR and comparison of Slug knockdown effect in COPD (red) and normal (blue) cells for the

genes coding for 5 transcription factors involved in somatic stem cell maintenance. Results are

Pearson correlations calculated with log2 (fold-change) and are presented as scatter plots with

regression line and R-square values.

Fig. 2 Genes involved in stem cell maintenance are more repressed by TGF-β in COPD bronchial

progenitors. Primary bronchial epithelial basal cells, normal and COPD, were grown on filters

without TGF-β or in presence of 1 ng/ml of TGF-β. At confluence, RNA lysates were prepared

and mRNA analyzed by RT-qPCR. GAPDH was used to normalize cDNA amounts between

samples and results were calculated as a ratio on GAPDH. Data shown are for n ≥ 5. **a**

Comparison of TGF-β effect on the expression of stem cell maintenance genes normal and COPD

cells. Results are presented as the fold-change induced by TGF-β on mRNA expression with

mean ±SEM. **b** Correlation in COPD progenitors between Slug knockdown and TGF-β effect on

the expression of the 5 genes coding for transcription factors involved in somatic stem cell

maintenance. Results are Pearson correlations calculated with log2 (fold-change) and are

presented as scatter plots with regression line and R-square values. Statistical significance is at P-

value < 5.00E-02 *, < 1.00E-03 *** as indicated. ns: non significant.

Table 1 Correlations between Slug and stem cell maintenance genes mRNA in COPD progenitors

treated with TGF-β

	Slug mRNA	
	no TGFβ	+ TGFβ
SOX2 mRNA	- 0.6566 (2.28E-01)	- 0.8951 (4.01E-02)
ELF5 mRNA	- 0.1203 (8.47E-01)	- 0.8976 (3.87E-02)
EPAS1 mRNA	- 0.4492 (4.47E-01)	- 0.7753 (1.23E-01)
PBX1 mRNA	- 0.6255 (2.59E-01)	- 0.7753 (1.23E-01)
STAT3 mRNA	- 0.1261 (8.39E-01)	0.1827 (7.68E-01)

361
362

Pearson correlation coefficients and associated P-value between expression levels.

Computation was performed on log2 of values.

Statistical significance was determined with a two-sided t-test.

Correlations are significant at P-value < 5.00E-02.

Additional File 1 Correlation Graphs corresponding to data in Table 1

Additional File 2 Characteristics of the study subjects.

Data are presented as Mean ±SD. Pack-year=1 year smoking 20 cigarettes per day. COPD,

chronic obstructive pulmonary disease; FEV1, forced expiratory volume in 1 s;

Additional File 3 Sequences of primers used for qPCR

Figure 1

a

b

Figure 2