The TMJ disc is a common ancestral feature in all mammals, as evidenced by the presence of a rudimentary disc during monotreme development

Neal Anthwal, Abigail S Tucker.

Centre for Craniofacial and Regenerative Biology, 27th Floor Guy's Tower, King's College London, London SE1 9RT

11
Abstract
The novel mammalian jaw joint, known in humans as the temporomandibular joint or TMJ, is
cushioned by a fibrocartilage disc. This disc is secondarily absent in therian mammals that
have lost their dentition, such as giant anteaters and some baleen whales. The disc is also
absent in all monotremes. However, it is not known if the absence in monotremes is
secondary to the loss of dentition, or if it is an ancestral absence. We use museum held
platypus and echidna histological sections to demonstrate that the developing monotreme jaw
joint forms a disc primordium that fails to mature and become separated from the mandibular
condyle. We then show that monotreme developmental anatomy is similar to that observed in
transgenic mouse mutants with reduced musculature. We therefore suggest that the absence of
the disc on monotremes is a consequence of the changes in jaw musculature associated with
the loss of adult teeth. Taken together, these data indicate that the ancestors of extant
monotremes likely had a jaw joint disc, and that the disc evolved in the last common ancestor
or all mammals.

26
27 **Keywords**
28 TMJ disc, monotreme, mammalian evolution, jaw joint, evo devo
29
30

31
Introduction

The temporomandibular joint (TMJ) is the one of the most used joints in the body, articulating the upper and lower jaw in mammals. A fibrous articular disc sits between the skeletal elements of the joint and acts as a cushion.

TMJ development occurs by the coming together of two membranous bones: the condylar process of the dentary bone in the mandible and the squamosal bone in the skull. The interaction of the condylar with the squamosal induces the formation of a glenoid (or mandibular) fossa on the latter (Wang et al., 2011). The articular disc sits between the two within a synovial capsule. The TMJ disc attaches to the superior head of the lateral pterygoid muscle anteriorly, and to ligaments posteriorly including the disco-mallear ligament that runs thought the capsule of the middle ear, joining the malleus to the TMJ disc. The TMJ articulates the jaw in all mammals, and is referred to as the squamosal dentary joint (SDJ) in those mammals without a fused temporal bone. In non-mammals the upper and lower jaw articulate via the endochondral quadrate and articular, known as the primary jaw joint (Wilson and Tucker, 2004). TMJ developmental anatomy reflects its evolutionary history as this novel, jaw joint forms after the development of the primary joint, which, in mammals, is integrated into the middle ear (Takechi and Kuratani, 2010; Anthwal et al., 2013; Maier and Ruf, 2016; Tucker, 2017). In recent years, a number of studies have advanced the understanding of middle ear evolution in the context of anatomical development (Luo, 2011; Anthwal et al., 2013, 2017; Urban et al., 2017; Wang et al., 2019), but little work has sought to understand the TMJ in an evolutionary and comparative developmental biology context. This is despite the crucial role that the formation of the TMJ has in mammalian evolution.

An important part of the TMJ is the disc that cushions its action. The origin of the disc is uncertain. The insertion of the lateral pterygoid muscle into the disc on the medial aspect, and the presence of the discomalleolar ligament has led to speculation that the disc represents a fibrocartilage sesamoid with a tendon of a muscle of mastication trapped by the novel mammalian jaw joint (Herring, 2003). However, studies in mice indicate that the disc develops from a region of flattened mesenchyme cells adjacent to, or possibly part of, the perichondrium of the developing condylar cartilage (Purcell et al., 2009, 2012; Hinton et al., 2015). Formation of the disc condensation is dependent on Ihh signalling from the cartilage (Shibukawa et al., 2007; Purcell et al., 2009; Yang et al., 2016), and Fgf signalling via Spry 1 and 2 genes from the adjacent muscles (Purcell et al., 2012). Therefore, the disc may have its origins in either a tendon, the novel secondary cartilage of the condylar process, or a combination of the two.

Interestingly the disc is absent in extant monotremes (Sprinz, 1964). Monotremes and therian mammals (marsupials and placentals) are evolutionary distant, with the common ancestor of the two subclasses being a mammal like reptile form around 160 million years ago (Kemp, 2005). Monotremes have a number of "reptile" like anatomical features such as a cloaca, external embryonic development in an egg, a straight cochlear in the inner ear and laterally protruding legs (Griffiths, 1978). The absence of a disc in both echidna and platypus suggests that the disc evolved after the split between monotremes and therian mammals, and is therefore a therian novelty. Alternatively, absence of the TMJ disc in extant monotremes might be due to a secondary loss of this structure, linked to changes in mastication with the reduction and loss of teeth. Extant adult monotremes are edentulous, possibly due to the evolution of electroreceptivity (Asahara et al., 2016). The juvenile platypus has rudimentary teeth that regress (Green, 1937), while the echidna shows only thickening of the dental epithelium during development. In contrast, fossil monotremes have a mammalian

tribosphenic dentition and were capable of chewing (Kemp, 2005). The presence or absence
of a disc in such fossils is difficult to ascertain due to lack of preservation of soft tissue. In
support of mastication playing a role in disc formation edentulous therian mammals, or those
lacking enamel, often lack a disc. These species include some (but not all) baleen whales (El
Adli and Deméré, 2015), giant ant eaters and sloths (Naples, 1999).
In order to address this uncertainty, we have examined the development of the TMJ in
monotremes and made comparison with mouse developmental models where muscle
development is perturbed.
89
Materials and Methods
Platypus *(Ornithorhynchus anatinus)* and short-beaked echidna *(Tachyglossus aculeatus)*
slides were imaged from the collections at the Cambridge University Museum of Zoology.
Details of samples imaged are in table 1. All museum samples have been studied in
previously published works (Watson, 1916; Green, 1937; Presley and Steel, 1978). Stages for
platypus are after Ashwell (Ashwell, 2012). Staging of echidna H.SP EC5 is estimated by
cross-referencing previous studies (Griffiths, 1978; Rismiller and McKelvey, 2003). CT cans
of adult platypus were a gift of Anjali Goswami, the Natural History Museum, London.
Mesp1Cre;Tbx1flox (Tbx1CKO) mice were derived as previously described (Anthwal et al.,
2015).
Tissue processing and histological staining: embryonic samples for histological sectioning
were fixed overnight at 4 °C in 4 % paraformaldehyde (PFA), before being dehydrated
through a graded series of ethanol and stored at -20˚C. For tissue processing, samples were
cleared with Histoclear II, before wax infiltration with paraffin wax at 60°C. Wax embedded
samples were microtome sectioned at 8 µm thickness, then mounted in parallel series on
charged slides.
For histological examination of bone and cartilage, the slides were then stained with
picrosirius red and alcian blue trichrome stain using standard techniques.
108

Species	Collection	ID	Estimated Stage	CRL
Ornithorhynchus anatinus	Cambridge	Specimen X	P6.5[*]	33mm
Ornithorhynchus anatinus	Cambridge	Specimen Delta	P10[*]	80mm
Ornithorhynchus anatinus	Cambridge	Specimen Beta	P80[*]	250mm
Tachyglossus aculeatus	Cambridge	Echidna H.SP EC5	P18[†]	83mm

109 *Table 1: Museum held specimens used in the current study. CRL – Crown rump length. [*]Estimate based on Ashwell, 2012.*
110 *[†]Estimate based on Griffiths, 1978 and Rismiller & McKelvey, 2003.*

111
112 **Results**
113 If the TMJ disc is a therian novelty then no evidence of a disc would be expected in extant
114 monotremes during development of the TMJ. The development of the jaw joint was therefore
115 examined in museum held histological sections of developing post-hatching platypus and
116 compared with the mouse.
117

Figure 1 Comparison of mouse (Mus musculus) and platypus (Ornithorhynchus anatinus) developing jaw joint reveals the presence of a jaw joint disc anlage in early post-hatching platypus despite absence of the disc in adults. A,B Histological sections of mouse jaw joint disc development at embryonic day 16.5 (A) and postnatal day 0 (B).C-D' Histological sections of estimated post hatching day 6.5 jaw joint at two different levels (C and D) Note that the separation between the disc anlage and condylar in D is probably a processing artefact. E,E'' Histological sections of estimated post hatching day 10 jaw joint. F Histological section of mature jaw joint in a juvenile platypus are estimated post hatching day 80. G µCT scan of jaw joint region of adult platypus. G.F. – glenoid fossa; Cdy. – condylar process; Cdy. Fibro. – condylar fibrocartilage; Synv. – synovial cavity of the jaw joint.

As other authors have previously described (Purcell et al., 2009; Hinton, 2014), in the embryonic day (E) 16.5 mouse, the disc anlage is observed as thickened later of mesenchyme connected to the superior aspect of the condylar cartilage (Figure 1A). At postnatal day (P) 0, the disc has separated from the condylar and sits within the synovial cavity of the jaw joint (Figure 1B). In a platypus sample estimated to be 6.5 days post hatching, the TMJ had been

initiated, but the joint cavity had not yet formed (Figure 1C,D). Close examination of the
superior surface of the condylar cartilage revealed a double layer of thickened mesenchyme in
the future fibrocartilage layer of the condylar (Figure 1 C',D'). The outer layer is similar to
that known to develop into the articular disc in therian mammals (Purcell et al., 2009). This
thickened mesenchyme persisted in older samples, estimated to be P10, where the synovial
cavity of the TMJ was beginning to form above (Figure 1 E,E'). In the most mature platypus
sample examined (around P80) the fibrocartilage layer of the condylar process was thick and
had a double-layered structure (Figure 1F). The outer layer was connected via a tendon to the
lateral pterygoid muscle. At this late stage of postnatal development, the platypus puggle
would have been expected to start leaving the burrow and to be eating a mixed diet, although
full weaning does not occur until around 205 days post hatching (Rismiller and McKelvey,
2003). In the mature platypus, the condylar process sits within a glenoid fossa (Figure 1 F,G),
which was not fully formed at earlier stages. A disc-like structure lying over the condylar and
connected to the adjacent muscles was therefore evident in the platypus postnatally but did
not lift off the condylar at any stage.
147
Next we examined the development of the TMJ in a derived young short-beaked echidna
puggle specimen with a crown-rump length of 83mm, which we estimate to be around P18.
The TMJ is not fully (Figure 2). The condylar process possessed a thick, doubled
fibrocartilage outer layer (Figure 2), much as was observed in the platypus (Figure 1D). The
outer fibrocartilage later was connected by connective tissue to the lateral pterygoid muscle
(Figure 2B'). Clear disc-like structures were therefore present during development in both
extant monotremes.
155

156
157 *Figure 2 Examination of the developing jaw joint reveals the presence of a jaw joint disc anlage in post-hatching day 18*
158 *short-beaked echidna (Tachyglossus aculeatus). A-B Histological staining at the forming jaw articulation in echidna young*
159 *estimated to be 18 days post hatching at two different level. Fibrocartilage disc anlage superior to the condylar and*
160 *connected by tendon lateral pterygoid muscle is observed. B' High-powered view of boxed region in B showing the*
161 *connection between the muscle and the developing disc. Cdy. – condylar process; m. lat. ptry. – lateral pterygoid muscle.*

162
Taken together, the developmental evidence suggests that extant monotremes initiate a layer
of fibrocartilage connected to the lateral pterygoid muscle, similar to the initiation of the TMJ
disc in therian mammals. However, unlike in therian mammals, the monotreme fibrocartilage
failed to separate from the condylar to form an articular disc in the TMJ. Interactions with
musculature, both mechanical (Habib et al., 2007; Purcell et al., 2012; Jahan et al., 2014;
Nickel et al., 2018) and molecular (Shibukawa et al., 2007; Gu et al., 2008; Purcell et al.,
2009, 2012; Kinumatsu et al., 2011; Michikami et al., 2012; Yasuda et al., 2012; Kubiak et
al., 2016), have been suggested to be responsible for the proper formation of the TMJ disc.
Lack of mechanical force in monotremes might therefore result in the disc remaining attached
to the condylar. In order to examine how changes in mechanical loading affect disc
development, we next examined disc development in the *Mesp1Cre;Tbx1flox* conditional
mutant mouse (*Tbx1CKO*). This mouse has a mesoderm specific deletion of the T-box

transcription factor *Tbx1*, resulting in hypomorphic muscle development (Grifone et al., 2008;
Aggarwal et al., 2010; Anthwal et al., 2015).
177
We used alcian blue / alizarin red stained histological sections to investigate the development
of the TMJ disc in *TbxCKO* mice at embryonic day 15.5. This is the stage when future disc
mesenchyme is first observed (see Figure 1A). In wildtype embryos, the future disc
mesenchyme was observed as a condensation attached to the superior surface of the condylar
fibrocartilage (Figure 3A). A distinct disc-like mesenchyme was also observed superior to the
condylar of the *Tbx1CKO* (Figure 3B). This mesenchyme and the fibrocartilage layer of the
condylar cartilage both appeared thicker in the *Tbx1CKO* compared to its wildtype littermate.
At E18.5, the wildtype TMJ disc had separated from the condylar process, and sat within a
synovial joint cavity (Figure 3C). In the *Tbx1CKO* an upper synovial cavity had formed,
similar to the WT, but the disc had failed to separate from the condylar (Figure 3D). Instead, a
thickened band of fibrocartilage was observed on the superior surface of the condylar process.
189
190

191
192 *Figure 3 Muscle-disc interactions are required for the maturation and separation of the jaw joint articular disc. A,B The disc*
193 *anlage is observed at E15.5 in both wildtype mice (A) and Mesp1Cre;Tbx1fl/fl mice with a hypomorphic muscle phenotype*
194 *(B). C,D By E18.5 the disc has separated from the condylar process in wildtype mice (C), but not in and Mesp1Cre;Tbx1fl/fl*
195 *mice.   Cdy.   –   condylar   process;   Cdy.   Fibro.   –   condylar   fibrocartilage*

Discussion

The absence of an articular disc in monotremes has been thought to be either a secondary loss related to the absence of mature dentition, or the disc being a later acquisition in the therian clade. The data presented here show that a mesenchyme similar to the TMJ disc is initiated in both platypus and echidna jaws during post-hatching development, but fails to mature and separated from the dentary condyle. In the light of the failure of the disc to fully separate in transgenic mouse models with hypomorphic muscle development, it seems likely that the disc has been secondarily lost in edentulous mammals, including monotremes.

The secondary jaw joint of some of the earliest mammals-like reptiles with a mandibular middle ear, such as Morganuconodon, were able to withstand the biomechanical stresses sufficient for feeding on the hard keratinised bodies of insects, while others such as Kuehneotherium could not (Gill et al., 2014). Later animals developed a range of mandibular movements during chewing, including rolling, yaw and front to back movements (Kemp, 2005; Luo et al., 2015; Grossnickle, 2017; Lautenschlager et al., 2017, 2018; Bhullar et al., 2019). It Is not clear if these species had evolved an articular disc, since fibrocartilage is rarely fossilised. Based on the presence of the first stages of disc formation during monotreme development it is likely that the common stem Jurassic mammal-like reptilian ancestor of both monotremes and therian mammals had a disc. The data presented here confirms an essential biomechanical component in disc development. Therefore, we are able to consider when during mammalian evolution these forces were able to act to enable disc formation. For example, it is probable that many late Triassic and early Jurassic mammaliaforms such a Hadrocodium (Luo, 2001) possessed an articular disc, since they possessed a well formed squamosal dentary joint and occluding teeth capable of chewing.

One hypothesis for the origin of the articular disc is that it formed from the tendon of a muscle of jaw closure of the primary jaw joint interrupted by the formation of the novel mammalian jaw joint (Herring, 2003). The tendons and skeleton of the front of the head are derived from the cranial neural crest, whereas much of the musculature is mesoderm derived (Santagati and Rijli, 2003; Yoshida et al., 2008). Interactions between the mesoderm and neural crest co-ordinate the muscular skeletal development of the head (Grenier et al., 2009). A striking piece of evidence for the tendon origin of the disc is the expression in the developing articular disc of *Scleraxis* (Purcell et al., 2012; Roberts et al., 2019), a specific regulator of tendon and ligament development (Schweitzer et al., 2001; Sugimoto et al., 2013). If the disc is derived from a tendon, then it may be thought of as a fibrocartilage sesamoid. Such sesamoids are found in joints and in tendons that are subject to compression, like the tendons that pass around bony pulleys such as the flexor digitorum profundus tendon in quadrupeds, the patella tendon and ligament (Benjamin and Ralphs, 1998), and the cartilago transiliens in crocodilians (Tsai and Holliday, 2011). Fibrocartilages also form at the enthesis of long bones. Interestingly, it has been demonstrated that much like the TMJ disc, enthesis fibrocartilage cells are derived from Hh responsive cells and that these cells are responsive to mechanical loading (Schwartz et al., 2015). To support the tendon origin of the TMJ disc, our data show that the formation of the disc is dependent on interactions between the skeletal and muscle components of the TMJ. Such tissue interaction is also a key process in the formation of tendons and ligaments (Eloy-Trinquet et al., 2009; Huang, 2017).

The mechanism by which the disc fails to separate from the condylar in monotremes is not yet clear. Hh signalling is known to be involved in both the initiation of the disc, and the later separation from the condylar (Purcell et al., 2009). It is still possible that the role in Hh in

separation of the disc is a therian innovation, and as such the reason that monotremes fail to
do so is a lack of the later Hh dependent developmental programme for disc separation.
However, the absence of the disc in therian edentates strongly suggests that the loss is
secondary. Furthermore, the failure of the disc to elevate off the condylar in *Tbx1CKO,* with
hypomorphic cranial musculature, suggests that the loss of discs in edentulous mammals is
due to changes in the developmental biomechanics of the muscle/bone interactions that
occurred as a consequence of loss of teeth, such as a reduction in size and power of the
muscles of mastication. The formation and maturation of the disc is unlikely to be directly
dependent on the presence of teeth. The TMJ disc forms normally during embryonic
development in mice quite some time before the eruption of the teeth during the third
postnatal week, while baleen whales vary in the presence or absence of TMJ discs, and indeed
TMJ synovial cavities (El Adli and Deméré, 2015). In addition, it is clear that movement of
the jaw is essential for maturation of the disc (see also (Habib et al., 2007)). Unfortunately,
due to the rarity of fresh material, it is not possible to further examine the mechanistic aspects
of TMJ development in edentulous monotreme species at the present time.
261
In conclusion, we demonstrate that during development, monotremes show evidence of
initiation of the fibrocartilage articular disc, despite all adult monotremes not having an
articular TMJ disc. The maturation and separation of the disc is dependent on biomechanical
interactions with the associated musculature, as demonstrated by the failure of disc maturation
and separation in mice mutants with hypomorphic cranial muscle. Therefore, toothed
ancestors of monotremes likely had a TMJ disc. Our research suggests that changes in the
cranial musculature that occurred as a consequence of a move towards edentulous dietary
niches resulted in absence of the TMJ in monotremes, a parallel loss occurring in edentulous
therian mammals (Figure 4). Finally, the presence of the disc anlage in monotremes indicates
that the mammal-like reptile ancestors of all modern mammals likely possessed a disc to
cushion the novel jaw articulation.
273

274
Figure 4 Maturation of the jaw joint articular disc in mammals is dependent on muscle interactions. In toothless mammals,
reduction in jaw musculature results in changes in muscle-disc interaction and so the disc does not separate from the
mandibular condyle to sit within the synovial joint capsule.

Acknowledgments
We would like to thank and acknowledge the following people. Anjali Goswami provided
µCT images of the adult platypus. Robert Asher provided access to samples held at the
Zoological Museum in Cambridge University. Peter Giere provided access to the Hill
Collection at the Berlin Museum für Naturkunde. Andrew Gillis provided assistance with
imaging.
286
287 **Funding**
288 This work was supported by the Wellcome Trust (102889/Z/13/Z).

290 **Bibliography**
291

Aggarwal, V. S., Carpenter, C., Freyer, L., Liao, J., Petti, M., and Morrow, B. E. (2010).
Mesodermal Tbx1 is required for patterning the proximal mandible in mice. *Dev. Biol.*
344, 669–681. doi:10.1016/j.ydbio.2010.05.496.
Anthwal, N., Joshi, L., and Tucker, A. S. (2013). Evolution of the mammalian middle ear and
jaw: adaptations and novel structures. *J. Anat.* 222, 147–60. doi:10.1111/j.1469-
7580.2012.01526.x.
Anthwal, N., Peters, H., and Tucker, A. S. (2015). Species-specific modifications of mandible
shape reveal independent mechanisms for growth and initiation of the coronoid.
Evodevo 6, 35. doi:10.1186/s13227-015-0030-6.
Anthwal, N., Urban, D. J., Luo, Z.-X., Sears, K. E., and Tucker, A. S. (2017). Meckel's cartilage
breakdown offers clues to mammalian middle ear evolution. *Nat. Ecol. Evol.* 1, 0093.
doi:10.1038/s41559-017-0093.
Asahara, M., Koizumi, M., Macrini, T. E., Hand, S. J., and Archer, M. (2016). Comparative
cranial morphology in living and extinct platypuses: Feeding behavior, electroreception,
and loss of teeth. *Sci. Adv.* 2. doi:10.1126/sciadv.1601329.
Ashwell, K. W. S. (2012). Development of the hypothalamus and pituitary in platypus
(Ornithorhynchus anatinus) and short-beaked echidna (Tachyglossus aculeatus). *J. Anat.*
221, 9–20. doi:10.1111/j.1469-7580.2012.01508.x.
Benjamin, M., and Ralphs, J. R. (1998). Fibrocartilage in tendons and ligaments--an
adaptation to compressive load. *J. Anat.* 193 (Pt 4, 481–94. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/10029181%0Ahttp://www.pubmedcentral.nih.g
ov/articlerender.fcgi?artid=PMC1467873.
Bhullar, B.-A. S., Manafzadeh, A. R., Miyamae, J. A., Hoffman, E. A., Brainerd, E. L., Musinsky,
C., et al. (2019). Rolling of the jaw is essential for mammalian chewing and tribosphenic
molar function. *Nature* 566, 528–532. doi:10.1038/s41586-019-0940-x.
El Adli, J. J., and Deméré, T. A. (2015). On the Anatomy of the Temporomandibular Joint and
the Muscles That Act Upon It: Observations on the Gray Whale, *Eschrichtius robustus*.
Anat. Rec. 298, 680–690. doi:10.1002/ar.23109.
Eloy-Trinquet, S., Wang, H., Edom-Vovard, F., and Duprez, D. (2009). Fgf signaling
components are associated with muscles and tendons during limb development. *Dev.*
Dyn. 238, 1195–206. doi:10.1002/dvdy.21946.
Gill, P. G., Purnell, M. A., Crumpton, N., Brown, K. R., Gostling, N. J., Stampanoni, M., et al.
(2014). Dietary specializations and diversity in feeding ecology of the earliest stem
mammals. *Nature* 512, 303–305. doi:10.1038/nature13622.
Green, H. L. H. H. (1937). VIII—The development and morphology of the teeth of

327 Ornithorhynchus *. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 228, 367–420.
328  doi:10.1098/rstb.1937.0015.
329 Grenier, J., Teillet, M.-A., Grifone, R., Kelly, R. G., and Duprez, D. (2009). Relationship
330  between neural crest cells and cranial mesoderm during head muscle development.
331  *PLoS One* 4, e4381. doi:10.1371/journal.pone.0004381.
332 Griffiths, M. (1978). *The Biology of the Monotremes*. London: Academic Press.
333 Grifone, R., Jarry, T., Dandonneau, M., Grenier, J., Duprez, D., and Kelly, R. G. (2008).
334  Properties of branchiomeric and somite-derived muscle development in Tbx1 mutant
335  embryos. *Dev. Dyn.* 237, 3071–8. doi:10.1002/dvdy.21718.
336 Grossnickle, D. M. (2017). The evolutionary origin of jaw yaw in mammals. *Sci. Rep.* 7, 45094.
337  doi:10.1038/srep45094.
338 Gu, S., Wei, N., Yu, L., Fei, J., and Chen, Y. (2008). Shox2-deficiency leads to dysplasia and
339  ankylosis of the temporomandibular joint in mice. *Mech. Dev.* 125, 729–42.
340  doi:10.1016/j.mod.2008.04.003.
341 Habib, H., Hatta, T., Rahman, O. I. F., Yoshimura, Y., and Otani, H. (2007). Fetal jaw
342  movement affects development of articular disk in the temporomandibular joint.
343  *Congenit. Anom. (Kyoto).* 47, 53–57. doi:10.1111/j.1741-4520.2007.00143.x.
344 Herring, S. W. (2003). TMJ anatomy and animals models. *J. Musculoskelet Neuronal Interact.*
345  3, 997–1003. doi:10.1016/j.biotechadv.2011.08.021.Secreted.
346 Hinton, R. J. (2014). Genes that regulate morphogenesis and growth of the
347  temporomandibular joint: A review. *Dev. Dyn.* 243, 864–74. doi:10.1002/dvdy.24130.
348 Hinton, R. J., Jing, J., and Feng, J. Q. (2015). Genetic Influences on Temporomandibular Joint
349  Development and Growth. *Craniofacial Dev.* 115, 85–109.
350  doi:10.1016/bs.ctdb.2015.07.008.
351 Huang, A. H. (2017). Coordinated development of the limb musculoskeletal system: Tendon
352  and muscle patterning and integration with the skeleton. *Dev. Biol.* 429, 420–428.
353  doi:https://doi.org/10.1016/j.ydbio.2017.03.028.
354 Jahan, E., Matsumoto, A., Rafiq, A. M., Hashimoto, R., Inoue, T., Udagawa, J., et al. (2014).
355  Fetal jaw movement affects Ihh signaling in mandibular condylar cartilage
356  development: the possible role of Ihh as mechanotransduction mediator. *Arch. Oral*
357  *Biol.* 59, 1108–18. doi:10.1016/j.archoralbio.2014.06.009.
358 Kemp, T. S. (2005). *The Origin and Evolution of Mammals*. Oxford, UK: Oxford University
359  Press.
360 Kinumatsu, T., Shibukawa, Y., Yasuda, T., Nagayama, M., Yamada, S., Serra, R., et al. (2011).
361  TMJ development and growth require primary cilia function. *J. Dent. Res.* 90, 988–94.
362  doi:10.1177/0022034511409407.
363 Kubiak, M., Ditzel, M., Kubiak, M., and Ditzel, M. (2016). A Joint Less Ordinary: Intriguing
364  Roles for Hedgehog Signalling in the Development of the Temporomandibular Synovial
365  Joint. *J. Dev. Biol.* 4, 25. doi:10.3390/jdb4030025.
366 Lautenschlager, S., Gill, P. G., Luo, Z.-X., Fagan, M. J., and Rayfield, E. J. (2018). The role of
367  miniaturization in the evolution of the mammalian jaw and middle ear. *Nature* 561,
368  533–537. doi:10.1038/s41586-018-0521-4.
369 Lautenschlager, S., Gill, P., Luo, Z.-X., Fagan, M. J., and Rayfield, E. J. (2017). Morphological
370  evolution of the mammalian jaw adductor complex. *Biol. Rev.* 92, 1910–1940.
371  doi:10.1111/brv.12314.
372 Luo, Z.-X. (2001). A New Mammaliaform from the Early Jurassic and Evolution of Mammalian
373  Characteristics. *Science (80-.).* 292, 1535–1540. doi:10.1126/science.1058476.

Luo, Z.-X. (2011). Developmental Patterns in Mesozoic Evolution of Mammal Ears. *Annu. Rev.*
Ecol. Evol. Syst. 42, 355–380. doi:10.1146/annurev-ecolsys-032511-142302.
Luo, Z.-X., Gatesy, S. M., Jenkins, F. A., Amaral, W. W., and Shubin, N. H. (2015). Mandibular
and dental characteristics of Late Triassic mammaliaform Haramiyavia and their
ramifications for basal mammal evolution. *Proc. Natl. Acad. Sci.* 112, E7101–E7109.
doi:10.1073/pnas.1519387112.
Maier, W., and Ruf, I. (2016). Evolution of the mammalian middle ear: a historical review. *J.*
Anat. 228, 270–83. doi:10.1111/joa.12379.
Michikami, I., Fukushi, T., Honma, S., Yoshioka, S., Itoh, S., Muragaki, Y., et al. (2012). Trps1 is
necessary for normal temporomandibular joint development. *Cell Tissue Res.* 348, 131–
40. doi:10.1007/s00441-012-1372-1.
Naples, V. L. (1999). Morphology, evolution and function of feeding in the giant anteater
(Myrmecophaga tridactyla). *J. Zool.* 249, S0952836999009036.
doi:10.1017/S0952836999009036.
Nickel, J. C., Iwasaki, L. R., Gonzalez, Y. M., Gallo, L. M., and Yao, H. (2018).
Mechanobehavior and Ontogenesis of the Temporomandibular Joint. *J. Dent. Res.* 97,
1185–1192. doi:10.1177/0022034518786469.
Presley, R., and Steel, F. L. D. (1978). The pterygoid and ectopterygoid in mammals. *Anat.*
Embryol. (Berl). 154, 95–110. doi:10.1007/BF00317957.
Purcell, P., Jheon, A., Vivero, M. P., Rahimi, H., Joo, A., Klein, O. D., et al. (2012). Spry1 and
spry2 are essential for development of the temporomandibular joint. *J. Dent. Res.* 91,
387–93. doi:10.1177/0022034512438401.
Purcell, P., Joo, B. W., Hu, J. K., Tran, P. V, Calicchio, M. L., O'Connell, D. J., et al. (2009).
Temporomandibular joint formation requires two distinct hedgehog-dependent steps.
Proc. Natl. Acad. Sci. U. S. A. 106, 18297–302. doi:10.1073/pnas.0908836106.
Rismiller, P. D., and McKelvey, M. W. (2003). Body mass, age and sexual maturity in short-
beaked echidnas, Tachyglossus aculeatus. *Comp. Biochem. Physiol. Part A Mol. Integr.*
Physiol. 136, 851–865. doi:10.1016/S1095-6433(03)00225-3.
Roberts, R. R., Bobzin, L., Teng, C. S., Pal, D., Tuzon, C. T., Schweitzer, R., et al. (2019). FGF
signaling patterns cell fate at the interface between tendon and bone. *Development*
146, dev170241. doi:10.1242/dev.170241.
Santagati, F., and Rijli, F. M. (2003). Cranial neural crest and the building of the vertebrate
head. *Nat. Rev. Neurosci.* 4, 806–818. doi:10.1038/nrn1221.
Schwartz, A. G., Long, F., and Thomopoulos, S. (2015). Enthesis fibrocartilage cells originate
from a population of Hedgehog-responsive cells modulated by the loading
environment. *Development* 142, 196–206. doi:10.1242/dev.112714.
Schweitzer, R., Chyung, J. H., Murtaugh, L. C., Brent, A. E., Rosen, V., Olson, E. N., et al.
(2001). Analysis of the tendon cell fate using Scleraxis, a specific marker for tendons
and ligaments. *Development* 128, 3855–66. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/11585810.
Shibukawa, Y., Young, B., Wu, C., Yamada, S., Long, F., Pacifici, M., et al. (2007).
Temporomandibular joint formation and condyle growth require Indian hedgehog
signaling. *Dev. Dyn.* 236, 426–34. doi:10.1002/dvdy.21036.
Sprinz, R. (1964). A Note On The Mandibular Intra-Articular Disc In The Joints Of Marsupialia
And Monotremata. *Proc. Zool. Soc. London* 144, 327–337. doi:10.1088/1751-
8113/44/8/085201.
Sugimoto, Y., Takimoto, A., Hiraki, Y., and Shukunami, C. (2013). Generation and

characterization of *ScxCre* transgenic mice. *genesis* 51, 275–283.
doi:10.1002/dvg.22372.

Takechi, M., and Kuratani, S. (2010). History of studies on mammalian middle ear evolution:
a comparative morphological and developmental biology perspective. *J. Exp. Zool. B.*
Mol. Dev. Evol. 314, 417–33. doi:10.1002/jez.b.21347.

Tsai, H. P., and Holliday, C. M. (2011). Ontogeny of the Alligator Cartilago Transiliens and Its
Significance for Sauropsid Jaw Muscle Evolution. *PLoS One* 6, e24935.
doi:10.1371/journal.pone.0024935.

Tucker, A. S. (2017). Major evolutionary transitions and innovations: the tympanic middle
ear. *Philos. Trans. R. Soc. B Biol. Sci.* 372, 20150483. doi:10.1098/rstb.2015.0483.

Urban, D. J., Anthwal, N., Luo, Z.-X., Maier, J. A., Sadier, A., Tucker, A. S., et al. (2017). A new
developmental mechanism for the separation of the mammalian middle ear ossicles
from the jaw. *Proc. R. Soc. B Biol. Sci.* 284, 20162416. doi:10.1098/rspb.2016.2416.

Wang, H., Meng, J., and Wang, Y. (2019). Cretaceous fossil reveals a new pattern in
mammalian middle ear evolution. *Nature* 576, 102–105. doi:10.1038/s41586-019-1792-
0.

Wang, Y., Liu, C., Rohr, J., Liu, H., He, F., Yu, J., et al. (2011). Tissue interaction is required for
glenoid fossa development during temporomandibular joint formation. *Dev. Dyn.* 240,
2466–2473. doi:10.1002/dvdy.22748.

Watson, D. M. S. (1916). The Monotreme Skull: A Contribution to Mammalian
Morphogenesis. *Philos. Trans. R. Soc. London. Ser. B, Contain. Pap. a Biol. Character*
207, 311–374. doi:10.2307/92025.

Wilson, J., and Tucker, A. S. (2004). Fgf and Bmp signals repress the expression of Bapx1 in
the mandibular mesenchyme and control the position of the developing jaw joint. *Dev.*
Biol. 266, 138–150. doi:10.1016/j.ydbio.2003.10.012.

Yang, L., Gu, S., Ye, W., Song, Y., and Chen, Y. P. (2016). Augmented Indian hedgehog
signaling in cranial neural crest cells leads to craniofacial abnormalities and dysplastic
temporomandibular joint in mice. *Cell Tissue Res.* 364, 105–115. doi:10.1007/s00441-
015-2306-5.

Yasuda, T., Nah, H. D., Laurita, J., Kinumatsu, T., Shibukawa, Y., Shibutani, T., et al. (2012).
Muenke Syndrome Mutation, FgfR3 P244R , Causes TMJ Defects. *J. Dent. Res.* 91, 683–
689. doi:10.1177/0022034512449170.

Yoshida, T., Vivatbutsiri, P., Morriss-Kay, G., Saga, Y., and Iseki, S. (2008). Cell lineage in
mammalian craniofacial mesenchyme. *Mech. Dev.* 125, 797–808.
doi:10.1016/j.mod.2008.06.007.