**Analysis of KSHV B lymphocyte lineage tropism in human tonsil reveals efficient infection of CD138+ plasma cells**

Farizeh Aalam*, Romina Nabiee*, Jesus Ramirez Castano and Jennifer E. Totonchy‡

Chapman University School of Pharmacy, Irvine, California, USA

*authors contributed equally to the manuscript and are listed alphabetically

‡Corresponding Author

Jennifer Totonchy

Chapman University School of Pharmacy

9410 Jeronimo Road

Irvine, CA

Phone: 714-516-5438

Email: totonchy@chapman.edu

**Running Title**: B cell tropism of KSHV

**Abstract**

Despite 25 years of research, the basic virology of Kaposi Sarcoma Herpesviruses (KSHV) in B lymphocytes remains poorly understood. This study seeks to fill critical gaps in our understanding by characterizing the B lymphocyte lineage-specific tropism of KSHV. Here, we use lymphocytes derived from 40 human tonsil specimens to determine the B lymphocyte lineages targeted by KSHV early during *de novo* infection in our *ex vivo* model system. We characterize the immunological diversity of our tonsil specimens and determine that overall susceptibility of tonsil lymphocytes to KSHV infection varies substantially between donors. We demonstrate that a variety of B lymphocyte subtypes are susceptible to KSHV infection and identify CD138+ plasma cells as a rare, but highly targeted cell type for *de novo* KSHV infection. Finally, we determine that the host immunological status influences the course of *de novo* infection in B lymphocytes. These results improve our understanding of KSHV transmission and the biology of early KSHV infection in a naïve human host, and lay a foundation for further characterization of KSHV molecular virology in B lymphocyte lineages.

**Introduction**

Kaposi Sarcoma-associated Herpesvirus (KSHV/HHV-8) is a lymphotrophic gamma-herpesvirus. In addition to its role in the pathogenesis of Kaposi Sarcoma (KS) (Chang et al., 1994), KSHV infection is associated with two lymphoproliferative disorders, multicentric Castleman disease (MCD) and primary effusion lymphoma (PEL) (Cesarman et al., 1995; Soulier et al., 1995), as well as a recently characterized inflammatory disorder KSHV inflammatory cytokine syndrome (KICS) (Uldrick et al., 2010). Although KSHV-associated lymphoproliferative disorders are rare, their

incidence has not declined as HIV treatment has improved (Powles et al., 2009; Bhutani

et al., 2015) suggesting that, in contrast to KS, immune reconstitution is not sufficient to

prevent KSHV-associated lymphoproliferative disease in people living with HIV/AIDS.

Moreover, the KSHV-associated lymphoproliferative diseases are uniformly fatal with

few effective treatment options (Carbone et al., 2014).

Despite the fact that KSHV is lymphotropic and causes pathological lymphoproliferation

*in vivo*, study of *de novo* KSHV infection in B lymphocytes has historically been difficult

(Kang and Myoung, 2017). Resting peripheral B cells and many established B cell-

derived cell lines are refractory to KSHV infection (Rappocciolo et al., 2008), but several

groups have recently been successful in infecting B lymphocytes derived from human

tonsil (Hassman et al., 2011; Myoung and Ganem, 2011a; b; Bekerman et al., 2013;

Totonchy et al., 2018; Nicol et al., 2016). KSHV DNA is detectable in human saliva and

salivary transmission is thought to be the primary route of person-to-person

transmission for KSHV (Casper et al., 2007; Pauk et al., 2000; Casper et al., 2004;

Vieira et al., 1997), making the oral lymphoid tissues a likely site for the initial infection

of B lymphocytes in a naïve human host. Thus, in addition to being susceptible to *ex*

*vivo* infection, tonsil lymphocytes represent a highly relevant model for understanding

early infection events in KSHV transmission.

The existing studies of KSHV infection in tonsil-derived B cells have explored a limited

number of cell surface markers including IgM, immunoglobulin lambda light chains and

activation markers on infected cells (Hassman et al., 2011; Nicol et al., 2016; Totonchy

et al., 2018). However, no studies to date have comprehensively explored the specific B

lymphocyte lineages targeted by KSHV infection in human tonsil specimens.

In this study, we performed KSHV infection of 40 human tonsil specimens from diverse

donors and utilized lineage-defining immunological markers by flow cytometry to

establish the primary B cell lineage tropism of KSHV. Our results demonstrate that the

susceptibility of tonsil-derived B lymphocytes to *ex vivo* KHSV infection varies

substantially from donor-to-donor, and that a variety of B cell lineages are susceptible to

KSHV infection. In particular, CD138+ plasma cells are highly targeted by KSHV

infection despite the fact that they are present at low frequencies in tonsil tissue. We

demonstrate that high susceptibility of plasma cells to KSHV infection cannot entirely be

explained by the presence of the CD138 heparin sulfate proteoglycan as an attachment

factor. Finally, we demonstrate that although the baseline T cell microenvironment does

not seem to influence susceptibility of tonsil specimens to KSHV infection, manipulation

of CD4/CD8 T cell ratios can alter the targeting of specific B cell lineages by KSHV.

These results provide new insights into early events driving the establishment of KSHV

infection in the human immune system and demonstrate that alterations in

immunological status can affect the dynamics of KSHV infection in B lymphocytes.

**Materials and Methods**

*Reagents and Cell Lines.* CDw32 L cells (CRL-10680) were obtained from ATCC and

were cultured in DMEM supplemented with 20% FBS (Sigma Aldrich) and

Penicililin/Streptomycin/L-glutamine (PSG/Corning). For preparation of feeder cells

CDw32 L cells were trypsinized and resuspended in 15 ml of media in a petri dish and

irradiated with 45 Gy of X-ray radiation using a Rad-Source (RS200) irradiator.

Irradiated cells were then counted and cyropreserved until needed for experiments.

Cell-free KSHV.219 virus derived from iSLK cells (Myoung and Ganem, 2011c) was a

gift from Javier G. Ogembo (City of Hope). Human tonsil specimens were obtained from

NDRI (www.ndri.org). Human fibroblasts were derived from primary human tonsil tissue

and immortalized using HPV E6/E7 lentivirus derived from PA317 LXSN 16E6E7 cells

(ATCC CRL-2203). Antibodies for flow cytometry were from BD Biosciences and

Biolegend and are detailed below.

*Isolation of primary lymphocytes from human tonsil.* De-identified human tonsil

specimens were obtained after routine tonsillectomy by NDRI and shipped overnight on

wet ice in DMEM+PSG. All specimens were received in the laboratory less than 24

hours post-surgery and were kept at 4°C throughout the collection and transportation

process.　Lymphocytes were extracted by dissection and maceration of the tissue in

RPMI media. Lymphocyte-containing media was passed through a 40µm filter and

pelleted at 1500rpm for 5 minutes. RBC were lysed for 5 minutes in sterile RBC lysing

solution (0.15M ammonium chloride, 10mM potassium bicarbonate, 0.1M EDTA). After

dilution to 50ml with PBS, lymphocytes were counted, and pelleted. Aliquots of $5(10)^7$ to

$1(10)^8$ cells were resuspended in 1ml of freezing media containing 90% FBS and 10%

DMSO and cryopreserved until needed for experiments.

*Infection of primary lymphocytes with KSHV.*　Lymphocytes were thawed rapidly at

37°C, diluted dropwise to 5ml with RPMI and pelleted. Pellets were resuspended in 1ml

RPMI+20%FBS+100µg/ml DNaseI+ Primocin 100µg/ml and allowed to recover in a low-

binding 24 well plate for 2 hours at 37°C, 5% $CO_2$. After recovery, total lymphocytes

were counted and Naïve B cells were isolated using Mojosort™ Naïve B cell isolation

beads (Biolegend 480068) or Naïve B cell Isolation Kit II (Miltenyi 130-091-150)

according to manufacturer instructions. Bound cells (non-naïve B and other

lymphocytes) were retained and kept at 37°C in RPMI+20% FBS+ Primocin 100μg/ml

during the initial infection process. 1(10)$^6$ Isolated naïve B cells were infected with iSLK-

derived KSHV.219 (dose equivalent to the ID20 at 3dpi on human fibroblasts) or Mock

infected in 400ul of total of virus + serum free RPMI in 12x75mm round bottom tubes via

spinoculation at 1000rpm for 30 minutes at 4°C followed by incubation at 37°C for an

additional 30 minutes. Following infection, cells were plated on irradiated CDW32 feeder

cells in a 48 well plate, reserved bound cell fractions were added back to the infected

cell cultures, and FBS and Primocin (Invivogen) were added to final concentrations of

20% and 100μg/ml, respectively. Cultures were incubated at 37°C, 5% $CO_2$ for the

duration of the experiment. At 3 days post-infection, cells were harvested for analysis by

flow cytometry.

*Flow cytometry staining and analysis of KSHV infected tonsil lymphocytes.* A proportion

of lymphocyte cultures at baseline or 3dpi representing ~500,000 cells were pelleted at

1400 rpm for 3 minutes into 96-well round bottom plates. Cells were resuspended in

100μl PBS containing (0.4ng/ml) fixable viability stain (BD Cat# 564406) and incubated

on ice for 15 minutes. Cells were pelleted and resuspended in 100μl cold PBS without

calcium and magnesium containing 2% FBS,0.5% BSA (FACS Block) and incubated on

ice for 10 minutes after which 100μl cold PBS containing 0.5% BSA and 0.1% Sodium

Azide (FACS Wash) was added. Cells were pelleted and resuspended in FACS Wash

containing B cell phenotype panel as follows for 15 minutes on ice: (volumes indicated

were routinely used for up to 0.5(10)^6 cells and were based on titration of the individual

antibodies on primary tonsil lymphocyte specimens) CD19-PerCPCy5.5 (2.5μl, BD

561295), CD20-PE-Cy7 (2.5ul, BD 560735), CD38-APC (10μl, BD 555462), IgD-APC-

H7 (2.5µl, BD 561305), CD138-v450 (2.5µl, BD 562098), CD27-PE (10µl BD 555441).

After incubation, 150µl FACS Wash was added and pelleted lymphocytes were washed

with a further 200µl of FACS Wash prior to being resuspended in 200µl FACS Wash for

analysis. Data was acquired on a BD FACS VERSE Flow Cytometer and analyzed

using FlowJo software. Readers should note that the BD FACS VERSE analysis

instrument lacks a 561nm laser so RFP lytic reporter expression from the KSHV.219

genome is not detectable in the PE channel. For baseline T cell frequencies 0.5e6 cells

from baseline uninfected total lymphocyte samples were stained and analyzed as above

with phenotype antibody panel as follows: CD95-APC (2µl, Biolegend 305611), CCR7-

PE (2µl, BD 566742), CD28-PE Cy7 (2µl, Biolegend 302925), CD45RO-FITC (3µl,

Biolegend 304204), CD45RA-PerCP Cy5.5 (2µl, 304121), CD4-APC H7 (2µl, BD

560158), CD19-V510 (3µl, BD 562953), CD8-V450 (2.5µl, BD 561426)

*KSHV neutralization via soluble Syndecan-1.* Infections were performed as described

above except KSHV.219 virus was pre-incubated for 30 minutes on ice with serum free

RPMI only or serum free RPMI containing recombinant human syndecan-1 protein

(rhCD138, BioVision, 7879-10) prior to being added to Naïve B lymphocytes. rhCD138

concentrations noted in the text indicate the final concentration of recombinant

syndecan-1 in the reconstituted total lymphocyte culture. Infection was analyzed at 3

days post-infection by flow cytometry for B cell lineages and KSHV infection as detailed

above. For experiments involving human fibroblasts virus was added to cells in serum

free media, cells were spinoculated for 30 minutes at 1000rpm, incubated at 37°C for 1

hour, then infection media was removed and replaced with complete media. At 3dpi

cells were harvested via trypsinization and analyzed for infection by flow cytometry.

*T cell depletion studies.* Infections were performed as described above except a sub-

population of total lymphocytes were depleted of either CD4 or CD8 T cells using

positive selection magnetic beads (MojoSort™ Human CD4 T Cell Isolation Kit

Cat#480009, MojoSort™ Human CD8 T Cell Isolation Kit Cat#480011). The resulting

depleted fractions or unmanipulated total lymphocytes were used to reconstitute naïve

B lymphocytes following infection rather than bound lymphocyte fractions as described

above.

*Statistical Analysis.* Data plots and statistical analysis were performed in R

software{RAlanguageanden:wf} using ggplot2{ggplotElegantGra:tn}

ggcorrplot{ggcorrplot:2018tg} and RColorBrewer (Neuwirth) packages. Specific

methods of statistical analysis and resulting values for significance and correlation are

detailed in the corresponding figure legends.

**Results**

**Variability of B cell lineages in human tonsil specimens.**

In order to explore the B cell lineages targeted by KSHV infection in human tonsil, we

procured a cohort of 40 de-identified human tonsil specimens from donors of diverse

age, sex and self-reported race (Figure 1A). Analysis of the baseline frequencies of

individual B cell lineages by multi-color flow cytometry (Supplemental Figure 1A)

revealed that the composition of individual human tonsil specimens is highly variable

(Figure 1B). This variation was independent of donor age for many lineages. However,

overall B cell frequencies declined with age as did germinal center, plasmablast and

transitional B cell populations while memory and naïve populations increased in

frequency with donor age (Figure 1C).

**Variable susceptibility of tonsil-derived B cells to *ex vivo* KSHV infection.**

Because of the heterogeneous nature of the samples, we predicted that each sample

would also have variable levels of susceptible B cell subtypes. Therefore, we employed

a method for normalizing infectious dose from donor-to-donor in order to obtain cross-

sectional data that was directly comparable (Figure 2A). For each sample we used

magnetic sorting to isolate untouched naïve B cells, which are a known susceptible cell

type (Totonchy et al., 2018), and infected 1 million naïve B cells with equivalent doses

of cell-free KSHV.219 virus. After infection, bound lymphocytes from the magnetic

separation were added back to each sample to reconstitute the total lymphocyte

environment. Infected cultures were incubated for three days and analyzed for both B

cell lineage markers and the GFP reporter present in the KSHV.219 genome to identify

infected cells. Overall, susceptibility of B cells to KSHV infection varied substantially

within the cohort (Figure 2B), but was not significantly correlated with age, sex or race

(Figure 2C).

**Specific targeting of individual B cell lineages by KSHV infection**

We next sought to establish the B lymphocyte tropism of KSHV in the human tonsil by

determining which B cell lineages are targeted for KSHV infection at early timepoints.

Because levels of individual B cell lineages are highly variable between samples (Figure

1B), we represented lineage-specific susceptibility data as the percentage of GFP+ cells

within each lineage so that data could be directly compared cross-sectionally within the

sample cohort. Our analysis of specific B cell lineages targeted for infection by KSHV

revealed that, although they represent a small proportion of the B cells within human

tonsil (Figure 1B), CD138+ plasma cells are infected at high frequencies at this early

timepoint. Indeed, CD19+ CD20- plasma cells displayed the highest within-lineage

susceptibility of any cell type with several replicates showing 100% infection of this

lineage at 3 dpi (Figure 3A). Other B cell lineages were susceptible to infection, but

were infected at relatively low within-population frequencies compared to plasma cells

(Figure 3A&B). Most B cell lineages showed linear correlation between within-lineage

infection and overall infection, while others like plasmablast, double negative, and

CD20- plasma cells showed no correlation to overall infection (Figure 3B). Pairwise

correlations revealed no significant effect of the baseline (pre-infection) frequency of

any B cell lineage on the susceptibility of that lineage to KSHV infection, indicating that

the B lymphocyte tropism of KSHV is dependent on cell-intrinsic factors and is not

simply a function of lineage population frequency (Figure 3C).

Taken together, our data demonstrate that (1) tonsil lymphocyte specimens are

immunologically heterogeneous, (2) susceptibility of tonsil lymphocytes to *ex vivo* KSHV

infection is highly donor-dependent and (3) diverse B cell lineages are specifically

targeted by KSHV early in infection. In particular, these data identify CD20- CD138+

plasma cells as a uniquely susceptible cell type for early KSHV infection in human

tonsil.

**CD138 as an attachment factor for KSHV infection of plasma cells**

Previous studies have shown that heparin sulfate proteoglycans (HSPG) of the

syndecan family can serve as an attachment factor facilitating KSHV entry via

interaction with the gH/gL glycoprotein complex (Hahn et al., 2009). In order to test

whether the high susceptibility of tonsil-derived plasma cells was due to increased

attachment via CD138 (syndecan-1), we attempted to selectively neutralize KSHV entry

by pre-treating cell-free virus particles with soluble recombinant CD138 (srCD138)

protein prior to infection. These experiments revealed a small, reproducible decrease in

overall KSHV infection of B lymphocytes with srCD138 treatment (Figure 4A). B cell

lineage analysis revealed decreased infection of plasma cell lineages in some samples,

but this decrease was highly variable with only 4 of 8 replicates showing a decrease.

KSHV infection of other B cell lineages was largely unaffected by srCD138 treatment

(Figure 4B). Moreover, KSHV infection of human fibroblasts was slightly decreased by

srCD138 treatment in 4 of 6 replicates (Figure 4C). Taken together, these results

suggest that, although srCD138 seems to weakly inhibit KSHV infection overall, the

effect is not B cell specific. Although plasma cell infection was inhibited by srCD138 in

some samples, the inconsistency between samples indicates that other factors exist

which influence plasma cell targeting and the high susceptibility of plasma cells in our

infection model cannot be explained by the use of plasma cell-expressed CD138 as an

attachment factor for KSHV entry.

**Immune status alters KSHV infection of B lymphocytes**

KSHV    lymphoproliferative    disorders    occur    primarily    in    the    context    of

immunosuppression, and other studies have shown interactions between T cells and

KSHV infected B cells in tonsil lymphocyte cultures (Myoung and Ganem, 2011a).

Therefore, we wanted to determine whether the immunological composition of the tonsil

lymphocyte environment would affect the establishment of KSHV infection in B

lymphocytes and specifically whether overall susceptibility or targeting of particular B

cell lineages is influenced by the presence or absence of T cells. Like B cell lineages,

levels of CD4+ and CD8+ T cell lineages vary considerably between tonsil donors

(Supplemental Figure 2A). However, unlike B lymphocyte subsets, the distribution of T

cell subsets are not generally correlated with donor age (Supplemental Figure 2B).

Moreover, CD4/CD8 T cell ratios are not correlated with donor age or overall

susceptibility of tonsil B lymphocytes to KSHV infection (Supplemental Figure 2C).

To determine whether manipulating the T cell environment would affect KSHV infection

in individual tonsil samples, we chose 11 samples with varying CD4/CD8 T cell ratios

and performed KSHV infections in which total lymphocytes, CD4-depleted total

lymphocytes or CD8-depleted total lymphocytes were added back following infection of

sorted naïve B cells. At 3dpi, we validated T cell depletions (data not shown) and

analyzed KSHV infection of B lymphocytes. Aggregated results for 11 individual tonsil

specimens show that neither CD4 nor CD8 depletion significantly altered overall KSHV

infection in tonsil-derived B cells (Figure 5A). However, due to the heterogeneous

nature of our tonsil samples (Figure 1B & Supplemental Figure 2A), we hypothesized

that the baseline T cell composition of each sample might influence the effect of T cell

depletion on KSHV infection. Indeed, when the change in KSHV infection in depleted

fractions is plotted against the baseline CD4/CD8 T cell ratio, we observe that KSHV

infection increased when depletions were performed in samples with high baseline

levels of CD4+ T cells (Figure 5B). We next analyzed the effect of T cell depletion on

KSHV infection of specific B cell lineages (Figure 5C). These data reveal that depletion

of CD4+ T cells increases infection of plasma cells as well as MZ-like and Transitional B

cell lineages and that the effect is dependent upon the baseline CD4/CD8 T cell ratio in

the sample with CD4+ T cell-rich samples showing the greatest effect. Interestingly,

CD8 depletions altered infection of different lineages compared to CD4 depletions, but

showed a similar dependence on the baseline level of CD4+ T cells. Taken together,

these results suggest that T cells influence the dynamics of KSHV infection in the B

lymphocyte compartment.

**Discussion**

*Ex vivo* infection of tonsil lymphocytes is emerging as a viable strategy for studying

early infection events for KSHV infection in B lymphocytes. However, the existing

literature on KSHV infection of tonsil lymphocytes is highly varied in both approach and

outcome. Hassman et. al. used cell free wild-type BCBL-1 derived KSHV virions to

infect total CD19+ B lymphocytes from tonsil specimens and used staining for LANA as

the only marker for infection, thus limiting their analysis to latently infected cells

(Hassman et al., 2011). Bekerman et. al. also used isolated CD19+ B cells as infection

targets, but employed a co-culture infection procedure using iSLK cells infected with the

recombinant KSHV.219 strain employing the GFP reporter as a marker for infection.

Nicol et. al. also employed a co-culture method to infect tonsil lymphocytes with

KSHV.219 but used Vero cells as producers and did not isolate CD19+ B cells prior to

infection. In two studies, Myoung et. al. used cell free KSHV.219 produced from Vero

cells to infect mixed lymphocyte cultures. With the exception of Bekerman et. al, all of

the above-mentioned studies employed some kind of activating agent (PHA stimulation

or CD40L stimulation) to manipulate the activation and/or proliferation of cells *in vitro*.

For our studies, we used cell-free, iSLK-derived KSHV.219 to infect Naïve B

lymphocytes followed by reconstitution of the total lymphocyte environment. We also

avoided activation of lymphocytes in both the isolation and culture procedures using our

previously-characterized CDw32 feeder cell system (Totonchy et al., 2018). To date, no

consensus has yet emerged on how to perform tonsil lymphocyte infection studies with

KSHV, and how differences in infection and culture procedure influences the resulting

data remains to be established.

Although previous studies in mixed lymphocyte cultures have explored limited surface

markers including immunoglobulins and activation markers (Nicol et al., 2016; Hassman

et al., 2011) and NK cell ligands (Bekerman et al., 2013), these studies essentially

treated all B cells as one population. Our current study is the first to use a

comprehensive panel of lineage-defining cell surface markers to carefully explore the

lineage-specificity of KSHV infection in B lymphocytes.

Our finding that KSHV efficiently targets CD138+ plasma cells early in infection of tonsil

B lymphocytes is particularly intriguing and relevant in the context of KSHV-mediated

lymphoproliferative diseases, which often have a plasma cell or plasmablast-like

phenotype (Chadburn et al., 2008; Carbone et al., 2010). Particularly for PEL, which

uniformly presents as a clonal CD138+ neoplasm, these results suggest that the

pathological cells may not be derived from KSHV-driven differentiation from less mature

lineages, but instead could be the result of modifications of terminally differentiated

plasma cells by direct infection. Recent studies have revealed intriguing interactions

between KSHV virology and cellular mediators of the unfolded protein response (UPR)

(Hu et al., 2016). The fact that the UPR is uniformly active in immunoglobulin-producing

cells, like plasma cells, suggests that these cells may provide unique advantages for

KSHV persistence. Certainly, our results highlight the virology of KSHV in primary

plasma cells as an area urgently requiring further study.

In this study, we were unable to establish that targeting of plasma cells was due to

enhanced virion binding via gH/gL interaction with the CD138 HSPG molecule, as was

suggested by a previous study (Hahn et al., 2009). However, we acknowledge that

selectively blocking virion binding to a specific HSPG in a primary cell system is

technically difficult, and we have no way to directly verify that our neutralization of gH/gL

binding sites using soluble CD138 was effective. Thus, the mechanisms underlying

KSHV targeting of plasma cells and other B lymphocyte lineages for infection remains to

be established. Studies in lymphoma cell lines have identified ephrin receptors as

critical factors in KSHV entry (Großkopf et al., 2019; Muniraju et al., 2019). Thus, our

future studies in this area will concentrate on the role of other heparin sulfate

proteoglycans and Ephrin family receptors in KSHV infection of tonsillar B lymphocyte

lineages.

Our characterization of immunological diversity of a large cohort of human tonsil

specimens will be of interest to the general immunology community. Although a few

studies have examined T cell (Petra et al., 2015) or B cell (Varon et al., 2017) subsets in

tonsil associated with particular disease states. To our knowledge, only one other study

has used multicolor flow cytometry to examine the immunological composition of both B

cells and T cells in a large cohort of human tonsil samples (Stanisce et al., 2018), and

this study was focused on comparing the microenvironments present in matched tonsils

and adenoids rather than comparison between donors based upon demographic data

as we have done here.

Our exploration of the impact of the T cell microenvironment on KSHV infection of B

lymphocytes presented in this study complements previous results from Myoung et. al.

showing that CD4+ T cells control lytic reactivation of KSHV in primary tonsillar B cells

(Myoung and Ganem, 2011a). We make the observation that manipulating the T cell

composition has a more profound effect on KSHV infection in tonsil specimens that

were CD4+ T cell rich at baseline. Moreover, although the specific B cell lineages

affected by depletion was different depending on whether CD4+ or CD8+ T cells were

experimentally depleted, the greater effect in T cell rich samples was consistent. Thus,

our data reveal that KSHV infection of B cells is sensitive to the overall immunological

microenvironment and highlight that there is complexity to this relationship that cannot

be adequately understood in the context of the current study. Future studies are

warranted to more carefully explore the contribution of donor-specific and context-

specific immunology to KSHV infection in tonsillar lymphocytes.

**Figure Legends**

**Figure 1: Variablility and age-dependence of B lymphocyte lineage distribution in**

**human tonsil.** (A) Donor demographics for human tonsil specimens used in this study.

(B) Frequency distributions of B cell lineages in tonsil specimens (see supplemental

figure 1A for lineage definitions) (C) Frequency of B cell lineages based on donor age.

Pearson correlation coefficients (r) with an absolute value greater than or equal to 0.4

are shown as red or blue inset text in the subset panels.

**Figure 2: Tonsil-derived B lymphocytes from diverse donors display variable**

**susceptibility to KSHV infection.** (A) Schematic of lymphocyte infection procedure.

(B) Frequency distribution for viable, CD19+GFP+ lymphocytes at 3dpi in 40 tonsil

specimens. (C) Infection frequency data as in (B) displayed with respect to donor age

(y-axis), sex (color), and self-reported race (shape).

**Figure 3: B lymphocyte lineage tropism of KSHV.** Within-lineage infection frequency

as a function of overall B cell infection frequency at 3dpi in for 40 tonsil specimens

shown (A) normalized to 100% or (B) scaled to each individual lineage population.

Pearson correlation coefficients (r) with an absolute value greater than or equal to 0.4

are shown in (B) as red inset text in the subset panels. (C) Correlation matrix showing

linear relationships (Pearson's correlation coefficient) between within-lineage infection

frequency and baseline (pre-infection) overall frequency of B lymphocytes and their

subsets.

**Figure 4: CD138 as an attachment factor for KSHV entry in plasma cells.** (A)

Purified KSHV virions were pre-treated with srCD138 protein and used for infection of B

lymphocytes. Viable, CD19+GFP+ lymphocytes were quantitated by flow cytometry at

3dpi. 8 experimental replicates of 5 unique tonsil specimens are shown where the

average infection rate was 1.8±0.5% in untreated controls. Data is represented as

change in GFP+ cells at each dose of srCD138 compared to untreated control and red

diamonds are the mean change at each dose. (B) Data as in (A) for within-subset GFP

quantitation. (C) Data as in (A) with E6/E7 transformed fibroblasts derived from human

tonsil as target cell type. 6 experimental replicates are shown where the average

infection rate was 25.5±16% in untreated controls.

**Figure 5: Manipulation of T cell microenvironment alters KSHV tropism for B cell**

**lineages.** Naïve B lymphocytes from 11 human tonsil specimens were infected with

KSHV.219 as in Figure 2A, total lymphocytes added back following infection were either

untreated or depleted of CD4+ or CD8+ T cells. Infections were analyzed at 3 days. (A)

Change in GFP+ B lymphocytes in CD4 or CD8 depleted samples compared to

controls. Colors indicate individual tonsil specimens. (B) infection data as in (A) plotted

against the baseline CD4/CD8 T cell ratio in the sample. (C) lineage-specific infection

data plotted against baseline CD4/CD8 T cell ratio. For B and C linear regression of the

data is shown as a blue line and Pearson correlation coefficients (r) are shown as red

text within panels only for r with an absolute value ≥ 0.6.

**Supplemental Figure 1: Gating schemes tonsil lymphocyte lineages.** Flow

cytometry data for a baseline uninfected sample from a 2-year-old male donor showing

representative gating and lineage definitions used in the study for (A) B cell lineages

based on (van Zelm et al., 2007) and (B) T cell lineages based on (Mahnke et al.,

2013).

**Supplemental Figure 2: Characterization of baseline T cell microenvironment in**

**human tonsils. (**A) Frequency distributions of T cell lineages in tonsil specimens (see

Supplemental Figure 1B for lineage definitions) (B) Frequency of T cell lineages based

on donor age. Pearson correlation coefficients (r) with an absolute value greater than or

equal to 0.4 are shown as red or blue inset text in the subset panels. (C) Correlations of

Donor age (left) or B lymphocyte susceptibility (right) plotted against sample baseline

CD4+/CD8+ T cell ratios.

**References**

Bekerman, E., D. Jeon, M. Ardolino, and L. Coscoy. 2013. A role for host activation-
induced cytidine deaminase in innate immune defense against KSHV. *PLoS Pathog.*
9:e1003748. doi:10.1371/journal.ppat.1003748.

Bhutani, M., M.N. Polizzotto, T.S. Uldrick, and R. Yarchoan. 2015. Kaposi Sarcoma–
Associated Herpesvirus-Associated Malignancies_ Epidemiology, Pathogenesis,
and Advances in Treatment. *Seminars in Oncology*. 42:223–246.
doi:10.1053/j.seminoncol.2014.12.027.

Carbone, A., E. Cesarman, A. Gloghini, and H.G. Drexler. 2010. Understanding
pathogenetic aspects and clinical presentation of primary effusion lymphoma

through its derived cell lines. *AIDS*. 24:479–490. doi:10.1097/QAD.0b013e3283365395.

Carbone, A., E. Vaccher, A. Gloghini, L. Pantanowitz, A. Abayomi, P. De Paoli, and S. Franceschi. 2014. Diagnosis and management of lymphomas and other cancers in HIV-infected patients. *Nat Rev Clin Oncol*. 11:223–238. doi:10.1038/nrclinonc.2014.31.

Casper, C., E. Krantz, S. Selke, S.R. Kuntz, J. Wang, M.-L. Huang, J.S. Pauk, L. Corey, and A. Wald. 2007. Frequent and asymptomatic oropharyngeal shedding of human herpesvirus 8 among immunocompetent men. *J. Infect. Dis.* 195:30–36. doi:10.1086/509621.

Casper, C., M. Redman, M.-L. Huang, J. Pauk, T.M. Lampinen, S.E. Hawes, C.W. Critchlow, R.A. Morrow, L. Corey, N. Kiviat, and A. Wald. 2004. HIV infection and human herpesvirus-8 oral shedding among men who have sex with men. *J. Acquir. Immune Defic. Syndr.* 35:233–238.

Cesarman, E., Y. Chang, P.S. Moore, J.W. Said, and D.M. Knowles. 1995. Kaposi's sarcoma-associated herpesvirus-like DNA sequences in AIDS-related body-cavity-based lymphomas. *N. Engl. J. Med.* 332:1186–1191. doi:10.1056/NEJM199505043321802.

Chadburn, A., E.M. Hyjek, W. Tam, Y. Liu, T. Rengifo, E. Cesarman, and D.M. Knowles. 2008. Immunophenotypic analysis of the Kaposi sarcoma herpesvirus (KSHV; HHV-8)-infected B cells in HIV+ multicentric Castleman disease (MCD). *Histopathology*. 53:513–524. doi:10.1111/j.1365-2559.2008.03144.x.

Chang, Y., E. Cesarman, M.S. Pessin, F. Lee, J. Culpepper, D.M. Knowles, and P.S. Moore. 1994. Identification of herpesvirus-like DNA sequences in AIDS-associated Kaposi's sarcoma. *Science*. 266:1865–1869.

Großkopf, A.K., S. Schlagowski, B.F. Hörnich, T. Fricke, R.C. Desrosiers, and A.S. Hahn. 2019. EphA7 Functions as Receptor on BJAB Cells for Cell-to-Cell Transmission of the Kaposi's Sarcoma-Associated Herpesvirus and for Cell-Free Infection by the Related Rhesus Monkey Rhadinovirus. *J Virol*. 93:1029. doi:10.1128/JVI.00064-19.

Hahn, A., A. Birkmann, E. Wies, D. Dorer, K. Mahr, M. Stürzl, F. Titgemeyer, and F. Neipel. 2009. Kaposi's sarcoma-associated herpesvirus gH/gL: glycoprotein export and interaction with cellular receptors. *J Virol*. 83:396–407. doi:10.1128/JVI.01170-08.

Hassman, L.M., T.J. Ellison, and D.H. Kedes. 2011. KSHV infects a subset of human tonsillar B cells, driving proliferation and plasmablast differentiation. *J Clin Invest*. 121:752–768. doi:10.1172/JCI44185.

Hu, D., V. Wang, M. Yang, S. Abdullah, D.A. Davis, T.S. Uldrick, M.N. Polizzotto, R.P.
Veeranna, S. Pittaluga, G. Tosato, and R. Yarchoan. 2016. Induction of Kaposi's
Sarcoma-Associated Herpesvirus-Encoded Viral Interleukin-6 by X-Box Binding
Protein 1. *J Virol.* 90:368–378. doi:10.1128/JVI.01192-15.

Kang, S., and J. Myoung. 2017. Primary lymphocyte infection models for KSHV and its
putative tumorigenesis mechanisms in B cell lymphomas. *J. Microbiol.* 55:319–329.
doi:10.1007/s12275-017-7075-2.

Mahnke, Y.D., T.M. Brodie, F. Sallusto, M. Roederer, and E. Lugli. 2013. The who's who
of T-cell differentiation: human memory T-cell subsets. *Eur. J. Immunol.* 43:2797–
2809. doi:10.1002/eji.201343751.

Muniraju, M., L.Z. Mutsvunguma, J. Foley, G.M. Escalante, E. Rodriguez, R. Nabiee, J.
Totonchy, D.H. Mulama, J. Nyagol, F. Wussow, A.K. Barasa, M. Brehm, and J.G.
Ogembo. 2019. Kaposi Sarcoma-Associated Herpesvirus Glycoprotein H Is
Indispensable for Infection of Epithelial, Endothelial, and Fibroblast Cell Types. *J*
*Virol.* 93:140. doi:10.1128/JVI.00630-19.

Myoung, J., and D. Ganem. 2011a. Active lytic infection of human primary tonsillar B
cells by KSHV and its noncytolytic control by activated CD4+ T cells. *J Clin Invest.*
121:1130–1140. doi:10.1172/JCI43755.

Myoung, J., and D. Ganem. 2011b. Infection of primary human tonsillar lymphoid cells
by KSHV reveals frequent but abortive infection of T cells. *Virology.* 413:1–11.
doi:10.1016/j.virol.2010.12.036.

Myoung, J., and D. Ganem. 2011c. Generation of a doxycycline-inducible KSHV
producer cell line of endothelial origin: maintenance of tight latency with efficient
reactivation upon induction. *J. Virol. Methods.* 174:12–21.
doi:10.1016/j.jviromet.2011.03.012.

Neuwirth, E. RColorBrewer: ColorBrewer Palettes.

Nicol, S.M., S. Sabbah, K.F. Brulois, J.U. Jung, A.I. Bell, and A.D. Hislop. 2016. Primary
B Lymphocytes Infected with Kaposi's Sarcoma-Associated Herpesvirus Can Be
Expanded In Vitro and Are Recognized by LANA-Specific CD4+ T Cells. *J Virol.*
90:3849–3859. doi:10.1128/JVI.02377-15.

Pauk, J., M.L. Huang, S.J. Brodie, A. Wald, D.M. Koelle, T. Schacker, C. Celum, S.
Selke, and L. Corey. 2000. Mucosal shedding of human herpesvirus 8 in men. *N.*
*Engl. J. Med.* 343:1369–1377. doi:10.1056/NEJM200011093431904.

Petra, D., K. Petra, K. Michaela, K. Daniela, H. Petr, K. Ladislav, K. Rami, H. Ondrej, K.
Zdenek, D. Pavla, K. Tomas, and F. Eva. 2015. Polyclonal, newly derived T cells
with low expression of inhibitory molecule PD-1 in tonsils define the phenotype of
lymphocytes in children with Periodic Fever, Aphtous Stomatitis, Pharyngitis and

469 Adenitis (PFAPA) syndrome. *Molecular Immunology*. 65:139–147.
470 doi:10.1016/j.molimm.2015.01.004.

471 Powles, T., D. Robinson, J. Stebbing, J. Shamash, M. Nelson, B. Gazzard, S. Mandelia,
472 H. Moller, and M. Bower. 2009. Highly Active Antiretroviral Therapy and the
473 Incidence of Non-AIDS-Defining Cancers in People With HIV Infection. *Journal of
474 Clinical Oncology*. 27:884–890. doi:10.1200/JCO.2008.19.6626.

475 Rappocciolo, G., H.R. Hensler, M. Jais, T.A. Reinhart, A. Pegu, F.J. Jenkins, and C.R.
476 Rinaldo. 2008. Human herpesvirus 8 infects and replicates in primary cultures of
477 activated B lymphocytes through DC-SIGN. *J Virol*. 82:4793–4806.
478 doi:10.1128/JVI.01587-07.

479 Soulier, J., L. Grollet, E. Oksenhendler, P. Cacoub, D. Cazals-Hatem, P. Babinet, M.F.
480 d'Agay, J.P. Clauvel, M. Raphael, and L. Degos. 1995. Kaposi"s sarcoma-
481 associated herpesvirus-like DNA sequences in multicentric Castleman"s disease.
482 *Blood*. 86:1276–1280.

483 Stanisce, L., E. Sims, C. Hou, Y. Koshkareva, J.P. Gaughan, I. Kuzin, and A. Bottaro.
484 2018. Differential cellular composition of human palatine and pharyngeal tonsils.
485 *Archives of Oral Biology*. 96:80–86. doi:10.1016/j.archoralbio.2018.08.020.

486 Totonchy, J., J.M. Osborn, A. Chadburn, R. Nabiee, L. Argueta, G. Mikita, and E.
487 Cesarman. 2018. KSHV induces immunoglobulin rearrangements in mature B
488 lymphocytes. *PLoS Pathog*. 14:e1006967. doi:10.1371/journal.ppat.1006967.

489 Uldrick, T.S., V. Wang, D. O'Mahony, K. Aleman, K.M. Wyvill, V. Marshall, S.M.
490 Steinberg, S. Pittaluga, I. Maric, D. Whitby, G. Tosato, R.F. Little, and R. Yarchoan.
491 2010. An interleukin-6-related systemic inflammatory syndrome in patients co-
492 infected with Kaposi sarcoma-associated herpesvirus and HIV but without
493 Multicentric Castleman disease. *CLIN INFECT DIS*. 51:350–358.
494 doi:10.1086/654798.

495 van Zelm, M.C., T. Szczepanski, M. van der Burg, and J.J.M. van Dongen. 2007.
496 Replication history of B lymphocytes reveals homeostatic proliferation and extensive
497 antigen-induced B cell expansion. *J Exp Med*. 204:645–655.
498 doi:10.1084/jem.20060964.

499 Varon, L.S., J. De Rosa, A.X.S. Machicote, L.A. Billordo, P.X.C. Baz, P.M.F.X. ndez,
500 I.K. Maciel, A.X.S. Blanco, and E.X.S.I. Arana. 2017. Characterization of tonsillar
501 IL10 secreting B cells and their role in the pathophysiology of tonsillar hypertrophy.
502 *Sci. Rep*. 1–9. doi:10.1038/s41598-017-09689-x.

503 Vieira, J., M.L. Huang, D.M. Koelle, and L. Corey. 1997. Transmissible Kaposi"s
504 sarcoma-associated herpesvirus (human herpesvirus 8) in saliva of men with a
505 history of Kaposi"s sarcoma. *J Virol*. 71:7083–7087.