Feral pigeon populations: their gene pool and links with local domestic breeds

Dimitri Giunchi[1, *, **], Nadia Mucci[2, *, **], Daniele Bigi[3], Chiara Mengoni[2], N. Emilio Baldaccini[1]

[1] Dipartimento di Biologia, Università di Pisa, Via Volta 6, 56126 Pisa

[2]Area per la Genetica della Conservazione (BIO-CGE), Dipartimento per il monitoraggio e la tutela dell'ambiente e per

la conservazione della biodiversità, Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA), Via Cà

Fornacetta, 9 40064 Ozzano dell'Emilia, Italy

[3] Department of Agricultural and Food Science (DISTAL), University of Bologna, Viale G. Fanin 46, 40127 Bologna,

Italy

* Corresponding authors, email: dimitri.giunchi@unipi.it, nadia.mucci@isprambiente.it

** D. Giunchi and N. Mucci contributed equally to the manuscript.

Abstract

Columba livia is a wild bird whose domestication has led to a large number of pigeon breeds. The occasional loss or

straying of domestic birds determined the origin of feral pigeons, which are now widespread all around the world. In

this study, we assumed that the main contribution to feral populations is provided by domestic breeds reared in the same

areas. We tested this hypothesis by analysing the variability of 12 microsatellite loci in nine Italian feral populations

sampled in areas with different intensities of breeding and selecting domestic breeds. We included in the analysis

samples belonging to domestic lineages commonly bred in Italy. Feral populations located in regions with a long-lasting

tradition of pigeon breeding showed a high level of admixture with domestic breeds, in particular with Racing Homer

and Piacentino. Ferals from Bolzano, Venice and Sassari were characterized by unique genetic components, almost all of

which are not shared by other feral populations and by the considered domestic breeds. Our results further emphasize the

complex origin of feral populations which can be properly investigated only by considering the pool of domestic pigeons bred in

the considered area and their past and present distribution.

Keywords: *Columba livia*, feral pigeons, domestic pigeons, genetic structure, microsatellite loci

Acknowledgments

We thank local regional authorities for providing permits to sample feral pigeons across Italy. We also thanks Andrea

Barbon, Pierluigi Berra, Danilo Pisu, Cecilia Soldatini and the staff of the wildlife recovery centres of the following

cities for their collaboration in collecting samples: Livorno (Gianluca Bedini, Renato Ceccherelli, Nicola Maggi),

Modena (Piero Milani), Reggio-Emilia (Sergio Mazzali), Sassari (Marco Muzzeddu). Thanks are due also to the

Federazione Colombofila Italiana and to the Federazione Italiana Allevatori Colombi (FIAC) for samples of domestic

breeds and for data regarding the rearing activity of Racing Homers in Italy. Wendy Doherty edited the language of the

manuscript.

Introduction

Feral pigeons are one of the most common inhabitants of cities all around the world, being a virtually cosmopolitan taxon (Lever 1987). The Neolithic domestication of the wild Rock Dove, dating back to about 6000 years BC (Sossinka 1982) and the subsequent selection of the various pigeon domestic breeds represent the initial steps of their origin. Indeed, feral pigeons originated from domestic pigeons released into the urban environment or escaped from farms (Johnston and Janiga 1995). This process began in the Old World and it is still ongoing in almost every place where domestic pigeons were introduced or bred (Lever 1987; Johnston 1994). Synanthropic wild Rock Doves seem to have contributed only marginally to the constitution of feral populations (Ballarini et al. 1989; Johnston and Janiga 1995).

The different ways pigeons established themselves in European and North-American urban habitats have been reviewed from a historical point of view by (Johnston and Janiga 1995; Haag-Wackernagel 1998; Baldaccini and Giunchi 2006). Given that nearly any domestic breed had (and still has) the potential to contribute to the feral gene pool, at least two main contributions to feral populations have been identified. Dovecotes had been rather widespread in several European countries until the XIX century [e.g. The British Isles (Ritchie 1920; Gompertz 1957); France (Van der Linden 1950); Italy (Giachetti 1914)). Pigeons breeding in dovecotes could revert to a free life in towns and sometimes they were even forced to leave their dovecotes e.g. during the French Revolution, when large dovecotes owned by aristocrats were destroyed (Van der Linden 1950). Urban dovecotes are now rare in Europe so they do not represent an important source of individuals for current feral populations. The main contribution to feral populations both in Europe and North America probably was and is still represented by homing pigeons that failed to return to their home loft (Goodwin 1960; Simms 1979; Stringham et al. 2012). In particular, (Simms 1979) suggested that juveniles homing pigeons lost at the beginning of their training period constituted an important component of feral populations at least in the UK.

In Italy, pigeon breeding has been an embedded activity since ancient times especially in Northern and Central regions, where several domestic breeds have been selected (McNeillie 1976; Bigi et al. 2016). It can be assumed that these local breeds have mostly contributed to the Italian feral populations, possibly with birds (both dovecote and wild individuals), imported mainly from Egypt (from the range of *C. l. gaddi* and *C. l. schimperi* subspecies) and Spain, used in pigeon shooting ranges till the first half of the last century (Ghigi in Toschi 1939).

In recent years, domestic pigeons have been the subject of several genetic investigations, mainly aimed at understanding their relationships and their geographic origins (Stringham et al. 2012; Shapiro et al. 2013; Biala et al. 2015; Bigi et al. 2016; Domyan and Shapiro 2017), whereas feral pigeons, on the contrary, received rather less attention. While a number of the above-mentioned studies on domestic breeds actually included one group of feral pigeons in their analyses (e.g. Stringham et al. 2012; Biala et al. 2015), only two studies were specifically focused on feral pigeons, with the aim of clarifying the pattern of genetic differentiation within and between urban areas (Jacob et al. 2015; Tang et al. 2018). To our knowledge, no study has systematically investigated the degree of influence of different domestic breeds on the genetic composition of feral populations.

In this paper, we hypothesized that the gene pool of feral pigeons living in a specific urban context should at least partially reflect the prevalent breeding activities of domestic breeds in the area, given the low dispersal propensity of these birds (Hetmanski 2007; Jacob et al. 2015). In particular, we would like to test the hypothesis put forward by (Stringham et al. 2012), that almost all feral populations should show strong affinities with racing breeds.

We compared the genetic pattern of nine Italian feral populations distributed in areas with different racing pigeon breeding

traditions with the aim of: 1) characterizing their genetic composition and structure; 2) testing their affinities with a number of

domestic pigeons commonly bred in Italy that could have contributed to the actual genetic pool (Bigi et al. 2016). For this reason,

we also included in the sampling a number of feral populations belonging to the Pianura Padana (Pavia, Reggio Emilia, Modena),

where the breeding of racing pigeons is traditionally widespread and where a significant number of Italian breeds originated

(Ghigi 1950; Bigi et al. 2016), and to Central (Pisa, Livorno) and Northeastern Italy (Venezia, Treviso, Bolzano), where racing

pigeon breeding is rather less common (Fig. 1) and to Sardinia (Sassari) where racing pigeon breeding is not present. Sardinia

possibly still hosts colonies of wild Rock Doves (Ragionieri et al. 1991; Baldaccini et al. 2000) which might have contributed to

the gene pool of feral populations in the area.

Material and Methods

A total of 194 samples were obtained from 9 urban areas (Fig. 1, Table 1) in North and Central Italy. About five

feathers were collected from each individual and conserved in ethanol 95% at -20 °C. All the procedures were

performed with the permission of the local authorities.

DNA extraction and microsatellite genotyping

DNA was extracted using the ZR Genomic DNA II (Zymo Research) and amplified at 12 loci microsatellite (*Cliμd01*,

Cliμt13, *Cliμd17*, *CliμT17*, *Cliμd16*, *Cliμd19* - (Traxler et al. 2000); *CliuD11*, *CliuT47*, *CliuT24*, *UUCli10*, *UUCli13*,

UUCli14n, *UUCli12*, *UUCli08* - (Stringham et al. 2012). Two independent replicates were performed in 8 μl of volume

containing 0.5 U of Hot Start Taq polymerase (Qiagen), 0.18 μM of each primer and 0.04 mg of Bovine Serum

Albumin Fraction V (Roche), with the following thermal protocol: (94 °C x 5'), 10 cycles at (94 °C x 40'') (55 °C-

0.5 °C x 40'') (72 °C x 60''), 35 cycles at (94 °C x 40'') (50 °C x 40'') (72 °C x 60''), and a final extension at 72 °C for

10'.

Data analysis

The number of alleles (*Na*), the number of effective (*Ne*) and private alleles, and the expected (*He*) and observed (*Ho*)

heterozygosity were obtained using GenAlEx 6.503 (Peakall and Smouse 2006, 2012; Smouse et al. 2015). Allelic

richness (*Ar*) was computed in Fstat to minimize the effect of a different sample size (Goudet 2001). Departure from

Hardy-Weinberg equilibrium was estimated with the exact test in Genepop on the web (Raymond and Rousset 1995)

(http://genepop.curtin.edu.au/) using 100 batches and 1000 iterations per batch.

Pairwise *Fst* computation and AMOVA test (Excoffier et al. 1992) were performed in Genetix 4.05.02 (Belkhir et al.

2002) and GenAlEx 6.503, respectively, to evaluate the significance of genetic differentiation between groups.

Genetic divergence among geographical groups was estimated using distance based on the Stepwise Mutation Model

(SSM) and the Infinite Allele Model (IAM).

Cavalli-Sforza chord distance (*Dc*) (Cavalli-Sforza and Edwards 1967) and the proportion of shared alleles (*Dps*)

(Bowcock et al. 1994) were computed in MSA (Dieringer and Schlotterer 2003) and the resulting networks were visualized in SplitTree 4.13.1 (Huson and Bryant 2006). *Da* distance (Nei 1973) and *Fst* distance (Latter 1972) with sample size bias correction, Goldestein distance $(\delta\mu)^2$ (Goldstein et al. 1995) and Shriver distance (*Dsw*) (Shriver et al. 1995) were computed and visualized in a Neighbour-Joining tree using Poptree on the web (http://poptree.med.kagawa-u.ac.jp/). A total of 10,000 bootstraps were used to reconstruct the tree topology.

Clustering was computed using a Bayesian model in Structure 2.3.4 (Pritchard et al. 2000; Falush et al. 2003; Hubisz et al. 2009) with no admixture and independent allele frequencies. A total of five independent runs for a number of subpopulations (K) from 1 to 10, was run with a burn-in period of 30,000 followed by 300,000 MCMC repetitions. Clumpp (Jakobsson and Rosenberg 2007) and Distruct (Rosenberg 2004) were used respectively to merge the five independent results for each K and to display the final data. A Discriminant Analysis of Principal Components (DAPC) that is not affected by Hardy-Weinberg disequilibrium was also performed to verify the substructure of feral populations using the Adegenet package for R software. A total of 80 PCs and four discriminant functions were retained to draw the plot.

All individuals belonging to the same urban area were characterized by identical coordinates. The hypothesis of Isolation by Distance among populations was tested in GenAlEx by means of the Mantel test calculated on the geographic distance and the *Fst* obtained by GenAlEx computation.

Relationship between feral pigeons and domestic breeds

Genotypes were aligned and compared with those obtained in (Bigi et al. 2016) to verify the affinity of feral populations with Italian domestic breeds. A total of 250 samples belonging to 10 Italian native breeds (Florentine, Italian Beauty Homer, Italian Owl, Italian Owl Rondone, Piacentino, Romagnol, Runt, Sottobanca, Triganino Schietto and Triganino Gazzo) and one international breed (Racing Homer) commonly bred in Italy, were included in the analysis (Table 2). *Dc, Dps, Da, Fst,* $(\delta\mu)^2$ and *Dsw* were computed as for the previous analysis on feral populations to identify the phylogenetic affinities. Cluster analysis was conducted in Structure using the models applied in the previous section with K ranging from 1 to 20. DAPC was calculated using the same number of PCs and discriminant functions used in the previous section. Cluster analysis and DAPC were run as reported in the previous paragraph.

After the preliminary results, five breeds (Florentine, Italian Owl, Italian Owl Rondone, Triganino Schietto and Triganino Gazzo) that did not show any genetic relationship with feral pigeon populations (ESM 1) were excluded from the analysis. The Florentine disappeared in Italy at the beginning of the last century (Giachetti 1914) and it has been reintroduced only recently, while the outdoor breeding of Italian Owls and of Triganino Schietto and Gazzo was interrupted from the first half of 20[th] century (Vaccari and Zambon 2014). Therefore only a scarce contribution of these five breeds to the present-day Italian feral populations can be expected.

Results

Genetic variability of feral populations

All samples were successfully genotyped. The descriptive statistics are listed in Table 3. The allelic number ranged

from 6.2 ± 0.5 SE (F_VE) to 8.5 ± 0.9 (F_RE) with an average value of 7.6 ± 0.8. The average number of effective

alleles decreased to 4.4 ±0.5. Mean values of expected (*He*) and observed (*Ho*) heterozygosity were 0.718 ± 0.062 and

0.726 ± 0.050, respectively. Allelic richness was computed on a minimum value of 14 samples and ranged from 5.6

(F_VE) to 7.6 (F_LE) with an average value of 6.9. A total of eight different private alleles were detected in Sardinian

ferals even if with a low frequency (< 10%).

No departure from Hardy Weinberg equilibrium was detected.

Phylogenetic trees built using *Da*, *Fst*, and *(δμ)²* distances did not show any significant structure (Fig. 2). Feral from

F_LE, F_PI, F_PV, F_MO and F_RE clustered together in all the trees, but the associated bootstrap values were so low

as to make the topology not significant. Acceptable bootstrap values were obtained only for the group constituted by

F_VE and F_TV. Additional results from other distance computations [Cavalli-Sforza chord distance (*Dc*), proportion

of shared alleles (*Dps*) and Shriver distance (*Dsw*)] were not shown as they did not produce a different structure. Nei's

genetic distances and *Fst* were not significant among F_MO, F_RE, F_LE and between F_PI and F_PV (ESM 2-3).

Mantel test evidenced a slightly significant correlation between the genetic and geographic distances ($r = 0.50$; $P = 0.03$,

ESM 4a); this correlation remained almost unchanged ($r = 0.46$; $P = 0.01$) when the Sardinian sample was removed

from the analysis (ESM 4b).

AMOVA did not result in a structured sampling with 90% of differences found within individuals and only 4% among

populations.

No defined genetic structure was identified by both multivariate and Bayesian analyses (Fig. 3a-c). DAPC analysis

showed an admixed pattern in Central Italy, whereas samples from Sassari (F_SS), Bolzano (F_BZ), and Venice (F_VE)

clustered separately from the others. The population from Treviso (F_TV) plotted in an intermediate position between

the remaining populations and F_VE.

The best value of ΔK computed in Structure Harvester identified four main clusters (Fig. 3b). Feral from Bolzano

(F_BZ), Venice (F_VE), and Sassari (F_SS) were associated with high *q* individual membership values to different

clusters while samples from F_PV, F_RE, F_MO, F_PI, and F_LE, were not differentiated and showed retraces of

admixture with F_BZ, F_VE and F_SS. The genetic composition of F_TV was mainly associated to Venice, but showed

traces of admixtures with the former five populations (Fig. 3c).

Relationship between feral pigeons and domestic breeds

Phylogenetic trees built using *Fst*, D*a* and *(δμ)²* distances did no show any significant structure (ESM 5) and feral

groups and the domestic lineages considered in the analysis were differently associated depending on the distance

computation considered. As in the previous section, Cavalli-Sforza chord distance (*Dc*), the proportion of shared alleles

(*Dps*) and Shriver distance (*Dsw*) were not informative and were not shown.

The DAPC of the nine feral populations and the six Italian breeds did not reveal a sharp separation between the two

groups (Fig. 4a). With the exception of domestic SB and RM, all the other breeds overlap with feral populations in the

plot. F_VE and F_SS did not show any tracks of an origin from the considered domestic lineages because in the plots
they did not overlap with any of them. The Bayesian analysis partially confirmed the results of DAPC and the *q*
individual membership values identified a relationship between domestic and local feral pigeons (Fig. 4c and 5). The
ΔK computation in Structure Harvester identified 8 main informative clusters (Fig. 4b). The evaluation of the estimated
membership coefficient for each population *(q)* at K = 8 identified the predominance of the domestic component
characterizing the Homers (light blue bar in Fig. 5) in feral populations from Lombardy, Emilia-Romagna and Tuscany,
particularly in F_RE, F_MO, and F_LE. Rather high percentages of domestic Piacentino (PC) and lower values of
domestic SB and RN were found in all populations, including F_BZ and F_SS, although the latter populations were
characterized by over 60% of a unique private component. The ferals from Venice (F_VE) showed a high percentage of
a unique genetic component, that was also detected in the other feral populations but that was almost irrelevant in
domestic lineages. As evidenced in Figure 5, the population of Venice did not show relevant traces of admixture with
the considered domestic populations.

Discussion

This paper represents one of the few studies dealing with the genetic structure of feral pigeon populations. Up to now,
only one study was specifically focused on clarifying the pattern of genetic differentiation of these birds between urban areas
(Jacob et al. 2015). Furthermore, while it is largely accepted that feral pigeons originated from domestic breeds (see for instance,
(Johnston and Janiga 1995), this study is the first one that systematically investigates the affinities between feral pigeon
populations and the domestic breeds commonly bred in the same area. To our knowledge only Biala et al. (2015) tried to quantify
the gene flow between several domestic breeds and feral pigeons. However, while Biala et al. (2015) used only one feral group,
composed of birds sampled in different towns, we sampled a significant number of true feral populations in order to investigate
the contribution of domestic breeds to their gene pool and how this contribution varies among populations located in different
areas.

On the whole, the populations analyzed in our study showed levels of genetic variability lower than those observed by (Jacob et
al. 2015). The low levels of variability indices which were found in the populations lacking genetic admixture allowed us to argue
that these values were influenced by the reduced incidence of gene flow among these populations.

As observed by (Jacob et al. 2015), also in our sample the genetic distance between feral populations was correlated with their
geographic distance, both considering all samples or only the peninsular ones. This result further confirms that the exchange rate
of individuals among cities is relatively rare (Johnston and Janiga 1995; Hetmanski 2007). It should be noted, however, that the
observed pattern of geographic differentiation is rather complex and only partially explained by the geographic distance between
populations. Indeed, while the present data do not support a well defined genetic structure, it is interesting to observe that Sardinia
and North-eastern populations (F_BZ, F_VE and F_TV to a lesser extent) tended to cluster separately. On the contrary,
populations belonging to Tuscany, Emilia-Romagna and Lombardy (F_PI, F_LE, F_MO, F_RE and F_PV) showed a high level
of admixtures almost independent of their geographic distance. The genetic distinction of populations like F_BZ, F_SS and F_VE,
can be attributed to their relative geographic isolation. Actually, Bolzano is located in an Alpine valley surrounded by habitats
mostly unsuitable for pigeons, while Sardinia is an island and it is known that pigeons do not like to fly over large water bodies
(Wagner et al. 1972). The effect of the surrounding water should probably be taken into account also for Venice: for instance, as
reported by (Soldatini et al. 2006), the number of Venetian pigeons involved in foraging flights outside the city is very low

considering the size of the population, which possibly confirm the low propensity of these pigeons to fly over the lagoon and thus

also to disperse inland.

However, the level of admixture of the remaining populations (F_PI, F_LE, F_MO, F_RE and F_PV) is quite difficult to explain

considering the above mentioned low rate of dispersal among cities (Johnston and Janiga 1995; Hetmanski 2007; Jacob et al.

2015). This pattern can be understood only when the domestic breeds are included in the analysis. Indeed, in these populations it

is possible to identify a significant component belonging to the domestic breeds considered in this study. In particular, the Racing

Homer and Piacentino components largely dominates the gene pool of these populations. These domestic components are still

detectable in F_SS, F_BZ, F_VE and F_TV, but at negligible percentages. This pattern can be explained by considering the long-

lasting and still ongoing tradition of keeping and selecting pigeon breeds especially in Emilia-Romagna and Lombardy (Ghigi

1950; McNeillie 1976). In particular, the Racing Homer component is quite evident mainly in populations located in areas where

Racing Homer breeding and racing is widespread (i.e. Emilia-Romagna and Lombardy, see Figure 1), while it is almost absent

where those activities are missing (i.e. Bolzano and Sardinia). Our data only partially confirm the hypothesis by Stringham et al.

(2012, but see also Goodwin 1960; Simms 1979) that Racing Homers constitute the most important component of feral pigeon

populations. Indeed, other breeds, such as Piacentino, can be dominant or co-dominant in the feral gene pool, as observed for

example in F_PI, F_PV and F_RE.

Our results further emphasize the complex origin of feral populations and suggest a past and probably ongoing flow of domestic

pigeons into feral populations in areas surrounded by a high number of pigeon fanciers. This seems to be confirmed by the

difference observed between F_VE and F_TV. These populations form a fairly separated cluster, which probably indicates that

they share a common origin. However, being located in a lagoon, Venice has no pigeon fanciers nearby and thus the genetic

contribution of the domestic breeds considered in this study to its feral population is very low. On the other hand, Treviso is

surrounded by farms that probably hosted and still host pigeon dovecotes, which facilitates the urban drift of domestic birds. It

should be noted that it is rather impossible to have detailed information regarding the actual distribution of pigeon breeding

around a given city and for this reason the above pattern is characterized by some unexplained variability that might be related to

the scale of our analysis.

The mechanisms leading to the admixture between domestic and feral pigeons have probably been and still are both pigeon

racing and feral pigeon foraging behaviour. As observed by Goodwin (1960) and Simms (1979) pigeon races are sources of

numbers of lost Racing Homers that flock together with ferals. Furthermore, the daily foraging flights of ferals towards the

surrounding crop fields (Johnston and Janiga 1995; Giunchi et al. 2012) may encourage farm dovecote individuals to join them.

As a partial support to this hypothesis, it should be noted that, birds living in cities mostly surrounded by an unsuitable foraging

habitat show both less propensity to perform foraging flight [i.e. Bolzano (Baldaccini et al. 2015) and Venice (Soldatini et al.

2006)] and a less relevant component of the studied domestic breeds in their gene pool.

Recent data on the monk parakeet *Myiopsitta monachus* and ring-necked parakeet *Psittacula krameri* suggest that a high degree

of admixture is not directly related to invasive success in an urban habitat and does not prevent the possibility of rapid adaptation

to the urban environment (Edelaar et al. 2015; Le Gros et al. 2016). In this regard, it would be interesting to a use genomic

approach in order to test whether the populations with a higher degree of admixtures, actually show higher frequencies of

phenotypic characters belonging to domestic breeds or whether the domestic phenotypes are quickly counter-selected by the

urban environment (see Johnston and Janiga 1995; Sol 2008)

As observed above, feral pigeons from Bolzano, Venice/ Treviso and Sassari were characterized by unique genetic components,

that are mainly not shared by other feral populations. Considering the geographic position of those populations, it can be

hypothesized that these unique components belong to domestic breeds not originated and/or reared in Italy (e.g. central-european

breeds for Bolzano, eastern breeds for Venice/Treviso). Concerning birds from Sassari, it should be noted that Sardinia still hosts

wild populations of Rock Doves (Ragionieri et al. 1991; Johnston 1992; Johnston and Janiga 1995). Johnston and Janiga (1995)

indicated that when wild and feral pigeons live in sympatry it is likely that they interbreed, thus it can be hypothesized that wild

pigeons may have contributed to F_SS. In this regard, it is interesting to observe that, contrary to Jacob et al. (2015) and Biala et

al. (2015), we found eight private alleles in feral samples, all belonging to the Sardinian population. Whether this is due to the

effect of domestic breeds not included in our sample and not affecting other feral populations or to the gene flow between wild

Rock Dove and feral pigeons as hypothesized by Ragionieri et al. (1991) could not be ascertained by our data and deserves

further investigations.

Our data emphasize the critical role of the sampling protocol when studying the relationship between feral pigeons and domestic

breeds. Indeed, papers studying this topic often did not consider single populations of feral pigeons, but mixed together pigeons

sampled in different cities (see e.g. Biala et al. 2015; Shao et al. 2019). Moreover the domestic breeds included in the analysis

were selected without taking into consideration the local tradition of pigeon breeding in the areas where feral pigeons were

sampled. This probably explains some of the inconsistencies in the results obtained for instance by Stringham et al. (2012) and

Shao et al. (2019), as the former suggested a strong relationship between Racing Homers and feral pigeons, which is not evident

in the latter study.

To conclude, our data provide the first detailed analysis of the relationships between feral pigeon populations and domestic

breeds, shedding light on the way feral populations originated and are probably maintained. Our results emphasize the complexity

of the feral gene pool whose composition shows high spatial variability possibly depending on both ecological and anthropic

factors. In particular, factors such as geographic isolation of feral populations along with the prevalent farming activities and the

local diversity of domestic pigeon breeds seem to play a central role in this regard. Last but not least, the first data on the

differentiation of Sardinian populations should not be overlooked because they could represent the first genetic evidence valuable

for the management and conservation of wild Rock Dove and feral populations where they occur in syntopy.

References

Baldaccini NE, Giunchi D (2006) Le popolazioni urbane di colombo: considerazioni sulla loro genesi e sulle metodologie di gestione. Biologia Ambientale 20:125–141

Baldaccini NE, Giunchi D, Mongini E, Ragionieri L (2000) Foraging flights of wild rock doves (Columba l. livia): a spatio-temporal analysis. Italian Journal of Zoology 67:371–377

Baldaccini NE, Lombardo D, Mongini E, Giunchi D (2015) I colombi della città di Bolzano: interventi di gestione e caratteristiche della popolazione

Ballarini G, Baldaccini NE, Pezza F (1989) Colombi in città. Aspetti biologici, sanitari e giuridici. Metodologie di controllo. Istituto Nazionale di Biologia della Selvaggina, Bologna

Belkhir K, Borsa P, Chikhi L, et al (2002) GENETIX 4.05, Windows TM Software for Population Genetics. Laboratory Genome, Populations, Interactions. Universite de Montpellier II, Montpellier, France.(in French)

Biala A, Dybus A, Pawlina E, Proskura WS (2015) Genetic diversity in eight pure breeds and urban form of domestic pigeon (Columba livia var. domestica) based on seven microsatellite loci. Journal of Animal and Plant

Sciences 25:1741–5

Bigi D, Mucci N, Mengoni C, et al (2016) Genetic investigation of Italian domestic pigeons increases knowledge about the long-bred history of *Columba livia* (Aves: Columbidae). Italian Journal of Zoology 83:173–182. https://doi.org/10.1080/11250003.2016.1172121

Bowcock AM, Ruiz-Linares A, Tomfohrde J, et al (1994) High resolution of human evolutionary trees with polymorphic microsatellites. Nature 368:455–457

Cavalli-Sforza LL, Edwards AW (1967) Phylogenetic analysis. Models and estimation procedures. American journal of human genetics 19:233–257

Dieringer D, Schlotterer C (2003) Microsatellite analyzer (MSA): a platform independent analysis tool for large microsatellite data sets. Molecular Ecology Notes 3:167–169

Domyan ET, Shapiro MD (2017) Pigeonetics takes flight: Evolution, development, and genetics of intraspecific variation. Developmental Biology 427:241–250. https://doi.org/10.1016/j.ydbio.2016.11.008

Edelaar P, Roques S, Hobson EA, et al (2015) Shared genetic diversity across the global invasive range of the monk parakeet suggests a common restricted geographic origin and the possibility of convergent selection. Molecular Ecology 24:2164–2176. https://doi.org/10.1111/mec.13157

Excoffier L, Smouse PE, Quattro JM (1992) Analysis of Molecular Variance Inferred From Metric Distances Among DNA Haplotypes: Application to Human Mitochondrial DNA Restriction Data. Genetics 131:479–491

Falush D, Stephens M, Pritchard J (2003) Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. Genetics 164:1567–1587

Ghigi A (1950) Piccioni Domestici e Colombicoltura. REDA, Roma

Giachetti C (1914) Piccioni Domestici - Francesco Battiato Editore. Catania

Giunchi D, Albores-Barajas Y, Balda NE, et al (2012) Feral Pigeons: Problems, Dynamics and Control Methods. In: Soloneski S, Larramendy ML (eds). InTech, Rijeka, pp 215–240

Goldstein DB, Linares AR, Cavalli-Sforza LL, Feldman MW (1995) An evaluation of genetic distances for use with microsatellite loci. Genetics 139:463–471

Gompertz T (1957) Some Observations on the Feral Pigeon in London (With Addendum: Two cases of polyneuritis in Feral Pigeons. *Derek Goodwin*). Bird Study 4:2–13. https://doi.org/10.1080/00063655709475863

Goodwin D (1960) Comparative ecology of feral pigeons in inner London. British Birds 53:201–212

Goudet J (2001) FSTAT, a program to estimate and test gene diversity and fixation indices (version 2.9. 3). http://www2. unil. ch/popgen/softwares/fstat. htm

Haag-Wackernagel D (1998) Die Taube: Vom heiligen Vogel der Liebesgöttin zur Strassentaube. Schwabe, Basel

Hetmanski T (2007) Dispersion asymmetry within a feral pigeon *Columba livia* population. Acta Ornithologica 42:23–31

Hubisz MJ, Falush D, Stephens M, Pritchard JK (2009) Inferring weak population structure with the assistance of sample group information. Molecular Ecology Resources 9:1322–1332

Huson DH, Bryant D (2006) Application of phylogenetic networks in evolutionary studies. Molecular Biology and Evolution 23:254–267

Jacob G, Prévot-Julliard A-C, Baudry E (2015) The geographic scale of genetic differentiation in the feral pigeon (*Columba livia*): implications for management. Biological Invasions 17:23–29. https://doi.org/10.1007/s10530-

014-0713-2

Jakobsson M, Rosenberg NA (2007) CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. Bionformatics 23:1801–1806

Johnston RF (1992) Evolution in the rock dove: skeletal morphology. The Auk 109:530–542

Johnston RF (1994) Geographic variation of size in feral pigeons. The Auk 111:398–404

Johnston RF, Janiga M (1995) The Feral Pigeons. Oxford University Press, London

Latter BDH (1972) Selection in finite populations with multiple alleles. III. Genetic divergence with centripetal selection and mutation. Genetics 70:475–490

Le Gros A, Samadi S, Zuccon D, et al (2016) Rapid morphological changes, admixture and invasive success in populations of Ring-necked parakeets (*Psittacula krameri*) established in Europe. Biological Invasions 18:1581–1598. https://doi.org/10.1007/s10530-016-1103-8

Lever C (1987) Naturalized birds of the world. Wiley, New York

McNeillie A (1976) Guide to the pigeons of the world. Elsevier-Phaidon, Oxford

Nei M (1973) Analysis of gene diversity in subdivided populations. PNAS 70:3321–3323. https://doi.org/10.1073/pnas.70.12.3321

Peakall R, Smouse PE (2006) GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research 6:288–295

Peakall R, Smouse PE (2012) GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. Bioinformatics 28:2537–2539

Pritchard JK, Stephens M, Peter D (2000) Inference of population structure using multilocus genotype data. Genetics 155:945–959

Ragionieri L, Mongini E, Baldaccini NE (1991) Problemi di conservazione in una popolazione di colombo selvatico (*Columba livia livia* Gmelin) della Sardegna. Supplemento alle Ricerche di Biologia della Selvaggina 18:35–46

Raymond M, Rousset F (1995) GENEPOP (version 1.2): population genetics software for exact tests and ecumenicism. Journal of Heredity 86:248–249

Ritchie J (1920) The Influence of Man on Animal Life in Scotland: a Study of Fauna Evolution. Cambridge University Press, Cambridge

Rosenberg N (2004) DISTRUCT: a program for graphical display for population structure. Molecular Ecology Notes 4:137–138

Shao Y, Tian H-Y, Zhang J-J, et al (2019) Genomic and phenotypic analyses reveal mechanisms underlying homing ability in pigeon. Molecular Biology and Evolution. https://doi.org/10.1093/molbev/msz208

Shapiro MD, Kronenberg Z, Li C, et al (2013) Genomic diversity and evolution of the head crest in the rock pigeon. Science 339:1063–1067. https://doi.org/10.1126/science.1230422

Shriver MD, Jin L, Boerwinkle E, et al (1995) A novel measure of genetic distance for highly polymorphic tandem repeat loci. Molecular Biology and Evolution 12:914–920

Simms E (1979) The public life of the street pigeon. Hutchinson, London

Smouse PE, Whitehead MR, Peakall R (2015) An informational diversity framework, illustrated with sexually

deceptive orchids in early stages of speciation. Molecular Ecology Resources 15:1375–1384

Sol D (2008) Artificial selection, naturalization, and fitness: Darwin's pigeons revisited. Biological Journal of the Linnean Society 93:657–665

Soldatini C, Mainardi D, Baldaccini EN, Giunchi D (2006) A temporal analysis of the foraging flights of feral pigeons (*Columba livia* f. *domestica*) from three Italian cities. Italian Journal of Zoology 73:83–92. https://doi.org/10.1080/11250000500502210

Sossinka R (1982) Domestication in birds. Avian Biology 6:373–403

Stringham SA, Mulroy EE, Xing J, et al (2012) Divergence, convergence, and the ancestry of feral populations in the domestic rock pigeon. Current Biology 22:302–308. https://doi.org/10.1016/j.cub.2011.12.045

Tang Q, Low GW, Lim JY, et al (2018) Human activities and landscape features interact to closely define the distribution and dispersal of an urban commensal. Evolutionary Applications 11.9:1598–1608. https://doi.org/10.1111/eva.12650

Toschi A (1939) Ricerche ed osservazioni sul Colombo selvatico (*Columba livia* L.). Ricerche di Zoologia Applicata alla Caccia 13:1–124

Traxler B, Brem G, Müller M, Achmann R (2000) Polymorphic DNA microsatellites in the domestic pigeon, *Columba livia* var. *domestica*. Molecular Ecology 9:366–368. https://doi.org/10.1046/j.1365-294x.2000.00874-2.x

Vaccari A, Zambon F (2014) Storia del Triganino Modenese. Notiziario della Federazione Italiana Allevatori Colombi 14:5–11

Van der Linden CG (1950) Le Pigeon Voyager. Payot, Paris

Wagner G, Galler SR, Schmidt-Koenig K, et al (1972) Topography and pigeon orientation. NASA, Washington, DC, pp 259–273

 Figure caption

Fig. 1 Sampling cities of feral pigeons (white dots), areas of origin of some of the Italian domestic breeds considered in
the analysis (arrows) and average number of juvenile Racing Homers reared in the different Italian regions in the period
2013-2018 (data provided by the Federazione Colombofila Italiana). Italian Racing Homer and Runt are not reported in
the map because their origin can not be ascribed to a well defined geographical area, even though the selection of the
Runt breed probably started in central Italy (Bigi et al. 2016).

Fig. 2. Phylogenetic trees drawn using three different distance-based models: *Da, Fst,* and *(δμ)²* distance. No
significant topology was supported in any of the three phylogenetic trees. The unique significant clustering was found
between F_TV and F_VE.

Fig. 3. Results of the Discriminant Analysis of Principal Components (DAPC) and of Bayesian computations on feral
pigeon samples. **(a)** DAPC evidenced a clear differentiation of F_BZ, F_SS, and F_VE, individuals from Central Italy
plotted together, while the position of F_TV was intermediate between F_VE and F_PV. **(b)** The ΔK computation
suggested that 4 clusters represented the best genetic subdivision of the sampling. **(c)** Bayesian analysis assigned F_BZ,
F_VE, and F_SS to unique and distinctive populations. The different colored bars in F_TV, F_PV, F_MO, F_RE, F_LE,
and F_PI describe admixed genetic compositions within these populations, thus evidencing a relevant genetic flow
among them.

Fig. 4. Discriminant Analysis of Principal Components e Bayesian computations in feral and domestic breeds. The
computations involved only pigeon lineages that could have contributed to the genetic composition of feral populations.
(a) DAPC showed that SB, and RM contributed marginally to the sampled feral populations, while IH, RH, PC and in
part RN clustered together with ferals from Lombardy, Emilia-Romagna and Tuscany. **(b)** The ΔK computation
suggested that 8 clusters represented the best genetic substructure of the sampling. **(c)** Bayesian analysis evidenced a
sharp distinction between feral and domestic samples. However, light blue, peach colored and light green bars found in
ferals suggested a probable origin of some individuals from IH, RH, PC and RN breeds.

Fig. 5. Bar chart showing population membership (Q) values at K=8. Admixed colored bars are representative of an
admixed genetic composition and origin. Identical colors indicate a common origin.

Table 1 Geographic origin of feral samples

City	Acronim	Region	No. samples
Bolzano	F_BZ	Trentino-Alto Adige	22
Treviso	F_TV	Veneto	24
Venice	F_VE	Veneto	22
Pavia	F_PV	Lombardy	20
Reggio Emilia	F_RE	Emilia-Romagna	25
Modena	F_MO	Emilia-Romagna	26
Pisa	F_PI	Tuscany	19
Leghorn	F_LE	Tuscany	14
Sassari	F_SS	Sardinia	22

314 **Table 2** Domestic breeds considered in the study. The breeds set off in bold (IO1, IO2, FL, TM1 and TM2)
315 were excluded from the analyses after preliminary investigations (see Material and Methods)

Domestic Breed	Acronim	No. samples
Italian Owl	**IO1**	**29**
Italian Owl Rondone	**IO2**	**20**
Florentine	**FL**	**20**
Piacentino	PC	25
Romagnol	RM	20
Runt	RN	19
Sottobanca	SB	26
Triganino Schietto	**TM1**	**26**
Triganino Gazzo	**TM2**	**20**
Racing Homer	RH	29
Italian Beauty Homer	IH	16

318 **Table 3** Variability indexes. Abbreviations: number of alleles (*Na*), number of effective alleles (*Ne*), allelic
319 richness (*Ar*), observed (*Ho*) and expected (*He*) heterozygosity. Except for *Ar*, values are averages ± SE

Population	*Na*	*Ne*	*Ar*	*Ho*	*He*
F_BZ	6.9 ± 0.7	3.9 ± 0.3	6.2	0.762 ± 0.037	0.727 ± 0.027
F_TV	7.8 ± 0.9	4.6 ± 0.6	6.9	0.705 ± 0.040	0.735 ± 0.036
F_VE	6.2 ± 0.5	3.4 ± 0.4	5.6	0.639 ± 0.065	0.651 ± 0.052
F_PV	7.9 ± 0.7	4.6 ± 0.5	7.2	0.710 ± 0.065	0.743 ± 0.038
F_RE	8.5 ± 0.9	4.5 ± 0.5	7.1	0.739 ± 0.048	0.736 ± 0.036
F_MO	8.3 ± 1.1	4.5 ± 0.6	7.0	0.714 ± 0.056	0.724 ± 0.046
F_PI	8.1 ± 0.8	5.0 ± 0.6	7.5	0.802 ± 0.124	0.758 ± 0.036
F_LE	7.6 ± 0.7	4.5 ± 0.6	7.6	0.708 ± 0.050	0.725 ± 0.044
F_SS	7.3 ± 0.8	4.6 ± 0.4	6.7	0.686 ± 0.069	0.733 ± 0.050
Global Mean ± SE	7.6 ± 0.8	4.4 ± 0.5	6.9 ± 0.6	0.718 ± 0.062	0.726 ± 0.050

326   **Figure 1**

332 **Figure 2**

(a)

(b)

(c)

K=4

Figure 3

349   **(a)**

352   **(b)**

355   **(c)**

358   **Figure 4**

Figure 5