Effect of nitrogen on fungal growth efficiency

Authors: D.P. Di Lonardo[1*], A. van der Wal [2], P. Harkes[3], W. de Boer[1,4]

[1]Department of Microbial Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Wageningen, the

Netherlands

[2]National Institute for Public Health and the Environment, Bilthoven, 3720 BA, The Netherlands

[3]Laboratory of Nematology, Wageningen University, Wageningen, the Netherlands

[4]Chairgroup of Soil Biology, Wageningen University, Wageningen, the Netherlands

*Corresponding author: Paolo Di Lonardo. Current address: Soil Biology Group, Wageningen

University & Research, P.O. Box 47, 6700 AA Wageningen, the Netherlands. E-mail:

paolo.dilonardo@wur.nl

Abstract

The contribution of fungi to carbon (C) and nitrogen (N) cycling is related to their growth efficiency

(amount of biomass produced per unit of substrate utilized). The concentration and availability of N

influences the activity and growth efficiency of saprotrophic fungi. When N is scarce in soils, fungi have

to invest more energy to obtain soil N, which could result in lower growth efficiencies. Yet, the effect of

N on growth efficiencies of individual species of fungi in soil has not been studied extensively. In this

study we investigated the influence of different concentrations of mineral N on the growth efficiency of

two common soil fungi, *Trichoderma harzanium* and *Mucor hiemalis* in a soil-like environment. We

hypothesized that a higher N availability will coincide with higher biomass production and growth

efficiency. To test this, we measured fungal biomass production as well as the respiration fluxes in sand

microcosms amended with cellobiose and mineral N at different C:N ratios. We found that for both fungal

species lower C:N ratios resulted in the highest biomass production as well as the highest growth

efficiency. This may imply that when N is applied concurrently with a degradable C source, a higher

amount of N will be temporarily immobilized into fungal biomass.

Keywords: C:N ratios; ergosterol; growth efficiency; respiration; saprotrophic fungi

1. Introduction

Fungi play a major role in terrestrial decomposition processes and they are important actors in soil organic matter (SOM) dynamics (van der Wal et al. 2013). They are thought to have a relevant contribution to the decomposition of the stable organic carbon (C) pool (Fontaine et al. 2007, 2011). On the other hand, filamentous fungi can promote the formation of soil macroaggregates (Willis et al. 2013) that promotes C sequestration by providing physical protection against decomposers and their degradative enzymes (Wilson et al. 2009).

It is generally assumed that soil microbial communities dominated by fungi have more efficient nitrogen (N) cycling than those dominated by bacteria (Wardle et al. 2004, Van Der Heijden et al. 2008). High N fertilizer additions have been indicated to be the cause of decrease in soil fungal biomass (de Vries et al. 2006, 2007), whereas the cessation of N-fertilizer use can cause a shift from bacterial to fungal dominated systems (de Vries et al. 2006, 2007, Postma-Blaauw et al. 2010). Increases in the abundance of fungi have been linked to a higher efficiency of N cycling and lower N losses from soils (de Vries et al. 2006, 2011, Gordon et al. 2008).

Growth efficiency is defined as the amount of biomass produced per unit of substrate utilized (Sinsabaugh et al. 2013, Mooshammer et al. 2014b). Microbial activity is highest when the C:N ratio of the substrate matches the demands of microbes (Hessen et al. 2004). According to the stoichiometric decomposition theory (Craine et al. 2007), decay processes are driven by the stoichiometry of substrates and adjustment in growth efficiencies may be the most important mechanism to cope with differences between substrate and biomass stoichiometry (Mooshammer et al. 2014b). It is expected that growth efficiencies are influenced by the C:N ratios of the organic substrates. When N is scarce in soils, fungi have to invest more energy in obtaining it, e.g. by producing extra-cellular enzymes, likely resulting in a low growth efficiencies. Generally, knowledge on microbial growth efficiency is of special interest for industrial applications (e.g., to obtain higher biomass or biosynthesized products). However, growth efficiencies have received increased attention of ecologists, due to its important implications for environmental processes (Geyer et al. 2016). Most of the ecological studies focused on soil microbial communities (e.g., Dijkstra et al., 2015; Geyer et al., 2018; Koranda et al., 2014; Mooshammer et al., 2014a; Sinsabaugh et al., 2013). However, fungal biomass production in soil microbial communities is not only affected by fungal growth responses but also by other processes such as predation and

competition. Hence, to have a basic understanding of the effect of substrate C:N ratio on fungal growth

responses in soil studies with single fungal species are needed.

In this study we investigated the influence of different concentrations of mineral N on the growth

efficiency of two common soil fungi, *Trichoderma harzanium* and *Mucor hiemalis* (De Boer et al. 2005,

Vinale et al. 2008) in a soil-like environment. As a carbon source we have chosen cellobiose as a model

compound for an easily degradable plant-derived carbohydrate (Martínez et al. 2005). We hypothesized

that higher nitrogen availability will coincide with higher fungal biomass production and growth efficiency.

2. Materials and Methods

Petri dishes (8.5 cm diameter) were filled with 60 g autoclaved, acid-washed quartz sand (granulation

0.1-0.5 mm; Honeywell Specialize Chemicals Seelze GmbH, Seelze, Germany). The lids of the Petri

dishes contained butyl rubber stoppers (Rubber BV, Den Haag, The Netherlands) to allow sampling the

gas from the headspace of the plates. The sand was amended with 10% (w/w) of a nutrient solution

containing (g l^{-1} demineralized water): KH_2PO_4, 0.10; K_2SO_4, 0.20; Yeast extract (Bacto™; Becton,

Dickinson and Company), 0.05; D-(+)-Cellobiose (Sigma-Aldrich), 5.0; MES (2-(N-morpholino)

ethanesulfonic acid, Sigma) 5.85. The latter compound was added because acid-washed sand has no

buffering capacity. To test the effect of different C:N ratios on fungal growth, the above described nutrient

solution received also NH_4NO_3 in different amounts. Three nutrient solutions were prepared: i) C-

cellobiose:N = 8:1, ii) C-cellobiose:N = 15:1 and iii) C-cellobiose:N = 50:1. The control treatment did not

receive any NH_4NO_3 addition. The pH of the nutrient solutions was adjusted to 6.5 with NaOH. Before

the addition of the nutrient solutions, sand was sterilized by two cycles of autoclaving (30 min at 121 ºC,

the second one after 24 hours). Next, it was dried at 120 ºC for two hours.

Fungal spores (10^4 spores g sand^{-1}) of *Trichoderma harzanium* and *Mucor hiemalis* were obtained from

pure cultures (Appendix S1) and mixed with the nutrient-containing sand. In total eight experimental

treatments were prepared: *T. harzanium* in sand with no nitrogen (TH No-N), with C:N=8 (TH 8:1), with

C:N=15 (TH 15:1) and with C:N=50 (TH 50:1); *M. hiemalis* in sand with no nitrogen (MH No-N), with

C:N=8 (MH 8:1), with C:N=15 (MH 15:1) and with C:N=50 (MH 50:1). Each treatment consisted of five

replicates. They were sealed with one layer of Diversified Biotech Petri Seal™ tape and one layer of

Parafilm, to avoid gas exchange with the external environment and maintaining air-tightness. Plates

were incubated in the dark at 20 °C.

During the 14-day incubation period, headspace CO_2 was sampled and measured (Appendix S1). CO_2

was analyzed at 2, 4, 7, 9, 11 and 14 days. At the end of the incubation period, soil was homogenized

by mixing, it was sampled and kept in aliquots in the freezer at -20 °C for ergosterol measurements,

DNA extractions and qPCR assays (Appendix S1). We calculated the growth efficiency for each fungus

on basis of the amount of ergosterol or ITS copy numbers per amount of CO_2 released. Growth

efficiencies were expressed as relative growth efficiencies where efficiencies of the C:N = 8 treatments

were set at 100%.

Differences in respiration, ergosterol, DNA copy numbers and growth efficiencies between treatments

were tested with one-way ANOVA followed by post-hoc Tukey's test, using IBM SPSS Statistics 22. In

some cases, due to unequal variances, Tukey's test was not possible and statistical comparisons were

performed by Tamhane's T2 test. We used linear regression analysis to test the relationship between

the different amounts of added N and ergosterol concentrations, and the DNA copy numbers.

3. Results and Disccussion

Growth efficiencies for both *M. hiemalis* and *T. harzanium*, as based on ergosterol and DNA copy

numbers, showed the same trend, namely an increase with decreasing C:N ratios (Fig. 1A and 1B, $P <$

0.05). We observed the same pattern for ergosterol production (Fig. S1A, $P < 0.05$). Furthermore, DNA

copy numbers for both fungal species showed an overall increase with an increasing amount of N (Fig.

S1B). However, only *M. hiemalis* grown in sand with C:N ratios 8 and 15 had significant higher DNA

copy numbers ($P < 0.05$). On the contrary, respiration fluxes did not increase with decreasing C:N ratios

(Fig. S2). Taken together these results indicate that more C-cellobiose was metabolized at lower C:N

ratio, implying that not all C-cellobiose has been metabolized in the treatments C:N = 15 and C:N = 50,

and certainly not in the control treatments, where there was no addition of mineral N. Our results are in

line with our hypothesis, namely that the highest growth efficiency is expected with higher nitrogen

availability. A similar growth efficiency pattern was observed for a litter-decomposing fungus grown on

maize litter, where the efficiencies increased accordingly with increasing N availability in the plant

material (Lashermes et al. 2016). In addition, our results suggest that when N becomes a limiting factor,

fungi invest extra energy to obtain N, for instance by recycling their cellular N via controlled autolysis

(Santamaria and Reyes 1988) or allocating N to essential metabolic processes (Wicklow 2006).

The linear regression analysis showed a significant positive linear relationship between the amount of

added N and ergosterol concentrations (R^2 = 0.8860; P < 0.0001 and R^2 = 0.8944; P < 0.0001, for *M.*

hiemalis and *T. harzanium* respectively; Fig. S3A) and between the amount of added N and DNA copy

numbers (R^2 = 0.7609; P < 0.0001 and R^2 = 0.3523; P = 0.006, for *M. hiemalis* and *T. harzanium*

respectively; Fig. S3B). Increase of fungal biomass can reduce N losses from soils (de Vries et al. 2006)

and, consequently, a higher fungal biomass in soils can be considered as an indicator of higher soil N

retention (de Vries et al. 2011). Applications of fertilizers to agricultural soils can result in N losses when

crops do not rapidly taken up the added N , and part of the N can be lost via leaching and denitrification

(de Vries and Bardgett 2012). Our study indicates that when N is applied concurrently with a degradable

C source, a higher amount of N is built into fungal biomass, thereby possibly reducing N losses (Simpson

et al. 2007, Liang and Balser 2011). Moreover, our study provides information on how nitrogen

influences fungal biomass production and growth efficiency. Knowledge on this effect of N is critical for

the assessment of soil C and N budgets.

Acknowledgments

We are thankful to Jan Geert Bruggink for help in the lab.

References

Boer W De, Folman LB, Summerbell RC, Boddy L. 2005. Living in a fungal world: Impact of

fungi on soil bacterial niche development. FEMS Microbiol Rev. 29:795–811,

doi:10.1016/j.femsre.2004.11.005.

Craine JM, Morrow C, Fierer N. 2007. Microbial nitrogen limitation increases decomposition.

Ecology. 88:2105–2113.

Dijkstra P, Salpas E, Fairbanks D, Miller EB, Hagerty SB, Groenigen KJ van, Hungate BA,

Marks JC, Koch GW, Schwartz E. 2015. High carbon use efficiency in soil microbial communities is

related to balanced growth, not storage compound synthesis. Soil Biol Biochem. 89:35–43,

doi:10.1016/j.soilbio.2015.06.021.

Fontaine S, Barot S, Barré P, Bdioui N, Mary B, Rumpel C. 2007. Stability of organic carbon in

deep soil layers controlled by fresh carbon supply. Nature. 450:277–80, doi:10.1038/nature06275.

Fontaine S, Henault C, Aamor A, Bdioui N, Bloor JMG, Maire V, Mary B, Revaillot S, Maron

PA. 2011. Fungi mediate long term sequestration of carbon and nitrogen in soil through their priming

effect. Soil Biol Biochem. 43:86–96, doi:10.1016/j.soilbio.2010.09.017.

Geyer K, Frey S, Dijkstra P, Sinsabaugh R. 2018. Clarifying the interpretation of carbon use

efficiency estimates in soil through methods comparison. Soil Biol Biochem. 20:8564,

doi:10.1016/J.SOILBIO.2018.09.036.

Geyer KM, Kyker-Snowman E, Grandy AS, Frey SD. 2016. Microbial carbon use efficiency:

accounting for population, community, and ecosystem-scale controls over the fate of metabolized

organic matter. Biogeochemistry. 127:173–188, doi:10.1007/s10533-016-0191-y.

Gordon H, Haygarth PM, Bardgett RD. 2008. Drying and rewetting effects on soil microbial

community composition and nutrient leaching. Soil Biol Biochem. 40:302–311,

doi:10.1016/j.soilbio.2007.08.008.

Harkes P, Verhoeven A, Sterken MG, Snoek LB, Elsen SJJ van den, Mooijman PJW, Quist

CW, Vervoort MTW, Helder J. 2017. The differential impact of a native and a non-native ragwort

species (Senecioneae) on the first and second trophic level of the rhizosphere food web. Oikos.

126:1790–1803, doi:10.1111/oik.04530.

Heijden MGA Van Der, Bardgett RD, Straalen NM Van. 2008. The unseen majority: Soil

microbes as drivers of plant diversity and productivity in terrestrial ecosystems. Ecol Lett. 11:296–310,

doi:10.1111/j.1461-0248.2007.01139.x.

Hessen D, Ågren G, Anderson T. 2004. Carbon sequestration in ecosystems: the role of

stoichiometry. Ecology. 85:1179–1192.

Koranda M, Kaiser C, Fuchslueger L, Kitzler B, Sessitsch A, Zechmeister-Boltenstern S,

Richter A. 2014. Fungal and bacterial utilization of organic substrates depends on substrate

complexity and N availability. FEMS Microbiol Ecol. 87:142–52, doi:10.1111/1574-6941.12214.

Lashermes G, Gainvors-Claisse A, Recous S, Bertrand I. 2016. Enzymatic Strategies and

Carbon Use Efficiency of a Litter-Decomposing Fungus Grown on Maize Leaves, Stems, and Roots.

Front Microbiol. 7:1–14, doi:10.3389/fmicb.2016.01315.

Liang C, Balser TC. 2011. Microbial production of recalcitrant organic matter in global soils:

implications for productivity and climate policy. Nat Rev Microbiol. 9:75–75, doi:10.1038/nrmicro2386-

c1.

Martínez AT, Speranza M, Ruiz-Dueñas FJ, Ferreira P, Camarero S, Guillén F, Martínez MJ,

Gutiérrez A, Río JC del. 2005. Biodegradation of lignocellulosics: microbial, chemical, and enzymatic

aspects of the fungal attack of lignin. Int Microbiol. 8:195–204.

Mooshammer M, Wanek W, Hämmerle I, Fuchslueger L, Hofhansl F, Knoltsch A, Schnecker J,

Takriti M, Watzka M, Wild B, Keiblinger KM, Zechmeister-Boltenstern S, Richter A. 2014a. Adjustment

of microbial nitrogen use efficiency to carbon:nitrogen imbalances regulates soil nitrogen cycling. Nat

Commun. 5:3694, doi:10.1038/ncomms4694.

Mooshammer M, Wanek W, Zechmeister-Boltenstern S, Richter A. 2014b. Stoichiometric

imbalances between terrestrial decomposer communities and their resources: Mechanisms and

implications of microbial adaptations to their resources. Front Microbiol. 5:1–10,

doi:10.3389/fmicb.2014.00022.

Nikolcheva LG, Bourque T, Bärlocher F. 2005. Fungal diversity during initial stages of leaf

decomposition in a stream. Mycol Res. 109:246–253, doi:10.1017/S0953756204001698.

Postma-Blaauw MB, Goede RGM de, Bloem J, Faber JH, Brussaard L. 2010. Soil biota

community structure and abundance under agricultural intensification and extensification. Ecology.

91:460–473, doi:10.1890/09-0666.1.

Ridder-Duine AS de, Smant W, Wal A van der, Veen J a. van, Boer W de. 2006. Evaluation of

a simple, non-alkaline extraction protocol to quantify soil ergosterol. Pedobiologia (Jena). 50:293–300,

doi:10.1016/j.pedobi.2006.03.004.

Santamaria F, Reyes F. 1988. Proteases produced during autolysis of filamentous fungi.

Trans Br Mycol Soc. 91:217–220, doi:10.1016/S0007-1536(88)80207-9.

Simpson AJ, Simpson MJ, Smith E, Kelleher BP. 2007. Microbially Derived Inputs to Soil

Organic Matter: Are Current Estimates Too Low? Environ Sci Technol. 41:8070–8076,

doi:10.1021/es071217x.

Sinsabaugh RL, Manzoni S, Moorhead DL, Richter A. 2013. Carbon use efficiency of microbial

communities: stoichiometry, methodology and modelling. Ecol Lett. 16:930–939,

doi:10.1111/ele.12113.

Vinale F, Sivasithamparam K, Ghisalberti EL, Marra R, Woo SL, Lorito M. 2008. Trichoderma–

plant–pathogen interactions. Soil Biol Biochem. 40:1–10, doi:10.1016/j.soilbio.2007.07.002.

Vries FT de, Bardgett RD. 2012. Plant–microbial linkages and ecosystem nitrogen retention:

lessons for sustainable agriculture. Front Ecol Environ. 10:425–432, doi:10.1890/110162.

Vries FT de, Bloem J, Eekeren N van, Brusaard L, Hoffland E. 2007. Fungal biomass in

pastures increases with age and reduced N input. Soil Biol Biochem. 39:1620–1630,

doi:10.1016/j.soilbio.2007.01.013.

Vries FT de, Groenigen JW van, Hoffland E, Bloem J. 2011. Nitrogen losses from two

grassland soils with different fungal biomass. Soil Biol Biochem. 43:997–1005,

doi:10.1016/j.soilbio.2011.01.016.

Vries FT de, Hoffland E, Eekeren N van, Brussaard L, Bloem J. 2006. Fungal/bacterial ratios

in grasslands with contrasting nitrogen management. Soil Biol Biochem. 38:2092–2103,

doi:10.1016/j.soilbio.2006.01.008.

Wal A van der, Geydan TD, Kuyper TW, Boer W de. 2013. A thready affair: linking fungal

diversity and community dynamics to terrestrial decomposition processes. FEMS Microbiol Rev.

37:477–494, doi:10.1111/1574-6976.12001.

Wardle D a, Bardgett RD, Klironomos JN, Setälä H, Putten WH van der, Wall DH. 2004.

Ecological linkages between aboveground and belowground biota. Science. 304:1629–33,

doi:10.1126/science.1094875.

White TJ, Bruns S, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal

ribosomal RNA genes for phylogenetics. PCR Protoc A Guid to Methods Appl. 315–322, doi:citeulike-

article-id:671166.

Wicklow D. 2006. The Fungal Community. Its Organization and Role in the Ecosystem. 3rd

Edition Edited by John Dighton, James F. White, and Peter Oudemans (Rutgers University). Taylor

& Francis Group, CRC Press, Boca Raton, FL. 2005. xx + 936 pp. 18.5 × 26 cm. $139.95. J Nat

Prod. 69:859–859, doi:10.1021/np0682244.

Willis A, Rodrigues BF, Harris PJC. 2013. The Ecology of Arbuscular Mycorrhizal Fungi. CRC

Crit Rev Plant Sci. 32:1–20, doi:10.1080/07352689.2012.683375.

Wilson GWT, Rice CW, Rillig MC, Springer A, Hartnett DC. 2009. Soil aggregation and carbon

sequestration are tightly correlated with the abundance of arbuscular mycorrhizal fungi: results from

long-term field experiments. Ecol Lett. 12:452–461, doi:10.1111/j.1461-0248.2009.01303.x.

 Figure caption

Fig. 1: Relative (%) fungal growth efficiencies in sand microcosms containing different C:N ratios after

14 days of incubation. Growth efficiency for each fungus is based on the amount of ergosterol (A) or ITS

copy numbers (B) per amount of CO_2 released. Growth efficiencies of the C:N = 8 treatments were set

at 100%. Statistically significant differences ($P < 0.05$) are indicated with different letters. MH: *M.*

hiemalis; TH: *T. harzanium*. No-N: no addition of N; 50:1 is C:N = 50; 15:1 is C:N = 15; 8:1 is C:N = 8.

Vertical bars represent standard errors (n = 5).