1 **Temperature change has the same effect as genetic variation on a morphological trait**
2 **involved in reproductive isolation between Drosophila sister species**
3

4 Alex E. Peluffo[1,*], Mehdi Hamdani[1], Alejandra Vargas-Valderrama[1], Jean R. David[2,3],
5 François Mallard[4], François Graner[5], Virginie Courtier-Orgogozo[1]
6
[1]Institut Jacques Monod, CNRS, Univ. de Paris, 75013 Paris, France.
[2]Institut Systématique Evolution Biodiversité (ISYEB), CNRS, MNHN, Sorbonne Université,
EPHE, 57 rue Cuvier, CP 50, 75005 Paris, France.
[3]Laboratoire Evolution, Génomes, Comportement, Biodiversité (EGCE), CNRS, IRD, Univ.
Paris-sud, Université Paris-Saclay, 91198 Gif-sur-Yvette, France.
[4]Institut de Biologie de l'École Normale Supérieure, CNRS UMR 8197, Inserm U1024, PSL
Research University, F-75005 Paris, France.
[5]Matière et Systèmes Complexes, CNRS UMR 7057, Univ. de Paris, Paris, France.
*Current address: Data Science Department, Pharnext S.A., 11 rue René Jacques, 92130
Issy-les-Moulineaux, France.
17
18 Corresponding authors: VCO: virginie.courtier@normalesup.org, tel: +33 1 57 27 80 43, fax:
19 +33 1 57 27 80 87; AEP: alex.peluffo@icloud.com, tel: +33 1 41 09 22 42
20
Author details
AEP: alex.peluffo@icloud.com ORCID 0000-0001-7216-4205
MH: m_hamdani@outlook.fr   no ORCID
AVV: alejandra.vargas-valderrama@inserm.fr ORCID 0000-0002-6003-3459
JRD: Jean.David@egce.cnrs-gif.fr no ORCID
FM: francois.mallard@ens.fr ORCID: 0000-0003-2087-1914
FG: francois.graner@univ-paris-diderot.fr ORCID : 0000-0002-4766-3579
VCO: virginie.courtier@normalesup.org ORCID: 0000-0002-9297-9230
29

30 **Author Contributions**
31 AEP designed the study, JRD collected fly strains, MH, AVV and AEP collected the data,
32 AEP and FG designed the analysis pipeline, FM wrote the ImageJ plugin to extract contour
33 pixels, AEP, FG and VCO analyzed the data, AEP, VCO and FG wrote the paper. All authors
34 have read and approved the manuscript.
35

Acknowledgments
We are grateful to São Tomé authorities for allowing us to collect flies. We thank David Stern
and Daniel Matute for fly strains. We thank the Courtier lab for helpful discussions. We also
thank Philippe Rinaudo for feedback on the statistical analyses. The research leading to this
paper has received funding from the European Research Council under the European
Community's Seventh Framework Program (FP7/2007–2013 Grant Agreement no. 337579)
to VCO and from the labex ''Who am I?'' (ANR-11-LABX-0071) and the Université de Paris
IdEx (ANR-18-IDEX-0001) funded by the French government through grant no. ANR-11-
IDEX-0005-02 to AEP.
45
46 DRYAD: doi:10.5061/dryad.kprr4xh1f. The temporary link until manuscript acceptance is
47 https://datadryad.org/stash/share/72YV3b0V2AELCpWuS_ULqMlZyORs9WILt6iMiInAkpE.
48

49 **Temperature change has the same effect as genetic variation on a morphological trait**
50 **involved in reproductive isolation between Drosophila sister species**
51
52 running title: shape plasticity to temperature & speciation
53

Abstract
Phenotypic plasticity, the capacity of one genotype to generate distinct phenotypes in
different environments, is usually thought to facilitate species divergence by opening
novel ecological niches to plastic individuals. Here we reveal a case of speciation
where this "plasticity first" scenario might not hold. Male genitalia are usually
extremely divergent between closely related species, but relatively constant within
one species. Under the lock-and-key hypothesis, rapid morphological evolution is
associated with a high match between male and female genitalia of the same species
and a low match between male and females of closely related species. Previous
studies have suggested plasticity of genitalia to be a proof against the lock-and-key
hypothesis since the environmentally triggered phenotypic change could modify the
"key". Here we examine the effect of temperature on the shape of the ventral
branches, a male genital structure involved in reproductive isolation, in the sister
species _Drosophila santomea_ and _D. yakuba_. We designed a semi-automatic
measurement pipeline that can reliably identify curvatures and landmarks based on
manually digitized contours of the ventral branches. With this method, we observed
that ventral branches are not plastic in _D. yakuba_ but that in _D. santomea_ temperature
change phenocopies interspecific genetic variation between both species for ventral
branches shape. Our results suggest that speciation of _D. santomea_ and _D. yakuba_
was associated with a gain of plasticity and that genitalia plasticity can be compatible
with the lock-and-key hypothesis.
(234 words)

76
77 **Keywords**
78 Plasticity, Speciation, Phenotypic accommodation, Genitalia, Reproductive isolation,
79 Drosophila, Landmark, Automatic Detection, Shape analysis
80
81 **Abbreviations**
82 **SE: Standard Error**
83 QTL: Quantitative Trait Locus
84 ST: spine thrust
85
86 **Data Accessibility Statement**
87 The images, contours, scripts, and measurement values are available on DRYAD (link not
88 shown here to keep anonymity).
89
90 **Conflict of Interest**
91 The authors declare that they have no competing interests.

Introduction

Phenotypic plasticity, the capacity for one genotype to generate multiple phenotypes in response to environmental variation, is a pervasive feature of biological systems (Debat & David, 2001; Klingenberg, 2019). The connection between plasticity and speciation is multifaceted (Lafuente & Beldade, 2019). On the one hand, plasticity can be heritable and modified by selection. On the other hand, plasticity can favor adaptation and speciation. As animals colonize novel habitats or face changing climate conditions, the phenotypic traits that are optimal for fitness are usually different from those experienced in the ancestral population. Waddington was among the first to suggest that organisms may solve this challenge by phenotypic plasticity first and later on by genetic fixation of what was previously an environmentally induced phenotypic trait (a process he called "genetic assimilation") (Waddington, 1942). According to several authors, the trait variations enabled by plasticity can initiate and accelerate the pace of adaptive evolution and promote morphological diversification. This central idea is at the basis of the "flexible stem hypothesis" (West-Eberhard, 2003; Schneider & Meyer, 2017), the "plasticity-first" model (Levis & Pfennig, 2016) or the "buying time" hypothesis, where plasticity allows the population to persist long enough for adaptive mutations to arise and become fixed (Pennisi, 2018). A key feature of all these views is that the phenotypic change triggered by the plastic response, which allows the colonization of the new niches, is a phenocopy, i.e., that the phenotypic change can be developmentally triggered by environmental variation or genetic variation interchangeably (Lafuente & Beldade, 2019). As we learn more about the genes mediating phenotypic plasticity (Gibert, 2017), it appears that similar phenotypic changes, either environmentally or genetically induced, can sometimes involve the same genetic loci. For example, the same enhancer of the gene *tan* contributes to both phenotypic plasticity in *Drosophila melanogaster* (Gibert *et al.*, 2016) and interspecific evolution between sister-species *D. santomea* and *D. yakuba* with respect to abdomen pigmentation (Jeong *et al.*, 2008).

Depending on the setting, plasticity can either accelerate, slow down, or have little effect on evolution and species divergence (Price *et al.*, 2003). Speciation, the process through which lineages diverge and become reproductively isolated, involves the accumulation over time of barriers limiting interbreeding, including divergence in ecological niches, behavioral isolation and genomic incompatibilities (Coyne & Orr, 2004). As early as 1844, anatomical differences in genitalia between closely related species were proposed to be an essential mechanism maintaining reproductive isolation, as the so-called "lock-and-key" hypothesis (Dufour, 1844; Masly, 2011). In animals with internal fertilization, genitalia are the most rapidly evolving organs in terms of morphology (Eberhard, 1988), suggesting that a significant part of the speciation process involves anatomical divergence in genitalia. Alternatively, genital evolution can be a by-product of other evolutionary processes occurring within single lineages, independently of speciation (such as sexual selection), and lead to reproductive isolation as a by-product, when individuals attempt to hybridize with other lineages (Masly, 2011).

The lock-and-key hypothesis, even in species where it seems applicable, has been challenged by a variety of observations, including the facts that (1) genitalia in females do not differ as much as in males, (2) closely related species with conspicuous genital differences can still often produce hybrids, (3) males with laser-ablated genital organs can still copulate with no observed defect, and (4) genitalia morphology can be sensitive to temperature or nutrition (Shapiro & Porter, 1989; Arnqvist & Thornhill, 1998; Andrade *et al.*, 2005; Masly, 2011; Simmons, 2014; LeVasseur-Viens *et al.*, 2015 and references therein). It

is thus possible that in some taxonomic groups interspecific differences in genital
morphology do not contribute to reproductive isolation.
To better comprehend the link between plasticity and speciation, careful
examinations of particular cases are essential, and genital traits involved in reproductive
isolation represent highly relevant model systems. How plastic are genitalia in general?
Surprisingly, few studies have examined genitalia after raising organisms in various
conditions. In the water strider Aquarius remigis, the mosquito *Aedes aegypti* and the fly *D.*
melanogaster, changes in larval crowding, nutrition conditions or temperature were found to
affect adult body size but had little effect on the size of the external genitalia (Wheeler *et al.*,
1993; Fairbairn, 2005; Shingleton *et al.*, 2009). However, in two other species, the mosquito
Anopheles albimanus and the fly *Drosophila mediopunctata*, the size and shape of the male
intromittent organ was found to vary with rearing temperature (Hribar, 1996; Andrade *et al.*,
2005). Overall, analysis of individuals sampled from the wild show that for a given arthropod
or mammal species, the genitalia are usually more or less the same size whereas adult body
size varies extensively (Eberhard *et al.*, 1998; Dreyer & Shingleton, 2011 and references
therein). These observations are concordant with the "lock-and-key hypothesis", where male
genitalia have to be of a particular size and shape to physically fit with the female genitalia.
They are also explained by the "one-size-fits-all" hypothesis, where females appear to prefer
males with genitalia of intermediate size (Eberhard *et al.*, 1998).
In order to analyze and quantify the possible link between plasticity, reproductive
isolation and interspecific divergence, we chose to examine the effect of temperature on a
male primary sexual trait likely involved in reproductive isolation between two Drosophila
sister species, *D. santomea* and *D. yakuba*. These two species form an attractive system
because their natural environment is relatively well characterized, they are known to
hybridize, and one of their most remarkable morphological difference is a primary sexual trait
that seem to be involved in a "lock-and-key" mechanism. *D. santomea* and *D. yakuba*
diverged approximately 0.5-1 million years ago (Turissini & Matute, 2017). They can be
crossed to generate fertile F1 females (Lachaise *et al.*, 2000). *D. santomea* is endemic to the
island of São Tomé, a volcanic island off the coast of Gabon (Lachaise *et al.*, 2000), while *D.*
yakuba is found in São Tomé and throughout sub-Saharan Africa (Lachaise *et al.*, 1988,
2000). In São Tomé, *D. santomea* lives in the mist forests at high elevations while *D. yakuba*
is found in open habitats associated with human presence, mostly at low elevations (Llopart
et al., 2005a; b) Both species co-occur at mid-elevation, around 1150 m, and hybrids have
been found consistently in this hybrid zone since its discovery in 1999 (Lachaise *et al.*, 2000;
Llopart *et al.*, 2005a; Comeault *et al.*, 2016; Cooper *et al.*, 2018). *D. santomea* being insular,
it is thought that this species originated from a common ancestor with *D. yakuba*, which
colonized the island about 0.5-1 million years ago (Cariou *et al.*, 2001; Llopart *et al.*, 2002;
Turissini & Matute, 2017), and that the present co-occurence of *D. santomea* and *D. yakuba*
in São Tomé reflects secondary colonization by *D. yakuba* from the African mainland, maybe
during the last 500 years when Portuguese colonised the island (Cariou *et al.*, 2001).
Analysis of genomic and mitochondrial DNA sequences indicate that gene flow occurred
between the *D. santomea* and *D. yakuba* more than 1,000 generations ago (Turissini &
Matute, 2017; Cooper *et al.*, 2019).
Multiple potential reproductive isolating mechanisms have been identified between
the two species, such as genetic incompatibilities (Coyne *et al.*, 2004; Moehring *et al.*, 2006),
ecological niche divergence (Matute *et al.*, 2009), mate discrimination (Lachaise *et al.*, 2000;
Coyne *et al.*, 2002), behavioral (Cande *et al.*, 2012), physiological (Matute, 2010) and
morphological differences (Lachaise *et al.*, 2000; Jeong *et al.*, 2008; Nagy *et al.*, 2018; Liu *et*

al., 2019). One reproductive isolating mechanism between *D. yakuba* and *D. santomea*
involves a difference in ventral branches shape in the male genitalia and is the most
conspicuous difference in male genitalia shape between the two species (Kamimura &
Mitsumoto, 2012b; Yassin & Orgogozo, 2013) (Figure 1). Ventral branches are located
below the aedeagus [i.e., the insect phallus (Rice et al., 2019)] and are only found in the *D.*
yakuba complex, which comprises D. teissieri, D. yakuba and *D. santomea* (Yassin and
Orgogozo 2013).
In *D. yakuba*, spiny ventral branches insert inside female protective pouches during mating.
In *D. santomea*, the male spines and female pouches are absent. These structures appear
to play important roles during copulation. When mating with *D. yakuba* males, *D. santomea*
females are wounded by the spines of the male ventral branches and they also live shorter
than females mating with conspecific males (Matute & Coyne, 2010; Kamimura, 2012;
Kamimura & Mitsumoto, 2012b). Compared to D. teissieiri females, D. santomea females
also survive less to interspecific copulation with *D. mauritiana* (Yassin & David, 2016).
Moreover, Kamimura and Mitsumoto (2012b) reported that "copulating pairs of D. santomea
males x D. yakuba females dislodge readily when disturbed", suggesting that the spines may
fasten genital coupling (Masly, 2011). We previously found that a major QTL on
chromosome 3L contributes to the ventral branches shape difference between *D. santomea*
and *D. yakuba* (Peluffo *et al.*, 2015).
In São Tomé the climate is very stable throughout the year, with only a 2.5°C- difference
between the average daily temperature of the warmest month (March) and of the coldest
one (July), and daily oscillations of about 5°C only (climate-data.org,
www.worldclim.org/bioclim). Based on temperature measurements at Monte Café (climate-
data.org), we estimate that the average temperature in the hybrid zone of Bom Sucesso
(1153m) varies between 15.5°C and 18°C throughout the year. In the wild, *D. santomea* flies
are thus likely developing mainly at temperatures around 18°C or lower.
In previous studies of ventral branch shape, flies were raised either at 21°C (Yassin
& Orgogozo, 2013) or 25°C (Kamimura, 2012; Kamimura & Mitsumoto, 2012b; Peluffo *et al.*,
2015). Here, we report that *D. santomea* males raised at 18°C develop spiny ventral
branches comparable to those of *D. yakuba* raised at 25°C. This is a surprising example
where organs potentially directly linked with reproductive isolation undergo a plastic
modification similar to the difference between two sister species. To better characterize the
morphological change in ventral branches shape, we developed a user-friendly method to
quantify contour curvatures and automatically detect spines using machine learning. We
used it to examine the plastic response of ventral branches development at 18°C and 25°C
both in newly collected wild strains and in strains kept in the lab for many years.
223
224
Material and Methods
Fly rearing and imaging
Fly strains (Table 1) were kept at 22˚C on standard yeast-cornmeal–agar medium in
uncrowded conditions before the beginning of the experiments. For each strain, roughly 20
individuals were transferred from the 22˚C stock to either 18˚C or 25°C, kept for a minimum
of two non-overlapping adult generations. Adult males were 5 to 7 days old when frozen at −
80°C for subsequent dissection. Dissection of genitalia was performed in 1X PBS at room
temperature. Each genitalia was mounted on standard glass slides in DMHF (Dimethyl

Hydantoin Formaldehyde, Entomopraxis) medium and kept overnight before imaging on an
Olympus IX83 inverted station at 40X.
235
236
237

Table 1. List of Isofemale lines used in this study. For each species, the most common name in the literature, the location, year of capture and reference to origin of the strain are given. All lines are indicated in the same order as in Figure 2.

Species	Name	Location	Year	Reference
D. yakuba	Ivory Coast	Ivory Coast	1955	Cornell National Drosophila Species Stock Center, Strain #14021-0261.00 (given by D. Stern)
D. yakuba	BM2015	São Tomé, Bom Sucesso Botanical Garden, 1150m	February 2015	This study
D. yakuba	Oku	Cameroun, Mt. Oku, 2000m	April 2016	This study
D. yakuba	Raphia	Cameroun, Mt. Oku, 1800m	April 2016	This study
D. santomea	STO.4	São Tomé, Obo Natural Reserve, submontane forest 1300-1450m	1998	(Lachaise *et al.*, 2000) Cornell National Drosophila Species Stock Center, Strain #14021-0271.00 (given by D. Stern)
D. santomea	STO Cago 1482	São Tomé 1482m	2001	(Llopart *et al.*, 2005b) This strain's original name is STO-LAGO 1482 (given by D. Stern)
D. santomea	Quija22	São Tomé, Quija River, 650m	2009	(Gavin-Smyth & Matute, 2013) This strain name is also Quija650.22 (given by D. Matute)
D. santomea	BM152	São Tomé Bom Sucesso Botanical Garden, 1150m	February 2015	This study
D. santomea	BM153	São Tomé Bom Sucesso Botanical Garden, 1150m	February 2015	This study
D. santomea	BM161	São Tomé Bom Sucesso Botanical Garden, 1150m	September 2016	This study
D. santomea	BM167	São Tomé Bom Sucesso Botanical Garden, 1150m	September 2016	This study
D. santomea	1563	EYFP lab strain derived from STO CAGO 1482, Insertion @ 3L:11.843.137	2001	(Stern *et al.*, 2017) (given by D. Stern)

Raw contour acquisition
All contours were digitized by the same person. Pictures were anonymized for manual contour acquisition so that the digitizer did not know the genotype. Digitization was skipped when the quality of the mounting was judged to be poor. For each picture, a custom ImageJ plugin was used to extract *x,y* coordinates (in pixels) of the contour. The plugin is designed to open all the pictures contained in a directory, allowing the user to manually draw a contour of the object of interest using the freehand tool of ImageJ.The raw contour is a series of points p_1, p_2, ..., p_n in a two-dimensional space *x,y* where *n* is the number of points over which the contour passes (usually $500 < n < 1000$). The contour is open and its endpoints are unimportant (Figure 1). It is analyzed (and twice smoothed) as follows.

Smoothed contour
The first layer of transformation is a rectangular smoothing filter over the raw contour to obtain the smoothed contour. At each point p_j with coordinates (x_j, y_j), we derive p'_j with coordinates (x'_j, y'_j) where $x'_j = \frac{1}{2\alpha n} \sum_{i=j-\alpha n}^{j+\alpha n} x_i$ and $y'_j = \frac{1}{2\alpha n} \sum_{i=j-\alpha n}^{j+\alpha n} y_i$.
Here the contour smoothing parameter *α*, to be adjusted via learning, describes the proportion of points (relative to the total number of points forming the contour) to include in the smoothing. This implies that the smoothed contour is *2αn* points shorter (*αn* on each side) than the raw contour.

Raw curvature of the smoothed contour
For each smoothed contour, the raw curvature *k* is computed with a sliding window of three points. For any set of three points *M*, *N*, *P* forming a triangle, the diameter of the circumscribed circle to this triangle, $2r = MP / \sin (MN,NP)$ can be computed as the product of the Euclidean distances divided by the cross product of the two (non-basal) sides, *r* is the curvature radius in *N*, and the curvature *k* in *N* is the inverse of *r* :

$$k = \frac{1}{r} = 2 \frac{|\overrightarrow{MN} \times \overrightarrow{NP}|}{MN \cdot NP \cdot PM}$$

The flatter the contour, the wider the circumscribed circle, the larger the radius r, and the smaller the curvature k. For each contour, the curvature profile is the curvature k_j computed over p'_j in $[p_{r1}, p_m]$ using its neighbouring points (M, N, $P = p'_{j-1}, p'_j, p'_{j+1}$) versus the curvilinear abscissa s_j of p'_j which is the sum of Euclidean distances from origin, $s_j = p'_1 p'_2 + p'_2 p'_3 + \ldots + p'_{j-1} p'_j$.

Refined curvature
We then use this first raw curvature estimation as information to refine the curvature in a second pass. In this second measure, the refined curvature k_j' is computed over an adaptive window of size $a = \frac{1}{|k|}$ for $k < 0.1$ and $a = 10$ otherwise: M, N, $P = p'_{j-a}, p'_j, p'_{j+a}$. This means that the curvature is computed over a larger distance where it is small (and curvature radius is large), which requires more smoothing, without losing the sharpness of curvature peak determination where the curvature is large.

Smoothed curvature

To improve curvature signal to noise ratio, for each point p'_j with coordinates (x'_j, y'_j) and refined curvature k'_j, we compute the smoothed curvature k''_j as a weighted moving average with triangular weights:

$$k''_j = \frac{\sum_{i=j-\beta n}^{j+\beta n} w_i \ k_i}{\sum_{i=j-\beta n}^{j+\beta n} w_i}$$

with $w_j = \beta n, \ldots, w_i = \beta n - |i-j|, \ldots, w_{j-\beta n} = w_{j+\beta n} = 0$ and where β is the smoothing parameter to adjust via learning. β describes the proportion of points (relative to the total number of contour points) to include in the smoothing. This implies that the smoothed curvature contour is $2\beta n$ points shorter (βn on each side) than the smoothed contour.

Landmark detection

Curvature around the start and end of the contour is noisy; it corresponds to a region of low curvature, at the beginning and end of the contour, outside of the region where we expect to find the five landmarks (Figure 1). In addition, the contour digitization by the user, which tends to start at a precise point and to end in a long stroke, results in a slight left-right asymmetry in the curvature profile. After superimposing all smoothed curvature profiles, we choose to exclude the first and last 20% of the smoothed contour. We find that the axis of symmetry (midline) is at position 0.475 instead of 0.5 for a symmetric profile. Landmarks are Bookstein's type 2 (local maxima of curvature) (Bookstein, 1992): maxima of the smoothed curvature for landmarks 1, 3, 5 and minima for landmarks 2 and 4. Having detected all minima and maxima, we first define landmark 3 as the maximum closest to the midline position, landmark 2 as the lowest minimum to the left of landmark 3 and landmark 1 as the maximum closest to landmark 2. Following the same logic, we define landmark 4 as the lowest minimum to the right of landmark 3 and landmark 5 as the maximum closest to landmark 4. Having detected all five landmarks, we found that there can seldom be more than one maximum between landmark 2 and landmark 4. In such situation, we allow resampling of landmark 3 to the highest maximum between landmarks 2 and 4. Finally, we exclude individuals that do not display all five landmarks after detection.

Spine thrust measure

Having detected all five landmarks, we quantify form using a measure previously introduced (Peluffo *et al.*, 2015), which is highly correlated to the Procrustes analysis principal component measure of inter-specific form variation, and which we called "spine thrust" (ST). ST is a measure of how much spines are elevated above the central ridge of the ventral branches and is computed as:

$$ST = \frac{1}{2}(Y_{L1} + Y_{L5}) - Y_{L3}$$

where Y_{L1}, Y_{L3} and Y_{L5} are the *Y* coordinate of landmarks 1, 3, 5, respectively. This measurement depends on the precise definition of *X* and *Y* axes. Here the *X*-axis is defined as the axis passing by landmarks 2 and 4 and oriented from 2 to 4, and with the *Y*-axis defined so that (X, Y) is an oriented orthonormal basis.

Machine learning

Detection of maxima and minima is a simple feature detection that relies on the derivative of the smoothed curvature profile. However, there are two parameters *α*, *β*, one for each smoothing filter (contour and curvature), which modulate the number and position of these

detected maxima and minima. It is possible to explore a set of values for α and β such that
the correlation between manually digitized landmarks and automatically detected landmarks
is optimized. Given that humans may introduce bias in the positioning of the landmarks, the
human output may not be optimal over the machine output. This is why we chose not to
quantify the learning success rate of our algorithm using the Area Under the Receiver
Operating Characteristic Curve but instead to search for combinations of parameter values
which yield the highest Pearson correlation value r^2 for ST measured over manually
digitized landmarks versus ST measured with automatically digitized landmarks.
344

Statistical analyses
All statistical analyses were performed using R version 3.4.3 (R Team 2016). We performed
two different sets of statistical analyses to investigate how ST changes across species, year
of collection and temperature. First, we fitted a standard multiple linear regression with
species, year of collection and temperature as numeric predictors using the standard R
function lm(). We chose the best model based on the variance explained provided by the
r^2 value. Table 2 presents the output of the lm() function using the R package jtools (v1.0.0)
(https://cran.r-project.org/web/packages/jtools/jtools.pdf) and its function export_summs()
with "scale" and "transform.response" set to "TRUE" which scales and centers the response
variable and reports standardized regression coefficients with their heteroskedasticity-robust
standard errors (see Table S1 for the raw lm() output). Second, we performed a regression
tree analysis and performed cross-validation using recursive partitioning with the regression
trees R package "rpart" version 4.1.13 (Therneau *et al.*, 2018) and the associated function
rpart() with the "anova" method and obtained the approximate r^2 from a 10-fold cross-
validation using the rsq.rpart() function. To confirm the importance of each factor on ST
change, we also performed random forest regression analysis using the R package
"RandomForest" version 4.6.14 and the randomForest() function in order. Both sets of
statistical analyses investigate the role of predictors in explaining a significant part of the
variance, multiple linear regression allows the use of interaction terms while regression trees
are easier to interpret (James *et al.*, 2013). In addition to these analyses, we systematically
plot distribution of ST across predictors (Fig. 2) showing individual values together with
mean, standard errors (which directly inform about two-by-two statistical significance
between groups), median, quartiles and estimates of the 95% confidence interval of the
medians, calculated as $\pm 1.58 \times \frac{IQR}{\sqrt{n}}$ where IQR is the interquartile range and n the number
of individuals for that IQR (Chambers *et al.*, 1983).
370

Results
Spine thrust (ST) can be measured semi-automatically
We previously reported that the shape of ventral branches in *D. santomea*, *D. yakuba* and
their hybrids can be characterized with a set of five manually detected landmarks, which
allows to calculate via simple arithmetic how much the lateral spines rise above the central
ridge, as a quantitative value named "spine thrust" (ST), expressed in micrometers (Peluffo
et al. 2015). The manual positioning of landmarks requires each point to be carefully
positioned on the exact feature for the ST measure to be exact. It can introduce between-
user and between-sample variability. In particular, the positioning of the three central
landmarks can be equivocal and may differ between users.
To use a less biased approach and automate the process, we decided to develop a
new measurement method that relies on manually digitized contours of the ventral branches,

which are easier to define than landmarks. The position contour of the ventral branches was
digitized by hand at an approximately four times faster rate than landmark detection,
because it can be done in a single stroke with a digital pen and the resulting ST measure is
barely sensitive to the exact pen position. We designed a pipeline that automatically
identifies the five landmarks based on the curvature of the manually digitized contours of the
ventral branches, and then calculates ST (Fig. 1). The typical rounded form of *D. santomea*
is then characterized by a null or negative value of ST (Figure 1, central panel) whereas the
spiny form of *D. yakuba* is characterized by a positive value of ST (Fig. 1, right panel). Note
that our method does not separate size and shape (Klingenberg, 2016), but considers
morphological form as a single quantifiable entity.
To assess repeatability, we digitized twice, at one month interval (at the beginning
and at roughly the midpoint of the digitizing effort), 30 individuals of the most characteristic
D. yakuba strain (Oku, sharp spines) and 31 individuals of the *D. santomea* strain which is
the most divergent from this *D. yakuba* strain (1563, extremely rounded shape and small
spines). We found a good correspondence   [2]between the two sets of automatic measures
(Fig. S2), indicating that our pipeline produces robust quantification of ventral branch form.
399
Learned α and β
401 We find that the same set of 30 *D. yakuba* and 31 *D. santomea* individuals is enough to
402 identify optimal parameter values for *α* (contour smoothing) and *β* (curvature smoothing).
403 We find that with $α = 0.025$ and $β = 0.055$ we obtain $r^2 = 0.91$ (Figure S1). Although a few
404 other combinations of *α* and *β* yield the same r^2 (Figure S1), we choose this set because it
405 is the one which applies the lowest degree of smoothing.
406
Strong interspecific difference in ST
In total, with our semi-automated method (and after removing $n = 71$ individuals
incorrectly dissected or mounted, 12% of total samples, with no apparent distribution bias),
we phenotyped 684 individuals raised at 18°C or 25°C throughout their development,
corresponding to four *D. yakuba* lines and seven *D. santomea* lines collected between 1998
and 2016 (Table 1). We checked all the automatically detected landmarks by eyes and found
that 30 individuals were incorrectly digitized, with a few landmarks either missing or
aberrantly positioned (see Fig. S3 for a sample of such individuals), and we excluded these
individuals (4% of 684) from subsequent analysis. These aberrant landmark profiles were
found in almost all the lines and at both temperatures, with no apparent distribution bias.
At 18°C and 25°C, for *D. santomea* and *D. yakuba*, in all 11 wild isofemale strains,
we observed within-strain variability in ST values (Figure 2A, for all groups, *n* per group is
between 26 and 31). At both temperatures, the mean ST of each of the seven *D. santomea*
strains is inferior to the mean ST of any of the four *D. yakuba* strain (Figure 2A). All *D.*
yakuba individuals have a positive ST, while most *D. santomea* strains have a mean ST
close to 0 (Fig. 2A). Accordingly, multiple linear regression analysis where the best fit model
is *ST ~ species x years x temperature* shows that the *species* independent variable explains
a significant part of the variance in ST ($p< 0.001$, Table 2). Overall, and despite within strain
variability as well as sensitivity to temperature variation, we confirm a morphological
difference of ventral branches between wild strains of *D. santomea* and *D. yakuba* using our
semi-automatic method of form quantification based on ST (Figure 2A, Table 2).
428

Table 2. Results of linear model fitting. This best model shows the contribution of each explanatory
variable, considered as a numerical value,and their interactions to the overall variance of ST in the full
D. santomea, D. yakuba dataset (shown in Figures 2A-B) at both temperatures (18°C and 25°C)
across all years using the standard R function lm(). The standardized effect values and their
heteroskedasticity-robust standard errors are reported together with the range of their *p*-values. For
example, species having an overall effect of 1.67 implies that going from *D. santomea* to *D. yakuba*
(all other things being equal) increases spine thrust absolute value by a relative (compared to the
other effects), dimension-less, value of 1.67. The raw effects together with the full output of the model
are provided in Table S1 (see methods). SE: standard error.

Factor *n* = 584, *r²* = 0.76	Effect	SE	*p*-value	Significance
Species	1.67	0.06	< 0.001	***
Years	0.54	0.06	< 0.001	***
Temperature	-0.29	0.05	< 0.001	***
Species x Years	-0.56	0.07	< 0.001	***
Species x Temperature	0.31	0.09	< 0.001	***
Years x Temperature	-0.23	0.09	< 0.05	*

439
440

Ventral branches of *D. santomea* are plastic to temperature

For *D. santomea*, in all strains but the oldest one collected in 1998, the mean ST is systematically smaller at 25°C compared to 18°C and standard errors do not overlap (Fig. 2). In contrast, no significant difference in mean ST between 25°C and 18°C is observed for *D. yakuba* strains, except for one strain collected in 2016 (*D. yakuba* Raphia) (Figure 2B). Multiple linear regression analysis supports a negative effect of temperature, as seen with *D. santomea* (p< 0.001, Table 2) and that effect is dependent on species (p< 0.001, Table 2). For the most recently collected wild strain of *D. santomea* (BM16.2), we compared the contours of the two most representative individuals raised at 18°C and 25°C, i.e., the two individuals with ST values closest to the median value of their group. We observed that the individual raised at 18°C has a more *D. yakuba*-like shape of ventral branches compared to the individual raised at 25°C (Fig. 3).

We find that the statistically significant effect of temperature on *D. santomea* only is also statistically dependent on the year at which the strain was collected (p < 0.05, Table 2). In order to interpret better our statistical analysis with multiple regression, we performed a 10-fold cross-validated regression tree analysis on the full data set (2 species, 11 strains, 584 individuals). The 10-fold cross-validated error rate is 0.3% and using an additive model of the shape *ST ~ species + years + temperature*. We found that the variance in the data set is first best partitioned by species and that temperature partitions the data set best for strains collected in 2015 and 2016 (Fig. 4, total variance explained as assessed by cross-validation r^2 is 0.77). To confirm those results, we also performed a random forest regression analysis with the same model as for the regression tree and found that the overall variance explained is $r^2 = 0.74$ and that the rank of importance of each independent variable is species > years > temperature. Altogether, our results show that in *D. santomea*, but not in *D. yakuba*, ventral branches are sensitive to temperature during development and that this effect is stronger in recently collected strains.

Temperature change phenocopies natural genetic variation for form

To compare the effects of temperature and of interspecific genetic variation on ventral branch form, we used our previous QTL mapping dataset of ventral branch form between *D. santomea* and *D. yakuba*, which comprises 365 *D. santomea* backcross individuals (Peluffo *et al.*, 2015). In this previous study, all flies were reared at 25°C as we found that this temperature was optimal to rear both species. The five landmarks were placed manually on images of the ventral branches. A generalized Procrustes analysis was performed on a set of 365 backcross progeny individuals and a larger dataset including the backcross progeny, F1 hybrids and parents. We found that, in both cases, the principal component PC1 explains an important part of the variance (58% in the full dataset and 41% in the backcross), that they are highly correlated (r² = 0.996) and that PC1 in the backcross is highly correlated to ST (spine thrust) (r² = 0.841) and not to centroid size (r² = 0.038).

This QTL mapping study revealed that a 2.7Mb locus on chromosome 3L explains 30% of the mean species difference in ST, meaning that replacing one *D. santomea* allele at this locus with a *D. yakuba* allele leads to an increase in ST of about 3 μm (Peluffo *et al.*, 2015). Pooling all the *D. santomea* lines examined in the present study, we find that a change in the raising temperature from 18°C to 25°C leads to an increase in ST of about 3.4 μm (Fig. 5). We conclude that the effect of temperature is as high as the effect of genetic variation at the major interspecific genetic locus.

While the ST density distribution of all *D. yakuba* individuals shows little overlap with the ST density distribution of all *D. santomea* individuals reared at 25°C, it overlaps more

489   with *D. santomea* individuals reared at 18°C (Fig. 5). Overall, our results show that
490   decreasing the temperature from 25°C to 18°C yield *D. santomea* males with spinier, *D.*
491   *yakuba*-like, ventral branches in the same way as introgressing a *D. yakuba* alleles in place
492   of a *D. santomea* allele at the major interspecific locus.
493
494
Discussion
A dataset-independent simultaneous quantification of shape and size
Our semi-automatic method, which relies on two simple layers of contour
transformation adjusted by regression based learning, is fast and allows the measure of form
variation through the simple outlining of ventral branches on 2D pictures. We note that in the
future, progress in edge detection algorithms (which for now introduce too much error to
measure with precision variations of the order of a few micrometers) might allow full
automation from pictures to form quantification.
Having drawn contours, we could also have relied on Fourier based analyses.
However, such methods require closed contours which in our case are difficult to draw since
the base of the ventral branches is a complex structure which cannot be easily delimited
from the cuticle of the ventral branches (Fig. 1). In addition, our method is more suitable for
contours in which very large and very small curvatures coexist. Furthermore, an important
limitation of morphometrics analyses on landmark data (e.g. Procrustes principal component
analysis) is that the PC values are dimensionless (Klingenberg, 2010) and may be difficult to
relate to physical features. With our simple measure of ST obtained from the automatically
detected landmarks, we are able to quantify and compare forms across studies. Importantly,
because we deal with absolute geometric measurements, our method simultaneously
analyzes shape and size, unlike most morphometric approaches (Claude, 2008;
Klingenberg, 2010). We believe this to be a strength in our case since both shape and size
of ventral branches probably contribute to the lock-and-key mechanism; e.g. spiny but short
D. yakuba ventral branches may not harm *D. santomea* females (Kamimura, 2012;
Kamimura & Mitsumoto, 2012b).
518
Effect of temperature on size and shape
In most insects and other ectotherms, adult body size typically increase with lower
temperatures (Angilletta Jr *et al.*, 2004). Bergmann's rule, which posits an increasing body
size with higher altitude, has been observed within the São Tomé island for the terrestrial
caecilian *Schistometopum thomense*, over a temperature range of 9°C (Measey & Van
Dongen, 2006). In contrast to other body parts, the genitalia of insects, and of *D.*
melanogaster in particular, have been reported as not, or little, plastic in response to
temperature or other types of environmental variation (Wheeler *et al.*, 1993; Eberhard, 2009;
Shingleton *et al.*, 2009). We find here that this is true also for *D. yakuba* but not for *D.*
santomea: changing the rearing temperature from 25°C to 18°C leads to an increase in
spine thrust in *D. santomea* male genitalia that phenocopies the morphological difference
observed between *D. santomea* and *D. yakuba*.
For each species, we find that strains raised in the same conditions display different
averages in ST, showing that the ventral branches form is influenced by genetic factors and
is able to evolve.
Plasticity of ventral branches form was detected for all the tested *D. santomea* strains
except the one that was maintained for the longest time in the laboratory. Furthermore, the
strains collected recently (in 2009, 2015 and 2016) display more pointed ventral branches at

18°C than the ones collected earlier. This suggests that as flies adapt to the laboratory
environment, the plasticity of ventral branches form towards temperature tends to be lost
and ventral branches tend to be more rounded. Recent studies show that *Drosophila* flies
can adapt to a laboratory environment in 20 generations only, which corresponds to about 8
months (Langmüller & Schlötterer, 2019).
Based on our experiments, we cannot fully rule out plasticity in *D. yakuba*. It is
possible that their genital morphology would be altered in external conditions outside of the
specific ones that we assayed here. In any case, we find that in our experimental conditions
the plasticity of genital form with respect to temperature is higher in *D. santomea* than *D.*
yakuba.
547
Laboratory observations should be complemented by analysis of wild-caught flies
Tests in the laboratory show that *D. santomea* flies appear to be poorly adapted to
high temperatures (Matute *et al.*, 2009). The optimal temperature for larval survival is 21°C
for *D. santomea* and 24°C for *D. yakuba.* Furthermore, when adult flies initially raised at
24°C are allowed to distribute themselves along a thermal gradient, they show a preference
for 23°C for *D. santomea* and between 26°C and 27°C for *D. yakuba* (Matute *et al.*, 2009).
These observations are in agreement with *D. santomea* being collected at higher altitudes
than *D. yakuba* in Sao Tomé. However, the reasons why the exact preferred temperature
values observed in the laboratory are different from the temperature values measured in the
geographic areas of the two species are unknown. Fly collections in Sao Tomé have mostly
been done on the north slopes of the island and in these areas *D. santomea* flies are found
at an altitude of 1150m or above (Lachaise *et al.*, 2000), which corresponds to temperatures
around 18°C or below (climate-data.org, www.worldclim.org/bioclim). However, we note that
on the southern slopes of the island a few *D. santomea* flies have also been collected at
lower altitudes (650m) in the dense mist forest near Rio Queijo (Matute & Coyne, 2010;
Nagy *et al.*, 2018). This suggests that *D. santomea* flies can also inhabit warmer regions of
the island and that they might be found across the native forest of Sao Tomé, which goes
down to sea level on the western slope of the island (Bell & Irian, 2019). Interestingly, this
type of coexistence is not unique on the island: two sister species of frogs closely match the
distribution of *D. santomea* and *D. yakuba*, respectively, with the endemic species
Hyperolius thomensis tied to wet forest habitats while its sister species *H. molleri* is in dry,
human-disturbed areas, and *H. thomensis* frogs have also been found in the southern forest
at 150 m (Bell & Irian, 2019).
It would be interesting to examine the genitalia of wild caught individual males of *D.*
santomea to check the form of their ventral branches at various altitudes. One possibility is
that at low altitudes in the southern part of the island *D. santomea* flies display rounded
ventral branches while in the hybrid zone with *D. yakuba* at 1150 m and at higher altitude*,*
where temperatures are 18°C or below, they have spinier ventral branches. Of note, *D.*
santomea flies have always been collected from traps and have never been observed
directly in their native environment: it is possible that they live in microenvironments whose
temperature is distinct from the one measured by climate stations (Feder *et al.*, 2000;
Negoua *et al.*, 2019).
580
Evolution of the plasticity of ventral branches form
To understand the relevance of this temperature sensitivity of genital form for the
past and present evolution of *D. santomea* and *D. yakuba*, more needs to be learnt about
their ecology and the plasticity of the ventral branches form of their closely related species,

D. teissieri. Ventral branches are only found in the three species of the *D. yakuba* complex,
D. santomea, *D. yakuba* and *D. teissieri* (Yassin & Orgogozo, 2013). Since ventral branch
form plasticity has not been studied in *D. teissieri*, it is unclear whether this plasticity to
temperature is an ancestral trait which has been lost in *D. yakuba* or if it is a novel trait which
evolved in *D. santomea* only. The species *D. teissieri* is not found in São Tomé but on the
mainland and a few islands of the African continent; it can hybridize with *D. yakuba* (Turissini
& Matute, 2017; Cooper *et al.*, 2018). In *D. teissieri* males, the spines are very long and no
layer of cuticle is present between them (Kamimura & Mitsumoto, 2012a; Yassin &
Orgogozo, 2013). In any case, even if the extent of ventral branch form plasticity in *D.
teissieri* was known, it would still be difficult to reconstruct ancestral trait states based on
only three species.
The female protective pouches, into which the spiny ventral branches of *D. yakuba*
males fit during copulation, were observed in *D. yakuba* but not in *D. santomea* females
raised at 21°C and 25°C (Kamimura & Mitsumoto, 2012b; Yassin & Orgogozo, 2013). It
would be interesting to check whether such pouches form in *D. santomea* females raised at
18°C, coinciding with the emergence of spiny ventral branches in males. Furthermore,
whether more pointed ventral branches in *D. santomea* males due to lower temperatures
affects copulation, reproduction and female physiology after mating is unknown.
If we assume that the São Tomé island species *D. santomea* arose from a *D.
yakuba*-like ancestor living on the African continent, one can hypothesize that regression in
ventral branch size and their plasticity evolved recently in the lineage leading to *D.
santomea*. Such a scenario is opposite to the most common view that posits that
morphological diversification tends to proceed through losses of plasticity, rather than gains
of plasticity ["flexible stem hypothesis" (West-Eberhard, 2003; Schneider & Meyer, 2017),
"plasticity-first" model (Levis & Pfennig, 2016) and "buying time" hypothesis (Pennisi, 2018)].
It is possible that the decrease in spine thrust that occurred during evolution in the lineage
leading to *D. santomea* was accompanied by a gain of ventral branches form plasticity
towards temperature. It is unclear whether the plasticity of ventral branches form to
temperature is adaptive. More knowledge about the ecology of *D. santomea* and its sister
species will be required to elaborate a convincing scenario to interpret the role of the ventral
branch form plasticity that we discovered.
616
Conclusion
Our data show that genitalia can be plastic to temperature and that this plasticity can evolve
coincidentally with speciation. Whereas the sensitivity of insect genitalia shape to
temperature or nutrition has been used previously as a proof against the lock-and-key
hypothesis (Arnqvist & Thornhill, 1998; Andrade *et al.*, 2005), our work suggests that
genitalia can be plastic without rejecting the lock-and-key hypothesis if the environmentally
induced changes do not hamper reproduction within each sister species lineage.
624
625
626
627
628   **Figures**
629
Figure 1. Landmark detection for ventral branches at 18°C and 25°C. For each
individual, a picture of the ventral branches is taken (top panel). The contour is digitized by
hand and smoothed (middle panel). The curvature along the contour is obtained by finite

differences, which are iterated for refining, and smoothed too (bottom panel). Smoothed
curvature (vertical axis), measured in inverse micrometers, is plotted along the contour,
starting from the leftmost point. The horizontal axis is the distance along the contour, called
the curvilinear abscissa, measured in micrometers. Here both axes are normalized by size
and represented in arbitrary units. In each plot the left dashed vertical line is the automatic
detection lower bound, the middle dashed line is the imputed global midpoint, the right
dashed line is the automatic detection upper bound (see methods). Red points represent
peaks and therefore curvature maxima whereas blue points represent cavities and therefore
curvature minima. Since *D. yakuba* is not sensitive to temperature, we only show one
characteristic shape. To understand where this genital structure is positioned within the male
genitalia, see Fig. 1 of Kamimura & Mitsumoto, 2012b.
644
Figure 2. Ventral branches form is sensitive to temperature in *D. santomea*. Isofemale
lines arranged by year of collection (see Table 1). (A) For each line, individual values (grey
points), median (thick black line), quartiles (colored boxplots), mean (colored points),
standard errors (black vertical segment over the colored points) and 95% confidence interval
estimates of the median (top and bottom notches) of automatically measured spine thrust
are shown. Each line was reared at 18°C (blue) or 25°C (red). (B) For each line, the same
mean and standard errors as in panel A are shown, together with the effect slope and
corresponding value of that effect (in μm).
653
654 **Figure 3. Difference in contour shape at 18°C versus 25°C within the same *D.**
655 ***santomea* isofemale strain collected in 2016**. Contours of the two individuals that have
656 the closest spine thrust value to the median value for D. santomea BM16.2 (right most strain
657 on Figure 2A) at 18°C and 25°C.
658
Figure 4. Regression tree for spine thrust measures of all *D. santomea* and *D. yakuba*
isofemale strains at both 18 and 25°C. Each node gives the spine thrust mean of all
samples included in that node and the proportion of the total dataset included in that node.
Below each node are two alternatives: to the left the condition is true and to the right the
condition is false. Note that the split between *D. santomea* and *D. yakuba* happens at the
top, thereby suggesting that neither temperature nor years have an effect on spine thrust
within *D. yakuba*.
666
667 **Figure 5. Density distribution of spine thrust values for *D. santomea* lines at 18 and**
668 **25°C and *D. yakuba* lines**. Density distribution inferred from *D. santomea* males raised at
669 18°C (blue) and 25°C (red), *D. yakuba* males at both temperatures (orange). Distributions
670 are inferred from the total data shown in Figure 2A.
671
672
673
674
675 **Supplementary Figures**
676
677 **Figure S1. Parameter adjustment for the machine detection algorithm**. Training the
678 algorithm relies on two layers of transformation that are each dependent on one parameter:
679 contour coordinate smoothing (horizontal axis) and curvature profile smoothing (vertical
680 axis). Training was performed using a set of 61 individuals, 31 *D. santomea* 1563 and 30 *D.*

yakuba Oku (see Table 1) for which we manually digitized both landmarks and contours. For each value of the two smoothing parameters, we performed linear regression of spine thrust from manually digitized landmarks against spine thrust derived from automatically digitized landmarks. The colors and values represent the r^2 from that regression. The value used for all detections is contoured in white.

Figure S2. Correlation between two sets of automatic measures from the same dataset. For the training dataset (31 *D. santomea* 1563 and 30 *D. yakuba* Oku), the same user digitized the same contours twice at one month interval and spine thrust was automatically measured. Each point represents one individual. The $y = x$ (black dashed line) and linear regression (full red line) are shown.

Figure S3. Representative samples of landmarks incorrectly identified with the machine detection algorithm. For each example, we show the smoothed contour, the corresponding curvature profile and identified landmarks.

Table S1. Raw output of R lm (formula = ST ~ Species * Year * Temperature).

References

Andrade, C.A., Hatadani, L.M. & Klaczko, L.B. 2005. Phenotypic plasticity of the aedeagus of Drosophila mediopunctata: effect of the temperature. *J. Therm. Biol.* **30**: 518–523.

Angilletta Jr, M.J., Steury, T.D. & Sears, M.W. 2004. Temperature, growth rate, and body size in ectotherms: fitting pieces of a life-history puzzle. *Integr. Comp. Biol.* **44**: 498–509.

Arnqvist, G. & Thornhill, R. 1998. Evolution of animal genitalia: patterns of phenotypic and genotypic variation and condition dependence of genital and non-genital morphology in water strider (Heteroptera: Gerridae: Insecta). *Genet. Res.* **71**: 193–212.

Bell, R.C. & Irian, C.G. 2019. Phenotypic and genetic divergence in reed frogs across a mosaic hybrid zone on São Tomé Island. *Biol. J. Linn. Soc.* **128**: 672–680.

Bookstein, F.L. 1992. *Morphometric tools for landmark data: Geometry and Biology*. Cambridge University Press, Cambridge, UK.

Cande, J., Andolfatto, P., Prud'homme, B., Stern, D.L. & Gompel, N. 2012. Evolution of Multiple Additive Loci Caused Divergence between Drosophila yakuba and D. santomea in Wing Rowing during Male Courtship. *PloS One* **7**: e43888.

Cariou, M.L., Silvain, J.F., Daubin, V., Da Lage, J.L. & Lachaise, D. 2001. Divergence between Drosophila santomea and allopatric or sympatric populations of D. yakuba using paralogous amylase genes and migration scenarios along the Cameroon volcanic line. *Mol. Ecol.* **10**: 649–660.

Chambers, J.M., Cleveland, W.S., Kleiner, B. & Tukey, P.A. 1983. *Graphical methods for data analysis.* Wadsworth & Brooks, Boston, USA.

Claude, J. 2008. *Morphometrics with R*. Springer Science & Business Media.

Comeault, A.A., Venkat, A. & Matute, D.R. 2016. Correlated evolution of male and female reproductive traits drive a cascading effect of reinforcement in Drosophila yakuba. In: *Proc. R. Soc. B*, p. 20160730. The Royal Society.

Cooper, B.S., Sedghifar, A., Nash, W.T., Comeault, A.A. & Matute, D.R. 2018. A maladaptive combination of traits contributes to the maintenance of a Drosophila hybrid zone. *Curr. Biol.* **28**: 2940–2947.

Cooper, B.S., Vanderpool, D., Conner, W.R., Matute, D.R. & Turelli, M. 2019. Wolbachia acquisition by Drosophila yakuba-clade hosts and transfer of incompatibility loci

between distantly related Wolbachia. *Genetics* **212**: 1399–1419.
Coyne, J.A., Elwyn, S., Kim, S.Y. & Llopart, A. 2004. Genetic studies of two sister species in
the Drosophila melanogaster subgroup, D. yakuba and D. santomea. *Genet. Res.* **84**:
11–26.
Coyne, J.A., Kim, S.Y., Chang, A.S., Lachaise, D. & Elwyn, S. 2002. Sexual isolation
between two sibling species with overlapping ranges: Drosophila santomea and
Drosophila yakuba. *Evolution* **56**: 2424–2434.
Coyne, J.A. & Orr, H.A. 2004. *Speciation*. Sinauer Associates, Sunderland, Massachusets.
Debat, V. & David, P. 2001. Mapping phenotypes: canalization, plasticity and developmental
stability. *Trends Ecol. Evol.* **16**: 555–561.
Dreyer, A.P. & Shingleton, A.W. 2011. The effect of genetic and environmental variation on
genital size in male Drosophila: canalized but developmentally unstable. *PLoS One*
6: e28278.
Dufour, L. 1844. Anatomie générale des diptères. *Ann. Sci. Nat.* **1**: 244–264.
Eberhard, W.G. 1988. *Sexual Selection and Animal Genitalia*. Harvard University Press.
Eberhard, W.G. 2009. Static allometry and animal genitalia. *Evol. Int. J. Org. Evol.* **63**: 48–
66.
Eberhard, W.G., Huber, B.A., Briceño, R.D., Salas, I. & Rodriguez, V. 1998. One size fits all?
Relationships between the size and degree of variation in genitalia and other body
parts in twenty species of insects and spiders. *Evolution* **52**: 415–431.
Fairbairn, D.J. 2005. Allometry for sexual size dimorphism: testing two hypotheses for
Rensch's rule in the water strider Aquarius remigis. *Am. Nat.* **166**: S69–S84. The
University of Chicago Press.
Feder, M.E., Roberts, S.P. & Bordelon, A.C. 2000. Molecular thermal telemetry of free-
ranging adult Drosophila melanogaster. *Oecologia* **123**: 460–465.
Gavin-Smyth, J. & Matute, D.R. 2013. Embryonic lethality leads to hybrid male inviability in
hybrids between Drosophila melanogaster and D. santomea. *Ecol. Evol.* **3**: 1580–
1589.
Gibert, J.-M. 2017. The flexible stem hypothesis: evidence from genetic data. *Dev. Genes*
Evol. **227**: 297–307.
Gibert, J.-M., Mouchel-Vielh, E., De Castro, S. & Peronnet, F. 2016. Phenotypic plasticity
through transcriptional regulation of the evolutionary hotspot gene tan in Drosophila
melanogaster. *PLoS Genet.* **12**: e1006218.
Hribar, L.J. 1996. Larval rearing temperature affects morphology of Anopheles albimanus
(Diptera: Culicidae) male genitalia. *J. Am. Mosq. Control Assoc.-Mosq. News* **12**:
295–297.
James, G., Witten, D., Hastie, T. & Tibshirani, R. 2013. Tree-based methods. In: *An*
introduction to statistical learning, pp. 303–335. Springer.
Jeong, S., Rebeiz, M., Andolfatto, P., Werner, T., True, J. & Carroll, S.B. 2008. The evolution
of gene regulation underlies a morphological difference between two Drosophila
sister species. *Cell* **132**: 783–793.
Kamimura, Y. 2012. Correlated evolutionary changes in Drosophila female genitalia reduce
the possible infection risk caused by male copulatory wounding. *Behav. Ecol.*
Sociobiol. **66**: 1107–1114.
Kamimura, Y. & Mitsumoto, H. 2012a. Genital coupling and copulatory wounding in
Drosophila teissieri (Diptera: Drosophilidae). *Can. J. Zool.* **90**: 1437–1440.
Kamimura, Y. & Mitsumoto, H. 2012b. Lock-and-key structural isolation between sibling
Drosophila species. *Entomol. Sci.* **15**: 197–201.
Klingenberg, C.P. 2010. Evolution and development of shape: integrating quantitative
approaches. *Nat. Rev. Genet.* **11**: 623.
Klingenberg, C.P. 2019. Phenotypic plasticity, developmental instability and robustness: the
concepts and how they are connected. *Front. Ecol. Evol.* **7**: 56.
Klingenberg, C.P. 2016. Size, shape, and form: concepts of allometry in geometric
morphometrics. *Dev. Genes Evol.* **226**: 113–137. Springer.
Lachaise, D., Cariou, M.-L., David, J.R., Lemeunier, F., Tsacas, L. & Ashburner, M. 1988.

Historical biogeography of the Drosophila melanogaster species subgroup. In:
789   *Evolutionary biology*, pp. 159–225. Springer.

790 Lachaise, D., Harry, M., Solignac, M., Lemeunier, F., Bénassi, V. & Cariou, M.L. 2000.
791   Evolutionary novelties in islands: Drosophila santomea, a new melanogaster sister
792   species from São Tomé. *Proc. Biol. Sci.* **267**: 1487–1495.

793 Lafuente, E. & Beldade, P. 2019. The genomics of developmental plasticity: recent progress
794   in animal models. *Front. Genet.* **10**: 720.

795 Langmüller, A.M. & Schlötterer, C. 2019. Low concordance of short-term and long-term
796   selection responses in experimental Drosophila populations. *bioRxiv* 759704.

797 LeVasseur-Viens, H., Polak, M. & Moehring, A.J. 2015. No evidence for external genital
798   morphology affecting cryptic female choice and reproductive isolation in Drosophila.
799   *Evolution* **69**: 1797–1807. Wiley Online Library.

800 Levis, N.A. & Pfennig, D.W. 2016. Evaluating 'plasticity-first'evolution in nature: key criteria
801   and empirical approaches. *Trends Ecol. Evol.* **31**: 563–574.

802 Liu, Y., Ramos-Womack, M., Han, C., Reilly, P., Brackett, K.L., Rogers, W., *et al.* 2019.
803   Changes throughout a Genetic Network Mask the Contribution of Hox Gene
804   Evolution. *Curr. Biol.*

805 Llopart, A., Elwyn, S., Lachaise, D. & Coyne, J.A. 2002. Genetics of a difference in
806   pigmentation between Drosophila yakuba and Drosophila santomea. *Evolution* **56**:
807   2262–2277.

808 Llopart, A., Lachaise, D. & Coyne, J.A. 2005a. An anomalous hybrid zone in Drosophila.
809   *Evol. Int. J. Org. Evol.* **59**: 2602–2607.

810 Llopart, A., Lachaise, D. & Coyne, J.A. 2005b. Multilocus analysis of introgression between
811   two sympatric sister species of Drosophila: Drosophila yakuba and D. santomea.
812   *Genetics* **171**: 197–210.

813 Masly, J.P. 2011. 170 years of "lock-and-key": genital morphology and reproductive isolation.
814   *Int. J. Evol. Biol.* **2012**.

815 Matute, D.R. 2010. Reinforcement of gametic isolation in Drosophila. *PLoS Biol.* **8**:
816   e1000341.

817 Matute, D.R. & Coyne, J.A. 2010. Intrinsic reproductive isolation between two sister species
818   of Drosophila. *Evol. Int. J. Org. Evol.* **64**: 903–920.

819 Matute, D.R., Novak, C.J. & Coyne, J.A. 2009. Temperature-based extrinsic reproductive
820   isolation in two species of Drosophila. *Evol. Int. J. Org. Evol.* **63**: 595–612.

821 Measey, G.J. & Van Dongen, S. 2006. Bergmann's rule and the terrestrial caecilian
822   Schistometopum thomense (Amphibia: Gymnophiona: Caeciliidae). *Evol. Ecol. Res.*
823   **8**: 1049–1059.

824 Moehring, A.J., Llopart, A., Elwyn, S., Coyne, J.A. & Mackay, T.F. 2006. The genetic basis of
825   postzygotic reproductive isolation between Drosophila santomea and D. yakuba due
826   to hybrid male sterility. *Genetics* **173**: 225–233.

827 Nagy, O., Nuez, I., Savisaar, R., Peluffo, A.E., Yassin, A., Lang, M., *et al.* 2018. Correlated
828   evolution of two copulatory organs via a single cis-regulatory nucleotide change.
829   *Curr. Biol.* **28**: 3450–3457.

830 Negoua, H., Chakir, M., David, J.R. & Capy, P. 2019. Climatic adaptation in Drosophila:
831   phenotypic plasticity of morphological traits along a seasonal cycle. In: *Annales de la*
832   *Société entomologique de France (NS)*, pp. 48–60. Taylor & Francis.

833 Peluffo, A.E., Nuez, I., Debat, V., Savisaar, R., Stern, D.L. & Orgogozo, V. 2015. A Major
834   Locus Controls a Genital Shape Difference Involved in Reproductive Isolation
835   Between Drosophila yakuba and Drosophila santomea. *G3 Bethesda Md*, doi:
836   10.1534/g3.115.023481.

837 Pennisi, E. 2018. *Buying time*. American Association for the Advancement of Science.

838 Price, T.D., Qvarnström, A. & Irwin, D.E. 2003. The role of phenotypic plasticity in driving
839   genetic evolution. *Proc. R. Soc. Lond. B Biol. Sci.* **270**: 1433–1440.

840 Rice, G., David, J.R., Kamimura, Y., Masly, J.P., Mcgregor, A.P., Nagy, O., *et al.* 2019. A
841   standardized nomenclature and atlas of the male terminalia of Drosophila
842   melanogaster. *Fly (Austin)* **13**: 51–64.

843   Schneider, R.F. & Meyer, A. 2017. How plasticity, genetic assimilation and cryptic genetic
844   variation may contribute to adaptive radiations. *Mol. Ecol.* **26**: 330–350.
845   Shapiro, A.M. & Porter, A.H. 1989. The lock-and-key hypothesis: evolutionary and
846   biosystematic interpretation of insect genitalia. *Annu. Rev. Entomol.* **34**: 231–245.
847   Annual Reviews 4139 El Camino Way, PO Box 10139, Palo Alto, CA 94303-0139,
848   USA.
849   Shingleton, A.W., Estep, C.M., Driscoll, M.V. & Dworkin, I. 2009. Many ways to be small:
850   different environmental regulators of size generate distinct scaling relationships in
851   Drosophila melanogaster. *Proc. R. Soc. B Biol. Sci.* **276**: 2625–2633.
852   Simmons, L.W. 2014. Sexual selection and genital evolution. *Austral Entomol.* **53**: 1–17.
853   Wiley Online Library.
854   Stern, D.L., Crocker, J., Ding, Y., Frankel, N., Kappes, G., Kim, E., *et al.* 2017. Genetic and
855   transgenic reagents for Drosophila simulans, D. mauritiana, D. yakuba, D. santomea,
856   and D. virilis. *G3 Genes Genomes Genet.* **7**: 1339–1347.
857   Therneau, T., Atkinson, B. & Ripley, B. 2018. *rpart: Recursive Partitioning and Regression*
858   *Trees. R Package. version 4.1-13*.
859   Turissini, D.A. & Matute, D.R. 2017. Fine scale mapping of genomic introgressions within the
860   Drosophila yakuba clade. *PLoS Genet.* **13**: e1006971.
861   Waddington, C.H. 1942. Canalization of development and the inheritance of acquired
862   characters. *Nature* **150**: 563.
863   West-Eberhard, M.J. 2003. *Developmental Plasticity and Evolution*. Oxford University Press.
864   Wheeler, D., Wong, A. & Ribeiro, J.M. 1993. Scaling of feeding and reproductive structures
865   in the mosquito Aedes aegypti L.(Diptera: Culicidae). *J. Kans. Entomol. Soc.* 121–
866   124.
867   Yassin, A. & David, J.R. 2016. Within-species reproductive costs affect the asymmetry of
868   satyrization in Drosophila. *J. Evol. Biol.* **29**: 455–460. Wiley Online Library.
869   Yassin, A. & Orgogozo, V. 2013. Coevolution between Male and Female Genitalia in the
870   Drosophila melanogaster Species Subgroup. *PLoS ONE* **8**: e57158.

	D. santomea		D. yakuba
	18°C	25°C	25°C

Peluffo et al. 2020

Table S1.

Call:
lm(formula = st ~ sp * year * temp, data = st_strains)

Residuals:
    Min   1Q Median   3Q   Max
-5.9035 -1.4710 -0.2689 1.3520 7.7223

Coefficients:
                 Estimate Std. Error t value Pr(>|t|)
(Intercept)   -7.505e+02 1.892e+02 -3.968 8.17e-05 ***
spYak   6.805e+02 2.138e+02   3.183 0.00154 **
year   3.760e-01 9.411e-02   3.995 7.30e-05 ***
temp   2.189e+01 8.710e+00   2.513 0.01224 *
spYak:year   -3.370e-01 1.064e-01 -3.167 0.00162 **
spYak:temp   -1.648e+01 9.816e+00 -1.679 0.09370 .
year:temp   -1.099e-02 4.333e-03 -2.536 0.01149 *
spYak:year:temp 8.301e-03 4.886e-03   1.699 0.08989 .

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.125 on 576 degrees of freedom
Multiple R-squared: 0.7556,	Adjusted R-squared: 0.7526
F-statistic: 254.3 on 7 and 576 DF, p-value: < 2.2e-16