Neural oscillations track the maintenance and

proceduralization of novel instructions

Oscillations in novel instructions implementation

Silvia Formica[a], Carlos González-García[a], Mehdi Senoussi[a], and Marcel Brass[a,b]

[a] Department of Experimental Psychology, Ghent University, Belgium

[b] School of Mind and Brain/Department of Psychology, Humboldt Universität zu Berlin,

Germany

Ghent University

Faculty of Psychology and Educational Sciences

Department of Experimental Psychology

Henri Dunantlaan 2

B-9000 Gent

Corresponding author: Silvia Formica (silvia.formica@ugent.be)

Abstract

Humans are capable of flexibly converting symbolic instructions into novel behaviors. Previous evidence and theoretical models suggest that the implementation of a novel instruction requires the reformatting of its declarative content into an action-oriented code optimized for the execution of the instructed behavior. While neuroimaging research focused on identifying the brain areas involved in such a process, the temporal and electrophysiological mechanisms remain poorly understood. These mechanisms, however, can provide information about the specific cognitive processes that characterize the proceduralization of information. In the present study, we recorded EEG activity while we asked participants to either simply maintain declaratively the content of novel S-R mappings or to proactively prepare for their implementation. By means of time-frequency analyses, we isolated the oscillatory features specific to the proceduralization of instructions. Implementation of the instructed mappings elicited stronger theta activity over frontal electrodes and suppression in mu and beta activity over central electrodes. On the contrary, activity in the alpha band, which has been shown to track the attentional deployment to task-relevant items, showed no differences between tasks. Together, these results support the idea that proceduralization of information is characterized by specific component processes such as orchestrating complex task settings and configuring the motor system that are not observed when instructions are held in a declarative format.

Keywords

Attention; Cognitive control; EEG oscillations; Instruction implementation; Motor preparation

Highlights

- Frontal theta power is increased during instructions implementation
- Attentional orienting in WM is analogous across maintenance and implementation
- Instructions implementation involves motor recruitment

1. Introduction

One peculiar aspect of human cognitive flexibility involves the ability to convert complex symbolic instructions into novel behaviors, even in the absence of prior practice (Cole, Laurent, & Stocco, 2013). Recent experimental evidence and theoretical proposals have put forward a serial heuristic model in which instructions are first encoded and maintained in a declarative format before being transformed into an action-oriented format (i.e., procedural representation) that allows executing the instructed task (Brass, Liefooghe, Braem, & De Houwer, 2017). Neuropsychological and behavioral evidence supports a dissociation between 'knowing' the content of an instruction and 'doing' the instructed cognitive or motor action, suggesting that maintaining its declarative content is insufficient to implement the instructed task, and that additional procedural codes are needed for optimal performance (Bhandari & Duncan, 2014; Duncan, Emslie, Williams, Johnson, & Freer, 1996; Liefooghe, Wenke, & De Houwer, 2012; Milner, 1963; Wenke, Gaschler, Nattkemper, & Frensch, 2009). Such immediate reformatting seems to occur only when the instruction is prepared for implementation, and not when participants merely maintain the information in a declarative manner, without the intention to implement (Liefooghe et al., 2012).

A growing number of fMRI studies has primarily focused on revealing which brain regions support the implementation of novel task sets (Demanet et al., 2016; González-García, Arco, Palenciano, Ramírez, & Ruz, 2017; Hartstra, Kühn, Verguts, & Brass, 2011; Hartstra, Waszak, & Brass, 2012; Palenciano, González-García, Arco, & Ruz, 2019; Ruge & Wolfensteller, 2010) and, more recently, on characterizing implementation-specific neural representations (Bourguignon, Braem, Hartstra, De Houwer, & Brass, 2018;

González-García, Formica, Wisniewski, & Brass, 2019; Muhle-Karbe, Duncan, De

Baene, Mitchell, & Brass, 2017; Palenciano, González-García, Arco, Pessoa, & Ruz,

2019; Ruge, Schäfer, Zwosta, Mohr, & Wolfensteller, 2019). These results consistently

point towards a crucial role of frontoparietal regions, and in particular of the prefrontal

cortex (PFC). Similarly, they provide evidence for distinct neural mechanisms supporting

implementing versus memorizing novel instructions (Brass et al., 2017). However, it is

still not clear to what extent declaratively representing information and proceduralization

involve processes that overlap or differ qualitatively. Crucially, investigating the oscillatory

dynamics of instruction implementation would allow to decompose the specific spectral

components that are involved in proceduralizing S-R mappings into action-oriented

codes, and to provide insights on their temporal unfolding. This would increase our

understanding of the functional significance of such processes, complementing the

knowledge on localization and representational format provided by fMRI research.

Specifically, we reasoned that both declarative maintenance and proactive

proceduralization of novel instructions likely engage certain shared processes, such as

selective attention towards the instructed content, whereas differences in brain activity

should reflect the reformatting process that is specific to the implementation of the

instruction. On the one hand, changes in the attentional focus are revealed by activity in

the alpha frequency range (8 – 14 Hz) (Hari & Salmelin, 1997; Klimesch, Sauseng, &

Hanslmayr, 2007; Senoussi, Moreland, Busch, & Dugué, 2019; van Ede, 2017).

Modulations of posterior alpha band oscillations are a well-established mechanism

associated with endogenous orienting of spatial attention and suppressing competition

from distractors, allowing for the enhancement of task-relevant information (Bonnefond &

Jensen, 2012; Jensen & Mazaheri, 2010; Sauseng et al., 2005). A substantial body of evidence reports decrease of alpha amplitude over posterior parietal regions contralateral to the attended spatial location, indicating mechanisms of orienting attention both to salient stimuli in the perceptual space and towards internally held representations (Capilla, Schoffelen, Paterson, Thut, & Gross, 2014; Gould, Rushworth, & Nobre, 2011; Mok, Myers, Wallis, & Nobre, 2016; Myers, Walther, Wallis, Stokes, & Nobre, 2015; Poch, Campo, & Barnes, 2014; Poch, Carretie, & Campo, 2017; Rihs, Michel, & Thut, 2007, 2009; Rohenkohl & Nobre, 2011; Schneider, Mertes, & Wascher, 2016; Thut, Nietzel, Brandt, & Pascual-Leone, 2006; van Dijk, Schoffelen, Oostenveld, & Jensen, 2008; Wallis, Stokes, Cousijn, Woolrich, & Nobre, 2015). Additionally, posterior alpha power has been observed to increase with the number of items relevant after attentional selection in a WM task (Jensen, 2002), also following retrospective selection (Manza, Hau, & Leung, 2014; Poch et al., 2017; Poch, Valdivia, Capilla, Hinojosa, & Campo, 2018). Therefore, we reasoned that alpha dynamics involved in the selection of internally maintained instructions should be independent of whether the task requires their implementation or mere maintenance.

On the other hand, we hypothesized that instruction implementation relies on a specific set of processes that result in the reformatting of the declarative content into action-oriented code. Therefore, we expected some oscillatory features to be specifically associated with proceduralization. First, the proactive reformatting of the instruction into a proceduralized action-bound representation should involve the allocation of cognitive control to allow the binding of stimulus and response. Ample evidence has associated control-related top-down mechanisms with larger theta (3 - 7 Hz) amplitude over mid-

frontal scalp electrodes (Cavanagh & Frank, 2014; Cohen & Donner, 2013). Recent computational models attribute to theta a central role in orchestrating long-range neural communication and in binding task-relevant information (Senoussi et al., 2020; Verbeke & Verguts, 2019; Verguts, 2017). At the same time, studies showed increased theta power for WM manipulation (Itthipuripat, Wessel, & Aron, 2013; Onton, Delorme, & Makeig, 2005). Therefore, we hypothesized stronger mid-frontal theta power during preparation for instruction implementation compared to maintenance, reflecting the proactive reformatting of the S-R mapping into a proceduralized action-bound representation. Nevertheless, behavioral evidence suggests that only a small set of mappings can be proceduralized and maintained in such a format prior to the moment of their actual implementation, hinting at a limit in the number of novel instructions that can be proceduralized at once (Liefooghe et al., 2012). As we expect theta power to track the occurrence of such reformatting, we reasoned that theta dynamics should not differ from declarative maintenance when the number of mappings to implement exceeds capacity limits.

Finally, we reasoned that proactively recoding the instruction for execution, and not merely maintaining it, should induce the preparation of the instructed motor plan (Everaert, Theeuwes, Liefooghe, & De Houwer, 2014; Meiran, Pereg, Kessler, Cole, & Braver, 2015a). Therefore, we looked at suppression in beta (15 – 30 Hz) and mu (8 – 12 Hz) frequency bands over central electrodes, oscillatory features traditionally associated with motor preparation and motor imagery (Cheyne, 2013; Pineda, 2005; Rhodes, Gaetz, Marsden, & Hall, 2018; Schneider, Barth, & Wascher, 2017; Tzagarakis, West, & Pellizzer, 2015). We expected this modulation to be larger while preparing to implement

the instructions, thus when an action-bound procedural representation is available, compared to the declarative maintenance of the mappings. Again, we predicted such difference to be reduced with a higher number of instructions, due to a limit in the number of action plans that can be prepared and maintained ready simultaneously.

To test our hypotheses, we recorded EEG activity while participants performed a task that encouraged a declarative maintenance of novel S-R mappings for recognition (Memorization task), and an Implementation task, intended to prompt their proactive reformatting for execution. In both cases, after the presentation of four mappings at the beginning of the trial, a retro-cue selected a subset of them as potential targets for the subsequent recognition or execution task. We reasoned that brain activity during the Cue-Target interval (CTI) would reflect the two-steps action described in a recent model of information prioritization in WM (Myers, Stokes, & Nobre, 2017). According to Myers and colleagues (2017), attention needs to be internally oriented towards the selected items, and the prioritized representations can then be reformatted into a behavior-optimized state. In the context of instruction implementation, the task-optimized representation is a procedural, action-bound code of the S-R mapping. On the other hand, successful maintenance is optimally achieved by means of a declarative representation of stimulus and response that does not entail any action plan. Therefore, during the CTI of the two tasks, we expected analogous features associated with attentional orienting, but instances of reformatting to be larger in preparation for instruction implementation.

2. Materials and Methods

2.1 Participants

Thirty-nine participants took part in the experiment (M_{age} = 21.74, *SD* = 4.50, 33 females) and received 30 euros as compensation. Sample size was not computed a priori: we aimed for thirty-five participants, which is on the upper bound of the sample size of studies investigating similar constructs (de Vries, Van Driel, Karacaoglu, & Olivers, 2018; Schneider et al., 2017; van Ede, Chekroud, Stokes, & Nobre, 2019a). All participants had normal or corrected-to-normal vision and thirty-three reported to be right-handed. Data from two participants were discarded due to low task performance (individual mean accuracy exceeded by 2.5 standard deviations the group mean accuracy in one of the two tasks and/or accuracy in response to catch trials in one of the tasks was below 60%); data from two additional participants were discarded following visual inspection because of excessive noise in the EEG recordings, resulting in a final sample size of thirty-five participants. All participants gave their informed consent prior to the beginning of the experiment, in accordance with the Declaration of Helsinki and the protocols of Ghent University.

2.2 Materials

The same set of stimuli was used as in previous studies on instruction implementation (Formica, González-García, & Brass, 2020; González-García, Formica, Liefooghe, & Brass, 2020). It consisted of 1550 pictures, grouped in two macro categories: animate (non-human animals) and inanimate (vehicles and musical instruments) (Brady, Konkle, Alvarez, & Oliva, 2013; Brodeur, Guérard, & Bouras, 2014; Griffin, Holub, & Perona, 2007; Konkle, Brady, Alvarez, & Oliva, 2010). All images had their background removed, were centered in a 200x200 pixels square and were converted to grayscale. Stimuli

presentation and response collection were performed using Psychopy toolbox (Pierce,

2007).

2.3 Experimental design

Participants performed the two tasks (from now on "Implementation task" and

"Memorization task") in a blocked fashion during one single session, and the order of the

two tasks was counterbalanced between participants. The structure of the trials for the

two tasks was identical up to the presentation of the target (Figure 1). Each trial started

with a red fixation cross presented for 2000ms (± 100ms, jittered) signaling the inter-trial

interval and allowing participants to blink if needed, followed by a white fixation cross for

250ms. Next, the encoding screen containing four S-R mappings (arranged in two rows)

appeared for 5 seconds. Mappings consisted of a new image associated with a bimanual

response: "index" referred to both index fingers (keys r and i) and "middle" referred to

both middle fingers (keys e and o). These bilateral responses were used instead of the

more traditional left and right options to avoid automatic motor activations elicited by the

mere presentation of lateralized response words (Bundt, Bardi, Abrahamse, Brass, &

Notebaert, 2015). Out of the four presented mappings, two contained images of animals

and two images of inanimate objects. Each image-response combination was presented

only once during the whole duration of the experiment. The rationale of this choice was

to maximize WM demands and minimize the retrieval of practiced S-R mappings from

long-term memory (Meiran, Pereg, Kessler, Cole, & Braver, 2015b). After a 750ms delay,

which is considered an interval sufficiently long to discard a contribution of iconic memory

(Souza & Oberauer, 2016), the retro-cue appeared and remained on the screen for

250ms. This consisted of a square presented centrally, with one, two or four (i.e., neutral

retro-cue) colored corner(s)[1]. The number of selected mappings is the second crucial

factor in our design, namely Load. Participants were instructed that the retro-cue signaled

which mapping(s) could be probed, with 100% validity (except in the case of catch trials,

see below). Importantly, when the retro-cue selected two mappings, these were always

on the same side of the screen. The subsequent cue-target interval (CTI) had a jittered

duration, lasting on average 1750ms (± 100ms). This interval largely exceeds the 300 –

500 ms considered necessary for retro-cues to benefit performance (Souza & Oberauer,

2016) and sufficiently long to allow time-frequency decomposition of low frequencies

(Cohen, 2014b). Finally, participants were presented with the target screen, which

differed depending on the task. In the Implementation task, the image of one of the cued

mappings was presented centrally and participants were required to press the associated

pair of keys (both index or both middle fingers). In the Memorization task, one mapping

(i.e., image and associated response) was presented centrally. The image in the

Memorization probe belonged to one of the selected mappings (except for catch trials,

see below). Participants had to report whether the presented mapping matched one of

those selected by the retro-cue, by pressing with both fingers of one hand for "yes" or

both fingers of the other hand for "no". The sides for "yes" and "no" were randomly

assigned on a trial basis to ensure that participants could not prepare any response during

the CTI, therefore encouraging the mere declarative maintenance of the instruction

memoranda. Labels with "yes" and "no" appeared at the bottom of the target screen

together with the mapping. In 50% of the trials, the target screen showed the same

---

[1] The physical properties of the retro-cue could potentially produce a combination of exogenous and endogenous orienting of attention in Load 1 and 2 (i.e., one side of it is colored and thus more salient). However, we are not strongly interested in this distinction and aim at using the retro-cue as the event to locate in time the onset of different demands between tasks.

mapping as in the encoding ("yes" response). In the other 50% of trials, one of the

selected images was presented associated with a different response with respect to

encoding ("no" response). Crucially, despite the similarities in trial structure, the

Memorization task relied on the declarative maintenance of the mappings as no response

could be prepared, whereas the Implementation task encouraged a proactive preparation

of the selected mappings for the execution of the required action plan. Both tasks included

catch trials (~25% for each task), in which the target screen displayed a different image

than the four displayed in the encoding screen. In this case, participants were required to

press the spacebar. This was done to discourage participants to adopt specific strategies,

such as memorizing only the mappings associated with one response option. Since we

were interested in the brain activity before the onset of the target, EEG recordings from

catch trials were analyzed together with regular trials. For each task, participants

completed 5 experimental blocks, for a total of 180 trials (60 per load, Figure 1).

Each of the two tasks was preceded by a practice session. This consisted of mini-blocks

of 12 trials, including all possible load conditions (i.e. number of mappings selected by

the retro-cue) and at least one catch trial. The only difference with the main task was the

presence of feedback at the end of each trial, signaling the accuracy of the response or

encouraging participants to respond faster in case no response was registered within the

maximum response time of 3 seconds. Performance was assessed at the end of each

mini-block: if accuracy was above 80%, practice was concluded, otherwise a new mini-

block started, up to a maximum of 3 blocks. S-R mappings used during the practice were

never presented again during the main task. The total duration of the experiment,

including cap preparation, practice, main tasks and breaks was approximately 150

minutes.

Figure1: Behavioral paradigm. Participants had to first encode four S-R mappings, for subsequent

recognition (Memorization task) or for a choice-reaction task (Implementation task). During the retention

interval, a Retro-Cue selected one, two or all four mappings as relevant for the upcoming probe. Participants

performed the two tasks separately in a block design.

2.4 EEG Recordings and pre-processing

Electrophysiological data were recorded using a BioSemi ActiveTwo system (BioSemi,

Amsterdam, Netherlands) with 64 Ag–AgCl electrodes arranged in the standard

international 10–20 electrode mapping (Klem, Lüders, Jasper, & Elger, 1999), with a

posterior CMS-DRL electrode pair. Two reference electrodes were positioned at the left

and right mastoids. Eye movements were registered with a pair of electrodes above and

below the left eye and two additional electrodes at the outer canthi of both eyes. EEG

signals were recorded at a 1024 Hz sampling rate.

EEG data were preprocessed using the Fieldtrip toolbox (Oostenveld, Fries, Maris, &

Schoffelen, 2010), running in MATLAB (MATLAB R2017b, The MathWorks, Inc., Natick,

Massachusetts, United States). First, the data were downsampled to 512 Hz and re-

referenced to the average of the mastoids. Then, a $0.5 - 45$ Hz band-pass FIR filter was

applied to the data, together with a Notch filter at 50 Hz and its harmonics. Data were

epoched relative to the onset of the retro-cue (from -1000 to 2500 ms) and demeaned to

the average of the whole epoch, to improve independent component analysis (ICA)

(Groppe, Makeig, & Kutas, 2009). Only trials in which the participant performed the

correct response were retained for subsequent analyses. Trials exhibiting movement

artifacts or excessive noise were removed following visual inspection of the data. Next,

eye movements artifacts were removed by means of ICA, using the EEGLAB (Delorme

& Makeig, 2004) runica algorithm as implemented in Fieldtrip. Components to discard

were selected based on their topography, the correlation between their time course and

the horizontal and vertical electrooculography, and their power spectrum. For most

participants, two components were removed (capturing blinks and horizontal eye

movements, respectively); in six participants blinks were reflected in two components,

leading to the removal of three IC, and in two participants only one component was

removed. Finally, data were visually inspected again to ensure successful cleaning and

one (N = 6), two (N = 2) or three (N = 1) excluded channels were interpolated by means

of spherical spline interpolation (Perrin, Pernier, Bertrand, & Echallier, 1989). This

cleaning procedure resulted in an average of 151.09 trials for the Implementation task

(*SD* = 12.29, 83.94%) and 143.06 trials for the Memorization task (*SD* = 14.07, 79.48%).

For each load condition, an average of 81.7% of trials were retained (Implementation,

Load 1: 52.14 (*SD* = 3.64) trials, Load 2: 50.26 (*SD* = 5.19) trials, Load 4: 48.68 trials (*SD*

= 5.74); Memorization, Load 1: 49.88 trials (*SD* = 4.84), Load 2: 47.86 trials (*SD* = 5.86),

Load 4: 45.31 trials (*SD* = 5.94).

2.5 Spectral Analysis

For each Task x Load condition (Memorization-Load 1, Memorization-Load 2,

Memorization-Load 4 and Implementation-Load 1, Implementation-Load 2,

Implementation-Load 4), time-frequency analysis was performed separately using

complex Morlet wavelet convolution, to estimate spectral power from 1 to 45 Hz in steps

of 1 Hz. The number of cycles in the wavelet was frequency-specific, ranging from 2 at 1

Hz and linearly spaced up to 7 cycles at 45 Hz, to achieve a good trade-off between

temporal and frequency precision (Cohen, 2014a). This analysis resulted in one time-

frequency spectrogram for each channel, condition and participant. Condition-specific

decibel normalization was then applied, using the time window between -500 and -200ms

before the onset of the retro-cue as baseline. To avoid data contamination due to the

smearing in time of the processing of the target, all statistical analyses are performed on

the time window from 0 to 1800ms with respect to retro-cue onset, therefore leaving

100ms gap between the end of our analyses window and the earliest jittered target onset.

2.6 Statistical Analyses

Reaction times (RTs), error rates (ER) and averaged alpha power (see below) were

separately entered in 2 (Task: Memorization vs Implementation) x 3 (Load: 1, 2, 4)

repeated measure ANOVAs, performed in JASP (Jasp Team, 2019).

To evaluate the statistical significance of differences between EEG time-courses or time-

frequency spectra, we adopted a cluster-based permutation approach (Maris &

Oostenveld, 2007), which is appropriate to assess the reliability of neural patterns over

neighboring data points. Moreover, this approach is robust against the multiple-

comparison problem, as the significance of clusters found in the observed group-level

data is estimated against a distribution of clusters obtained by randomly permuting the

assignement of participants to each group. First, depending on the number of conditions

to be compared, we performed either a two-sided t-test (two conditions) or an F-test (three

conditions), at an α level of 0.05, between all data points of the observed conditions.

Then, we considered as cluster a group of adjacent data points with same sign

significance and as cluster-size the sum of all t- or F-values in the cluster. Next, we used

5,000 permutations of participant-level data to estimate a distribution of cluster sizes

under the null hypothesis that there were no differences between conditions. The *P*-value

for each cluster in the observed group-level data corresponds to the proportion of

permutations in which the largest cluster size was larger than the size of the considered

observed cluster. Again, we used a significance alpha level of 0.05, therefore only

observed clusters whose size was larger than the size of the largest cluster in at least

95% of permutations are reported. This approach provides a statistically sound procedure

to compare two conditions and establish the existence of a difference between them,

reliably accounting for multiple comparisons. However, because clusters found through

this procedure reflect the result of the second level inference between the observed and

the null permuted distributions, no strong conclusions can be drawn with respect to their

boundaries, such as the exact latency or frequency range of the effect (Sassenhagen &

Draschkow, 2019).

2.7 Contralateral alpha suppression

Consistent findings reported alpha desynchronization over posterior regions contralateral

to the attended spatial location, indicating top-down anticipatory mechanisms of

attentional orienting in the perceptual or internal space (Bonnefond & Jensen, 2012;

Capilla et al., 2014; Gould et al., 2011; Jensen & Mazaheri, 2010; Mok et al., 2016; Myers

et al., 2015; Rihs et al., 2007, 2009; Rohenkohl & Nobre, 2011; Sauseng et al., 2005;

Thut et al., 2006; van Dijk et al., 2008; Wallis et al., 2015).

To estimate the suppression of alpha oscillations contralateral to the cued hemifield, we

averaged the time-frequency spectra corresponding to Load 1 and Load 2, separately for

the two tasks. Trials with neutral retro-cues (Load 4) were excluded from this analysis

because they did not induce attentional orienting towards one of the two visual hemifields.

Based on previous literature, two pairs of electrodes were selected in right (P8, PO8) and

left (P7, PO7) posterior parietal regions (de Vries, van Driel, & Olivers, 2019a; Gould et

al., 2011; Schneider et al., 2017; van Ede, Chekroud, Stokes, & Nobre, 2019b). Power

spectra were extracted from these electrode pairs, averaged in the alpha frequency range

(8 – 14 Hz), and collapsed between retro-cues pointing to the left and right hemifield, in

order to extract one power time series for the contralateral hemisphere and one for the

ipsilateral hemisphere. The two time series were compared by means of cluster-based

permutation testing, separately for the two tasks (see Statistics section). Additionally, the

cluster-based permutation approach was also used to compare the contralateral alpha

suppression between the two tasks. Namely, for each task the ipsilateral power time

series was subtracted to the contralateral one, separately for each participant, and the

resulting difference waves were compared. This analysis was performed to test whether

the deployment of alpha oscillations to orient attention towards relevant mental

representations differed between tasks.

2.8 Load-dependent alpha increase

Posterior alpha power has been observed to increase with the number of items relevant

after attentional selection in a WM task (Fukuda, Mance, & Vogel, 2015; Jensen, 2002),

also following retrospective selection (Manza et al., 2014; Poch et al., 2017, 2018).

Observing such modulation in our data would be evidence in favor of an effective selection

of the relevant mappings. To investigate the increase in alpha power relative to the

number of retained items, we extracted the average power for the frequencies ranging

from 8 to 14 Hz, separately for each condition and task, from the electrode pair Pz and

POz (Jensen, 2002), across the whole time window from 0 to 1800ms with respect to the

onset of the retro-cue. This operation led to one averaged alpha power value for each

Task-Load combination per participant, that were entered in a 2 (Task) x 3 (Load)

repeated measure ANOVA.

2.9 Mid-frontal theta in Implementation

The two tasks differ to the extent they require the reformatting of the encoded information

in an action-oriented code. We hypothesized such transformation to rely on a top-down

mechanism supported by low frequency oscillations. To test our predictions, time-

frequency spectra were extracted from the CTI (0 – 1800 ms) in the frequency range 3 –

7 Hz, averaged from channels Fz and AFz. The choice of these two channels was guided

by both their widespread use in the literature concerning midfrontal theta (Berger et al.,

2019; Cooper et al., 2019; Onton et al., 2005; Popov et al., 2018; Senoussi et al., 2020)

and a condition-independent channel selection procedure. Namely, we fitted the power

spectra from all channels, computed across all trials, regardless of the Load condition

and the Task, with the FOOOF toolbox (Haller et al., 2018). This method allows to

decompose different components of the power spectrum (i.e., the aperiodic (1/f pattern)

and periodic components) and thus to better capture the oscillatory profile of

electrophysiological data. This analysis confirmed that, across conditions, channels Fz

and AFz exhibited the largest peaks in the power spectra between 3 and 7 Hz (see

Supplementary materials for details).

We tested the main effect of Task by comparing the time-frequency spectra of

Implementation and Memorization (averaged across levels of Loads) by means of cluster-

based permutation test. Similarly, we tested the main effect of Load by contrasting the

spectra for each level of Load, averaged across Task condition. We then tested for a

significant effect of the interaction between our two factors by running a cluster-based

permutation test on the three time-frequency spectra resulting from pairwise subtraction

of the two tasks for each Load level (i.e., Implementation 1 minus Memorization 1,

Implementation 2 minus Memorization 2 and Implementation 4 minus Memorization 4).

2.10 Motor-related mu and beta suppression in Implementation

Analogously, we hypothesized that action-oriented procedural representations should

entail a motor plan, reflected in stronger signatures of proactive motor preparation. We

expected differences in two frequency bands associated with motor preparation, namely

mu (8 -12 Hz) and beta (15 – 30 Hz) (Cheyne, 2013; Pineda, 2005). To test for this, we

extracted time-frequency spectra averaged across electrodes C3 and C4 from the CTI (0

– 1800 ms). Next, we again used a cluster-based permutation approach outlined for mid-

frontal theta, to look for significant clusters of different activity between tasks, loads, and

their interaction. We selected electrodes C3 and C4 because we expected the effect to

be maximal over motor and pre-motor cortices[2] (Marchesotti, Bassolino, Serino, Bleuler,

& Blanke, 2016; McFarland, Miner, Vaughan, & Wolpaw, 2000). Moreover, we averaged

across the two of them because the goal of this comparison was to investigate pre-

movement preparatory oscillatory activity, rather than the effector-specific M1 activation

(Neuper & Pfurtscheller, 2001; Pfurtscheller, Brunner, Schlögl, & Lopes da Silva, 2006;

Pfurtscheller & Neuper, 1997). In our study, responses were bimanual in the

Implementation task and lateralized in the Memorization task. However, we reasoned that

our test is independent of such difference, as before target onset responses from both

hands had to be prepared in the two tasks.

3. Results

3.1 Behavioral results

Concerning behavioral performance, we expected valid retro-cues to have a beneficial

effect on reaction times and error rates in both tasks (Souza & Oberauer, 2016). Repeated

---

[2] Electrodes selection for analyses in the mu and beta frequency ranges was based on the literature and on the clearer assumptions on the brain areas causing the expected oscillations (i.e., pre-SMA and M1). Moreover, we chose to not rely on an independent electrode selection as we did for theta because the FOOOF toolbox results in higher frequencies are more susceptible to muscular artifacts and therefore less accurate.

measures ANOVA on RTs confirmed a significant main effect of Load[3] ($F_{1.44, 49.06}$ =

104.65, $p < 0.001$, $\eta^2_p = 0.76$): RTs increased with the number of mappings selected by

the retro-cue (Load 1: M = 978ms, SD = 196, Load 2: M = 1168ms, SD = 150, Load 4: M

= 1266ms, SD = 142). The main effect of Task was also significant ($F_{1, 34}$ = 237.20, $p <$

0.001, $\eta^2_p = 0.87$), with slower RTs in Memorization (M = 1355ms, SD = 185) than

Implementation (M = 907ms, SD = 160). The interaction between the two factors also

resulted to be significant ($F_{1.32, 44.88}$ = 4.42, $p = 0.031$, $\eta^2_p = 0.11$). More specifically, the

effect of Load was larger in the Implementation task ($F_{1.44, 49.06}$ = 93.20, $p < 0.001$)

compared to the Memorization task ($F_{1.44, 49.06}$ = 45.50, $p < 0.001$) (Figure 2a).

Error rates showed a significant main effect of Task ($F_{1, 34}$ = 11.08, $p = 0.002$, $\eta^2_p = 0.25$)

and of Load ($F_{1.67, 57.91}$ = 25.65, $p < 0.001$, $\eta^2_p = 0.43$). Participants were significantly

more accurate in the Implementation task (M = 0.14, SD = 0.09) compared to the

Memorization task (M = 0.19, SD = 0.10), and were more accurate when less items were

selected by the retro-cue (Load 1: M = 0.12, SD = 0.07, Load 2: M = 0.17, SD = 0.10,

Load 4: M = 0.20, SD = 0.12). The interaction between the two factors was not significant

($p = 0.266$) (Figure 2b).

Regarding catch trials, participants could successfully detect a new image in both the

Memorization task (error rate in catch trials: M = 0.13, SD = 0.08) and the Implementation

task (M = 0.09, SD = 0.07). Nevertheless, they were significantly less accurate (t_{34} = 3.16,

$p = 0.003$, $d = 0.53$) in the Memorization task.

---

[3] Mauchly's test revealed that the assumption of sphericity is violated ($p < 0.05$). Greenhouse-Geisser correction is applied here and in all results where the sphericity assumption is violated.

Such differences in performance between Implementation and Memorization are not

surprising and can be at least partially attributed to the task-specific processes taking

place after the occurrence of the target (Formica et al., 2020; Muhle-Karbe et al., 2017).

Specifically, the Memorization task requires the comparison of the presented mapping to

the encoded ones, the identification of the response side, and the execution of the

intended response. On the contrary, the Implementation task only involves the

identification of the target stimulus and the execution of the associated motor response.

Figure 2: Behavioral results. **a**) Reaction times (ms). **b**) Error rates. Both measures confirmed a benefit

in performance (i.e., faster RTs and less errors) when the Retro-Cue selected less mappings. In each

boxplot, the thick line inside box plots depicts the second quartile (median) of the distribution (n = 35). The

bounds of the boxes depict the first and third quartiles of the distribution. Whiskers denote the 1.5

interquartile range of the lower and upper quartile. Dots represent individual subjects' scores.

3.2 EEG Results

3.2.1 Contralateral alpha power decrease

As hypothesized, we observed a stronger reduction in alpha power in contralateral

compared to ipsilateral electrodes across loads 1 and 2 both in Implementation ($P =$

0.013, cluster-corrected) and Memorization ($P = 0.003$, cluster-corrected) tasks (see

Supplementary material for the same analysis performed separately for the two load

conditions). The significant clusters spanned the time window around 600 – 800ms.

Notably, we did not find any difference between the two tasks (no cluster survived multiple

comparison correction, even at a cluster threshold of $P = 0.1$), supporting our hypothesis

that the mechanisms allowing for attentional orienting are engaged to a similar extent in

the two tasks (Figure 3a). To qualitatively confirm our electrodes selection and the

location of the significant effect, contralateral minus ipsilateral half-topographies are

reported separately for each task (Figure 3b).

3.2.2 Load-dependent alpha increase

We predicted alpha power to also track the number of items selected by the retro-cue to

be relevant for the ongoing task. The repeated measures ANOVA on the averaged alpha

power over Pz – POz in the whole CTI (0 – 1800ms) revealed a significant main effect of

Load ($F_{2, 68} = 24.81$, $p < 0.001$, $\eta^2_p = 0.42$) (Figure 3c). Specifically, alpha power values

increased with the number of retained items (Load 1 vs Load 2: $t = -3.24$, $p = 0.006$

Bonferroni-corrected; Load 2 vs Load 4: $t = -3.38$, $p < 0.001$ Bonferroni-corrected), both

in Implementation (Load 1: $M = 0.83$, $SD = 1.83$, Load 2: $M = 1.12$, $SD = 1.83$, Load 4: M

$= 1.42$, $SD = 1.92$) and Memorization (Load 1: $M = 0.77$, $SD = 1.78$, Load 2: $M = 1.22$,

$SD = 1.87$, Load 4: $M = 1.79$, $SD = 1.90$). Crucially, neither the effects of Task nor the

interaction Task x Load were significant (ps > 0.1). Scalp topographies of alpha power

across the entire CTI are reported separately for each Task and Load condition (Figure

3d).

Figure 3: Alpha power dynamics. **a**) Time courses of the difference waves (contralateral minus ipsilateral time course) of alpha power activity between contralateral and ipsilateral electrode clusters (PO8 - P8 and

PO7 - P7). Both difference waves show a significant deflection from 600 to 800ms after the onset of the

retro-cue. Horizontal lines above the x-axis indicate significant temporal clusters (blue: Implementation, red:

Memorization), obtained with cluster-based permutation testing). Shading represents the s.e.m., calculated

across participants (n = 35). **b**) Half topographies showing differences in alpha power for contralateral minus

ipsilateral electrodes collapsed across hemispheres. White dots indicate the a-priori selected electrodes

used in the cluster-based permutation analysis. **c**) Alpha power values averaged in the time window 0 –

1800 ms after Retro-Cue onset from electrodes Pz and POz. Error bars represents the s.e.m. calculated

across participants (n = 35). **d**) Topographies of each Task x Load condition. White dots indicate the a-

priori selected electrodes (Pz and POz) from which activity was extracted for the ANOVA.

3.2.3 Mid-frontal theta in Implementation

We expected to find stronger theta power over mid-frontal electrodes in the

Implementation task compared to the Memorization task. Supporting this prediction, we

found a significant cluster (P = 0.04, corrected for multiple comparisons) when comparing

the power estimates of the two tasks in the frequency range 3 – 7 Hz from electrodes Fz

and AFz (Figure 4). On the contrary, the test aimed at detecting a main effect of Load

approached but did not reach significance (P = 0.09).

Figure 4: Theta dynamics in Implementation compared to Memorization. a) Time-frequency power spectrum extracted from the electrodes Fz and AFz. **b**) Averaged power time course (3 – 7 Hz) from the electrode Fz and AFz. Thicker outline indicates timepoints belonging to the significant cluster. **c)** Each topography represents the averaged activity in the frequency range 3 – 7 Hz, in 200 ms time windows within the CTI (0 – 1800 ms). Black dots indicate the electrodes used for the statistical analysis.

Moreover, our corollary hypothesis concerning theta dynamics predicted such effect to be modulated by the number of mappings to be implemented or memorized. Specifically, we expected that in Load 4 Implementation and Memorization would not differ, in line with behavioral findings showing that 4 novel instructions cannot be proceduralized at once and would therefore be maintained in a declarative format in both conditions (Liefooghe et al., 2012). However, the effect of the interaction resulted to be not significant (P = 0.53). Despite the lack of a significant interaction, and thus the impossibility to draw any strong conclusion from post-hoc tests, we explored the effect of Task across different Load levels by means of pairwise comparisons. We observed a significant cluster only in the contrast

Implementation 2 vs Memorization 2 (P = 0.02), whereas no cluster was observed for

Load 4, and no cluster survived correction in Load 1 (P = 0.23).

One potential confound when investigating activity in low frequency bands is that this

might be reflecting sustained ERPs. To rule out this possibility we compared across tasks

the Contingent Negative Variation (CNV), a slow negative potential over centro-frontal

electrodes during preparation for an upcoming target (Walter, Cooper, Aldridge,

McCallum, & Winter, 1964). The CNV did not show a differential pattern between

Implementation and Memorization (see Supplementary material), suggesting that our

frontal activity is better described by low frequency oscillations, rather than a sustained

ERP. Furthermore, at the lowest estimated frequency of 2 Hz, the wavelet was two-cycles

long, i.e. spanning a 1s interval, which is considered to be selective to true oscillations

and able to disentangle them from non-oscillatory signal (Cohen, 2014a). Finally,

interpreting this frontal activity as the correlate of target expectancy would not account for

the effect specific to Load 2 observed in time-frequency analyses.

3.2.4 Motor-related mu and beta suppression in Implementation

We hypothesized that proceduralization would induce increased recruitment of motor and

premotor regions, in preparation for the instructed upcoming movement, compared to

Memorization. Specifically, we predicted a suppression of oscillations in the mu (8 – 12

Hz) and beta (15 – 30 Hz) frequency ranges, which have been shown to reflect motor

cortex activity (Cheyne, 2013; Pineda, 2005; Rhodes et al., 2018; Schneider et al., 2017;

Tzagarakis et al., 2015). To test for this hypothesis, we again used cluster-based

permutations and we tested the effects of Task, Load, and their interaction, separately for

each frequency range of interest (i.e. mu and beta).

In line with our prediction, we found significant clusters when contrasting the two Tasks

both in the mu ($P = 0.01$) and beta ($P = 0.03$) frequency ranges. This finding supports the

idea that proceduralization involves a recruitment of motor areas for the activation of the

instructed motor plan (Figure 5).

Figure 5: Mu and Beta dynamics in Implementation compared to Memorization. a) Time-frequency

power spectrum from electrodes C3 and C4. **b)** Averaged power time course in the Mu ($8 - 12$ Hz)

frequency range from electrodes C3 and C4: thicker outline indicates timepoints belonging to the significant

cluster. Topographies in 200 ms time windows: black dots indicate the electrodes used for statistical

analyses. **c)** Averaged power time course in the Beta ($15 - 30$ Hz) frequency range from electrodes C3 and

C4: thicker outline indicates timepoints belonging to the significant cluster. Topographies in 200 ms time

windows: black dots indicate the electrodes used for statistical analyses.

Similarly, the effect of Load resulted to be significant in both mu (P < 0.001) and beta (P

< 0.001) range, suggesting that these motor-related oscillatory features are also sensitive

to the number of items being handled.

As for theta, our secondary hypothesis was that the differences in mu and beta dynamics

between tasks would be modulated by the number of mappings. This hypothesis would

be in line with a capacity limit in the amount of instructions that can be proceduralized at

once, and thus converted in an action plan. However, in neither of the two frequency

bands the cluster-based permutations tests revealed significant clusters for an effect of

such interaction (mu: P = 0.14, beta: P = 0.57). As exploratory follow-up analyses we

performed pairwise comparisons separately for each Load level (i.e., Implementation 1

vs Memorization 1, Implementation 2 vs Memorization 2, Implementation 4 vs

Memorization 4). Although interpretations of these results have to be cautious and do not

allow to draw any strong statistical conclusion, it is interesting to observe that again only

in Load 2 the two tasks showed clusters of different activity (beta: P1 < 0.001, P2 = 0.02,

P3 = 0. 21, mu: P1 = 0.003, P2 = 0.006, P3 = 0.042). On the contrary, comparisons in

Load 1 (beta: P = 0.19, mu: no cluster) and Load 4 (beta: P = 0.35, mu: P = 0.13) yielded

no significant clusters.

4. Discussion

In the present study, we investigated the spectral and temporal dynamics underlying

novel instruction implementation. In line with our predictions and the framework proposed

by Myers and colleagues (2017), we observed an analogous unfolding of early attentional

mechanisms across tasks, reflected in similar patterns of alpha power dynamics.

Crucially, Implementation and Memorization also showed differences in frontal theta, and

central mu and beta oscillations, suggesting that Implementation-specific processes go

beyond the attentional prioritization of the relevant items. These mechanisms are likely

involved in the reformatting of the selected S-R mappings into a behavior-optimized,

action-bound process requiring the exertion of cognitive control and resulting in the

preparation of the instructed motor plan (van Ede, 2020).

An underlying assumption of our study is that the intention to implement the instructed

mappings is a crucial necessary factor to trigger the reformatting from declarative to

procedural state (Liefooghe et al., 2012). This exclusive role of the intention to implement

has been questioned by some recent behavioral evidence, suggesting an automatic

reformatting of the instruction independent of task demands (Liefooghe & De Houwer,

2017). This might be due to the response set used in the specific task, or to other

idiosyncratic factors. Despite these inconsistencies in the behavioral measures, the

dissociation between Implementation and Memorization is well established at the neural

level (Demanet et al., 2016; González-García et al., 2019; Muhle-Karbe et al., 2017).

Here, we show such differences are reflected in oscillatory mechanisms.

4.1 Alpha-mediated attentional orienting is analogous in Implementation and

Memorization

Activity in the alpha frequency band has been consistently associated with attentional

processing (van Ede, 2017). In particular, posterior alpha power decreases over posterior

electrodes contralateral to the attended hemifield, both in perceptual and in WM tasks

(Jensen & Mazaheri, 2010; Poch et al., 2017; Sauseng et al., 2005). In our experiment,

participants were encouraged to focus on the selected items and discard the unselected

ones. Coherently, in both tasks we found after the retro-cue a significant suppression of

alpha oscillations contralateral to the attended hemifield. The most influential model

proposed to account for this phenomenon is referred to as *Gating by Inhibition* (Jensen &

Mazaheri, 2010). According to this framework, top-down alpha modulation of sensory

cortices allows for the inhibition of irrelevant inputs, contributing to the creation of a

functional network optimized to perform the task (Mazaheri et al., 2014; Van Diepen,

Foxe, & Mazaheri, 2019). Our clusters of alpha lateralization extended around 600 –

800ms after retro-cue. This dynamic is consistent with previous findings on lateralized

retrospective orienting of internal attention, showing modulations within 500 and 1000ms

(Mok et al., 2016; Poch et al., 2017, 2018; Wallis et al., 2015; Wolff, Ding, Myers, &

Stokes, 2015; Wolff, Jochim, Akyürek, & Stokes, 2017). The occurrence of the selection

process is further supported by load-dependent centro-posterior alpha power modulation.

The amplitude of alpha power has been observed to be sensitive to the number of items

retained during a WM task (Fukuda et al., 2015; Jensen, 2002). In both our tasks, alpha

increased with load. Crucially, we observed no task-dependent differences in the

dynamics of alpha oscillations, suggesting that the information provided by the retro-cue

is used to analogously orient attention and select the items that are likely to be probed,

independently of task demands.

4.2 Low frequencies oscillations for S-R binding in Implementation

We predicted low frequency oscillations over frontal sensors to be associated with the

proactive reformatting of the S-R mapping(s) into an action-oriented code. The crucial

contribution of the PFC in quickly converting symbolic instructions into task-sets for

prospective action has been widely established in several fMRI studies (Brass, Wenke, Spengler, & Waszak, 2009; Cole, Bagic, Kass, & Schneider, 2010; Dumontheil, Thompson, & Duncan, 2011; González-García et al., 2017; Hartstra et al., 2011; Palenciano, González-García, Arco, & Ruz, 2019; Ruge & Wolfensteller, 2010). In particular, declarative maintenance and implementation of novel mappings can be dissociated using univariate and multivariate analyses in prefrontal regions, such as the inferior frontal junction (IFJ) and the inferior frontal sulcus (IFS) (Bourguignon et al., 2018; Demanet et al., 2016; Hartstra et al., 2011; Muhle-Karbe et al., 2017). Activation of these areas has been interpreted as reflecting the creation and the activation of procedural condition-action rules from the instructed mappings, a process that is thought to involve the manipulation of the available information and the exertion of cognitive control (Brass et al., 2017). Analogously, the need for top-down control over complex goal-directed cognitive operations is traditionally linked with oscillations in the theta frequency range over prefrontal cortices (Cavanagh & Frank, 2014; Cooper et al., 2019). Moreover, recent influential models of large-scale brain interactions attribute to theta the crucial role of orchestrating information exchange between distant areas by synchronizing their firing pattern (Fries, 2005, 2015; Lisman & Jensen, 2013; McLelland & VanRullen, 2016; Verguts, 2017). We hypothesized the Implementation task in our experiment to require a more extensive allocation of cognitive control, given the need to manipulate the declarative representations of the selected S-R mappings into their procedural counterparts, leading to increased power in low frequencies over frontal sensors. Supporting this prediction, our Implementation and Memorization tasks differed in the

extent to which they engaged oscillations in the theta frequency range over prefrontal

regions.

Previous studies show mixed findings with respect to the role of midfrontal theta power in

dissociating maintenance and manipulation of WM items. Berger and colleagues (2019)

reported no differences in theta power between retention and mental spatial mirroring

(i.e., manipulation) of a grid of colored dots. On the contrary, the study by Itthipuripat and

colleagues (2013) used a sequence updating paradigm and revealed increased theta for

both contrasts Manipulation vs Maintenance and Manipulation (correct trials) vs

Manipulation (incorrect trials). Interestingly, they found low theta ($2 - 4$ Hz) to be sensitive

to manipulation demands, whereas high theta ($5 - 7$ Hz) increased in both contrasts

(Itthipuripat et al., 2013). Although the exact boundaries of our significant cluster cannot

be strongly interpreted due to the nature of the statistical approach we adopted (see

statistical Methods section), the effect of Task we find in our study appears to be

predominant in the lower frequencies. One possibility is that different frequencies within

the theta band serve different cognitive functions. This view has recently been gaining

support. Evidence in favor of a more fine-grained characterization of theta oscillations

include variations in theta peak frequency depending on task demands and difficulty

(Senoussi et al., 2020), and differences in the functional significance of high (posterior)

and low (anterior) hippocampal theta (Goyal et al., 2020).

A functional role has been attributed to theta oscillations in the prioritization of relevant

information in visual WM. Oscillations in the delta/low theta ($2 - 6$ Hz) range have been

observed during switches of internal attention from one search template to another for

prospective use (de Vries et al., 2018; de Vries, van Driel, & Olivers, 2019b; Senoussi et

al., 2019). Such involvement has also been causally tested by applying rhythmic (5 Hz)

and arrhythmic transcranial magnetic stimulation to the lateral PFC during a change

detection task with retro-cues (Riddle et al., 2020). Rhythmic stimulation caused an

entrainment of the ongoing theta oscillations and benefitted performance, compared with

the arrhythmic train of pulses (Lakatos, Gross, & Thut, 2019; Riddle et al., 2020; Thut,

Schyns, & Gross, 2011; Thut, Veniero, et al., 2011).

In our study, the difference in theta activity observed between tasks suggests that its

function might go beyond the exertion of short-lasting top-down control over attentional

prioritization, but also encompasses the reformatting of the prioritized memoranda into

behavior-guiding representations. This is possible for the Implementation task, whereas

in the Memorization task the prioritized representations can only be used as a template

to guide the upcoming task (de Vries, Slagter, & Olivers, 2019; Olivers & Eimer, 2011;

Olivers, Peters, Houtkamp, & Roelfsema, 2011).

Therefore, the stronger theta activity in Implementation likely reflects the inherently more

extensive reformatting process of proceduralization, which we assume involves a theta-

driven binding between the stimulus and its corresponding motor plan (Combrisson et al.,

2017; Engel, Roelfsema, Fries, Brecht, & Singer, 1997; Verguts, 2017). The Memorization

task also requires the prioritization and the binding of stimulus and response, but this

would engage theta dynamics to a lesser extent, given the impossibility to activate a motor

plan.

Although strong conclusions cannot be made with respect to timing due to the temporal

smearing inherent to time-frequency analyses, attentional orienting and theta-mediated

binding seem to be occurring simultaneously. The temporal extents of the two clusters

are, in fact, largely overlapping. Therefore, our results hint at a parallel engagement of

the processes that underlie these frequencies.

4.3 Motor preparation in Implementation is reflected in mu and beta suppression

Previous fMRI studies showed that the proactive proceduralization of novel instructions

is accompanied by increased neural activity in pre-motor areas, suggesting motor

preparation (Bourguignon et al., 2018; Hartstra et al., 2011, 2012; Muhle-Karbe et al.,

2017; Ruge & Wolfensteller, 2010). In current views, the recruitment of motor areas is

thought to boost instruction implementation by enabling mental simulation of the overt

application of the instructed mappings (Brass et al., 2017). Motor imagery has been

associated with suppression in mu and beta bands over motor cortices (Cheyne, 2013;

McFarland et al., 2000; Pfurtscheller et al., 2006; Pfurtscheller & Neuper, 1997; Pineda,

2005), which we therefore predicted to occur while the participant is preparing to

implement a specific motor plan, as opposed to the declarative maintenance of the

mapping. Moreover, the engagement of motor areas during implementation is consistent

with the creation of a functional network optimized for the rapid and efficient execution of

the mappings.

In line with our hypothesis, we found these markers of motor preparation to be higher in

Implementation compared to Memorization, already during the CTI. While in both tasks

participants had to perform an overt motor response at the end of the trial, only during

Implementation they could proactively start preparing their response to the upcoming

target. This rules out the possibility that the observed differences reflect more general

mechanisms of unspecific motor energization, temporal estimation or target expectancy

(Nobre & Van Ede, 2018; Van Elswijk, Kleine, Overeem, & Stegeman, 2007; Wiener,

Parikh, Krakow, & Coslett, 2018). Overall, our results suggest that preparing to execute

novel mappings involves the activation of a specific motor plan corresponding to the

instructed response, to a much larger extent then the motor engagement observed in

Memorization.

Concerning the timing of the implementation-induced motor engagement, the temporal

extent of the significant cluster suggests a later onset with respect to alpha and theta

dynamics. Although strong conclusions cannot be drawn, based on this interpretation of

our permutation results and theoretical models such as Myers' (2017), we are inclined to

speculate that motor preparation is initiated following attentional selection of the relevant

mappings. From this perspective, theta would act as an overarching mechanism to bind

and control the task-optimized representations. Future studies should investigate the

exact timing properties and the interplay between these processes.

4.4 Capacity limits of proceduralization

As a corollary hypothesis, we predicted that the capacity limits of proceduralization

observed in behavioral studies (Liefooghe et al., 2012) would be reflected in stronger

proceduralization-related oscillatory features with fewer mappings. However, we did not

observe a significant interaction of Task and Load in none of our frequency bands of

interest. It is possible that our design was not powered enough to detect such effects.

Despite the not significant interaction, for exploratory purposes we looked at the effect of

Task separately for each Load. Interestingly, we observed task-related differences only

in Load 2. The interpretation of these results requires caution and can only be speculative

given the exploratory nature of these analyses. Yet, they could reflect that relevant

oscillations become crucial when in need to coordinate multiple bindings, but still within

capacity limitations. From this perspective, Load 4 would show no differences because

too many S-R mappings cannot be simultaneously recoded into action-oriented

representations (Liefooghe et al., 2012), and Load 1 would instead allow a more ballistic

implementation of the prepared motor plan, without the need to carefully coordinate

different competing procedural representations, as is the case in Load 2. These

exploratory results pave the way for future research investigating more accurately the

capacity limits of proceduralization: how they are overcome with extensive practice and

how they reflected in power modulations.

5. Conclusions

Our study shed light for the first time on the oscillatory dynamics associated with the

retrospective prioritization and proceduralization of novel instructions. We showed that

alpha-mediated mechanisms of attentional orienting are in place to prioritize the relevant

items, independently from the upcoming task demands. Conversely, other neural features

were sensitive to the task. The implementation of novel mappings, as opposed to their

declarative maintenance, is characterized by increased power in low frequencies and by

stronger suppression of mu and beta activity. The former is consistent with the purported

role of theta oscillations in cognitive control for the successful binding of stimulus and

response into a behavior-optimized procedure, and the latter to reflect motor preparation

of the instructed motor responses. Overall, our results support the idea that under optimal

task conditions, proceduralization can occur proactively in preparation for the upcoming

response execution demands.

Future research should investigate how proceduralization affects the underlying neural

representation of the instructed mapping, under the assumption that it is recoded from a

declarative to a procedural format, and how prefrontal regions mediate this reformatting

process. Moreover, future studies should explore whether capacity limits of

proceduralization (Liefooghe et al., 2012) are caused by concurrent activation of

interfering motor plans or intrinsic to the neural mechanisms supporting the recoding of

instructions into procedures, such as the theta-mediated binding.

Acknowledgements

S.F. and C.G.G. were supported by Special Research Fund of Ghent University

BOF.GOA.2017.0002.03. C.G.G. was additionally supported by the European Union's

Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie

grant agreement no. 835767. M.S. was supported by Research Foundation Flanders

(FWO) grant number G012816N. MB is funded by an Einstein Strategic Professorship of

the Einstein Foundation Berlin.

Data and code availability

772 Data and code will be made available upon publication.

REFERENCES

Berger, B., Griesmayr, B., Minarik, T., Biel, A. L., Pinal, D., Sterr, A., & Sauseng, P.
(2019). Dynamic regulation of interregional cortical communication by slow brain
oscillations during working memory. *Nature Communications*, *10*(1), 1–11.
https://doi.org/10.1038/s41467-019-12057-0

Bhandari, A., & Duncan, J. (2014). Goal neglect and knowledge chunking in the
construction of novel behaviour. *Cognition*, *130*(1), 11–30.
https://doi.org/10.1016/j.cognition.2013.08.013

Bonnefond, M., & Jensen, O. (2012). Alpha oscillations serve to protect working
memory maintenance against anticipated distracters. *Current Biology*, *22*(20),
1969–1974. https://doi.org/10.1016/j.cub.2012.08.029

Bourguignon, N. J., Braem, S., Hartstra, E., De Houwer, J., & Brass, M. (2018).
Encoding of Novel Verbal Instructions for Prospective Action in the Lateral
Prefrontal Cortex: Evidence from Univariate and Multivariate Functional Magnetic
Resonance Imaging Analysis. *Journal of Cognitive Neuroscience*, *26*(3), 1–15.
https://doi.org/10.1162/jocn_a_01270

Brady, T. F., Konkle, T., Alvarez, G. A., & Oliva, A. (2013). Real-world objects are not
represented as bound units: Independent forgetting of different object details from
visual memory. *Journal of Experimental Psychology: General*, *142*(3), 791–808.
https://doi.org/10.1037/a0029649

Brass, M., Liefooghe, B., Braem, S., & De Houwer, J. (2017). Following new task
instructions: Evidence for a dissociation between knowing and doing. *Neuroscience*

and Biobehavioral Reviews, *81*(June), 16–28.

https://doi.org/10.1016/j.neubiorev.2017.02.012

Brass, M., Wenke, D., Spengler, S., & Waszak, F. (2009). Neural Correlates of

Overcoming Interference from Instructed and Implemented Stimulus-Response

Associations. *Journal of Neuroscience*, *29*(6), 1766–1772.

https://doi.org/10.1523/JNEUROSCI.5259-08.2009

Brodeur, M. B., Guérard, K., & Bouras, M. (2014). Bank of Standardized Stimuli (BOSS)

Phase II: 930 New Normative Photos. *PLoS ONE*, *9*(9), e106953.

https://doi.org/10.1371/journal.pone.0106953

Bundt, C., Bardi, L., Abrahamse, E. L., Brass, M., & Notebaert, W. (2015). It wasn't me!

Motor activation from irrelevant spatial information in the absence of a response.

Frontiers in Human Neuroscience, *9*(OCT).

https://doi.org/10.3389/fnhum.2015.00539

Capilla, A., Schoffelen, J.-M., Paterson, G., Thut, G., & Gross, J. (2014). Dissociated α-

Band Modulations in the Dorsal and Ventral Visual Pathways in Visuospatial

Attention and Perception. *Cerebral Cortex*, *24*(2), 550–561.

https://doi.org/10.1093/cercor/bhs343

Cavanagh, J. F., & Frank, M. J. (2014). Frontal theta as a mechanism for cognitive

control. *Trends in Cognitive Sciences*, *18*(8), 414–421.

https://doi.org/10.1016/j.tics.2014.04.012

Cheyne, D. O. (2013). MEG studies of sensorimotor rhythms: A review. *Experimental*

Neurology, *245*, 27–39. https://doi.org/10.1016/j.expneurol.2012.08.030

Cohen, M. X. (2014a). *Analyzing neural time series data: theory and practice*. MIT

Press.

Cohen, M. X. (2014b). Analyzing Neural Time Series Data.

Cohen, M. X., & Donner, T. H. (2013). Midfrontal conflict-related theta-band power

reflects neural oscillations that predict behavior. *Journal of Neurophysiology*,

110(12), 2752–2763. https://doi.org/10.1152/jn.00479.2013

Cole, M. W., Bagic, A., Kass, R., & Schneider, W. (2010). Prefrontal dynamics

underlying rapid instructed task learning reverse with practice. *Journal of*

Neuroscience, *30*(42), 14245–14254. https://doi.org/10.1523/JNEUROSCI.1662-

10.2010

Cole, M. W., Laurent, P., & Stocco, A. (2013). Rapid instructed task learning: A new

window into the human brain's unique capacity for flexible cognitive control.

Cognitive, Affective, & Behavioral Neuroscience, *13*(1), 1–22.

https://doi.org/10.3758/s13415-012-0125-7

Combrisson, E., Perrone-Bertolotti, M., Soto, J. L., Alamian, G., Kahane, P., Lachaux, J.

P., … Jerbi, K. (2017). From intentions to actions: Neural oscillations encode motor

processes through phase, amplitude and phase-amplitude coupling. *NeuroImage*,

147, 473–487. https://doi.org/10.1016/j.neuroimage.2016.11.042

Cooper, P. S., Karayanidis, F., McKewen, M., McLellan-Hall, S., Wong, A. S. W.,

Skippen, P., & Cavanagh, J. F. (2019). Frontal theta predicts specific cognitive

control-induced behavioural changes beyond general reaction time slowing.

NeuroImage, *189*(January), 130–140.

https://doi.org/10.1016/j.neuroimage.2019.01.022

de Vries, I. E. J., Slagter, H. A., & Olivers, C. N. L. (2019). Oscillatory Control over

Representational States in Working Memory. *Trends in Cognitive Sciences*.

https://doi.org/10.1016/j.tics.2019.11.006

de Vries, I. E. J., Van Driel, J., Karacaoglu, M., & Olivers, C. N. L. (2018). Priority

Switches in Visual Working Memory are Supported by Frontal Delta and Posterior

Alpha Interactions. *Cerebral Cortex*, *28*(11), 4090–4104.

https://doi.org/10.1093/cercor/bhy223

de Vries, I. E. J., van Driel, J., & Olivers, C. N. L. (2019a). Decoding the status of

working memory representations in preparation of visual selection. *NeuroImage*,

191, 549–559. https://doi.org/10.1016/j.neuroimage.2019.02.069

de Vries, I. E. J., van Driel, J., & Olivers, C. N. L. (2019b). Decoding the status of

working memory representations in preparation of visual selection. *NeuroImage*,

191, 549–559. https://doi.org/10.1016/J.NEUROIMAGE.2019.02.069

Delorme, A., & Makeig, S. (2004). EEGLAB : an open source toolbox for analysis of

single-trial EEG dynamics including independent component analysis. *Journal of*

Neuroscience Methods, *134*, 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Demanet, J., Liefooghe, B., Hartstra, E., Wenke, D., De Houwer, J., & Brass, M. (2016).

There is more into 'doing' than 'knowing': The function of the right inferior frontal

sulcus is specific for implementing versus memorising verbal instructions.

NeuroImage, *141*, 350–356. https://doi.org/10.1016/j.neuroimage.2016.07.059

Dumontheil, I., Thompson, R., & Duncan, J. (2011). Assembly and use of new task rules

in fronto-parietal cortex. *Journal of Cognitive Neuroscience*, *23*(1), 168–182.

https://doi.org/10.1162/jocn.2010.21439

Duncan, J., Emslie, H., Williams, P., Johnson, R., & Freer, C. (1996). Intelligence and

the Frontal Lobe: The Organization of Goal-Directed Behavior. *Cognitive*

Psychology, *30*(3), 257–303. https://doi.org/10.1006/COGP.1996.0008

Engel, A. K., Roelfsema, P. R., Fries, P., Brecht, M., & Singer, W. (1997). Role of the

temporal domain for response selection and perceptual binding. *Cerebral Cortex*,

7(6), 571–582. https://doi.org/10.1093/cercor/7.6.571

Everaert, T., Theeuwes, M., Liefooghe, B., & De Houwer, J. (2014). Automatic motor

activation by mere instruction. *Cognitive, Affective and Behavioral Neuroscience*,

14(4), 1300–1309. https://doi.org/10.3758/s13415-014-0294-7

Formica, S., González-García, C., & Brass, M. (2020). The effects of declaratively

maintaining and proactively proceduralizing novel stimulus-response mappings.

Cognition, *201*, 104295. https://doi.org/10.1016/j.cognition.2020.104295

Fries, P. (2005). A mechanism for cognitive dynamics: neuronal communication through

neuronal coherence. *Trends in Cognitive Sciences*, *9*(10), 474–480.

https://doi.org/10.1016/J.TICS.2005.08.011

Fries, P. (2015, October 7). Rhythms for Cognition: Communication through Coherence.

Neuron. Cell Press. https://doi.org/10.1016/j.neuron.2015.09.034

Fukuda, K., Mance, I., & Vogel, E. K. (2015). α power modulation and event-related

slow wave provide dissociable correlates of visual working memory. *Journal of*

Neuroscience, *35*(41), 14009–14016. https://doi.org/10.1523/JNEUROSCI.5003-

14.2015

González-García, C., Arco, J. E., Palenciano, A. F., Ramírez, J., & Ruz, M. (2017).

Encoding, preparation and implementation of novel complex verbal instructions.

NeuroImage, *148*(November 2016), 264–273.

https://doi.org/10.1016/j.neuroimage.2017.01.037

González-García, C., Formica, S., Liefooghe, B., & Brass, M. (2020). Attentional

prioritization reconfigures novel instructions into action-oriented task sets.

Cognition, *194*, 104059. https://doi.org/10.1016/j.cognition.2019.104059

González-García, C., Formica, S., Wisniewski, D., & Brass, M. (2019). Frontoparietal

action-oriented codes support novel task set implementation. *BioRxiv*, 830067.

https://doi.org/10.1101/830067

Gould, I. C., Rushworth, M. F., & Nobre, A. C. (2011). Indexing the graded allocation of

visuospatial attention using anticipatory alpha oscillations. *Journal of*

Neurophysiology, *105*(3), 1318–1326. https://doi.org/10.1152/jn.00653.2010

Goyal, A., Miller, J., Qasim, S., Watrous, A. J., Stein, J. M., Inman, C. S., … Jacobs, J.

(2020). Functionally distinct high and low theta oscillations in the human

hippocampus. *BioRxiv*, 498055. https://doi.org/10.1101/498055

Griffin, G., Holub, A., & Perona, P. (2007). Caltech-256 Object Category Dataset.

Retrieved from https://authors.library.caltech.edu/7694/

Groppe, D. M., Makeig, S., & Kutas, M. (2009). Identifying reliable independent

components via split-half comparisons. *NeuroImage*, *45*(4), 1199–1211.

https://doi.org/10.1016/j.neuroimage.2008.12.038

Haller, M., Donoghue, T., Peterson, E., Varma, P., Sebastian, P., Gao, R., … Voytek, B.

(2018). Parameterizing neural power spectra. *BioRxiv*, 299859.

https://doi.org/10.1101/299859

Hari, R., & Salmelin, R. (1997, January 1). Human cortical oscillations: A neuromagnetic

view through the skull. *Trends in Neurosciences*. https://doi.org/10.1016/S0166-

2236(96)10065-5

Hartstra, E., Kühn, S., Verguts, T., & Brass, M. (2011). The implementation of verbal

instructions: An fMRI study. *Human Brain Mapping*, *32*(11), 1811–1824.

https://doi.org/10.1002/hbm.21152

Hartstra, E., Waszak, F., & Brass, M. (2012). The implementation of verbal instructions:

Dissociating motor preparation from the formation of stimulus-response

associations. *NeuroImage*, *63*(3), 1143–1153.

https://doi.org/10.1016/j.neuroimage.2012.08.003

Itthipuripat, S., Wessel, J. R., & Aron, A. R. (2013). Frontal theta is a signature of

successful working memory manipulation. *Experimental Brain Research*, *224*(2),

255–262. https://doi.org/10.1007/s00221-012-3305-3

Jasp Team. (2019). JASP (Version 0.11.1)[Computer software]. Retrieved from

https://jasp-stats.org/

Jensen, O. (2002). Oscillations in the Alpha Band (9-12 Hz) Increase with Memory Load

during Retention in a Short-term Memory Task. *Cerebral Cortex*, *12*(8), 877–882.

https://doi.org/10.1093/cercor/12.8.877

Jensen, O., & Mazaheri, A. (2010). Shaping Functional Architecture by Oscillatory Alpha

Activity: Gating by Inhibition. *Frontiers in Human Neuroscience*, *4*(November), 1–8.

https://doi.org/10.3389/fnhum.2010.00186

Klem, G. H., Lüders, H. O., Jasper, H. H., & Elger, C. (1999). The ten-twenty electrode

system of the International Federation. The International Federation of Clinical

Neurophysiology. *Electroencephalography and Clinical Neurophysiology.*

Supplement, *52*, 3–6.

Klimesch, W., Sauseng, P., & Hanslmayr, S. (2007). EEG alpha oscillations: The

inhibition–timing hypothesis. *Brain Research Reviews*, *53*(1), 63–88.

https://doi.org/10.1016/J.BRAINRESREV.2006.06.003

Konkle, T., Brady, T. F., Alvarez, G. A., & Oliva, A. (2010). Conceptual distinctiveness

supports detailed visual long-term memory for real-world objects. *Journal of*

Experimental Psychology: General, *139*(3), 558–578.

https://doi.org/10.1037/a0019165

Lakatos, P., Gross, J., & Thut, G. (2019, September 23). A New Unifying Account of the

Roles of Neuronal Entrainment. *Current Biology*. Cell Press.

https://doi.org/10.1016/j.cub.2019.07.075

Liefooghe, B., & De Houwer, J. (2017). Automatic effects of instructions do not require

the intention to execute these instructions. *Journal of Cognitive Psychology*, *30*(1),

108–121. https://doi.org/10.1080/20445911.2017.1365871

Liefooghe, B., Wenke, D., & De Houwer, J. (2012). Instruction-based task-rule

congruency effects. *Journal of Experimental Psychology: Learning, Memory, and*

Cognition, *38*(5), 1325–1335. https://doi.org/10.1037/a0028148

Lisman, J. E., & Jensen, O. (2013). The Theta-Gamma Neural Code. *Neuron*, *77*(6),

1002–1016. https://doi.org/10.1016/j.neuron.2013.03.007

Manza, P., Hau, C. L. V., & Leung, H.-C. (2014). Alpha Power Gates Relevant

Information during Working Memory Updating. *Journal of Neuroscience*, *34*(17),

5998–6002. https://doi.org/10.1523/JNEUROSCI.4641-13.2014

Marchesotti, S., Bassolino, M., Serino, A., Bleuler, H., & Blanke, O. (2016). Quantifying

the role of motor imagery in brain-machine interfaces. *Scientific Reports*, *6*(1),

24076. https://doi.org/10.1038/srep24076

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-

data. *Journal of Neuroscience Methods*, *164*, 177–190.

https://doi.org/10.1016/j.jneumeth.2007.03.024

Mazaheri, A., van Schouwenburg, M. R., Dimitrijevic, A., Denys, D., Cools, R., &

Jensen, O. (2014). Region-specific modulations in oscillatory alpha activity serve to

facilitate processing in the visual and auditory modalities. *NeuroImage*, *87*, 356–

362. https://doi.org/10.1016/J.NEUROIMAGE.2013.10.052

McFarland, D. J., Miner, L. A., Vaughan, T. M., & Wolpaw, J. R. (2000). Mu and Beta

Rhythm Topographies During Motor Imagery and Actual Movements. *Brain*

Topography, *12*(3), 177–186. https://doi.org/10.1023/A:1023437823106

McLelland, D., & VanRullen, R. (2016). Theta-Gamma Coding Meets Communication-
through-Coherence: Neuronal Oscillatory Multiplexing Theories Reconciled. *PLoS*
Computational Biology, *12*(10), e1005162.

Meiran, N., Pereg, M., Kessler, Y., Cole, M. W., & Braver, T. S. (2015a). Reflexive
activation of newly instructed stimulus–response rules: evidence from lateralized
readiness potentials in no-go trials. *Cognitive, Affective, & Behavioral*
Neuroscience, *15*(2), 365–373. https://doi.org/10.3758/s13415-014-0321-8

Meiran, N., Pereg, M., Kessler, Y., Cole, M. W., & Braver, T. S. (2015b). The power of
instructions: Proactive configuration of stimulus–response translation. *Journal of*
Experimental Psychology: Learning, Memory, and Cognition, *41*(3), 768–786.
https://doi.org/10.1037/xlm0000063

Milner, B. (1963). Effects of Different Brain Lesions on Card Sorting: The Role of the
Frontal Lobes. *Archives of Neurology*, *9*(1), 90–100.
https://doi.org/10.1001/archneur.1963.00460070100010

Mok, R. M., Myers, N. E., Wallis, G., & Nobre, A. C. (2016). Behavioral and Neural
Markers of Flexible Attention over Working Memory in Aging. *Cerebral Cortex*,
26(4), 1831–1842. https://doi.org/10.1093/cercor/bhw011

Muhle-Karbe, P. S., Duncan, J., De Baene, W., Mitchell, D. J., & Brass, M. (2017).
Neural Coding for Instruction-Based Task Sets in Human Frontoparietal and Visual
Cortex. *Cerebral Cortex (New York, N.Y. : 1991)*, *27*(3), 1891–1905.
https://doi.org/10.1093/cercor/bhw032

Myers, N. E., Stokes, M. G., & Nobre, A. C. (2017). Prioritizing Information during

Working Memory: Beyond Sustained Internal Attention. *Trends in Cognitive*

Sciences, *21*(6), 449–461. https://doi.org/10.1016/j.tics.2017.03.010

Myers, N. E., Walther, L., Wallis, G., Stokes, M. G., & Nobre, A. C. (2015). Temporal

Dynamics of Attention during Encoding versus Maintenance of Working Memory:

Complementary Views from Event-related Potentials and Alpha-band Oscillations.

Journal of Cognitive Neuroscience, *27*(3), 492–508.

https://doi.org/10.1162/jocn_a_00727

Neuper, C., & Pfurtscheller, G. (2001). Event-related dynamics of cortical rhythms:

Frequency-specific features and functional correlates. *International Journal of*

Psychophysiology, *43*(1), 41–58. https://doi.org/10.1016/S0167-8760(01)00178-7

Nobre, A. C., & Van Ede, F. (2018). Anticipated moments: Temporal structure in

attention. *Nature Reviews Neuroscience*, *19*(1), 34–48.

https://doi.org/10.1038/nrn.2017.141

Olivers, C. N. L., & Eimer, M. (2011). On the difference between working memory and

attentional set. *Neuropsychologia*, *49*(6), 1553–1558.

https://doi.org/10.1016/j.neuropsychologia.2010.11.033

Olivers, C. N. L., Peters, J., Houtkamp, R., & Roelfsema, P. R. (2011, July). Different

states in visual working memory: When it guides attention and when it does not.

Trends in Cognitive Sciences. https://doi.org/10.1016/j.tics.2011.05.004

Onton, J., Delorme, A., & Makeig, S. (2005). Frontal midline EEG dynamics during

working memory. *NeuroImage*, *27*(2), 341–356.

https://doi.org/10.1016/J.NEUROIMAGE.2005.04.014

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2010). FieldTrip: Open Source

Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological

Data. *Computational Intelligence and Neuroscience*, *2011*.

https://doi.org/10.1155/2011/156869

Palenciano, A. F., González-García, C., Arco, J. E., Pessoa, L., & Ruz, M. (2019).

Representational organization of novel task sets during proactive encoding. *The*

Journal of Neuroscience, 0725–19. https://doi.org/10.1523/jneurosci.0725-19.2019

Palenciano, A. F., González-García, C., Arco, J. E., & Ruz, M. (2019). Transient and

Sustained Control Mechanisms Supporting Novel Instructed Behavior. *Cerebral*

Cortex, *29*(9), 3948–3960. https://doi.org/10.1093/cercor/bhy273

Perrin, F., Pernier, J., Bertrand, O., & Echallier, J. F. (1989). Spherical splines for scalp

potential and current density mapping. *Electroencephalography and Clinical*

Neurophysiology, *72*(2), 184–187. https://doi.org/10.1016/0013-4694(89)90180-6

Pfurtscheller, G., Brunner, C., Schlögl, A., & Lopes da Silva, F. H. (2006). Mu rhythm

(de)synchronization and EEG single-trial classification of different motor imagery

tasks. *NeuroImage*, *31*(1), 153–159.

https://doi.org/10.1016/J.NEUROIMAGE.2005.12.003

Pfurtscheller, G., & Neuper, C. (1997). Motor imagery activates primary sensorimotor

area in humans. *Neuroscience Letters*, *239*(2–3), 65–68.

https://doi.org/10.1016/S0304-3940(97)00889-6

Pineda, J. A. (2005). The functional significance of mu rhythms: Translating 'seeing' and

'hearing' into 'doing'. *Brain Research Reviews*, *50*(1), 57–68.

https://doi.org/10.1016/j.brainresrev.2005.04.005

Poch, C., Campo, P., & Barnes, G. R. (2014). Modulation of alpha and gamma

oscillations related to retrospectively orienting attention within working memory.

European Journal of Neuroscience, *40*(2), 2399–2405.

https://doi.org/10.1111/ejn.12589

Poch, C., Carretie, L., & Campo, P. (2017). A dual mechanism underlying alpha

lateralization in attentional orienting to mental representation. *Biological*

Psychology, *128*(March), 63–70. https://doi.org/10.1016/j.biopsycho.2017.07.015

Poch, C., Valdivia, M., Capilla, A., Hinojosa, J. A., & Campo, P. (2018). Suppression of

no-longer relevant information in Working Memory: An alpha-power related

mechanism? *Biological Psychology*, *135*(February), 112–116.

https://doi.org/10.1016/j.biopsycho.2018.03.009

Popov, T., Popova, P., Harkotte, M., Awiszus, B., Rockstroh, B., & Miller, G. A. (2018).

Cross-frequency interactions between frontal theta and posterior alpha control

mechanisms foster working memory. *NeuroImage*, *181*(July), 728–733.

https://doi.org/10.1016/j.neuroimage.2018.07.067

Rhodes, E., Gaetz, W. C., Marsden, J., & Hall, S. D. (2018). Transient Alpha and Beta

Synchrony Underlies Preparatory Recruitment of Directional Motor Networks.

Journal of Cognitive Neuroscience, *30*(6), 867–875.

https://doi.org/10.1162/jocn_a_01250

Riddle, J., Scimeca, J. M., Cellier, D., Dhanani, S., Esposito, M. D., Riddle, J., …

    Esposito, M. D. (2020). Causal Evidence for a Role of Theta and Alpha Oscillations

    in the Control of Working Memory Report Causal Evidence for a Role of Theta and

    Alpha Oscillations in the Control of Working Memory. *Current Biology*, 1–7.

    https://doi.org/10.1016/j.cub.2020.02.065

Rihs, T. A., Michel, C. M., & Thut, G. (2007). Mechanisms of selective inhibition in visual

    spatial attention are indexed by α-band EEG synchronization. *European Journal of*

    Neuroscience, *25*(2), 603–610. https://doi.org/10.1111/j.1460-9568.2007.05278.x

Rihs, T. A., Michel, C. M., & Thut, G. (2009). A bias for posterior α-band power

    suppression versus enhancement during shifting versus maintenance of spatial

    attention. *NeuroImage*, *44*(1), 190–199.

    https://doi.org/10.1016/j.neuroimage.2008.08.022

Rohenkohl, G., & Nobre, A. C. (2011). Alpha oscillations related to anticipatory attention

    follow temporal expectations. *Journal of Neuroscience*, *31*(40), 14076–14084.

    https://doi.org/10.1523/JNEUROSCI.3387-11.2011

Ruge, H., Schäfer, T. A., Zwosta, K., Mohr, H., & Wolfensteller, U. (2019). Neural

    representation of newly instructed rule identities during early implementation trials.

    ELife, *8*, 1–25. https://doi.org/10.7554/eLife.48293

Ruge, H., & Wolfensteller, U. (2010). Rapid formation of pragmatic rule representations

    in the human brain during instruction-based learning. *Cerebral Cortex*, *20*(7), 1656–

    1667. https://doi.org/10.1093/cercor/bhp228

Sassenhagen, J., & Draschkow, D. (2019). Cluster-based permutation tests of

MEG/EEG data do not establish significance of effect latency or location. *Psychophysiology*, (August 2018), 1–8. https://doi.org/10.1111/psyp.13335

Sauseng, P., Klimesch, W., Stadler, W., Schabus, M., Doppelmayr, M., Hanslmayr, S., … Birbaumer, N. (2005). A shift of visual spatial attention is selectively associated with human EEG alpha activity. *European Journal of Neuroscience*, *22*(11), 2917–2926. https://doi.org/10.1111/j.1460-9568.2005.04482.x

Schneider, D., Barth, A., & Wascher, E. (2017). On the contribution of motor planning to the retroactive cuing benefit in working memory: Evidence by mu and beta oscillatory activity in the EEG. *NeuroImage*, *162*(April), 73–85. https://doi.org/10.1016/j.neuroimage.2017.08.057

Schneider, D., Mertes, C., & Wascher, E. (2016). The time course of visuo-spatial working memory updating revealed by a retro-cuing paradigm. *Scientific Reports*, *6*(1), 21442. https://doi.org/10.1038/srep21442

Senoussi, M., Moreland, J. C., Busch, N. A., & Dugué, L. (2019). Attention explores space periodically at the theta frequency. *Journal of Vision*, *19*(5), 1–17. https://doi.org/10.1167/19.5.22

Senoussi, M., Verbeke, P., Desender, K., De Loof, E., Talsma, D., Verguts, T., & Dunantlaan, H. (2020). Theta oscillations shift towards optimal frequency for cognitive control. *BioRxiv*, 2020.08.30.273706. https://doi.org/10.1101/2020.08.30.273706

Souza, A. S., & Oberauer, K. (2016). In search of the focus of attention in working memory: 13 years of the retro-cue effect. *Attention, Perception, & Psychophysics*,

78(7), 1839–1860. https://doi.org/10.3758/s13414-016-1108-5

Thut, G., Nietzel, A., Brandt, S. A., & Pascual-Leone, A. (2006). Alpha-band

electroencephalographic activity over occipital cortex indexes visuospatial attention

bias and predicts visual target detection. *The Journal of Neuroscience : The Official*

Journal of the Society for Neuroscience, *26*(37), 9494–9502.

https://doi.org/10.1523/JNEUROSCI.0875-06.2006

Thut, G., Schyns, P. G., & Gross, J. (2011). Entrainment of Perceptually Relevant Brain

Oscillations by Non-Invasive Rhythmic Stimulation of the Human Brain. *Frontiers in*

Psychology, *2*(JUL), 170. https://doi.org/10.3389/fpsyg.2011.00170

Thut, G., Veniero, D., Romei, V., Miniussi, C., Schyns, P., & Gross, J. (2011). Rhythmic

TMS causes local entrainment of natural oscillatory signatures. *Current Biology*,

21(14), 1176–1185. https://doi.org/10.1016/j.cub.2011.05.049

Tzagarakis, C., West, S., & Pellizzer, G. (2015). Brain oscillatory activity during motor

preparation: Effect of directional uncertainty on beta, but not alpha, frequency band.

Frontiers in Neuroscience, *9*(JUN), 1–13. https://doi.org/10.3389/fnins.2015.00246

Van Diepen, R., Foxe, J. J., & Mazaheri, A. (2019). The functional role of alpha-band

activity in attentional processing: The current zeitgeist and future outlook. *Current*

Opinion in Psychology. https://doi.org/10.1016/J.COPSYC.2019.03.015

van Dijk, H., Schoffelen, J.-M., Oostenveld, R., & Jensen, O. (2008). Prestimulus

oscillatory activity in the alpha band predicts visual discrimination ability. *The*

Journal of Neuroscience : The Official Journal of the Society for Neuroscience,

28(8), 1816–1823. https://doi.org/10.1523/JNEUROSCI.1853-07.2008

van Ede, F. (2017). Mnemonic and attentional roles for states of attenuated alpha

oscillations in perceptual working memory: A review. *European Journal of*

Neuroscience, 1–7. https://doi.org/10.1111/ejn.13759

van Ede, F. (2020). Visual working memory and action: Functional links and bi-

directional influences. *Visual Cognition*, 1–13.

https://doi.org/10.1080/13506285.2020.1759744

van Ede, F., Chekroud, S. R., Stokes, M. G., & Nobre, A. C. (2019a). Concurrent visual

and motor selection during visual working memory guided action. *Nature*

Neuroscience, *22*(3), 477–483. https://doi.org/10.1038/s41593-018-0335-6

van Ede, F., Chekroud, S. R., Stokes, M. G., & Nobre, A. C. (2019b). Concurrent visual

and motor selection during visual working memory guided action. *Nature*

Neuroscience, *22*(3), 477–483. https://doi.org/10.1038/s41593-018-0335-6

Van Elswijk, G., Kleine, B. U., Overeem, S., & Stegeman, D. F. (2007). Expectancy

induces dynamic modulation of corticospinal excitability. *Journal of Cognitive*

Neuroscience, *19*(1), 121–131. https://doi.org/10.1162/jocn.2007.19.1.121

Verbeke, P., & Verguts, T. (2019). Learning to synchronize: How biological agents can

couple neural task modules for dealing with the stability-plasticity dilemma. *PLOS*

Computational Biology, *15*(8), e1006604.

https://doi.org/10.1371/journal.pcbi.1006604

Verguts, T. (2017). Binding by random bursts: A computational model of cognitive

control. *Journal of Cognitive Neuroscience*. https://doi.org/10.1162/jocn

Wallis, G., Stokes, M., Cousijn, H., Woolrich, M., & Nobre, A. C. (2015). Frontoparietal

and Cingulo-opercular Networks Play Dissociable Roles in Control of Working

Memory. *Journal of Cognitive Neuroscience*, *27*(10), 2019–2034.

https://doi.org/10.1162/jocn_a_00838

Walter, W. G., Cooper, R., Aldridge, V. J., McCallum, W. C., & Winter, A. L. (1964).

Contingent negative variation : An electric sign of sensori-motor association and

expectancy in the human brain. *Nature*, *203*(4943), 380–384.

https://doi.org/10.1038/203380a0

Wenke, D., Gaschler, R., Nattkemper, D., & Frensch, P. A. (2009). Strategic influences

on implementing instructions for future actions. *Psychological Research*

Psychologische Forschung, *73*(4), 587–601. https://doi.org/10.1007/s00426-009-

0239-x

Wiener, M., Parikh, A., Krakow, A., & Coslett, H. B. (2018). An intrinsic role of beta

oscillations in memory for time estimation. *Scientific Reports*, *8*(1), 1–17.

https://doi.org/10.1038/s41598-018-26385-6

Wolff, M. J., Ding, J., Myers, N. E., & Stokes, M. G. (2015). Revealing hidden states in

visual working memory using electroencephalography. *Frontiers in Systems*

Neuroscience, *9*(September), 1–12. https://doi.org/10.3389/fnsys.2015.00123

Wolff, M. J., Jochim, J., Akyürek, E. G., & Stokes, M. G. (2017). Dynamic hidden states

underlying working-memory-guided behavior. *Nature Neuroscience*, *20*(6), 864–

871. https://doi.org/10.1038/nn.4546

1162