A sister lineage of the *Mycobacterium tuberculosis* complex discovered in the African
Great Lakes region

Authors:

Jean Claude Semuto Ngabonziza[1,2,3#], Chloé Loiseau[4,5#], Michael Marceau[6#], Agathe Jouet[7],
Fabrizio Menardo[4,5], Oren Tzfadia[2], Rudy Antoine[6], Esdras Belamo Niyigena[1], Wim Mulders[2],
Kristina Fissette[2], Maren Diels[8], Cyril Gaudin[7], Stéphanie Duthoy[7], Willy Ssengooba[9],
Emmanuel André[10], Michel K Kaswa[11], Yves Mucyo Habimana[12], Daniela Brites[4,5], Dissou
Affolabi[13], Jean Baptiste Mazarati[14], Bouke Catherine de Jong[2], Leen Rigouts[2,3], Sebastien
Gagneux[4,5]*, Conor Joseph Meehan[2,15]*, Philip Supply[6]*

Affiliations:

[1]National Reference Laboratory Division, Department of Biomedical Services, Rwanda
Biomedical Center, Kigali, Rwanda. [2]Mycobacteriology Unit, Department of Biomedical
Sciences, Institute of Tropical Medicine, Antwerp, Belgium. [3]Department of Biomedical
Sciences, Antwerp University, Antwerp, Belgium. [4]Swiss Tropical and Public Health Institute,
Basel, Switzerland. [5]University of Basel, Basel, Switzerland. [6]Univ. Lille, CNRS, Inserm, CHU
Lille, Institut Pasteur de Lille, U1019 - UMR 9017 - CIIL - Center for Infection and Immunity of
Lille, F-59000 Lille, France. [7]Genoscreen, Lille, France. [8]BCCM/ITM, Mycobacterial culture
collection, Institute of Tropical Medicine, Antwerp, Belgium. [9]Department of Medical
Microbiology, College of Health Sciences, Makerere University, Kampala, Uganda.
[10]Laboratory of Clinical Bacteriology and Mycology, Department of Microbiology and Immunology,
KU Leuven, Leuven, Belgium. [11]National Tuberculosis Program, Kinshasa, DR Congo.
[12]Tuberculosis and Other Respiratory Diseases Division, Institute of HIV/AIDS Disease
Prevention and Control, Rwanda Biomedical Center, Kigali, Rwanda. [13]Laboratoire de
Référence des Mycobactéries, Cotonou, Benin. [14]Department of Biomedical Services,
Rwanda Biomedical Center. [15]School of Chemistry and Biosciences, University of Bradford,
Bradford, UK.

[#]**contributed equally**

[*]**Joint senior authors**

Materials and correspondence: Conor Meehan: c.meehan2@bradford.ac.uk, Philip Supply:
philip.supply@ibl.cnrs.fr, Sebastien Gagneux: sebastien.gagneux@swisstph.ch

Abstract

The human- and animal-adapted lineages of the *Mycobacterium tuberculosis* complex

(MTBC) are thought to have clonally expanded from a common progenitor in Africa.

However, the molecular events that accompanied this emergence remain largely unknown.

Here, we describe two MTBC strains isolated from patients with multidrug-resistant

tuberculosis, representing an as-yet-unknown lineage, named Lineage 8 (L8), seemingly

restricted to the African Great Lakes region. Using genome-based phylogenetic

reconstruction, we show that L8 is a sister clade to the known MTBC lineages. Comparison

with other complete mycobacterial genomes indicate that the divergence of L8 preceded the

loss of the *cobF* genome region - involved in the cobalamin/vitamin B12 synthesis - and gene

interruptions in a subsequent common ancestor shared by all other known MTBC lineages.

This discovery further supports an East African origin for the MTBC and provides additional

molecular clues on the ancestral genome reduction associated with adaptation to a

pathogenic lifestyle.

Introduction

Tuberculosis (TB), caused by members of the *Mycobacterium tuberculosis* complex (MTBC), is among the ancient scourges of humankind[1], and remains the leading cause of mortality globally due to an infectious disease[2]. Intense research has been dedicated to decipher the evolutionary history of the MTBC and to understand the causes underlying the worldwide spread of TB[3–5]. Current genome data show that the MTBC is comprised of the five human-adapted lineages representing *M. tuberculosis sensu stricto* (L1–4, and L7), two other human-adapted lineages traditionally referred to as *M. africanum* (L5-6) and at least nine animal-adapted lineages[6]. Africa is the only continent where all MTBC lineages are present, suggesting that the MTBC emerged from a common ancestor therein and then clonally expanded to the rest of the world following human migrations[3,7–10]. However, the genomic traits of this common ancestor and the region from which this expansion took place in Africa remain unknown. Whole genome sequencing (WGS) analyses showed that rare human TB bacilli with a smooth colony morphotype, highly restricted to the Horn of Africa and named *Mycobacterium canettii* (alias smooth tubercle bacilli or STB) represent early evolutionary branching lineages that predate the emergence of the most recent common ancestor (MRCA) of the MTBC (or of the rest of the MTBC, if *M. canettii* is considered to be part of the complex)[4,11,12] Indeed, whereas known MTBC strains differ by no more than ~2,000 Single Nucleotide Polymorphisms (SNPs)[13], *M. canettii* strains are 10 to 25-fold more genetically diverse and separated by at least 14,000 SNPs from the hitherto known MTBC MRCA[4,12]. Moreover, *M. canettii* strains are less virulent and possess highly mosaic genomes, possibly reflecting adaptation to an environmental reservoir favouring active lateral gene flow[4,14–16]. These biological differences support the existence of lineages that reflect intermediate stages in the evolution from *M. canettii* towards the obligate MTBC pathogens.

Here, we describe two exceptional strains representing a new clade, diverging before the

MRCA of the other MTBC lineages. These two strains were discovered in two independent

analyses, and were both multidrug-resistant (MDR; i.e. resistant to at least rifampicin and

isoniazid). One was isolated from a TB patient in Rwanda through an ongoing MDR-TB

diagnostic trial in Africa. The second isolate was from a patient in Uganda, and was

discovered upon screening publicly available draft genome datasets, where it was

misclassified as an *M. bovis* strain. We used PacBio and Illumina WGS to reconstruct the full

circular genome of the Rwandan strain. We used these data and the available Illumina

sequencing data of the Ugandan strain to reconstitute the phylogeny of this novel lineage,

which we named Lineage 8 (L8), and further investigate molecular and evolutionary events

associated with the emergence of the MTBC.

Results

Patient with the L8 TB strain in Rwanda

The strain in Rwanda was isolated from a male patient, aged 77 years, HIV-negative, resident

of Rulindo district bordering with the Southwest of Uganda, and who had lived in Uganda

previously. The patient was diagnosed with rifampicin-resistant TB based on standard Xpert

MTB/RIF testing (Xpert; Cepheid, Sunnyvale, CA, USA), which probes for mutations in the

rifampicin resistance-determining region of the *rpoB* gene of MTBC strains[17]. The results of

the assay showed a rare delayed probe B reaction (~3% prevalence in Rwanda)[18], presumed

(and later confirmed; see below) to be due to the rifampicin resistance-associated

Asp435Tyr mutation[19].

Per routine practice, the patient was initiated on standard short-course MDR-TB treatment

(i.e. 9-month WHO-endorsed MDR-TB regimen, including moxifloxacin, kanamycin,

protionamide, ethambutol, clofazimine, high dose isoniazid and pyrazinamide)[20]. However,

the patient developed hypotension, and eventually died due to probable cardiac failure,

after 20 days of treatment. Phenotypic drug-susceptibility testing (DST) confirmed resistance

to both rifampicin and isoniazid, and susceptibility to other anti-TB drugs including

ethambutol, fluoroquinolones, and second-line injectables.

Growth characteristics and biochemical properties of the Rwandan strain

The strain from Rwanda was grown in 12.5 days on Mycobacterial Growth Indicator Tubes.

Colonies were observed on the fifth week after initial inoculation on Löwenstein-Jensen

medium, indicating a slow grower phenotype with rough colonies (Figure 1). The strain also

displayed archetypal biochemical characteristics of *M. tuberculosis sensu stricto*, including

niacin production combined with urease hydrolysis (Table 1).

Genotypic resistance and SNP profile of the Rwandan strain by Deeplex Myc-TB

Following the MDR-TB diagnosis, the strain was included in the first set of tests for an

ongoing MDR-TB diagnostic trial "DIAgnostics for MDR-TB in Africa (DIAMA) Clinicaltrials.gov,

NCT03303963", evaluating a new targeted deep sequencing assay, called Deeplex-MycTB.

Deeplex-MycTB testing confirmed the presence of the *rpoB* Asp435Tyr mutation conferring

rifampicin resistance, along with the *inhA* Ser94Ala mutation conferring isoniazid resistance,

consistently with the MDR phenotype identified by phenotypic DST (**Figure 2**). This strain

also harboured two alleles in phylogenetic positions in *embB* (Ala378) and *gidB* (Ala205) not

associated with resistance to ethambutol or streptomycin, which were both shared by

several MTBC lineages (L1, 5, 6, 7, and animal lineages) and *M. canettii*[21]. In addition, nine

other - so far uncharacterized - SNPs were identified in six of the 18 gene targets

interrogated by the assay (**Figure 2**). Moreover, this test detected an atypical spoligotype

pattern, 1111100000000000000000000000001110000000 (**Figure 2**), which was further

confirmed by conventional membrane-based spoligotyping testing. This spoligotype pattern

was unique in the global spoligotype database that comprises 111,637 MTBC isolates from

131 countries[22].

WGS analysis and phylogenetic position of the Rwandan and Ugandan strains

Results from WGS analysis of the Rwandan strain using Illumina sequencing confirmed all

Deeplex-MycTB findings.

The strain isolated in Uganda was discovered independently upon screening global MTBC

genome datasets publicly available on NCBI/EBI, comprising approximately 20,000 genomes

in total. From subsequent processing with our WGS analysis pipeline, we found 1 genome

that did not classify in any of the 7 human-adapted lineages or 9 animal-adapted ecotypes

known at the time, but had been misclassified as *M. bovis* isolated from a human patient[23].

These WGS data revealed a similar spoligotype

1111100000000000000000000000000001111000111, characterized by the presence of

spacers 1 to 5 and 34 to 37 (vs 34-36 in the Rwandan strain) with all intervening spacers

missing. Moreover, the Ugandan strain also shared the same *rpoB* Asp435Tyr and *inhA*

Ser94Ala mutations and the same sequence alleles in *embB* and *gidB*. The Ugandan strain

contained an additional *katG* Ser315Thr mutation conferring isoniazid resistance, as well as

the *embA* C-11A and *embB* Asp328Tyr mutations, and two *pncA* missense mutations,

predictive of pyrazinamide resistance. Moreover, only three of the nine aforementioned

uncharacterized SNPs detected by Deeplex-MycTB were shared between both strains.

To further assess the relationships between both strains and in comparison to other MTBC

strains, a maximum likelihood phylogeny was inferred from 241 MTBC genomes, including

representatives of all known human- and animal- adapted lineages[6] and using a *M. canettii*

strain as an outgroup. *M. canettii* represents the closest outgroup to the MTBC including L8,

as shown by subsequent comparative analysis of a complete L8 genome (see below), and

previous observations of ~2.0 Mb larger genomes and substantially lower average nucleotide

identities of phylogenetically closest non-tuberculous mycobacterial species such as *M.*

marinum and *M. kansasii*[4,24,25]. This reconstruction revealed a unique phylogenetic position

of the two new genomes from Rwanda and Uganda (**Figure 3**), representing a newly

characterised monophyletic clade in which none of the known MTBC genomes are

contained. A core genome-based phylogeny was also constructed from representatives of

the MTBC lineages as well as *M. canettii*, *M. marinum* and *M. kansasii* (Supplementary figure

1). This phylogeny confirmed the placement of the L8 Rwandan strain as being a new clade

between *M. canettii* and the other lineages of the MTBC. Based on these phylogenies, this

clade shares a MRCA with the rest of the MTBC, thus representing a new sister clade to the

known MTBC, which we named Lineage 8 (L8). Comprehensive SNP analysis identified a total

of 189 SNPs separating both L8 genomes, which is within the range of zero to 700 SNPs

found between any two strains within any of the lineages 1 to 7 of the MTBC[13]. On average,

two strains of the MTBC including L8 differed by 1443 SNPs (corresponding to 0.04% of the

genome, excluding repetitive/'problematic' regions), thus at least an order of magnitude

lower than the SNP distance separating any MTBC strain from *M. canettii*[4, 13].

The absence of any matching pattern in the global spoligotype database, as well as the lack

of detection of this clade in previous large WGS datasets of MTBC strains from global

sources, suggests that L8 is rare and seemingly geographically restricted to the African Great

Lakes region. Specifically, the L8 spoligotype signature and the 3 SNPs shared by both L8

strains were not detected in any of 115 MTBC genomes from a previous drug resistance

survey in Uganda[26], nor in 380 rifampicin-resistant strains from Rwanda collected between

1991 and 2018, from routine drug resistance surveillance as well as various drug resistance

surveys[27–29]. Furthermore, among 14 other available isolates out of 27 from Uganda and

Rwanda tested by Gene Xpert MTB/RIF that showed the same delayed probe B as L8, none

displayed the L8 signatures when tested by Deeplex Myc-TB or by classical spoligotyping.

Likewise, none of > 1,500 clinical samples from TB patients tested by Deeplex-MycTB from a

recent nationwide drug resistance survey performed in the Democratic Republic of Congo

displayed the L8 spoligotype signature or the specific SNPs.

Defining features of a complete L8 genome

To further assess the sister position of L8 and split from the remaining MTBC inferred from

SNP analysis, the Rwandan strain was subjected to long read-based PacBio sequencing.

Comparison of the obtained assembly with 36 available complete genomes of MTBC

members, comprising L1-L4 (including H37Rv), *M. africanum* (L6) and *M. bovis* strains,

showed a highly syntenic organization, with no major structural rearrangement between

both groups. Although the assembled L8 genome of 4,379,493 bp was within the 4.34-4.43

Mb size range of the other MTBC genomes, it was 30 kb smaller than the 4.41-Mb mean size

of genomes of *M. tuberculosis sensu stricto*[30]. However, the largest part of this gap was

accounted by the absence of three genomic regions in L8, corresponding to regions of

difference (RDs) known to be variably present or absent in other MTBC

(sub)lineages[31,32](Supplementary Table 2). These include a 9.3-kb PhiRv1 prophage region

(RD3), as well as 10.0-kb and 8.5-kb segments corresponding to RD14 and RD5, comprising

the *plcABC* gene cluster and the *plcD* gene regions, respectively[31]. In L8, each of the two

latter regions only contained one copy of the IS*6110* insertion sequence, devoid of direct

repeats (DRs) that normally flank IS*6110* copies after transposition, indicating that these

deletions in L8 resulted from recombination between two adjacent IS*6110* copies with loss

of the intervening sequences[33]. These mobile DNA-related deletions, which also arose

independently in several other MTBC branches[31,34], probably occurred after the divergence

of L8 from the other MTBC lineages. Apart from these three deletions and two dozen

repetitive/multicopy genes (IS6110-related, PE/PPE-, or Mce-encoding), we only found 5

non-repetitive genes, included in two small segments (3.4 and 4.4 kb), which were

undetected in the complete L8 genome while being present in reference MTBC genomes

(Supplementary Table 2).

Conversely, a particular 4.4 kb genome region was present in the genomes of both L8 strains

and in *M. canettii*, but absent in the 36 available complete genomes of MTBC members

(Supplementary Table 3). This region comprises the *cobF* gene (**Figure 4**), encoding the

precorrin 6A synthase involved in the cobalamin/vitamin B12 synthesis, along with two other

genes, respectively encoding a PE-PGRS protein family member and a protein of unknown

function. This region shared by the L8 and the *M. canettii* genomes is also present in the

phylogenetically proximal non-tuberculous mycobacterial species *M. marinum* and *M.*

kansasii (Supplementary Table 3). Via BLAST analysis, we further confirmed the systematic

absence of *cobF* in any of 6,456 quality draft genome assemblies available as of January 2020

from the NCBI, from strains belonging to lineages 1 to 7 or the animal lineages of the MTBC.

Moreover, we thereby determined that the junction between the sequences flanking the

cobF deletion was at the same nucleotide position in all but 6 of these genomes, resulting in

the truncation of *rv0943c* and *rv0944* genes as seen in the complete MTBC genomes (Fig. 4).

Consistent with the clonal evolution of the MTBC with negligible, if any, horizontal gene

transfer between strains[1,14,32,35] this perfect conservation of this sequence junction suggests

that *cobF* was lost in the MRCA of the other MTBC lineages, after its divergence from L8. The

6 exceptions were 3 strains from lineage 4.3 and 3 from lineage 3 that showed slightly larger

deletions, including the 5' region of *rv0943c* or the 5' region of *rv0943c*, *rv0944* and the 5'

region of *rv0945*, respectively, suggesting probable subsequent deletion events in particular

sub-branches of these lineages.

In contrast, none of the almost 900 other genes specifically identified in the *M. canettii*

genomes, and absent in the other MTBC genomes, were found in the L8 genome. The latter

finding thus supports the close relationship with the previously known MTBC lineages

indicated by the SNP-based phylogeny, as well as the outgroup position of *M. canettii*

relatively to the MTBC including L8.

Further evidence for the early branching of L8 relative to the rest of the MTBC comes from

examination of interrupted coding sequences (ICDSs). These ICDSs correspond to frameshifts

or in-frame stop codons detected in genes originally intact in a common progenitor, thus

putatively representing so-called molecular scars inherited during progressive

pseudogenization of the MTBC genomes[36,37]. Four orthologues of MTBC ICDSs were

previously found to be intact in the genomes of *M. canettii* strains, as well as in *M. marinum*

and *M. kansasii*[4]. One of these four orthologues (*pks8*), which belongs to a multigene family

encoding polyketide synthases involved in the biosynthesis of important cell envelope

lipids[38], was also intact in the genomes of both L8 strains (Figure 5 and Supplementary Table

4). Moreover, we found an additional orthologue of MTBC ICDSs (i.e. *rv3899c-rv3900c*),

coding for a conserved hypothetical protein, which was intact in the genomes of *M. canettii*,

M. kansasii, M. marinum and both L8 strains (Supplementary Table 4). These two molecular

scars were also likely acquired by the other MTBC lineages after their divergence from the

common progenitor shared with L8.

The assembled L8 genome also included 48 out of 50 genes (the exceptions are rv3513c

encoding the probable fatty-Acid-Coa ligase FadD18 and the PhiRv1 region; see above)

present in MTBC members but not found in any of the *M. canettii* genomes, including a

number of genes putatively acquired through horizontal gene transfer by the common

ancestor of the MTBC after its separation from *M. canettii*[4] (Supplementary Table 5). This

observation additionally supports both the close relationship with the previously known

MTBC lineages and the outgroup position of *M. canettii* relatively to the MTBC including L8.

Likewise, consistent with the rough colony morphotype of the Rwandan strain, both L8

strains displayed the single polyketide-synthase-encoding *pks5* gene configuration shared by

all MTBC members, instead of the dual *pks5* conformation found in *M. canettii* strains

involved in the smooth colony phenotype of the latter strains[15]. Thus, the recombination

between the two *pks5* genes and the loss of the intervening *pap* gene, thought to have

resulted in surface remodelling and incremental gain of virulence after the phylogenetic

separation from *M. canettii*[15], already existed in the common progenitor of L8 and the rest

of the MTBC. Moreover, both L8 strains also contained the intact TbD1 and RD9 regions,

shared by the other "ancestral" *M. tuberculosis* lineages (L1, L7) but subsequently lost by the

so-called "modern" lineages of *M. tuberculosis* (TbD1 lost in L2-4), *M. africanum* (L5 and L6)

and the animal lineages (RD9)[31].

In contrast to the highly clonal population structure of the MTBC, *M. canettii* strains are

highly recombinogenic, as apparent from mosaic sequence arrangements in their genomes

and functional DNA transfer between *M. canettii* strains mediated by a distributive conjugal

transfer (DCT)-like mechanism[4,39]. However, no significant genome-wide recombination

signal was detected by ClonalFrameML analysis[40] between L8 and other MTBC strains

(Supplementary Figure 2). In particular, and in contrast to the numerous recombination

segments in *M. canettii*[4], the complete L8 genome only contained 26 possible recombination

segments, yet the longest of these was 607bp and the average length was 142bp.

Discussion

The discovery of L8 provides unique insights into an ancestor of the MTBC that existed after the *pks5*-recombination-mediated surface remodelling, which occurred after separation of the MTBC MRCA from the *M. canettii* clade, but preceded the loss of the *cobF* region and gene interruptions in a later common ancestor of the other MTBC lineages. The seeming restriction of this lineage to the African Great Lakes region represents new evidence supporting an origin for the MTBC in the eastern part of the African continent. These findings reinforce results from previous work suggesting an East- rather than a West African origin of the MTBC[3,4,9,11,41].

A distinct ecological niche, linked to a potential environmental reservoir, has been hypothesized to explain the marked geographic restriction of *M. canettii* strains to the Horn of Africa, the lower persistence of these strains in infection models as well as their genome mosaicism implying multiple DNA recombination events within the *M. canettii* strain pool[4,12]. However, although our analysis is limited to two genomes identified to date, our results suggest that L8 is as clonal as the rest of the MTBC[3,14,34,42]. Moreover, the observation that both L8 strains share two uncommon rifampicin- and isoniazid-resistance conferring mutations in *rpoB* and *inhA* suggests that multidrug resistance was already acquired in the common ancestor of these two strains. Isoniazid and rifampicin were introduced in TB treatments in the African Great Lakes region in the late fifties and 1983, respectively (Dr. Armand Van Deun, personal communication). These shared MDR-defining mutations, and the detection of these isolates in human patients in both cases (with reported absence of previous TB history for the Rwandan patient), suggest that these patients were infected with an already-resistant strain, which was exposed to drug selective pressure already decades ago and had been likely circulating in the community for some time. Overall, this pattern

thus suggests human-to-human transmission rather than infection from a non-human

source. While based on only two initial strains, these results are consistent with the

presumed scenario of a human rather than a zoonotic origin for the MTBC[31,43].

Of note, given the above timeline of introduction of rifampicin and isoniazid in both

countries, the ~100 SNPs distance separating these two strains from their MRCA would imply

a rapid molecular clock for L8, in the range of the upper bound of 2.2 SNPs/genome/year

most recently estimated for other MTBC clades[44]. However, this mutation rate cannot be

confirmed until additional L8 samples are uncovered.

Remarkably, the absence of other L8 strains in datasets from Uganda, Rwanda and DRC,

together comprising more than 2,000 strains, suggests that L8 is rare even within the African

Great Lakes region. Such scarcity is compatible with selective sweeps of later branching

MTBC strains, introduced more recently into the region. Similar scenarios have also been

proposed to explain the slow apparent replacement of MTBC L5 and L6 by L4 in West

Africa[45–47] and the restriction of L7 to Northern Ethiopia[48].

Loss-of-function linked to the deletion of *cobF* is a plausible candidate molecular event

involved in such a replacement scenario for L8. Indeed, loss-of-function appears to be an

important mechanism driving the pathoadaptive evolution of the TB pathogen, as shown for

the role of the loss of lipo-oligosaccharide production (via recombination in the *pks5* locus)[15]

in the evolution towards increased virulence from *M. canettii* to MTBC strains. Likewise, loss

of secretion of PPE-MPTR and PE_PGRS proteins by the type VII secretion system ESX-5 (via

mutations of the *ppe38* locus) has been involved in the hypervirulence of recent branches of

L2 (alias "modern" Beijing) strains[49]. The loss of the *cobF* region in the other MTBC lineages,

inferred from comparative genomics with *M. canettii* and non-tuberculous mycobacteria[4],

was previously hypothesised to reflect enhanced adaptation to an intracellular parasitic

lifestyle[50]. Indeed, the cobalamin/vitamin B12 synthesis pathway, of which the *cobF*-

encoded precorring-6a synthase is a component, represents a highly complex and energy

consuming process with about 30 enzymatic steps[51]. While the absence of this component

may not entirely ablate cobalamin biosynthesis[52,53], its loss might have resulted in gain of

fitness and reflect enhanced pathogenic professionalisation, by economical reliance upon

the mammalian host environment as source of vitamin B12. As an additional plausible but

not necessarily mutually exclusive hypothesis, such selective sweep of L8 might have been

(further) enhanced by the loss of theTbD1 region in later branching MTBC strains. This

region, which we also found intact in L8, as is the case in the "ancestral" *M. tuberculosis*

lineages L1 and L7, encodes members of the mycobacterial membrane protein families

MmpL. Very recent findings indicate that the loss of this region in later branching MTBC

strains was also associated with a gain of virulence, and the deletion of TbD1 at the origin of

the "modern" *M. tuberculosis* lineages L2/L3/L4 has therefore been suggested as a key driver

for their global epidemic spread[54]. If true, more recently emerged or introduced *cobF*- and

TbD1-deleted strains might conceivably have largely outcompeted L8 strains. This hypothesis

could be explored by assessing the growth and the virulence/fitness in cellular and animal

models, of recombinant *cobF*- and/or TbD1-knock-out, as well as *cobF*- and/or TbD1-knock-

in strains, derived from the available Rwandan L8 strain and other MTBC strains,

respectively.

In conclusion, our genomic data, on an as-yet-unknown ancestral stage between the MTBC

and the putative progenitor pool of *M. canettii*-like mycobacteria, thus suggest further

experiments to examine candidate molecular events potentially involved in the

pathoadaptive evolution of *M. tuberculosis*. The discovery of such rare strains raises the

possibility for the existence of further extant strains, especially in Eastern Africa,

representing other clades further closing the biological gap between the MTBC and *M.*

canettii.

Methods

Rwandan patient recruitment and ethics statement

The patient in Rwanda was recruited into, and gave informed consent for, the DIAMA study which was approved by the Rwanda National Ethical committee (IRB 00001497 of IORG0001100; Ref No.0069/RNEC/2017).

Phenotypic characterization

We studied conventional mycobacterial growth and biochemical characteristics including colony morphology, niacin production, nitrate reduction, p-nitro benzoic acid growth inhibition, catalase production, urea hydrolysis, tween 80 hydrolysis, and thiophene carboxylic acid hydrazide growth inhibition[55]. For comparative purpose, a reference set of the seven known human-adapted MTBC lineages[56], together with *M. canettii* (BCCM/ITM500199), *M. bovis* (BCCM/ITM500324), *M. bovis BCG* (BCCM/ITM500101), and *M. orygis* (BCCM/ITM501026) strains were processed with the novel strain isolated in Rwanda. Moreover, phenotypic drug-susceptibility testing to first- and second-line anti-TB drugs was done using the proportion method[57]. The strain isolated in Uganda was not available for phenotypic characterization.

Targeted and whole genome sequencing of the Rwandan strain

Targeted sequencing was performed by using the Deeplex-MycTB assay[58] (Genoscreen, France). Briefly, this assay relies on a 24-plexed PCR amplification of mycobacterial species identification (*hsp65*), genotyping (spoligotyping and phylogenetic single nucleotide polymorphisms (SNPs)) and 18 *M. tuberculosis* complex drug resistance-associated gene targets. This test and short-read Illumina-based WGS were performed on the Rwandan strain as follows. A bead beating method was used to extract DNA from colonies (Supplementary method 1). Libraries of Deeplex-MycTB amplicons or genome fragments were constructed

using the Nextera XT kit and sequenced on an Illumina MiSeq platform with paired end, 150-

bp read lengths (Illumina, CA, USA). DNA extraction suitable for PacBio SMRT sequencing

was performed using the Genomic DNA Buffer Set (Qiagen Inc, Germantown, Maryland,

USA) (Supplementary method 2). Sequencing was performed on a PacBio RS II using the

SMRT technology.

Deeplex-MycTB analysis and spoligotyping

Analysis of the Deeplex-MycTB sequencing data, including SNP calling and spoligotype

identification, was performed by read mapping on *M. tuberculosis* H37Rv sequence

references, using a parameterized web application (GenoScreen)[58]. Membrane-based

spoligotyping was performed as described previously[59].

Illumina whole genome sequencing analysis

Raw genomic reads from the newly sequenced L8 genome from Rwanda and the L8 genome

from Uganda (SAMN02567762) were processed as previously described[60]. Briefly, the reads

were trimmed with Trimmomatic v0.33.22[61] and reads larger than 20 bp were kept. The

software SeqPrep (https://github.com/jstjohn/SeqPrep) was used to identify and merge any

overlapping paired-end reads. The resulting reads were aligned to the reconstructed

ancestral sequence of the MTBC[62] using the mem algorithm of BWA v0.7.13[63]. Duplicated

reads were marked using the MarkDuplicates module of Picard v2.9.1

(https://github.com/broadinstitute/picard) and local realignment of reads around InDels was

performed using the RealignerTargetCreator and IndelRealigner modules of GATK v3.4.0[64].

SNPs were called with Samtools v1.2 mpileup[65] and VarScan v2.4.1[66] using the following

thresholds: minimum mapping quality of 20, minimum base quality at a position of 20,

minimum read depth at a position of 7X, maximum strand bias for a position 90%.

The spoligotype pattern of the strain from Uganda was extracted *in silico* from the raw reads

using kvarQ[67].

Phylogenetic reconstruction

The maximum likelihood phylogeny was inferred with RAxML v.8.2.8[68] using an alignment

containing only polymorphic sites and the branch lengths of the tree were rescaled using

invariant sites (rescaled_branch_length = (branch_length * alignment_length) /

(alignment_length +invariant_sites))[44,69].

A position was considered polymorphic if at least one genome had a SNP at that position.

Deletions and positions not called according to the minimum threshold of 7x were encoded

as gaps. We excluded positions with more than 20% missing data, positions falling in PE-

PGRS genes, phages, insertion sequences and in regions with at least 50 bp identity to other

regions in the genome. We also excluded variable positions falling in drug resistance-related

genes. The phylogeny was computed using the general time-reversible model of sequence

evolution (-m GTRCAT -V options), 100 bootstrap inferences and *M. canettii* (SRR011186)

was used as an outgroup to root the phylogeny.

Whole genome *de novo* assembly, annotation and comparative genomics

Raw PacBio reads obtained from the Rwandan strain were assembled with Canu v1.6[70], using

default settings and an expected genome size of 4.4 Mbp, typical of MTBC strains. After

discarding 60,272 reads below minimal quality parameters, 106,681 reads were used for the

assembly. Based on the expected genome size, the average coverage depth was estimated at

186x using raw reads, and 39x and 38x using corrected and trimmed reads, respectively. The

obtained unique contig of 4,387,285 bp was circularized with Circlator v1.5.5[71] using default

settings, resulting in an assembly of 4,379,493 bp. Additional sequence verification and

correction was then performed by mapping Illumina reads obtained from the same strain,

using pacbio-utils version 0.2[72] (https://github.com/douglasgscofield/PacBio-utilities) and snippy

version 4.4[73] (https://github.com/tseemann/snippy). Alignments of the final assembly were

performed against an ensemble of complete genome sequences available from 38 strains of

tubercle bacilli. This set included 34 *M. tuberculosis* strains from lineages 1, 2, 3 and 4

(comprising H37Rv), *M. africanum* L6 GM041182, *M. bovis* AF2122/97, as well as the closest

STB-A (CIPT 140010059) and most distant (STB-K) *M. canettii* strains (Supplementary Table

1). Comparative alignments and genome annotation were performed based on BLAST

searches and analysis of gene synteny, using Artemis and Artemis comparison tool[74], as well

as a custom Multiple Annotation of Genomes and Differential Analysis (MAGDA) software

previously used for annotation of *M. canettii* and *Helicobacter pylori* genomes[4,75].

Comparisons with orthologues from *M. canettii* STB-D, -E, -G, -H, -I, and -J in addition to STB-

A and -K, and from *M. marinum* type strain M and *M. kansasii* genomes were additionally

done using the Microscope platform v3.13.3[76]. When applicable, annotations were

transferred from those of *M. tuberculosis* or *M. canettii* orthologs in the

TubercuList/Mycobrowser database, using BLAST matches of > 90% protein sequence

identity, an alignable region of >80% of the shortest protein length in pairwise comparisons

and visual inspection of the gene synteny. Genome completeness was assessed using

CheckM[77] using the lineage-specific workflow and *Mycobacterium* as the genus. The PacBio

assembled Rwandan strain was found to have 98.74% completeness and 0% contamination

or strain heterogeneity, making it suitable for further analyses.

ACT comparison files were generated using MAUVE 2015-02-25 software to visualize the

genome-wide distribution of SNP densities between the assembled L8 genome from Rwanda

and *M. tuberculosis* H37Rv and *M. canettii* STB-A and STB-K genomes. Recombination

between L8 and other MTBC lineages or *M. canettii* was assessed from a progressive MAUVE

alignment of the PacBio assembled L8 genome and previously published closed genomes[73]

using ClonalFrameML[40]. To further assess the phylogenetic placement of the Lineage 8

strain, a core gene alignment was constructed using the completed genomes of the

Rwandan L8 strain, representatives of the MTBC lineages 1-4 and 6, *M. canettii, M. bovis, M.*

marinum (accession GCF_000419315.1) and *M. kansassii* (accession GCF_000157895.3). The

GFF files of each genome were input to roary[78] with an 80% identity cut-off, as has been

done for previous genus-level mycobacterial core alignments[79]. A phylogenetic tree was

constructed from the core gene superalignment using RAxML-NG v 0.9.0[80] under the

GTR+Gamma model of evolution with 20 starting trees. Bootstrapping was run until

autoMRE converged with a value of 0.03 (50 replicates).

Accession codes

The complete genome sequence of the L8 strain from Rwanda was deposited in the NCBI

repository under project PRJNA598991 with SRR10828835 and SRR10828834 accession

codes for Illumina- and PacBio-derived genome sequences, respectively.

References

1.   Gagneux, S. Ecology and evolution of Mycobacterium tuberculosis. *Nature Reviews*

Microbiology **16**, 202–213 (2018).

2.   World Health Organization. *Global Tuberculosis Report 2018*. (2018).

3.   Comas, I. *et al.* Out-of-Africa migration and Neolithic coexpansion of Mycobacterium

tuberculosis with modern humans. *Nature genetics* **45**, 1176–82 (2013).

4.   Supply, P. *et al.* Genomic analysis of smooth tubercle bacilli provides insights into

ancestry and pathoadaptation of Mycobacterium tuberculosis. *Nature genetics* **45**,

172–9 (2013).

5.   Bos, K. I. *et al.* Pre-Columbian mycobacterial genomes reveal seals as a source of New

World human tuberculosis. *Nature* **514**, 494–7 (2014).

6.   Brites, D. *et al.* A New Phylogenetic Framework for the Animal-Adapted

Mycobacterium tuberculosis Complex. *Frontiers in Microbiology* **9**, 2820 (2018).

7.   Rutaihwa, L. K. *et al.* Multiple Introductions of Mycobacterium tuberculosis Lineage 2–

Beijing Into Africa Over Centuries. *Frontiers in Ecology and Evolution* **7**, 112 (2019).

8.   O'Neill, M. B. *et al.* Lineage specific histories of Mycobacterium tuberculosis dispersal

in Africa and Eurasia. *Molecular Ecology* **28**, 3241–3256 (2019).

9.   Wirth, T. *et al.* Origin, spread and demography of the Mycobacterium tuberculosis

complex. *PLoS pathogens* **4**, e1000160 (2008).

10.   Gagneux, S. *et al.* Variable host-pathogen compatibility in Mycobacterium

tuberculosis. *Proceedings of the National Academy of Sciences of the United States of*

America **103**, 2869–73 (2006).

11.   Gutierrez, M. C. *et al.* Ancient origin and gene mosaicism of the progenitor of

Mycobacterium tuberculosis. *PLoS pathogens* **1**, e5 (2005).

12.   Blouin, Y. *et al.* Progenitor "Mycobacterium canettii" clone responsible for lymph

467   node tuberculosis epidemic, Djibouti. *Emerging infectious diseases* **20**, 21–8 (2014).

468   13.   Coscolla, M. & Gagneux, S. Consequences of genomic diversity in Mycobacterium

469   tuberculosis. *Seminars in Immunology* **26**, 431–444 (2014).

470   14.   Boritsch, E. C. *et al.* Key experimental evidence of chromosomal DNA transfer among

471   selected tuberculosis-causing mycobacteria. *Proceedings of the National Academy of*

472   *Sciences of the United States of America* **113**, 9876–81 (2016).

473   15.   Boritsch, E. C. *et al.* pks5-recombination-mediated surface remodelling in

474   Mycobacterium tuberculosis emergence. *Nature Microbiology* **1**, 15019 (2016).

475   16.   Mortimer, T. D. & Pepperell, C. S. Genomic signatures of distributive conjugal transfer

476   among mycobacteria. *Genome biology and evolution* **6**, 2489–500 (2014).

477   17.   Helb, D. *et al.* Rapid detection of Mycobacterium tuberculosis and rifampin resistance

478   by use of on-demand, near-patient technology. *Journal of Clinical Microbiology* **48**,

479   229–237 (2010).

480   18.   Ng, K. C. S. *et al.* Automated algorithm for early identification of rifampicin-resistant

481   tuberculosis transmission hotspots in Rwanda [abstract]. *The International Journal of*

482   *Tuberculosis and Lung Disease* **22**, 605 (2018).

483   19.   Torrea, G. *et al.* Variable ability of rapid tests to detect Mycobacterium tuberculosis

484   rpoB mutations conferring phenotypically occult rifampicin resistance. *Scientific*

485   *Reports* **9**, (2019).

486   20.   Trébucq, A. *et al.* Treatment outcome with a short multidrug-resistant tuberculosis

487   regimen in nine African countries. *The International Journal of Tuberculosis and Lung*

488   *Disease* **22**, 17–25 (2018).

21.   Coll, F. *et al.* A robust SNP barcode for typing Mycobacterium tuberculosis complex

strains. *Nature Communications* **5**, (2014).

22.   Couvin, D., David, A., Zozio, T. & Rastogi, N. Macro-geographical specificities of the

prevailing tuberculosis epidemic as seen through SITVIT2, an updated version of the

Mycobacterium tuberculosis genotyping database. *Infection, Genetics and Evolution*

(2018). doi:10.1016/j.meegid.2018.12.030

23.   Wanzala, S. I. *et al.* Retrospective Analysis of Archived Pyrazinamide Resistant

Mycobacterium tuberculosis Complex Isolates from Uganda-Evidence of Interspecies

Transmission. *Microorganisms* **7**, (2019).

24.   Stinear, T. P. *et al.* Insights from the complete genome sequence of Mycobacterium

marinum on the evolution of Mycobacterium tuberculosis. *Genome Research* **18**, 729–

741 (2008).

25.   Wang, J. *et al.* Insights on the emergence of Mycobacterium tuberculosis from the

analysis of Mycobacterium kansasii. *Genome biology and evolution* **7**, 856–70 (2015).

26.   Ssengooba, W. *et al.* Whole genome sequencing to complement tuberculosis drug

resistance surveys in Uganda. *Infection, genetics and evolution: journal of molecular*

epidemiology and evolutionary genetics in infectious diseases **40**, 8–16 (2016).

27.   Carpels, G. *et al.* Drug resistant tuberculosis in sub-Saharan Africa: an estimation of

incidence and cost for the year 2000. *Tubercle and lung disease: the official journal*

of the International Union against Tuberculosis and Lung Disease **76**, 480–6 (1995).

28.   Umubyeyi, A. N. *et al.* Results of a national survey on drug resistance among

pulmonary tuberculosis patients in Rwanda. *The international journal of tuberculosis*

and lung disease: the official journal of the International Union against Tuberculosis

and Lung Disease **11**, 189–94 (2007).

29.   World Health Organization & WHO. WHO | Global tuberculosis report 2016. *WHO*

(2019).

30. Yang, T. *et al.* Pan-genomic study of Mycobacterium tuberculosis reflecting the

primary/secondary genes, generality/individuality, and the interconversion through

copy number variations. *Frontiers in Microbiology* **9**, (2018).

31. Brosch, R. *et al.* A new evolutionary scenario for the Mycobacterium tuberculosis

complex. *Proceedings of the National Academy of Sciences of the United States of*

America **99**, 3684–9 (2002).

32. Tsolaki, A. G. *et al.* Functional and evolutionary genomics of Mycobacterium

tuberculosis: Insights from genomic deletions in 100 strains. *Proceedings of the*

National Academy of Sciences of the United States of America **101**, 4865–4870 (2004).

33. Brosch, R. *et al.* Genomic analysis reveals variation between Mycobacterium

tuberculosis H37Rv and the attenuated M. tuberculosis H37Ra strain. *Infection and*

Immunity **67**, 5768–5774 (1999).

34. Hirsh, A. E., Tsolaki, A. G., DeRiemer, K., Feldman, M. W. & Small, P. M. Stable

association between strains of Mycobacterium tuberculosis and their human host

populations. *Proceedings of the National Academy of Sciences of the United States of*

America **101**, 4871–6 (2004).

35. Chiner-Oms *et al.* Genomic determinants of speciation and spread of the

Mycobacterium tuberculosis complex. *Science Advances* **5**, (2019).

36. Deshayes, C. *et al.* Detecting the molecular scars of evolution in the Mycobacterium

tuberculosis complex by analyzing interrupted coding sequences. *BMC evolutionary*

biology **8**, 78 (2008).

37. Smith, N. H., Hewinson, R. G., Kremer, K., Brosch, R. & Gordon, S. V. Myths and

misconceptions: the origin and evolution of Mycobacterium tuberculosis. *Nature*

538  *Reviews Microbiology* **7**, 537–544 (2009).

539 38. Etienne, G. *et al.* Identification of the polyketide synthase involved in the biosynthesis

540  of the surface-exposed lipooligosaccharides in mycobacteria. *Journal of Bacteriology*

541  **191**, 2613–2621 (2009).

542 39. Boritsch, E. C. & Brosch, R. Evolution of Mycobacterium tuberculosis: New Insights

543  into Pathogenicity and Drug Resistance. in *Tuberculosis and the Tubercle Bacillus,*

544  *Second Edition* **4**, 495–515 (American Society of Microbiology, 2016).

545 40. Didelot, X. & Wilson, D. J. ClonalFrameML: efficient inference of recombination in

546  whole bacterial genomes. *PLoS computational biology* **11**, e1004041 (2015).

547 41. Hershberg, R. *et al.* High Functional Diversity in Mycobacterium tuberculosis Driven by

548  Genetic Drift and Human Demography. *PLoS Biology* **6**, e311 (2008).

549 42. Supply, P. *et al.* Linkage disequilibrium between minisatellite loci supports clonal

550  evolution of Mycobacterium tuberculosis in a high tuberculosis incidence area.

551  *Molecular Microbiology* **47**, 529–538 (2003).

552 43. Mostowy, S., Cousins, D., Brinkman, J., Aranaz, A. & Behr, M. A. Genomic Deletions

553  Suggest a Phylogeny for the *Mycobacterium tuberculosis* Complex. *The Journal of*

554  *Infectious Diseases* **186**, 74–80 (2002).

555 44. Menardo, F., Duchêne, S., Brites, D. & Gagneux, S. The molecular clock of

556  Mycobacterium tuberculosis. *PLoS pathogens* **15**, e1008067 (2019).

557 45. Niobe-Eyangoh, S. N. *et al.* Genetic biodiversity of Mycobacterium tuberculosis

558  complex strains from patients with pulmonary tuberculosis in Cameroon. *Journal of*

559  *Clinical Microbiology* **41**, 2547–2553 (2003).

560 46. Godreuil, S. *et al.* First molecular epidemiology study of Mycobacterium tuberculosis

561  in Burkina Faso. *Journal of Clinical Microbiology* **45**, 921–927 (2007).

47.   Groenheit, R. *et al.* The Guinea-Bissau Family of Mycobacterium tuberculosis Complex

Revisited. *PLoS ONE* **6**, e18601 (2011).

48.   Comas, I. *et al.* Population Genomics of Mycobacterium tuberculosis in Ethiopia

Contradicts the Virgin Soil Hypothesis for Human Tuberculosis in Sub-Saharan Africa.

Current Biology **25**, 3260–3266 (2015).

49.   Ates, L. S. *et al.* Mutations in ppe38 block PE_PGRS secretion and increase virulence of

Mycobacterium tuberculosis. *Nature Microbiology* **3**, 181–188 (2018).

50.   Boritsch, E. C. *et al.* A glimpse into the past and predictions for the future: the

molecular evolution of the tuberculosis agent. *Molecular Microbiology* **93**, 835–852

(2014).

51.   Martens, J. H., Barg, H., Warren, M. & Jahn, D. Microbial production of vitamin B12.

Applied Microbiology and Biotechnology **58**, 275–285 (2002).

52.   Gopinath, K. *et al.* A vitamin B $_{12}$ transporter in *Mycobacterium tuberculosis*. *Open

Biology* **3**, 120175 (2013).

53.   Minias, A., Minias, P., Czubat, B. & Dziadek, J. Purifying Selective Pressure Suggests

the Functionality of a Vitamin B12 Biosynthesis Pathway in a Global Population of

Mycobacterium tuberculosis. *Genome Biology and Evolution* **10**, 2326–2337 (2018).

54.   Bottai, D. *et al.* TbD1 deletion as a driver of the evolutionary success of modern

epidemic Mycobacterium tuberculosis lineages. *Nature Communications* **11**, 1–14

(2020).

55.   LEÃO, S. C. *et al.* Practical handbook for the phenotypic and genotypic identification of

mycobacteria. (2004).

56.   Borrell, S. *et al.* Reference set of Mycobacterium tuberculosis clinical strains: A tool for

research and product development. *PLOS ONE* **14**, e0214088 (2019).

57.   Kent, P. & Kubica, G. *Public Health Mycobacteriology: A Guide for the Level III*

587   *Laboratory US Department of Health and Human Services, Centres for Disease Control.*

588   (1985).

589   58.   Makhado, N. A. *et al.* Outbreak of multidrug-resistant tuberculosis in South Africa

590   undetected by WHO-endorsed commercial tests: an observational study. *The Lancet*

591   *Infectious Diseases* **18**, 1350–1359 (2018).

592   59.   van der Zanden, A. G. M. *et al.* Improvement of differentiation and interpretability of

593   spoligotyping for Mycobacterium tuberculosis complex isolates by introduction of

594   new spacer oligonucleotides. *Journal of clinical microbiology* **40**, 4628–39 (2002).

595   60.   Menardo, F. *et al.* Treemmer: a tool to reduce large phylogenetic datasets with

596   minimal loss of diversity. *BMC Bioinformatics* **19**, 164 (2018).

597   61.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

598   sequence data. *Bioinformatics (Oxford, England)* **30**, 2114–20 (2014).

599   62.   Comas, I. *et al.* Human T cell epitopes of Mycobacterium tuberculosis are

600   evolutionarily hyperconserved. *Nature genetics* **42**, 498–503 (2010).

601   63.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler

602   transform. *Bioinformatics (Oxford, England)* **25**, 1754–60 (2009).

603   64.   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for

604   analyzing next-generation DNA sequencing data. *Genome research* **20**, 1297–303

605   (2010).

606   65.   Li, H. A statistical framework for SNP calling, mutation discovery, association mapping

607   and population genetical parameter estimation from sequencing data. *Bioinformatics*

608   *(Oxford, England)* **27**, 2987–93 (2011).

66.   Koboldt, D. C. *et al.* VarScan 2: somatic mutation and copy number alteration

discovery in cancer by exome sequencing. *Genome research* **22**, 568–76 (2012).

67.   Steiner, A., Stucki, D., Coscolla, M., Borrell, S. & Gagneux, S. KvarQ: targeted and

direct variant calling from fastq reads of bacterial genomes. *BMC Genomics* **15**, 881

(2014).

68.   Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

69.   Duchene, S. *et al.* Inferring demographic parameters in bacterial genomic data using

Bayesian and hybrid phylogenetic methods. *BMC evolutionary biology* **18**, 95 (2018).

70.   Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer

weighting and repeat separation. *Genome research* **27**, 722–736 (2017).

71.   Hunt, M. *et al.* Circlator: automated circularization of genome assemblies using long

sequencing reads. *Genome Biology* **16**, 294 (2015).

72.   Scofield, D. G. GitHub - douglasgscofield/PacBio-utilities: Collection of utilities for

working with PacBio-based assemblies. Available at:

https://github.com/douglasgscofield/PacBio-utilities. (Accessed: 20th December

2019)

73.   Seemann, T. GitHub - tseemann-snippy Rapid bacterial SNP calling and core genome

alignments. Available at: https://github.com/tseemann/snippy. (Accessed: 20th

December 2019)

74.   Carver, T. *et al.* Artemis and ACT: viewing, annotating and comparing sequences

stored in a relational database. *Bioinformatics (Oxford, England)* **24**, 2672–6 (2008).

75.   Li, H. *et al.* East-Asian Helicobacter pylori strains synthesize heptan-deficient

lipopolysaccharide. *PLoS Genetics* **15**, (2019).

76.   Médigue, C. *et al.* MicroScope-an integrated resource for community expertise of

gene functions and comparative analysis of microbial genomic and metabolic data.

Briefings in bioinformatics **20**, 1071–1084 (2019).

77.   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:

Assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. *Genome Research* **25**, 1043–1055 (2015).

78.   Page, A. J. *et al.* Roary: Rapid large-scale prokaryote pan genome analysis.

Bioinformatics **31**, 3691–3693 (2015).

79.   Fedrizzi, T. *et al.* Genomic characterization of Nontuberculous Mycobacteria. *Scientific*

Reports **7**, (2017).

80.   Kozlov, A. M., Darriba, D., Flouri, T., Morel, B. & Stamatakis, A. RAxML-NG: a fast,

scalable and user-friendly tool for maximum likelihood phylogenetic inference.

Bioinformatics (Oxford, England) **35**, 4453–4455 (2019).

Acknowledgements

Calculations were partially performed at sciCORE (http://scicore.unibas.ch/) scientific computing core facility at University of Basel. This work was supported by EDCTP2 grant DRIA2014-326—DIAMA of the European Union, the Belgian General Directorate for Development Cooperation (PhD fellowship to JCSN), grant ANR-16-CE35-0009 from Agence Nationale de la Recherche, the Swiss National Science Foundation (grants 310030_188888, IZRJZ3_164171, IZLSZ3_170834 and CRSII5_177163), and the European Research Council (309540-EVODRTB). The views and opinions of authors expressed herein do not necessarily state or reflect those of EDCTP. P.S., C.G. and S.D. declare the following competing interests: P.S. was a consultant of Genoscreen; C.G. and S.D. were employees of the same company. The other authors declare no competing interests. The funders had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

Author contributions

S.G., P.S., C.M., B.C.d.J., L.R. and J.C.S.N. designed the study. P.S., J.C.S.N., C.L., M.M., C.M. and S.G. analyzed data and wrote the manuscript, with comments from all authors. A.J. and C.M. performed the assembly of sequences. M.M. annotated the L8 genome. C.L., F.M., D.B. and A.J. performed SNP analyses and phylogenetic reconstruction. M.M., A.R. and P.S. conducted comparative analyses of complete mycobacterial genomes, with support from C.L. and O.T. J.C.S.N., E.B.N., I.M.H., J.B.M., W.M., K.F. and M.D. performed and/or analyzed data from mycobacterial isolation, growth assays, phenotypic characterization and/or molecular tests. S.D., C.G., J.C.S.N., E.B.N., E.A. and M.K.K. conducted targeted deep sequencing analyses. S.D., C.G. and W.S. L. Majlessi, F.S., C. Locht and C. Leclerc conducted and/or analyzed immune assays. J.T., A. Criscuolo and S.B. conducted MLST, recombination and/or phylogenetic analyses. L.F. conducted histopathological analyses. V.K., M.O. and C.P.

670   created bioinformatics tools and analyzed data. M.F. isolated STB strains. T.S. and T.P.S.

671   conducted core genome and NeighborNet analyses.

672
673   **Figure 1.** Microscopic image of L8 on Dubos agar medium showing typical rough colonies

674   (read at 100x).

Figure 2. Deeplex-MycTB results identifying a multidrug resistant strain from Rwanda with

an atypical genotypic profile in the *M. tuberculosis* complex. Target gene regions are

grouped within sectors in a circular map according to the tuberculous drug resistance with

which they are associated. The two sectors in red indicate regions where rifampicin and

isoniazid resistance associated mutations are detected. The multiple sectors in blue refer to

regions where as yet uncharacterized mutations are detected, while sectors in green

indicate regions where no mutation or only mutations not associated with resistance (shown

in gray around the map) were detected. Green lines above gene names represent the

reference sequences with coverage breadth above 95%. Limits of detection (LOD) of

potential heteroresistance (reflected by subpopulations of reads bearing a mutation),

depending on the coverage depths over target sequence positions, are represented by grey

(LOD 3%) and orange zones (variable LOD >3%–80%; only seen in extremities of a few

targets, such as the two *rrs* regions) above the reference sequences within the sectors.

Information on an unrecognized spoligotype, an equivocal SNP-based and on mycobacterial

species identification, based on *hsp65* sequence best match, are shown in the centre of the

circle. *AMI, amikacin; BDQ, bedaquiline; CAP, capreomycin; CFZ, clofazimine; EMB,

ethambutol; ETH, ethionamide; FQ, fluoroquinolones; KAN, kanamycin; LIN, linezolid; INH,

isoniazid; PZA, pyrazinamide; RIF, rifampin; SM, streptomycin; SIT, spoligotype international

type.

Figure 3 . Maximum likelihood phylogeny of 241 MTBC genomes, inferred from 43,442

variable positions. The scale bar indicates the number of substitutions per polymorphic site.

Branches corresponding to human-adapted strains are coloured and branches corresponding

to animal-adapted strains are depicted in black. The phylogeny is rooted on *M. canettii* and

bootstrap values are shown for the most important splits.

Figure 4. Aligned genome segments showing the *cobF* gene region in *M. tuberculosis* L8, M.

canettii CIPT140010059 (alias STB-A), M. kansasii ATCC12478 and M. marinum M strains, and

the corresponding deletion in *M. tuberculosis* H37Rv, M. bovis AF2122/97, and *M. africanum*

GM041182. Coding sequences of this region are shown in green, and flanking coding

sequences in red. Sequences flanking the deletion point in truncated genes in *M.*

tuberculosis, *M. africanum* and *M. bovis*, and in the cobF region present in *M. canettii*, *M.*

kansasii and *M. marinum* are indicated in red and black, respectively. Dashed lines

correspond to missing segment parts relatively to the longest segment found in *M. marinum*.

Figure 5. Aligned genome segments showing the interrupted coding sequences *pks8/17* in

M. tuberculosis H37Rv and *M. bovis* AF2122/97 gene region, and complete *pks8* genes in L8,

M. canettii CIPT140010059 (alias STB-A) and *M. kansasii* ATCC12478. *pks8/17* and *pks8*

coding sequences are shown in green, and flanking genes in red. Sequences flanking the 1-

nucleotide deletion and resulting in a frameshift in *M. tuberculosis* complex strains are

indicated. Dashed lines correspond to missing segment parts relatively to the longest

segment found in *M. canettii.*

Table 1. Standard biochemical characteristics of selected mycobacterial species or *M. tuberculosis* complex lineages/subspecies versus L8

Mycobacterial species/lineage	BCCM/ITM REF N°	Niacin production	Nitrate reduction	Urease hydrolysis	Tween hydrolysis	Catalase production	Arylsulfatase
L8	500961	+	+	+	weak +	-	-
L1	500941	+	+	+	-	weak +	-
L2	500945	+	-	+	-	-	-
L3	500947	+	+	+	-	-	-
L4	500951	+	+	+	+	-	-
L5	500953	+	-	-	+	-	-
L6	500957	+	-	weak +	weak +	-	-
L7	500959	+	-	weak +	-	-	-
M. bovis	500324	-	-	weak +	-	-	-
M. bovis BCG	500101	-	-	+	-	weak +	-
M. orygis	501026	+	-	+	-	-	-
M. canettii	500199	-	-	+	+	weak +	-
M. fortuitum	500465	-	+	+	-	weak +	+

+: positive reaction; -: negative