The role of maternal effects on offspring performance in familiar and novel environments

Authors: Milan Vrtílek[1]*, Pierre J. C. Chuard[2]*, Maider Iglesias-Carrasco[3], Michael D. Jennions[3] and Megan L. Head[3]

Affiliations:

[1]The Czech Academy of Sciences, Institute of Vertebrate Biology, Brno, Czech Republic

[2]Department of Biological Sciences, Bishop's University, Sherbrooke, Canada

[3]Research School of Biology, The Australian National University, Canberra, Australia

* These authors share joint first authorship on this work

Correspondence: Milan Vrtílek, The Czech Academy of Sciences, Institute of Vertebrate Biology, Květná 8, 603 65 Brno, Czech Republic; phone: +420 543 422 557, fax: +420 543 211 346, e-mail: vrtilek@ivb.cz

Conflict of interest

We declare that we do not have any conflict of interest.

Running title: Maternal effects in novel environment

Version: 21 January 2020

Figures: 0

Tables: 3 (+2 Supplementary Tables)

Word count: 5702 words

Abstract:

Stressful conditions, like novel host environment, can stimulate mothers to

produce offspring with phenotypes that better suit the conditions they are

likely to experience (i.e. adaptive maternal effects). However, mothers

might vary in their ability to adjust their offspring's phenotype in response

to environmental cues. This could generate a maternal-by-environment

interaction (M×E) that inflates estimates of genotype-by-environment

effects (G×E). Our main aim in this study was to test for M×E with a

paternal full-sib/half-sib breeding design in the seed beetle *Callosobruchus*

maculatus, where we split and reared offspring from the same mother on

two different bean host types – original and novel. Partitioning of offspring

phenotypic variation indicated that maternal effects were very small for all

the measured life-history traits, prohibiting a formal test for M×Es. There

was also little evidence that the degree of maternal preference to oviposit

on one host type predicted her offspring's relative performance on this

host. Additive genetic variance for most traits was, however, relatively high

on both hosts. While there was higher heritability in the novel host, there

was no evidence for G×Es and most cross-host genetic correlations were

highly positive. This suggests that offspring from the same family ranked

similarly for performance on both host types. Our results point to a genetic

basis of host adaptation in the seed beetle, rather than non-genetic

maternal effects. Even so, we encourage researchers to test for potential

M×Es because, due to a lack of testing, it remains unclear how often they

arise.

Keywords: genetic correlation, host adaptation, maternal effects, seed

beetle

Introduction

Phytophagous insects are a valuable model for studying life-history evolution, especially in the context of trade-offs arising from adaptation to a preferred host (Agrawal *et al.*, 2010). Antagonistic pleiotropy is predicted to prevent populations of parasites or herbivores from becoming 'masters of all trades' and adapting perfectly to multiple hosts (Agrawal *et al.*, 2010). This prediction is based on the assumption that antagonistic pleiotropy leads to a trade-off in performance on different hosts (Joshi and Thompson, 1995). At the same time, phytophagous insects exhibit a variety of non-genetic maternal effects that can also influence life-history traits (Fox and Dingle, 1994; Fox *et al.*, 1997; Van Asch *et al.*, 2010; Cahenzli and Erhardt, 2013). Adaptive maternal effects that result in cross-generational plasticity are considered to be one way that individuals can perform well under new environmental conditions before selection acts to favour advantageous genetic variation (Fox *et al.*, 1997; Cahenzli *et al.*, 2015). Maternal effects can have far reaching consequences, including niche breadth expansion through shifts in phenology (Van Asch *et al.*, 2010), and even speciation when they facilitate host shifts (Forbes *et al.*, 2017).

Environmental challenges often lead mothers to alter investment in their offspring. This can either facilitate adaptation to new hosts (Fox *et al.*, 1997), or be detrimental to offspring (Magalhães *et al.*, 2011). Maternal

effects can moderate population dynamics (Benton, St Clair, & Plaistow,

2008), and sometimes even alter the likelihood of population extinction or

persistence (Bernardo, 1996). In addition, non-genetic maternal effects can

influence offspring in an environment-specific manner (Bernardo, 1996;

Fox *et al.*, 1997; Räsänen and Kruuk, 2007; Vega-Trejo *et al.*, 2018). It is

therefore no surprise that environment-specific maternal effects (M×E)

could falsely inflate estimates of a Genotype-by-Environment interaction

(G×E) (Vega-Trejo *et al.*, 2018). In the same way that maternal effects have

been shown to inflate estimates of genetic effects (Kruuk and Hadfield,

2007), M×E interactions may result in variation in the response of different

genotypes across environments being incorrectly treated as evidence for a

G×E interaction (Via and Lande, 1985; Saltz *et al.*, 2018). In such cases, a

genetic trade-off in the degree of adaptation to different hosts would be

artificially inflated.

In this study, we investigated the presence of maternal effects, as

well as the potential for an interaction between maternal effects and her

offspring's rearing environment on general offspring life-history traits, as

well as her daughters' egg laying preferences on different hosts in the seed

beetle, *Callosobruchus   maculatus*   (Coleoptera: Chrysomelidae).

Callosobruchus maculatus is a common pest to legume storages worldwide.

The larvae feed on multiple legume species (family Fabaceae) with varying

degrees of success (Gompert and Messina, 2016; Price *et al.*, 2017;

Messina *et al.*, 2018). Females lay individual eggs on host beans shortly

after copulation (Mitchell, 1975). Host choice is crucial as the larvae cannot

move between individual beans. Incorrect decisions on where a female

chooses to lay her eggs inevitably lead to lower quality or even non-viable

offspring (Mitchell, 1975; Messina and Fry, 2003; Messina *et al.*, 2007).

Larvae hatch 4-5 days after oviposition and burrow into the bean to feed

on the endosperm. Depending on the temperature and host species, adults

typically emerge within 26-36 days of oviposition (Fox *et al.*, 2003;

Messina, 2004a). *Callosobruchus maculatus* shows profound sexual

dimorphism in life-history traits with males emerging earlier, being smaller

and living shorter than females (Guntrip *et al.*, 1997; Fox, Bush, *et al.*,

2004).

We used a full-sib/half-sib split brood design to tease apart the role

of additive genetic and maternal effects on *C. maculatus* life-history traits

when larvae develop on two host types – original (cowpea beans) and

novel (mung beans). We hypothesised that:

1) There would be strong maternal effects on offspring life history

traits (Fox, 1993a; Messina and Fry, 2003). This led us to predict that

offspring from a single female (maternal full-siblings) are more similar to

118 each other than to their half-siblings (shared sire, different mother) or to

119 offspring from families sired by other males than expected based solely on

120 genetic relatedness.

121  2) Maternal effects are environment-specific (M×E). Specifically, we

122 predicted there would be a negative correlation between full-sibling

123 performance on the two host types, even after controlling for a negative

124 genetic correlation due to antagonistic pleiotropy.

125  3) Novel host type would be challenging for offspring. Here, we

126 predicted that offspring developing in the novel host would suffer reduced

127 performance compared to those developing in the original host type (i.e.

128 lower larval survival, longer larval development, lower body mass at

129 emergence, and a shorter adult lifespan).

130  4) Maternal host preference would predict offspring performance.

131 We predicted here that offspring would perform better on the host type

132 preferred by their mother when she laid her eggs.

135 **Methods**

Experiment overview. We estimated genetic and maternal effects for

four life-history traits and host egg laying preference using a half-sib/full-

sib split brood design with two different host types - original vs. novel. We

used cowpea (*Vigna unguiculata*, Fabaceae) as the original and mung bean

(*V. radiata*) as the novel host species in our experiment. Cowpea and mung

are both suitable hosts for *C. maculatus* (Fox and Messina, 2018) but

populations kept on specific hosts for many generations show better

performance on their usual host beans compared to novel hosts (Messina,

2004a).

Origin and maintenance of stock beetles. Beetles used as the

parental generation for our experiment were obtained from a large stock

population originally sourced from the University of Western Australia

(Perth, Australia) in 2017 where they had been bred on cowpea for at least

90 generations. We maintained this stock in our lab at the Australian

National University on cowpea for another 9 generations before we began

our experiment. Our stock was maintained in four, regularly mixed (every 5

generations) populations of over 500 individuals, each kept on cowpea at

25-26 °C.

Establishing the parental generation (P). To obtain virgin males and females for the parental generation (P), we exposed approximately 2000 un-infested cowpea beans to stock beetles for a period of 48 h. Each bean was then placed in an individual Eppendorf tube (with a pinhole for airflow) and had 5-8 eggs on its surface (a density which is usual for our stock). Once isolated, we monitored these beans until adults began to emerge. We collected virgin beetles each morning and used them in our experiment on the same day. We knew that beetles were virgins as they were either the only beetle to have emerged that day, or all beetles that had emerged were of the same sex. Every evening, we discarded any beetles that had emerged during that day. Using the parental beetles that emerged each morning, we mated males and females according to a full-sib/half-sib breeding design: each male was sequentially mated with four random females over the day and the mating order was noted (similar to Fox *et al.* (2003)). We aimed for a total of 89 families (sires) with 356 dams. Pairs that did not copulate within 30 min were separated for half an hour before being placed together for another mating attempt. Each female was weighed prior to mating to the nearest 0.001 mg (Cubis Ultra-Micro balance, Sartorius Lab Instruments GmbH., Goettingen, DEU). All matings took place over six days ('day mated', range 1-6, see *Partitioning of Phenotypic Variance*). There is therefore a positive correlation between

parental age (egg to adulthood) and 'day mated', but it is imperfect

because parents emerged from eggs initially laid over two days (i.e. 48

hours; see above). There is no correlation between 'day mated' and adult

parental age because all matings occurred on the day that the parents

emerged.

Maternal host preference. Once females had mated, we conducted

choice trials to determine their preferred host type for egg laying. We

considered females ready to lay eggs once the pair dismounted (Wilson

and Hill, 1989). For the host preference trials, we mixed cowpea (original

host) and mung beans (novel host) in covered Petri dish (ø 5.5 cm). We

used proportionally more mung than cowpea beans (8:4) because cowpea

beans have an approximately 1.6-times larger surface area (Paukku and

Kotiaho, 2008) and are 4-times heavier than mung beans (in our study the

average ± SD bean mass was 294±45 mg for cowpea, and 72±8 mg for

mung bean). Females were left to lay eggs for two hours, after which they

were removed and the number of eggs on each host bean type was

counted. Relative preference was calculated as the proportion of the eggs

that a female laid on the original host (cowpea). The values for relative

host preference therefore ranged between 0 and 1.

Generation of offspring (F1). Directly after the host preference trial,

we transferred individual dams to plastic containers (ø 4 cm, height 6 cm)

with 10-13 mung beans and left them to lay eggs for 14-18 h. We then

moved dams to new containers with 10-13 cowpea beans for 9-10 h. This

difference in laying time was required because initial trials showed that

more time was necessary for females to lay a sufficient number of eggs on

the novel host (mung beans), than on cowpea (the original host). Females

were presented with mung beans first to prevent them from laying all their

eggs on what we expected to be the preferred host type (cowpea)

(Messina and Slade, 1997). Maternal age and/or laying order effects are

unlikely to have influenced our results due to the short period of time we

allowed females to lay eggs (i.e. <24 h). Previous studies have shown no

effect of female age on offspring quality during the first 24 h of laying (Fox

and Dingle, 1994).

Once females had laid eggs on both host types, we collected up to 10

individual beans of each host type for each female. If a female had laid

eggs on fewer than 10 beans of a given type, we used them all. We ensured

that each bean had only one egg laid on it by haphazardly scraping off

surplus eggs with a scalpel. We weighed beans within 24 h after oviposition

to the nearest 0.001 mg to measure the resources available to the larvae.

Beans with an egg were then placed individually in Eppendorf tubes with

perforated lids and incubated at 26 °C.

We started regular monitoring of F1 emergence on day 24 post-

oviposition. However, we missed the onset of emergence of 40 beetles on

mung (1.2% of the total sample) by approximately 1 day as they emerged

sooner than expected. To be conservative, data from these early emerging

beetles was removed from the final dataset, because their emergence

could have been on that day or the day before. This uncertainty might not

only affect estimates of larval development duration, but also that of any

correlated traits such as body mass, as we did not provide adults with food.

Measurement of offspring (F1) traits. When an offspring emerged,

the date was recorded, and it was sexed and weighed (to the nearest 0.001

mg), and the bean was removed from the tube. After weighing, the beetle

was returned to its tube and checked daily for survival. We also tested the

host preference of two newly emerged female offspring per dam raised on

each host type (total N = 4/dam). To do this we mated each daughter with

a randomly selected male from the stock and then ran an egg laying

preference trial identical to that described above for their mothers. We

monitored survival until the death of all emerged offspring (day 79 post-

oviposition). At that time, we also censused larval (egg-to-emergence)

survival, assuming that larvae in beans from which a beetle had not yet

emerged had died.

Sample sizes. Our sample size was determined a priori based on

power analysis from Lynch and Walsh (1998) to create 89 families (89 sires

with 356 dams) and try to obtain 10 offspring per dam per host type (i.e. 7

120 offspring). The final sample size is lower due to unsuccessful matings

or too few eggs being laid. We analysed data for sons and daughters

separately due to their marked sexual dimorphism which invariably leads

to strong interactions between sex and other effects. We only included

offspring of dams that produced offspring reared on both host types, and

we removed offspring with extreme values (>3 standard deviations (SD)

from the mean). We collected data on four offspring life-history traits:

(egg-to-emergence) 'larval survival' (dead or emerged; N = 2 877),

'duration of larval development' (number of days between oviposition and

offspring emergence; N = 1 221 daughters and 1 189 sons), 'body mass'

(weight at emergence in mg; N = 1 228 daughters and 1 200 sons) and

'adult lifespan' (number of days between offspring emergence and death;

N = 1 226 daughters and 1 197 sons). In addition, we measured the host

egg laying preferences of 514 daughters. The slightly lower sample sizes

(<1%) for larval development or adult lifespan compared to body mass is

due to exclusion of response variable outliers and dams producing

offspring of certain sex on a single host type only. Prior to any statistical

analysis, but after all data has been collected, we registered this project on

the   Open   Science   Forum   webpage:

https://osf.io/ft7eq/?view_only=0bab0a33bb4246adb64c919601a72757

The final analytical approach we used was different, however, from the

original plan and we explain our changes in the Annotated registration

section in Supplementary Information.

Partitioning of phenotypic variance. To determine the strength of

maternal effects on offspring traits in *C. maculatus* (Prediction 1) and

whether maternal effects depend on the offspring rearing environment,

(i.e. the existence of a M×E interaction - Prediction 2), we built a 'minimal

model' containing only fixed effects directly attributable to the

experimental design. These were 'host type' (original/novel), 'bean mass'

(standardized within each host type to zero mean and 1 SD variance), 'dam

mating order' (i.e. dam position in sire's mating sequence, 1 to 4), and 'day

mated' (1-6). We did not fit 'day mated' as a random effect because it

resulted in overparametrized models that produced unstable and weak

variance estimates. As random effects, we used terms that reflected the

nested breeding design. We defined between-sire (family) variance as 'sire

ID', between-dam variance as 'dam ID' nested inside sire ID, and between-

individual variation as 'residual' variance.

We fitted a bivariate model for each life-history trait to allow the

components of offspring phenotypic variance to differ between the two

host types. It means that for each random factor we estimated two values

for variance in each model: one for offspring on cowpeas and the other for

offspring on mung beans. The bivariate model then allowed us to estimate

the covariance between offspring traits on the two host types. The

response variables for the bivariate model, therefore, consisted of two sub-

traits created from the single column of trait values. The first column (sub-

trait) contained trait values recorded for offspring raised on cowpea with

the values for those raised on mung bean left blank. The other column

contained values from offspring raised on mung bean and rows for cowpea

offspring were left blank.

Daughter host preference data was fitted differently, because we

wanted to account for variation in the total number of eggs laid and not

just fit the preference ratio (values 0-1) as a response. We again used a

bivariate model, with the number of eggs on cowpea in the first column of

the response, and the total number of eggs laid in the second column. To

estimate variance components separately for offspring from each host

type, we specified interactions with random effects: that is between host

type and sire (HOST:sire ID); between host type and dam (HOST:dam ID);

and between host type and the residual term.

Based on the outputs from the minimal models for each trait, we

then performed the phenotypic variance partitioning and calculated the

maternal-effects proportion (m^2) and the cross-environmental maternal-

effects correlation (M×E) (r_M), narrow-sense heritability (h^2), evolvability

(CV_A) and the cross-environmental correlation for additive genetic effects

(G×E) (r_G). We used maternal effects proportion, $m^2 = V_M/V_{total} = (V_{dam}-$

$V_{sire})/(V_{sire}+V_{dam}+V_{residual})$, to quantify the importance of maternal effects

(Messina & Fry, 2003). The correlation between maternal effects for the

two host types (M×E) was $r_M = COV_{M\ original-novel}/\sqrt{(V_{M\ original})}*\sqrt{(V_{M\ novel})}$,

where $COV_{M\ original-novel} = COV_{dam\ original-novel} - COV_{sire\ original-novel}$. We calculated

heritability as $h^2 = V_A/V_{total} = 4*V_{sire}/(V_{sire}+V_{dam}+V_{residual})$ (Lynch and Walsh,

1998), assuming dominance and epistatic effects were negligible.

Evolvability was defined by Houle (1992) as $CV_A = \sqrt{V_A}/\mu = \sqrt{(4*V_{sire})}/\mu$,

where μ is the mean of a trait (Garcia-Gonzalez *et al.*, 2012). Evolvability

standardizes additive genetic variance over the trait mean and is therefore

useful to compare the potential evolutionary response among traits, unlike

heritability which is conditional on the amount of residual phenotypic

variation (Hansen et al., 2011). The correlation between additive genetic

effects on the two host types (G×E) was $r_G = COV_{sire\ original-novel}/\sqrt{(V_{sire}}$

$_{original})*\sqrt{(V_{sire\ novel})}$.

Testing for host effect and dam host egg laying preference on

offspring phenotype. To determine the effect of novel host type on

offspring (Prediction 3) and to test whether offspring performed better on

the host type that was preferred by their mother (Prediction 4), we built a

'full model'. The full model always included all the terms outlined above for

the minimal model, as well as, 'dam host preference' (i.e. the relative

preference ratio: 0-1 values). We included the interaction between host

preference and host type 'HOST:dam host preference' to test if the effect

of the strength of the preference for the original host on offspring traits

differs depending on the host type. We also included interactions between

host type and bean mass 'HOST:bean mass', as well as host type and day

mated 'HOST:day mated' as fixed effects to test for potential host-specific

influences.

Model fitting. We fitted the models using the package MCMCglmm

(ver. 2.28) (Hadfield, 2010) in R software (ver. 3.5.3). The continuous

response variables - duration of larval development, body mass and adult

lifespan; were fitted using Gaussian family and identity link, while larval

survival (binomial distribution) was fitted as a categorical variable (with

implicit logit link). All continuous variables, including the response, were

standardized to zero mean and variance in units of SD (standardised $y = (y -$

$\mu)/SD$). Note that standardisation allows for direct among-trait comparison

by transforming response variable to SD scale (Matsumura *et al.*, 2012).

Daughter host preference was tested as bivariate vector with the number

of eggs on original host and total number of eggs using a binomial

distribution (specified as 'multinomial2' in the MCMCglmm) (Hadfield,

2010).

To fit the MCMC models, we used locally uninformative inverse-

Wishart priors for the random-effects variance (G) with expected variance-

covariance matrix at limit 'V' = 1 on the diagonal (i.e. two values – one per

each host), and the degree of freedom 'nu' = 1.002. We also added

parameter-expanded priors 'α mu' = 0 and 'α V' = 1 000 on diagonal to

increase mixing and effective sample size (Hadfield, 2010). Priors for

residual variance (R) were set as V = 1 on diagonal and nu = 1.002. In the

analysis of survival, the residual variance had to be fixed to 1, so that the

binomial mixed-effects model could be estimated (Hadfield, 2010). For

fixed effects, we retained default priors with variance = 0 and mean = 10^{10}.

To generate posterior distributions from the minimal model, we ran 120

000 iterations with burn-in of 20 000 and thinning interval of 100, so that

we obtained an effective sample size of 1 000. In the full model, we

increased number of iterations to 240 000, burn-in 40 000 and thinning

interval to 200 for the Gaussian traits. For daughter host preference, the

setup was 550 000 iterations with burn-in of 50 000 and 500 for the

thinning interval. For model of larval survival, we increased the iterations

to 1 200 000 with a burn-in of 200 000 and thinning interval of 1 000 as we

were aiming for an autocorrelation <0.1 at the lag corresponding to the

thinning interval (Wilson et al., 2010). Model convergence was assessed by

visual examination of the traces using function 'plot(mcmc.model)'.

Estimates of heritability in larval survival and daughter host egg laying

preference on the observed data scale were computed from the latent

scale variance estimates using the package 'QGglmm' (de Villemereuil *et*

al., 2016). We always interpreted the outcome of the complete model

(without simplification) and assessed the importance of individual model

terms based on their credible intervals (CrI).

Results

Prediction 1 and 2 – testing for maternal effects and M×E interaction. We observed very weak maternal effects for all the measured traits. Posterior estimates of variation due to maternal effects ($V_M = V_{dam} - V_{sire}$) ranged between -0.271-0.437 in the minimal models (Table 1). Note that negative values for variance are not meaningful, so the real maternal effects were close to zero. As a consequence, posterior estimates of maternal effects proportion (m^2) credible intervals did not exclude zero (Table 2). Given the low maternal effects whose estimates were even negative in some cases, it was not possible to calculate cross-environmental correlation of maternal effects (r_M), and to test for M×E interactions. Although we extracted the cross-environmental maternal effects covariance, to calculate r_M we also needed the product of the square-root of maternal effects variances (V_M) from the two host types as the denominator. A square-root is, however, not defined for negative values on real number scale.

Table 1. Variance-covariance structure and fixed-effects estimates based

on the minimal models.

Components of phenotypic variance related to additive genetic

effects - heritability, evolvability and G×E. We observed higher heritability

for 3 offspring life-history traits in the novel host, but evolvability was

similar for the two host types. We found medium to high heritability on the

novel host ($h^2 = 0.471$-0.621) for development duration of sons, body mass

of daughters and adult lifespan of both sexes (Table 2). The adult lifespan

of daughters also had medium heritability in offspring from the original

host type ($h^2 = 0.367$), but the credible interval was wide and came close to

including zero (Table 2). Despite a trend for higher heritability on the novel

host type, the estimates of evolvability (variance due to additive genetic

effects standardised over trait mean) were rather similar for the two host

types. Comparing evolvability among traits across sexes and host types, it

was highest for body mass at emergence ($CV_A = 8.4$-12.6 %, Table 2).

We recorded highly positive correlations of additive genetic effects

(r_G) for offspring performance on the two host types (Table 2). The only

exceptions were for the larval survival and development duration of sons,

which exhibited negative genetic correlation but with wide credible

intervals overlapping zero (Table 2). The within-family larval survival and

416   development duration of sons on the novel host did not correspond to that

417   on the original host, potentially suggesting a Genotype-by-Environment

418   interaction (G×E).

Table 2. Phenotypic variance partitioning.

Predictions 3 and 4 – testing for host type suitability and host

preference-performance. Unexpectedly, given our predictions, offspring

generally performed better on the novel host (mung) than on the original

host (cowpea) (Table 1, 3; see Supplementary Table 1 for the list of average

trait estimates for both host types). We did not find a positive relationship

between the strength of the dam host egg laying preference and the

performance of their offspring on either host type. The one exception was

for daughter body mass: the daughters of dams that more strongly

preferred the cowpea host emerged heavier when raised on cowpea,

whereas we observed no such effect for daughters raised on mung beans

(Table 3). Daughters reared on the original cowpea host also exhibited a

higher egg laying preference for cowpeas than those reared on mung

beans (raw average host preference ratio of 0.677 versus 0.516; where 1 is

a strict preference for cowpeas).

Effect of experimental variables on offspring traits - bean size,

mating order and mating date. Host-type specific relative bean size (zero-

centered per host type) and mating date were important predictors of

offspring traits. For both host types, offspring that developed in larger

beans had a poorer outcome: increased development duration for both

sexes, reduced body mass in daughters and lower adult lifespan in sons.

The effects of mating day were, however, mixed (Table 3). Daughters of

later mated families took longer to develop, but sons emerged at a larger

size, and both sexes subsequently lived longer. Daughters from later mated

families also showed a stronger egg laying preference for the novel host

type.

Table 3. Fixed-effects estimates from the full models.

Discussion

Summary. Our main aim was to test for environment-specific

maternal effects on offspring life-history traits, that is Maternal-by-

Environment interactions (M×E). We used a full-sib/half-sib split brood

design with seed beetles (*C. maculatus*) reared on two types of host:

original – cowpea; and novel – mung bean. Maternal effects proved to be

negligible on both host types, making it meaningless to estimate maternal

effect correlations across the two hosts and thereby test for M×E

interactions. Instead, we found that additive genetic variance played an

important role, particularly in the novel environment. Contrary to our

predictions, developing on the novel host type did not lower offspring

performance for any of the measured traits. On the original host type,

daughters whose mothers preferred to lay eggs on this host were heavier

than those of mothers that preferred the novel host type. Daughters

reared on the original host also preferred to lay eggs on this host, while

those from the novel host did not exhibit a preference. Below we discuss

our findings in more detail.

Missing maternal effects. We were unable to formally test for a

difference in the relative importance of maternal effects in different

environments, because maternal effects were very weak in both host

types. Intriguingly, previous studies on *C. maculatus* have reported strong

maternal effects on male development duration (Fox, 1993a), female

fecundity, adult body mass and lifespan (Messina and Fry, 2003). However,

these non-genetic maternal effects in experimental studies with *C.*

maculatus mostly seem to stem from variation in maternal age (Fox and

Dingle, 1994; Fox *et al.*, 2003; Lind *et al.*, 2015). Older *C. maculatus* females

have previously been shown to produce offspring with lower egg-to-adult

survival, longer larval development (Fox *et al.*, 2003; Lind *et al.*, 2015) and a

shorter lifespan (Lind *et al.*, 2015). In our study, neither male nor female

adult parental age varied as we collected parental beetles and mated them

on the day of emergence. While mating day, which is imperfectly

correlated with lifetime age (i.e. zygote to emergence), had notable effects

on several offspring life-history traits and daughters' host egg laying

preference, removing it from the model did not increase estimates of

maternal effect proportions (Supplementary Table 2). This suggests that

mating day acts like a 'block' factor that accounts for experimental

variation that affected the entire family sired by a given sire (and not to

lifetime age *per se*) due to stochastic daily fluctuations of unknown origin.

Non-genetic maternal effects due to variation in age were therefore

eliminated by mating dams of the same age to the single sire in each

family.

Heritability of some life-history traits was higher in the novel

environment, but evolvability was similar between hosts and the genetic

correlations between hosts were highly positive. Heritability of fitness-

related traits is traditionally assumed to be low as additive genetic

variation is expected to be depleted by natural selection (Mousseau and

Roff, 1987; Teplitsky *et al.*, 2009). Here, we recorded considerably higher

heritability for a number of life-history traits in the novel host environment

(mung beans), especially son's development duration, daughter's body

mass and adult lifespan. Higher heritability can be the result of higher

additive genetic variance, reduced residual variance, or both, because it is

defined by a ratio. The traits with non-zero heritability in our current study

showed increased additive genetic variance rather than reduced residual

variance in the novel environment. This suggests there was larger standing

genetic variation for these traits in the novel environment. However,

similar values of trait evolvability, the additive genetic variance

standardised over trait mean, indicated that the traits are likely to respond

at a similar rate to selection on both host types. We also found strong

positive genetic correlations indicating parallel reaction norms for offspring

from different families when reared on the two host types. It is important

to bear in mind, however, that quantitative genetic studies are ultimately

inconclusive about whether or not there is genetic trade-off in adaptation

to different hosts (Gompert and Messina, 2016) and any results based on

inspection of genetic correlation should be viewed with caution.

The novel host type appeared to be better for offspring. Offspring

life-history traits differed between the two host types in an unexpected

way with offspring performing better on the novel host type. While

offspring survival was almost 90 % on cowpeas (the original host), it was

even higher on the novel host (mung bean), exceeding 90 %. There was no

obvious cost as the duration of larval development was shorter on mung

beans, and the beetles that emerged were larger and lived longer as adults.

It is not straightforward, however, to tell which host species is generally

better for *C. maculatus*, because it depends on other factors such as the

level of competition (Messina, 2004a; Fox and Messina, 2018),

temperature (Stillwell *et al.*, 2007) and the population's evolutionary

history (Messina, 2004a).

In the present study, we eliminated any effect of larval competition

by rearing one larva per bean. We therefore reduced the fitness advantage

that cowpeas provide due to their larger size when there is larval

competition (Fox and Savalli, 1998; Messina, 2004a; Fox and Messina,

2018). In our experiment, the original host type was cowpeas. Although

adult lifespan on mung beans can be longer than that on cowpea (Fox,

Bush, *et al.*, 2004), cowpea strains are overall more viable than those from

mung beans (Messina, 2004a). Currently, we still lack a good explanation

for the observed positive effect of a novel host type on the measured life-

history traits, but we offer some speculative suggestions.

First, the seemingly higher performance of *C. maculatus* on the novel

host type could be caused by female avoidance of this host. We only

analysed data from offspring of dams that laid eggs on both hosts. Our

interpretation of the data would change, if the offspring of dams that only

lay eggs on cowpea are of inferior quality. In total about 10 % of dams (36

in total) produced offspring on one host only, and of these 21 dams

showed strict cowpea preference in the host egg-laying preference trials.

There is therefore a possibility that the dams producing lower quality

offspring avoided mung as a host. Unfortunately, including the data on

offspring from the excluded dams will not resolve the issue, because we do

not know how their offspring would perform on mung beans.

Second, we note that the lower performance of offspring on cowpea

might have been because, following the preference trial, females were

provided with mung beans to lay eggs, and only after that with cowpea.

Eggs laid late in the laying sequence have previously been shown to be of

lower quality in some insects (Rossiter, 1996). The lifetime fecundity of *C.*

maculatus is 60-80 eggs and depends on female body size, and mate and

host availability (Mitchell, 1975; Wilson and Hill, 1989; Messina and Fry,

2003). The dams in our experiment laid around 30 eggs per 24 h and were

therefore probably still far from nearing the end of egg laying in our

experiment. In addition, past studies suggest there is no negative effect of

maternal age on offspring performance on the first day of female

reproductive life (Fox and Dingle, 1994). We therefore suggest that the

sequence of host type presentation for egg laying is very unlikely to explain

poorer offspring performance on cowpeas.

Host preference of *C. maculatus* was only weakly related to

offspring performance on each host species. We tested two relationships

with regard to female host egg laying preference – whether dam host

preference correlated with offspring performance on each host, and if

rearing host type affected a daughter's host preference. Host egg laying

preference often predicts offspring performance on that host in

phytophagous insects (Gripenberg *et al.*, 2010). However, despite the

critical role of oviposition choice in *C. maculatus*, for all the traits we

measured, the strength of the dam preference for cowpea only correlated

with daughter body mass. Specifically, dams that more strongly preferred

cowpea produced heavier daughters on cowpea but not on mung.

Previous studies show that strains of *C. maculatus* maintained for >

40 generations on mung bean or cowpea exhibit a strong preference for

their recent host type (Messina, 2004b). (Fox, Stillwell, *et al.*, 2004) used

different host strains to demonstrate that there is no immediate effect of

the current host type (i.e. switching a strain to another host for 1

generation) on the host preference. Variation in host preference in *C.*

maculatus is mainly due to additive genetic effects, but is also influenced

by dominance and epistasis (Fox, 1993b; Fox, Stillwell, *et al.*, 2004).

However, we found that while daughters reared on cowpea preferred

cowpea over a novel host (mung bean), those reared on mung beans

showed no preference for either host type. This suggests that there might

be a G×E interaction affecting host egg laying preferences and more

generations would be necessary to see whether the host preference

evolves after the host switch (as in Fox, Stillwell, *et al.* (2004) or Messina

(2004b)).

Conclusion. Maternal effects were low for all measured traits in our study

for offspring reared on either a novel or familiar host. We therefore did not

test for Maternal-by-Environment interactions. A next step in the

exploration of the role of M×E interaction on estimates of genetic variation

and G×E interactions in *C. maculatus* would be to run tests using a less

favourable novel host as the species we used are both of high quality to *C.*

maculatus (Gompert *et al.*, 2015). By creating more challenging conditions

for offspring, maternal effects might emerge as an important factor

affecting variation in offspring performance (Fox *et al.*, 1997; Räsänen and

Kruuk, 2007). Despite being unable to test for an interaction between

maternal effects and environment (M×E), we still provided a way to

formally test for it. Studies partitioning phenotypic variation across

environments should not only account for variation in performance of

different genotypes between environments (G×E), which is a common

practice, but also control for potential M×E.

Acknowledgements

We thank Timothée Bonnet and Loeske E. B. Kruuk (RSB ANU) for

very helpful discussions about phenotypic variation partitioning. We are

thankful to Carolina Dolan, Krish S. Sanghvi and Elroy Kwan Au (RSB ANU)

for their help with the experiment and collecting data. MV and PJCC both

received 6 months Endeavour fellowship from Australian Government.

Author contribution

MV, PJCC, MLH and MDJ conceived and designed the study; MV, PJCC

and MIC collected data; MV analysed the data and drafted the manuscript;

and PJCC, MIC, MLH and MDJ revised it. All authors approved the final

version of the manuscript.

Conflict of interest

621 We declare that we do not have any conflict of interest.

Data archiving

624 Data with accompanying R code are stored at OSF (link:

625 https://osf.io/ft7eq/?view_only=0bab0a33bb4246adb64c919601a72757)

626 and FigShare (link: https://figshare.com/s/7878e68e31590dce594d).

Supplementary Information

629 Supplementary information is available at *bioRxiv* website along the

630 manuscript file.

References

Agrawal AA, Conner JK, Rasmann S (2010). Tradeoffs and negative

correlations in evolutionary ecology. In: *Evolution since Darwin: the*

first 150 years, Sinauer, pp 243–268.

Van Asch M, Julkunen-Tiito R, Visser ME (2010). Maternal effects in an

insect herbivore as a mechanism to adapt to host plant phenology.

Funct Ecol **24**: 1103–1109.

Bernardo J (1996). Maternal effects in animal ecology. *Am Zool* **36**: 83–105.

Cahenzli F, Erhardt A (2013). Transgenerational acclimatization in an

herbivore-host plant relationship. *Proc R Soc B Biol Sci* **280**.

Cahenzli F, Wenk BA, Erhardt A, Rosenheim JA (2015). Female butterflies

adapt and allocate their progeny to the host-plant quality of their own

larval experience. *Ecology* **96**: 1966–1973.

Forbes AA, Devine SN, Hippee AC, Tvedte ES, Ward AKG, Widmayer HA, *et*

al. (2017). Revisiting the particular role of host shifts in initiating insect

speciation. *Evolution (N Y)* **71**: 1126–1137.

Fox CW (1993a). Maternal and genetic influences on egg size and larval

performance in a seed beetle. *Heredity* **73**: 509–517.

Fox CW (1993b). A quantitative genetic analysis of oviposition preference

and larval performance on two hosts in the bruchid beetle,

Callosobruchus maculatus. Evolution **47**: 166–175.

Fox CW, Bush ML, Roff DA, Wallin WG (2004). Evolutionary genetics of

lifespan and mortality rates in two populations of the seed beetle,

Callosobruchus maculatus. Heredity **92**: 170–181.

Fox CW, Bush ML, Wallin WG (2003). Maternal age affects offspring

lifespan of the seed beetle. *Funct Ecol* **17**: 811–820.

Fox CW, Dingle H (1994). Dietary mediation of maternal age effects on

offspring performance in a seed beetle (Coleoptera: Bruchidae). *Funct*

Ecol **8**: 600–606.

Fox CW, Messina FJ (2018). Evolution of larval competitiveness and

associated life-history traits in response to host shifts in a seed beetle.

J Evol Biol **31**: 302–313.

Fox CW, Savalli UM (1998). Inheritance of environmental variation in body

size: Superparasitism of seeds affects progeny and grandprogeny body

size via a nongenetic maternal effect. *Evolution* **52**: 172.

Fox CW, Stillwell RC, Amarillo-Suárez AR, Czesak ME, Messina FJ (2004).

Genetic architecture of population differences in oviposition

behaviour of the seed beetle Callosobruchus maculatus. *J Evol Biol* **17**:

1141–1151.

Fox CW, Thakar MS, Mousseau TA (1997). Egg size plasticity in a seed

beetle: an adaptive maternal effect. *Am Nat* **149**: 149–163.

Garcia-Gonzalez F, Simmons LW, Tomkins JL, Kotiaho JS, Evans JP (2012).

Comparing evolvabilities: Common errors surrounding the calculation

and use of coefficients of additive genetic variation. *Evolution* **66**:

2341–2349.

Gompert Z, Jahner JP, Scholl CF, Wilson JS, Lucas LK, Soria-Carrasco V, *et al.*

(2015). The evolution of novel host use is unlikely to be constrained by

trade-offs or a lack of genetic variation. *Mol Ecol* **24**: 2777–2793.

Gompert Z, Messina FJ (2016). Genomic evidence that resource-based

trade-offs limit host-range expansion in a seed beetle. *Evolution* **70**:

1249–1264.

Gripenberg S, Mayhew PJ, Parnell M, Roslin T (2010). A meta-analysis of

preference-performance relationships in phytophagous insects. *Ecol*

Lett **13**: 383–393.

Guntrip J, Sibly RM, Holloway GJ (1997). The effect of novel environment

and sex on the additive genetic variation and covariation in and

between emergence body weight and development period in the

cowpea weevil, *Callosobruchus maculatus* (Coleoptera, Bruchidae).

Heredity **78**: 158–165.

Hadfield JD (2010). MCMC methods for multi-response generalized linear

mixed models: The MCMCglmm R package. *J Stat Softw* **33**.

Houle D (1992). Comparing evolvability and variability of quantitative traits.

Genetics **130**: 195–204.

Joshi A, Thompson JN (1995). Trade-offs and the evolution of host

specialization. *Evol Ecol* **9**: 82–92.

697 Kruuk LEB, Hadfield JD (2007). How to separate genetic and environmental

698   causes of similarity between relatives. *J Evol Biol* **20**: 1890–1903.

699 Lind MI, Berg EC, Alavioon G, Maklakov AA (2015). Evolution of differential

700   maternal age effects on male and female offspring development and

701   longevity. *Funct Ecol* **29**: 104–110.

702 Lynch M, Walsh B (1998). *Genetics and Analysis of Quantitative Traits*.

703   Sinauer, pp. 980.

704 Magalhães S, Blanchet E, Egas M, Olivieri I (2011). Environmental effects on

705   the detection of adaptation. *J Evol Biol* **24**: 2653–2662.

706 Matsumura S, Arlinghaus R, Dieckmann U (2012). Standardizing selection

707   strengths to study selection in the wild: A critical comparison and

708   suggestions for the future. *Bioscience* **62**: 1039–1054.

709 Messina FJ (2004a). Predictable modification of body size and competitive

710   ability. *Evolution* **58**: 2788–2797.

711 Messina FJ (2004b). How labile are the egg-laying preferences of seed

712   beetles? *Ecol Entomol* **29**: 318–326.

713 Messina FJ, Fry JD (2003). Environment-dependent reversal of a life history

714   trade-off in the seed beetle *Callosobruchus maculatus*. *J Evol Biol* **16**:

715   501–509.

Messina FJ, Lish AM, Gompert Z (2018). Variable responses to novel hosts

by populations of the seed beetle *Callosobruchus maculatus*

(Coleoptera: Chrysomelidae: Bruchinae). *Environ Entomol* **47**: 1194–

1202.

Messina FJ, Morrey JL, Mendenhall M (2007). Why do host-deprived seed

beetles 'dump' their eggs? *Physiol Entomol* **32**: 259–267.

Messina FJ, Slade AF (1997). Inheritance of host-plant choice in the seed

beetle *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Ann*

Entomol Soc Am **90**: 848–855.

Mitchell R (1975). The evolution of oviposition tactics in the bean weevil,

Callosobruchus maculatus (F.). *Ecology* **56**: 696–702.

Mousseau TA, Roff DA (1987). Natural selection and the heritability of

fitness components. *Heredity* **59**: 181–197.

Paukku S, Kotiaho JS (2008). Female oviposition decisions and their impact

on progeny life-history traits. *J Insect Behav* **21**: 505–520.

Price TN, Leonard A, Lancaster LT (2017). Warp-speed adaptation to novel

hosts after 300 generations of enforced dietary specialisation in the

seed beetle *Callosobruchus maculatus* (Coleoptera: Chrysomelidae:

Bruchinae). *Eur J Entomol* **114**: 257–266.

Räsänen K, Kruuk LEB (2007). Maternal effects and evolution at ecological

time-scales. *Funct Ecol* **21**: 408–421.

Rossiter M (1996). Incidence and consequences of inherited environmental

effects. *Annu Rev Ecol Syst* **27**: 451–476.

Saltz JB, Bell AM, Flint J, Gomulkiewicz R, Hughes KA, Keagy J (2018). Why

does the magnitude of genotype-by-environment interaction vary?

Ecol Evol **8**: 6342–6353.

Stillwell RC, Wallin WG, Hitchcock LJ, Fox CW (2007). Phenotypic plasticity

in a complex world: Interactive effects of food and temperature on

fitness components of a seed beetle. *Oecologia* **153**: 309–321.

Teplitsky C, Mills JA, Yarrall JW, Merilä J (2009). Heritability of fitness

components in a wild bird population. *Evolution* **63**: 716–726.

Vega-Trejo R, Head ML, Jennions MD, Kruuk LEB (2018). Maternal-by-

environment but not genotype-by-environment interactions in a fish

without parental care. *Heredity* **120**: 154–167.

Via S, Lande R (1985). Genotype-environment interaction and the evolution

of phenotypic plasticity. *Evolution* **39**: 505–522.

de Villemereuil P, Schielzeth H, Nakagawa S, Morrissey MB (2016). General

methods for evolutionary quantitative genetic inference from

generalized mixed models. *Genetics* **204**: 1281–1294.

Wilson K, Hill L (1989). Factors affecting egg maturation in the bean weevil

Callosobruchus maculatus. Physiol Entomol **14**: 115–126.

Tables

Table 1. Variance-covariance structure and main effects estimates

based on the minimal models. Phenotypic variance decomposition based

on these estimates is presented in Table 2. (A) gives model summary for

larval survival and life history in male offspring (sons); (B) is for life history

of female offspring (daughters) and their host preference. Estimates whose

credible interval does not overlap zero are emphasised in bold. Note that

residual variance for both host types in the binomial model of larval

survival was set to 1 by default. The host type effect needs to be added to

the intercept (value for original host type) to obtain an estimate for the

effect in the novel host type.

Table 2. Phenotypic variance partitioning. Estimates of maternal

effects proportion (m^2), heritability (h^2), cross-environmental correlations

(r_M and r_G) and evolvability (CV_A) in sons (A) and daughters (B). Heritability

and evolvability estimates for larval survival and daughter host preference

(asterisk) were recalculated to the observed data scale using the package

'QGglmm' (de Villemereuil *et al.,* 2016). Bold emphasis is put on the

estimates with credible intervals not overlapping zero.

Table 3. Fixed-effects estimates from the full models. (A) gives larval

survival and life-history traits in male offspring (sons); (B) is for life history

of female offspring and their host preference. Bold emphasis is put on

estimates with credible intervals not overlapping zero. The host type effect

needs to be added to the intercept (value for original host type) to obtain

an estimate for the effect in the novel host type. The same applies to the

host-specific interactions.

Table 1. Variance-covariance structure and fixed-effects estimates based on the minimal models.

785   (A)

| | Trait | Larval survival | | Duration of larval development | | Adult body mass | | Adult lifespan | |
| | Sex (Sample size) | juveniles (N = 2877) | | sons (N = 1189) | | sons (N = 1200) | | sons (N = 1195) | |
	Parameter	Estimate	CrI	Estimate	CrI	Estimate	CrI	Estimate	CrI
Variance-covariance	V_{sire} orig	0.002	(0.000,0.361)	0.001	(0.000,0.101)	0.001	(0.000,0.103)	0.053	(0.000,0.120)
	V_{dam} orig	0.403	(0.000,0.946)	0.107	(0.000,0.207)	**0.129**	(0.044,0.221)	**0.110**	(0.037,0.205)
	V_{res} orig	1	-	**0.941**	(0.857,1.136)	**0.855**	(0.754,0.968)	**0.736**	(0.687,0.886)
	V_{sire} novel	0.004	(0.000,1.083)	**0.081**	(0.008,0.131)	0.001	(0.000,0.110)	**0.107**	(0.022,0.226)
	V_{dam} novel	0.005	(0.000,0.654)	**0.057**	(0.004,0.127)	**0.138**	(0.075,0.238)	**0.138**	(0.051,0.258)
	V_{res} novel	1	-	**0.420**	(0.367,0.482)	**0.500**	(0.438,0.575)	**0.642**	(0.570,0.741)
	COV_{sire} orig-novel	0.000	(-0.179,0.266)	-0.002	(-0.040,0.032)	0.000	(-0.008,0.083)	0.047	(-0.002,0.130)
	COV_{dam} orig-novel	-0.001	(-0.140,0.425)	0.017	(-0.025,0.073)	**0.096**	(0.043,0.181)	**0.090**	(0.035,0.178)
Main effects	Intercept (orig HOST)	**2.567**	(2.203,3.083)	**0.302**	(0.160,0.476)	**-0.449**	(-0.602,-0.228)	**-0.351**	(-0.543,-0.17)
	novel HOST	**0.502**	(0.133,0.945)	**-0.814**	(-0.920,-0.659)	**0.665**	(0.556,0.785)	**0.496**	(0.355,0.594)
	bean mass	**-0.193**	(-0.303,-0.013)	**0.086**	(0.026,0.121)	-0.022	(-0.078,0.019)	**-0.080**	(-0.127,-0.022)
	mating order	0.026	(-0.186,0.139)	0.042	(-0.015,0.102)	0.041	(-0.033,0.112)	0.051	(-0.012,0.133)
	day mated	0.088	(-0.091,0.282)	**0.167**	(0.097,0.240)	**0.119**	(0.062,0.219)	**0.110**	(0.010,0.193)

788   (B)

	Trait	Host preference		Duration of larval development		Adult body mass		Adult lifespan	
	Sex (sample size)	daughters (N = 514)		daughters (N = 1221)		daughters (N = 1228)		daughters (N = 1224)	
	Parameter	Estimate	CrI	Estimate	CrI	Estimate	CrI	Estimate	CrI
Variance-covariance	V_{sire} orig	0.000	(0.000,0.006)	0.001	(0.000,0.082)	0.001	(0.000,0.122)	**0.082**	(0.012,0.186)
	V_{dam} orig	0.000	(0.000,0.005)	**0.092**	(0.016,0.197)	**0.133**	(0.077,0.275)	**0.104**	(0.038,0.229)
	V_{res} orig	**0.034**	(0.023,0.048)	**0.799**	(0.713,0.931)	**0.674**	(0.596,0.782)	**0.721**	(0.656,0.865)
	V_{sire} novel	0.000	(0.000,0.008)	0.018	(0.000,0.075)	**0.088**	(0.004,0.162)	**0.116**	(0.047,0.233)
	V_{dam} novel	0.000	(0.000,0.006)	**0.078**	(0.025,0.143)	**0.117**	(0.070,0.229)	**0.093**	(0.011,0.166)
	V_{res} novel	**0.033**	(0.024,0.055)	**0.397**	(0.351,0.459)	**0.440**	(0.384,0.492)	**0.660**	(0.589,0.767)
	COV_{sire} orig-novel	0.000	(-0.002,0.003)	0.000	(-0.011,0.045)	0.035	(-0.004,0.110)	**0.077**	(0.015,0.160)
	COV_{dam} orig-novel	0.000	(-0.001,0.002)	0.037	(-0.004,0.093)	**0.146**	(0.069,0.205)	**0.064**	(0.012,0.142)
Main effects	Intercept (orig HOST)	**-0.426**	(-0.567,-0.329)	**0.428**	(0.306,0.614)	**-0.582**	(-0.757,-0.377)	**-0.215**	(-0.404,-0.043)
	novel HOST	**-0.254**	(-0.339,-0.172)	**-1.001**	(-1.104,-0.876)	**0.988**	(0.862,1.073)	**0.537**	(0.433,0.671)
	bean mass	0.007	(-0.036,0.049)	**0.083**	(0.036,0.121)	**-0.079**	(-0.119,-0.034)	-0.021	(-0.070,0.034)
	mating order	0.016	(-0.025,0.059)	0.020	(-0.012,0.101)	0.032	(-0.054,0.087)	-0.045	(-0.112,0.026)
	day mated	**-0.059**	(-0.099,-0.007)	**0.242**	(0.168,0.298)	0.088	(-0.004,0.176)	0.049	(-0.033,0.171)

791   **Table 2.** Phenotypic variance partitioning.

792   (A)

| Trait | Larval survival | | | Duration of larval development | | Adult body mass | | Adult lifespan | |
| Sex | juveniles | | | sons | | sons | | sons | |
Parameter	Estimate	CrI	Estimate*	Estimate	CrI	Estimate	CrI	Estimate	CrI
V_M orig	0.437	(-0.237,0.970)	-	0.088	(-0.086,0.212)	0.110	(-0.046,0.214)	0.062	(-0.065,0.185)
m^2 orig	0.158	(-0.144,0.500)	-	0.079	(-0.070,0.191)	0.114	(-0.046,0.199)	0.064	(-0.065,0.193)
V_M novel	-0.271	(-1.161,0.549)	-	-0.016	(-0.097,0.115)	0.084	(-0.008,0.243)	0.043	(-0.143,0.228)
m^2 novel	-0.303	(-0.523,0.314)	-	0.002	(-0.180,0.193)	0.203	(-0.008,0.326)	0.046	(-0.160,0.236)
r_M	-	-	-	-	-	-	-	-	-
h^2 orig	0.006	(0.000,0.862)	0.029	0.001	(0.000,0.364)	0.002	(0.000,0.394)	0.215	(0.000,0.484)
h^2 novel	1.431	(0.000,1.962)	0.095	**0.472**	(0.116,0.938)	0.004	(0.000,0.593)	**0.471**	(0.072,0.922)
r_G	-0.499	(-0.961,0.850)	-	-0.228	(-0.935,0.607)	0.977	(-0.458,0.999)	**0.989**	(0.468,0.999)
CV_A orig	-	-	-	0.008	(0.000,0.020)	**0.087**	(0.001,0.183)	**0.027**	(0.008,0.044)
CV_A novel	-	-	-	**0.019**	(0.009,0.025)	**0.126**	(0.003,0.171)	**0.036**	(0.016,0.050)

(B)

Trait	Host preference			Duration of larval development		Adult body mass		Adult lifespan	
Sex	daughters			daughters		daughters		daughters	
Parameter	Estimate	CrI	Estimate*	Estimate	CrI	Estimate	CrI	Estimate	CrI
V_M orig	0.000	(-0.005,0.007)	-	0.090	(-0.042,0.200)	0.095	(-0.012,0.281)	0.030	(-0.120,0.199)
m^2 orig	0.000	(-0.140,0.169)	-	0.087	(-0.046,0.207)	0.142	(-0.014,0.292)	0.031	(-0.118,0.201)
V_M novel	0.000	(-0.009,0.006)	-	0.037	(-0.031,0.138)	0.079	(-0.071,0.209)	-0.090	(-0.193,0.102)
m^2 novel	0.001	(-0.161,0.159)	-	0.070	(-0.064,0.253)	0.081	(-0.121,0.289)	-0.119	(-0.214,0.093)
r_M	-	-	-	-	-	-	-	-	-
h^2 orig	0.004	(0.000,0.572)	0.004	0.002	(0.000,0.344)	0.003	(0.000,0.518)	**0.367**	(0.017,0.700)
h^2 novel	0.005	(0.000,0.644)	0.004	0.003	(0.000,0.550)	**0.482**	(0.068,0.947)	**0.621**	(0.247,0.980)
r_G	0.832	(-0.872,0.999)	-	0.727	(-0.727,0.999)	**0.937**	(0.004,0.999)	**0.876**	(0.394,0.999)
CV_A orig	-	-	-	0.008	(0.000,0.018)	**0.084**	(0.002,0.121)	**0.025**	(0.013,0.040)
CV_A novel	-	-	-	**0.012**	(0.002,0.019)	**0.091**	(0.044,0.134)	**0.032**	(0.018,0.038)

799 **Table 3.** Fixed-effects estimates from the full models.

800 (A)

Trait	Larval survival		Duration of larval development		Adult body mass		Adult lifespan	
Sex (Sample size)	juveniles (N = 2877)		sons (N = 1189)		sons (N = 1200)		sons (N = 1197)	
Parameter	Estimate	CrI	Estimate	CrI	Estimate	CrI	Estimate	CrI
Host								
Intercept (orig HOST)	**2.542**	(2.259,3.114)	**0.261**	(0.145,0.474)	**-0.392**	(-0.602,-0.244)	**-0.405**	(-0.548,-0.182)
novel HOST	**0.554**	(0.101,0.920)	**-0.810**	(-0.925,-0.646)	**0.653**	(0.570,0.783)	**0.483**	(0.351,0.580)
bean mass (orig)	-0.177	(-0.327,0.045)	**0.094**	(0.007,0.170)	-0.060	(-0.146,0.009)	**-0.090**	(-0.160,-0.010)
Interaction: bean mass (novel)	-0.102	(-0.376,0.197)	-0.026	(-0.111,0.089)	0.029	(-0.031,0.157)	0.063	(-0.083,0.122)
Dam properties								
dam mating order	-0.012	(-0.178,0.153)	0.043	(-0.016,0.098)	0.046	(-0.035,0.111)	0.060	(-0.017,0.128)
day mated	0.164	(-0.092,0.370)	0.097	(-0.005,0.200)	**0.131**	(0.044,0.248)	**0.126**	(0.012,0.220)
Interaction: day mated (novel)	-0.198	(-0.513,0.212)	0.081	(-0.028,0.233)	-0.017	(-0.128,0.081)	-0.034	(-0.154,0.072)
dam preference for cowpea (orig)	-0.043	(-0.219,0.216)	-0.031	(-0.136,0.057)	0.012	(-0.104,0.087)	0.006	(-0.099,0.082)
Interaction: dam preference (novel)	0.105	(-0.179,0.463)	0.051	(-0.071,0.163)	0.045	(-0.069,0.137)	0.008	(-0.115,0.097)

803   (B)

| Trait | Host preference | | Duration of larval development | | Adult body mass | | Adult lifespan | |
| Sex (Sample size) | daughters (N = 514) | | daughters (N = 1221) | | daughters (N = 1228) | | daughters (N = 1226) | |
Parameter	Estimate	CrI	Estimate	CrI	Estimate	CrI	Estimate	CrI
Host								
Intercept (orig HOST)	**-0.457**	(-0.553,-0.337)	**0.412**	(0.277,0.600)	**-0.558**	(-0.737,-0.369)	**-0.188**	(-0.409,-0.042)
novel HOST	**-0.228**	(-0.333,-0.166)	**-0.983**	(-1.094,-0.868)	**0.981**	(0.863,1.074)	**0.571**	(0.429,0.679)
bean mass (orig)	-0.004	(-0.054,0.062)	**0.111**	(0.053,0.203)	**-0.125**	(-0.198,-0.055)	-0.073	(-0.143,0.002)
Interaction: bean mass (novel)	0.013	(-0.074,0.088)	-0.044	(-0.167,0.024)	0.103	(-0.008,0.171)	0.047	(-0.018,0.176)
Dam properties								
dam mating order	0.017	(-0.026,0.057)	0.036	(-0.017,0.102)	0.025	(-0.049,0.090)	-0.033	(-0.116,0.022)
day mated	**-0.059**[#]	(-0.122,0.000)	**0.255**	(0.148,0.352)	**0.127**	(0.019,0.240)	0.126	(-0.001,0.220)
Interaction: day mated (novel)	0.017	(-0.077,0.091)	-0.011	(-0.127,0.107)	-0.085	(-0.197,0.020)	-0.074	(-0.220,0.029)
dam preference for cowpea (orig)	-0.001	(-0.051,0.064)	-0.029	(-0.125,0.048)	**0.103**	(0.012,0.203)	-0.014	(-0.110,0.078)
Interaction: dam preference (novel)	-0.022	(-0.110,0.059)	0.029	(-0.062,0.143)	**-0.096**	(-0.196,-0.009)	-0.003	(-0.132,0.085)

804   [#] marginally important (upper limit of the Credible Interval = -0.0003839)