The role of maternal effects on offspring performance in familiar and novel environments

Authors: Milan Vrtílek[1]*, Pierre J. C. Chuard[2]*, Maider Iglesias-Carrasco[3], Michael D. Jennions[3] and Megan L. Head[3]

Affiliations:

[1]The Czech Academy of Sciences, Institute of Vertebrate Biology, Brno, Czech Republic

[2]Department of Biological Sciences, Bishop's University, Sherbrooke, Canada

[3]Research School of Biology, The Australian National University, Canberra, Australia

* These authors share joint first authorship on this work

Correspondence: Milan Vrtílek, The Czech Academy of Sciences, Institute of Vertebrate Biology, Květná 8, 603 65 Brno, Czech Republic; phone: +420 543 422 557, fax: +420 543 211 346, e-mail: vrtilek@ivb.cz

Conflict of interest: We declare that we do not have any conflict of interest.

Running title: Environment-specific maternal effects

Version: November 26, 2020

Figures: 1

Tables: 4 (+1 Supplementary Table)

Word count: 6 736 words

Abstract

Maternal effects are an important evolutionary force that may either facilitate adaptation to a new environment or buffer against unfavourable conditions. The degree of variation in traits expressed by siblings from different mothers is frequently sensitive to environmental conditions. This could generate a Maternal-by-Environment interaction (M×E) that could inflate estimates of Genotype-by-Environment effects (G×E). We aimed to test for environment-specific maternal effects (M×E) using a paternal full-sib/half-sib breeding design in the seed beetle *Callosobruchus maculatus*, where we split and reared offspring from the same mother on two different bean host types – original and novel. Our quantitative genetic analysis indicated that maternal effects were very small on both host types for all the measured life-history traits. There was also little evidence that maternal oviposition preference for a particular host type predicted her offspring's performance on that host. Further, additive genetic variance for most traits was relatively high on both hosts. While there was higher heritability for offspring reared in the novel host, there was no evidence for G×Es, and most cross-host genetic correlations were positive. This suggests that offspring from the same family ranked similarly for performance on both host types. Our results point to a genetic basis of host adaptation in the seed beetle, rather than maternal effects. Even so, we encourage researchers to test for

48   potential M×Es because, due to a lack of testing, it remains unclear how

49   often they arise.

51   **Keywords:** genetic correlation, host adaptation, maternal effects, seed

52   beetle, life history

Introduction

Maternal effects modify phenotypes that undergo selection and are therefore a potentially important evolutionary force (Mousseau and Fox, 1998; Räsänen and Kruuk, 2007; Moore *et al.*, 2019). They may either facilitate adaptation to novel environments (Fox and Savalli, 2000; Leftwich *et al.*, 2019), or buffer against changing conditions through trans-generational phenotypic plasticity (Shama *et al.*, 2014). Maternal effects have also been shown to have a range of other implications, such as moderating population dynamics (Plaistow and Benton, 2009), and increasing niche breadth (Van Asch *et al.*, 2010). Maternal effects, which can arise due to both genetic and environmental variation among mothers, can be defined as 'the causal influence of the maternal genotype or phenotype on the offspring phenotype' (Wolf and Wade, 2009). Genetic maternal effects might increase total heritability and facilitate evolution (Wilson *et al.*, 2005; Räsänen and Kruuk, 2007), as opposed to phenotypic (environmental) maternal effects. Teasing these two sources of maternal effects apart is often challenging, however (Kruuk and Hadfield, 2007). Irrespective of their genetic basis, maternal effects increase the degree of similarity between siblings and may thus bias estimates of direct genetic effects upward if ignored (Räsänen and Kruuk, 2007; McAdam *et al.,* 2014). This can lead to

an overestimation of the likely response to selection (McGlothlin and

Galloway, 2014).

Maternal effects are often environment sensitive (Rossiter, 1998;

Galloway, 2005; Räsänen and Kruuk, 2007; McAdam *et al.,* 2014). It is

unclear, however, whether challenging or favourable conditions will result

in stronger maternal effects (Charmantier and Garant, 2005; Rowiński and

Rogell, 2017). A harsher environment is typically assumed to increase

variation among offspring from different mothers (Charmantier and Garant,

2005; Räsänen and Kruuk, 2007). This may, for example, stem from

differences in maternal ability to provision offspring (Parichy and Kaplan,

1992). On the other hand, marine sticklebacks showed weaker maternal

effects under stressful conditions (warmer temperature) (Shama *et al.*,

2014). Similarly higher population densities reduced offspring differences

attributable to variation in maternal age in soil mites (Plaistow and Benton,

2009). Environment-dependent variation in maternal effects like this

represents a form of Maternal-by-Environment interaction, M×E (Figure 1B).

Another form of M×E occurs, irrespective of change in variance across

environments, if individual mothers differ in how they affect their offspring

in an environment-specific manner (Figure 1C). In an experiment with great

tits for instance (Berthouly *et al.*, 2008), mothers supplemented with

carotenoids raised smaller nestlings than the (un-supplemented) control

mothers. When the brood size was increased, however, the result was

opposite and the offspring of carotenoid-supplemented mothers grew faster

compared to controls (Berthouly *et al.*, 2008). A similar interaction was also

observed for offspring immune response (Berthouly *et al.*, 2008). The level

of maternal carotenoids thus affects the fate and relative performance of

the great tit offspring depending on the brood size. This interaction would

cause cross-over of the maternal effects reaction norms because of the

change in rank of maternal siblings between the two environments (Figure

1C). In general, a quantitative genetic analysis of potential M×E can help us

to identify the mechanisms that increase similarity between siblings within

the same environment beyond genetic relatedness. That analysis would be

especially useful if such mechanisms themselves are undescribed, or their

role is not fully understood. Many cases probably involve a mixture of

changes in ranking and variances (as in Figure 1D), but the change in relative

phenotype of offspring (i.e. shift in the rank) due to maternal effects is less

often accounted for. In general, ignoring significant levels of M×E might bias

estimates of additive genetic effects in a host-specific way and inflate G×E

as shown by Vega-Trejo *et al.* (2018). This could partially explain

inconsistency in G×E estimation across different studies (Saltz *et al.*, 2018).

Phytophagous insects are a valuable model group to study life-history

evolution, especially when interested in trade-offs arising from

specialization to a preferred host (Via and Lande, 1985; Agrawal, 2000). At

the same time, phytophagous insects exhibit a variety of maternal effects

that influence key life-history traits (Fox and Dingle, 1994; Rossiter, 1996;

Fox and Czesak, 2000; Van Asch *et al.*, 2010), including host-specific

maternal effects (Fox *et al.*, 1997; Cahenzli and Erhardt, 2013). In the current

study, we investigated the presence of maternal effects, and the potential

for an interaction between maternal effects and offspring rearing host

environment, on general offspring life-history traits, as well as daughters'

egg laying preferences for different host types in the seed beetle,

Callosobruchus maculatus (Coleoptera: Chrysomelidae). This beetle is a

common pest of stored legumes worldwide. The larvae feed on multiple

legume species (family Fabaceae) with varying degrees of success (Gompert

and Messina, 2016; Price *et al.*, 2017; Messina *et al.*, 2018). Females lay eggs

on host beans shortly after copulation (Mitchell, 1975). Host choice is crucial

as larvae cannot move between beans. Incorrect decisions about where a

female chooses to lay her eggs inevitably lead to lower quality, or even dead,

offspring (Mitchell, 1975; Messina and Fry, 2003; Messina *et al.*, 2007).

Larvae hatch 4-5 days after oviposition and burrow into the bean to feed on

the endosperm. A single seed commonly harbours multiple larvae. The

variable level of larval competition within seeds results in different sized

adult beetles, which affects their condition, female reproductive

performance and egg size (Fox and Savalli, 1998). Depending on the

temperature and host species, adults typically emerge within 26-36 days of

oviposition (Fox *et al.*, 2003; Messina, 2004a). *Callosobruchus maculatus*

shows large sexual dimorphism in life-history traits with males emerging

earlier, being smaller and having a shorter lifespan than females (Guntrip *et*

al., 1997; Fox, Bush, *et al.*, 2004; Iglesias-Carrasco *et al.*, 2020).

We used a full-sib/half-sib split brood design to tease apart the role of

additive genetic and maternal effects on *C. maculatus* life-history traits

when larvae develop on two host types – original (cowpea) and novel (mung

bean). We hypothesised that:

1) There are strong maternal effects on offspring life-history traits

(Fox, 1993; Messina and Fry, 2003). This led us to predict that a considerable

component of offspring phenotypic variation, beyond that due to additive

genetic effects, would be due to offspring having different mothers.

2) Maternal effects are environment-specific (M×E). Specifically, we

predicted that the maternal effects variance would differ between offspring

host environments (Figure 1B), and/or that the rank of offspring from the

same mother will change between the two host types decreasing the cross-

environmental maternal correlation (Figure 1C, D).

3) Novel host type will be more challenging for offspring. Hence,

offspring developing in the novel host would suffer reduced performance

compared to those developing in the original host type (i.e. lower larval

survival, longer larval development, lower body mass at emergence, and

shorter adult lifespan).

4) Maternal host preference will predict offspring performance. We

predicted that offspring would perform better on the host type preferred by

their mother when she laid her eggs.

Methods

Experiment overview. We aimed to estimate genetic and maternal effects

for four life-history traits and host egg laying preference. To do this we used

a half-sib/full-sib split brood design with offspring reared on two different

host types - original or novel. We created the parental generation for our

experiment by mating beetles from our stock population. We mated parents

and tested mother egg laying preference for the two host types. We then let

them lay eggs on both host types and measured offspring life-history traits

and the daughters' host preference. We used animal mixed-effects models

to test our predictions.

Origin and maintenance of stock beetles. Beetles used as the parental

generation for our experiment were obtained from a large stock population

originally sourced from the University of Western Australia (Perth, Australia)

in 2017 where they had been bred on cowpea (*Vigna unguiculata*, Fabaceae)

for at least 90 generations. We maintained this stock in our lab at the

Australian National University on cowpea for another 9 generations before

we began this experiment. Our stock was maintained in four, regularly mixed

(every 5 generations) populations of over 500 individuals, each kept on

cowpea at 25-26 °C.

Establishing the parental generation (P). To obtain virgin males and females

for the parental generation (P), we exposed approximately 2000 un-infested

cowpea beans to stock beetles for a period of 48 h. Each bean, which had 5-

8 eggs on its surface (a density which is usual for our stock), was then placed

in an individual Eppendorf tube (with a pinhole for airflow). Once isolated,

we monitored these beans until adults began to emerge. We collected virgin

beetles each morning and used them in our experiment on the same day.

We knew that beetles were virgins as they were either the only beetle to

have emerged that day, or all beetles that had emerged were of the same

sex. Every evening, we discarded any extra beetles that had emerged during

that day to aid in the collection of virgins the following day. Using the beetles

that emerged each morning, we mated males and females according to a

full-sib/half-sib breeding design: each male was sequentially mated with

four random females over the day and the mating order was noted (similar

to Fox *et al.* (2003)). Pairs that did not copulate within 30 min were

separated for half an hour before being placed together for another mating

attempt. Each female was weighed (to the nearest 0.001 mg) prior to mating

(Cubis Ultra-Micro balance, Sartorius Lab Instruments GmbH., Goettingen,

Germany). All matings took place over six days ('day mated', range 1-6, see

Model comparison and partitioning of phenotypic variance). There is

therefore a positive correlation between parental developmental duration

(egg to adulthood) and 'day mated', but it is imperfect because parents

emerged from eggs initially laid over two days (i.e. 48 hours; see above).

There is no correlation between 'day mated' and adult parental age because

all matings occurred on the day that the parents emerged.

Maternal host preference. We used mung bean (*V. radiata*) as the novel

host type. We define the host as novel in that it was 'unlikely to have been

experienced by the study population within an evolutionary timescale'

(Rowiński and Rogell, 2017), approximately 100 generations in our case (see

also Kawecki (1995). Cowpea and mung are both suitable hosts for *C.*

maculatus (Messina, 2004a; Fox and Messina, 2018), but populations kept

on a specific host for multiple generations usually show better performance

on their usual host than a novel host (Messina, 2004a).

Once females had mated, we conducted choice trials to determine

their preferred host type for egg laying. We considered females ready to lay

eggs once the pair dismounted (Wilson and Hill, 1989). For the host

preference trials, we mixed cowpeas (original host) and mung beans (novel

host) in covered Petri dish (∅ 5.5 cm). We used proportionally more mung

than cowpea beans (8:4) to roughly equalise the available surface area of

each bean type (similar to 10:5 used by Messina and Slade (1997)). Cowpea

beans have an approximately 1.6-times larger surface area (Paukku and

Kotiaho, 2008) and are 4-times heavier than mung beans (in our study the

average ± SD bean mass was 294±45 mg for cowpea, and 72±8 mg for mung

bean). Females were left to lay eggs for two hours, after which they were

removed and the number of eggs on each bean was counted. Relative

preference was calculated as the proportion of the eggs that a female laid

on the original host type (cowpea). The values for relative host preference

therefore ranged between 0 and 1.

Generation of offspring (F1). Directly after the host preference trial, we

transferred individual dams to plastic containers (ø 4 cm, height 6 cm) with

10-13 mung beans and left them to lay eggs for 14-18 h. We then moved

dams to new containers with 10-13 cowpea beans for 9-10 h. This difference

in laying time was required because initial trials showed that more time was

necessary for females to lay a sufficient number of eggs on the novel host

(mung beans), than on cowpea (the original host). Females were presented

with mung beans first to prevent them from laying all their eggs on what we

expected to be the preferred host type (cowpea) (Messina and Slade, 1997).

Maternal age and/or laying order effects are unlikely to have influenced

differences between mung and cowpea-reared offspring due to the short

period of time we allowed females to lay eggs (i.e. <24 h). Previous studies

have shown no effect of female age (hence laying order) on offspring quality

during the first days of laying after emergence (Wasserman and Asami, 1985;

Fox, 1993; Fox and Dingle, 1994; Iglesias-Carrasco *et al.*, 2018). Further, one

of the authors (MLH) found, in a separate experiment looking at oviposition

preferences and offspring traits over the same time frame, that the order in

which host beans are presented does not have any observable effect on

offspring survival, development duration, or body mass at emergence

(Zhang *et al.*, in preparation).

Once females had laid eggs on both host types, we collected up to 10

beans of each host type for each female. If a female had laid eggs on fewer

than 10 beans of a given type, we used them all. We ensured that each bean

had only one egg laid on it by scraping off surplus eggs with a scalpel. We

then weighed beans, to the nearest 0.001 mg, within 24 h of oviposition to

measure the resources available to the larvae. Beans with an egg were then

placed individually in Eppendorf tubes with perforated lids and incubated at

26 °C.

We started regular monitoring of F1 emergence on day 24 post-

oviposition. However, we missed the onset of emergence of 40 beetles on

mung (1.3% of all emerged offspring) by approximately 1 day as they

emerged sooner than expected. Uncertainty in the date of their emergence

could introduce noise into the estimates of larval development duration,

body mass at emergence and adult lifespan, so we removed these beetles

from the final dataset. Including these early emerging beetles did not

qualitatively change our results.

Measurement of offspring (F1) traits. We recorded the date of offspring

emergence from the bean, sex and body mass at emergence (to the nearest

0.001 mg), and then removed the bean from the tube. After weighing, the

beetle was returned to its tube and checked daily for survival. We also tested

the host preference of two newly emerged female offspring per dam raised

on each host type (total N = 4/dam). To do this we mated each daughter with

a randomly selected male from the stock and then ran an egg laying

preference trial identical to that described above for their mothers. We

monitored survival until the death of all emerged offspring (day 79 post-

oviposition). At that time, we also censused larval (egg-to-emergence)

survival, assuming that larvae in beans from which a beetle had not yet

emerged had died.

Sample sizes. Our target sample size was determined *a priori* based on Lynch

and Walsh (1998). We created 89 families (89 sires with 356 dams) and

aimed for 10 offspring per dam, per host type (i.e. 7 120 offspring). The final

sample size (N = 3 431) is lower due to unsuccessful matings or too few eggs

being laid. Despite not reaching our target sample size our final sample size

is still higher than that in other similar studies of maternal effects (the

studies in a meta-analysis by Moore *et al.* (2019), for example, have a mean

sample size of 1 370, and a much lower median of 555). We analysed data

for sons and daughters separately due to their large sexual dimorphism

which invariably leads to strong interactions between sex and other effects.

We excluded offspring of dams that did not lay eggs during host preference

trials (285 individuals) and data from 13-24 daughters and 12-27 sons with

missing, or extreme values (>3 standard deviations (SD) from the mean),

with the exact number depending on the focal trait. We analysed four

offspring life-history traits: (egg-to-emergence) 'larval survival' (dead or

emerged; N = 3 146), 'duration of larval development' (number of days

between oviposition and offspring emergence; N = 1 420 daughters and 1

384 sons), 'body mass' (weight at emergence in mg; N = 1 431 daughters and

1 399 sons) and 'adult lifespan' (number of days between offspring

emergence and death; N = 1 425 daughters and 1 389 sons). In addition, we

measured the host egg laying preferences of 665 daughters. The analysed

offspring came from 82 sires and 245-249 dams for the life-history traits, and

239 dams for host preference analysis. The proportion of offspring reared

on the original host type was 52-57 % (see Supplementary Table S1). Prior to

any statistical analysis, but after all data had been collected, we registered

this   project   on   the   Open   Science   Forum   webpage:

https://osf.io/ft7eq/?view_only=0bab0a33bb4246adb64c919601a72757

The final analytical approach we used was different, however, from the

original plan – partly due to feedback from reviewers. We explain our

changes in the *Annotated registration* section in the Supplementary

Information.

Overview of data analysis approach. Our main objective was to test for

maternal effects on the offspring traits (Prediction 1) and their potential

interaction with offspring rearing environment (Prediction 2). First, we built

a 'minimal model' containing only fixed effects directly attributable to the

experimental design. To test whether maternal effects are important, we

then created a set of candidate models with different random effects

structures (Table 1). We used the best-selected model for partitioning of

phenotypic variance (Table 2). Finally, we assess the effect of host type

(Prediction 3) and dam host egg laying preference (Prediction 4) on offspring

traits with a 'full' model containing additional fixed effects (Table 3).

Model specification, model comparison and partitioning of phenotypic

variance. We started data analysis with a minimal mixed-effects model that

contained, in the fixed-effects part, only variables ensuing from the

experimental design. These were 'host type' (original/novel), 'bean mass'

(standardized to zero mean and 1 SD variance, within each host type, to look

at the effect of size within each host type), 'dam mating order' (i.e. dam

position in a sire's mating sequence, integer 1 to 4), and 'day mated' (integer

1-6). We did not fit 'day mated' as a random effect because of the low

number of levels (Bolker *et al.*, 2009). We specified the random effects

structure using an animal model approach (Kruuk, 2004; Wilson *et al.*, 2010),

where variance-covariance estimates between relatives are computed from

a pedigree.

We fitted the models using Bayesian framework with package

MCMCglmm (ver. 2.29) (Hadfield, 2010) in R software (ver. 4.0.0; R Core

Team, 2020). The continuous response variables (duration of larval

development, body mass and adult lifespan) were fitted using Gaussian

family and identity link, while larval survival (dead or emerged) was fitted

using 'categorical' family (binomial distribution with logit link). Daughter

host preference was treated as a bivariate vector with the number of eggs

on the original host and the total number of eggs collected in the egg laying

preference trial using 'multinomial2' family (binomial distribution and log-

odds ratio link function) (Hadfield, 2010).

To test for the best random effects structure, we performed model

comparison on a set of 7 candidate models (Table 1) for each trait using the

Deviance Information Criterion (DIC) (Spiegelhalter *et al.*, 2002). DIC is a

Bayesian alternative to Akaike Information Criterion (AIC) that takes into

account the fit of the model but penalizes for model complexity, with lower

DIC values indicating a better model (Hadfield, 2010; Wilson *et al.*, 2010).

Model comparison with DIC should be performed cautiously for non-

Gaussian response variables (Hadfield, 2010; Wilson *et al.*, 2010), so we did

not conduct model selection for larval survival or host preference.

The simplest model 'G' included only additive genetic effects specified

by the random effect of 'animal'. We then also included dam identity to

estimate maternal effects in addition to additive genetic effects ('G+M'

model). These two models do not, however, take into account possible host-

specific differences in respective variance components (this is shown in

Figure 1A). We thus formulated more realistic models that estimated the

variance component (including residual variance) separately for each host

type (as in Figure 1B). These were specified through random-effect

interactions (Hadfield, 2010) with host type 'idh(HOST):animal' for 'G_{sep}', or

with 'idh(HOST):dam' for 'M_{sep}' models (Table 1). This is similar to fitting two

separate host-specific models. To account for potential change in the

ranking of additive genetic or maternal effects (Figure 1C), we fitted models

that also estimated cross-environmental covariance ('G_{cov}' or 'M_{cov}'). When

specifying variance-covariance matrix of the random-effect interaction we

used 'unstructured' form, e.g. 'us(HOST):animal' (Hadfield, 2010). As a

result, we not only obtained separate variance estimates for each host type,

but also the corresponding covariance to compute the cross-environmental

correlations (Figure 1D) – the proxies for Genotype-by-Environment (G×E),

or Maternal-by-Environment (M×E) interactions (Lynch and Walsh, 1998). In

the models estimating separate variances per host type, we always specified

host-specific residual variance (as 'rcov=~idh(HOST):units') (Hadfield, 2010).

We then performed phenotypic variance partitioning based on the

outputs of the best-selected minimal model for each trait. We calculated the

maternal-effects proportion (m^2), narrow-sense heritability (h^2), evolvability

(coefficient of additive genetic variation, CV_A) and the cross-environmental

correlation for additive genetic (r_G – potential G×E interaction) and for

maternal effects (r_M – potential M×E interaction). We used maternal effects

proportion, $m^2 = \dfrac{V_M}{V_P} = \dfrac{V_{dam}}{(V_{animal} + V_{dam} + V_{residual})}$, to quantify the

importance of maternal effects (as in e.g. Messina and Fry (2003); Vega-

Trejo *et al.* (2018); Moore *et al.* (2019)). The V_M contains both genetic and

non-genetic maternal effects, and potentially also genetic dominance, as we

did not separate them with our experimental design. We calculated

heritability as $h^2 = \dfrac{V_A}{V_P} = \dfrac{V_{animal}}{(V_{animal} + V_{dam} + V_{residual})}$ (Lynch and Walsh, 1998),

assuming epistatic effects were negligible. Evolvability was defined by

(Houle, 1992) as $CV_A = \dfrac{\sqrt{V_A}}{\mu} = \dfrac{\sqrt{V_{animal}}}{\mu}$, where μ is the mean of a trait.

Evolvability standardizes additive genetic variance over the trait mean and

is therefore useful to compare the potential evolutionary response among

traits, unlike heritability which is conditional on the amount of residual

phenotypic variation (Hansen *et al.*, 2011). The correlation between additive

genetic effects on the two host types (G×E interaction) was $r_G =$

$\dfrac{COV_{animal\,(original-nov\quad)}}{\sqrt{V_{animal\,(original)}V_{animal\,(novel)}}}$. The same formula was used for maternal

effects correlation between the two hosts (potential M×E interaction) using

the dam-related covariance across the two host types. Estimates of V_A, V_P

and h^2 for larval survival and host preference were back-transformed to the

original data scale using the function 'QGparams' (from QGglmm package,

ver.0.7.4; de Villemereuil *et al.* (2016)).

Testing for host effect and dam host egg laying preference on offspring

phenotype. To determine the effect of novel host type on offspring

performance (Prediction 3) and to test whether offspring performed better

on the host type that was preferred by their mother (Prediction 4), we built

a 'full model'. The full mixed-effects model always included all the terms

outlined above for the minimal model with the random effects from the

best-selected minimal model (Table 1). We then included 'dam host

preference' (i.e. the relative preference ratio: 0-1 values) and the interaction

between preference and host type 'HOST:dam host preference' to test if the

effect of the strength of the preference for the original host on offspring

traits differs depending on the host type. We also included interactions

between host type and bean mass 'HOST:bean mass', as well as host type

and day mated 'HOST:day mated' as fixed effects to test for potential host-

specific effects.

Model fitting. To fit the MCMC models, we used non-informative priors for

the random-effects variance (G) with expected variance-covariance matrix

at limit 'V' = 1, and the degree of freedom 'nu' = 0.002 for each random term

(with nu = 1.002 in models with the random effects estimated separately per

host type) (Wilson *et al.*, 2010). To further improve mixing and effective

sample size, we specified parameter-expanded priors as 'α mu' = 0 and 'α V'

= 1 000 (Hadfield, 2010). Priors for residual variance (R) were set as V = 1 and

nu = 0.002 (nu = 1.002 when residual variance was estimated separately for

each host type). In the analysis of survival, the residual variance had to be

fixed to 1, so that the binomial mixed-effects model could be estimated

(Hadfield, 2010). For fixed effects (B), we retained default priors with mean

= 0 and variance = 10^{10}. To generate posterior distributions from the minimal

and full models, we ran 360 000 iterations with burn-in of 60 000 and

thinning interval of 300, so that we obtained an effective sample size of 1

000 for each estimated model term. For larval survival, we increased the

iterations to 5 500 000 with a burn-in of 500 000 and thinning interval of

5000 as we were aiming for an autocorrelation <0.1 at the lag corresponding

to the thinning interval (Wilson *et al.*, 2010). Model convergence was

evaluated by visual examination of the traces using function

'plot(mcmc.model)'. We assessed the importance of individual model terms

based on their credible intervals (CrI).

Results

Prediction 1 and 2 – testing for maternal effects and M×E interaction. The

model comparison suggests very weak maternal effects for all the measured

traits. The models incorporating maternal effects performed poorly (Table

1), and the best model contained only additive genetic effects estimated for

each host type and the cross-environmental covariance (the 'G_{cov}' model).

This shows that either additive genetic variance, residual variance or both

differed between host types and that the cross-environmental genetic

covariance differed from zero. Estimate of the cross-environmental

covariance has to be examined directly to see whether the correlation differs

from +1 indicating a potential Genotype-by-Environment interaction (see

below).

Table 1. Model comparison.

Components of phenotypic variance related to additive genetic effects -

heritability, evolvability and G×E. We observed higher heritability for

offspring life-history traits in the novel host type (h^2 = 0.365-0.649), than on

the original host (h^2 = 0.210-0.459) (Table 2). However, the additive genetic

variance was substantially higher in the novel host only for adult lifespan of

sons. For the other traits, we recorded a similar magnitude of additive

genetic variance in the two hosts, but higher residual phenotypic variance in

the original host type, thereby reducing trait heritability (Table 2). The

presence of similar levels of additive genetic variance in both hosts is also

apparent in the estimates of evolvability (variance due to additive genetic

effects standardised over trait means) which were comparable for the two

host types (Table 2). When comparing evolvability among traits (across sexes

and host types) we found that it was highest for adult lifespan (CV_A = 16.8-

20.8 %, Table 2). For the non-Gaussian traits (larval survival and host egg

laying preference), estimates of additive genetic variance and heritability

obtained from the G_{cov} model were very low (h^2 = 0.001-0.060, Table 4A)

when compared to the Gaussian life-history traits.

We recorded highly positive correlations between offspring

performance on the two host types for additive genetic effects (r_G) (Table 2).

The only exception was for the development duration of sons, which

exhibited only a weak genetic correlation between hosts (Table 2). The

credible interval (-0.279, 0.533) suggests a Genotype-by-Environment

interaction (G×E).

Table 2. Phenotypic variance partitioning.

Predictions 3 and 4 – testing for host type suitability and host preference-

performance. Unexpectedly, given our predictions, offspring generally

performed better on the novel host (mung) than on the original host

(cowpea) (Table 3; Supplementary Table S1). We recorded very high larval

survival overall, however, so that survival on mung beans was only

marginally better than on cowpea (Table 4B). We did not find a positive

relationship between the strength of the dam host preference for egg laying

and the performance of their offspring on either host type. The one

exception was for daughter body mass: daughters of dams that more

strongly preferred the cowpea emerged heavier when raised on cowpea, but

not when raised on mung beans (Table 3). Daughters reared on the original

cowpea host also exhibited a higher egg laying preference for cowpeas than

those reared on mung beans. Raw average host preference ratio of

daughters reared on cowpea was 0.681 versus 0.517 for those from mung

(where 1 is a strict preference for cowpeas). The host preference ratio of

their mothers (reared on cowpea) was 0.724.

Table 3. Fixed-effects estimates from the full models.

Effect of experimental variables on offspring traits - bean size, mating date

and mating order. Host-specific relative bean size (zero-centered per host

type) and mating date were important predictors of offspring traits. For both

host types, offspring that developed in larger beans had a poorer outcome:

increased development duration in daughters, as well as reduced body mass

and lower adult lifespan in both sexes. The effects of mating day on offspring

performance were, however, mixed (Table 3). Offspring of later mated

families took longer to develop, but sons emerged at a larger size and lived

longer. Daughters from later mated families also showed a modest decrease

in the preference for the original host type (Table 4B). Finally, if a dam was

later in the sire's mating sequence this marginally extended the

development duration of her daughters.

Table 4. Phenotypic variance partitioning and full model inference for

the non-Gaussian traits.

Discussion

Summary. Our main aim was to test for environment-specific maternal effects on offspring life-history traits and their correlation, that is, Maternal-by-Environment interactions (M×E). We used a full-sib/half-sib split brood design with seed beetles (*C. maculatus)* reared on two types of host: original – cowpea; and novel – mung bean. Maternal effects, however, proved to be negligible for all of the measured traits, on both host types. Instead, we found that additive genetic variance played an important role in determining phenotypic variation. Contrary to our predictions, offspring reared on the novel host type performed better for all the measured traits. Daughters whose mothers preferred to lay eggs on the original host were heavier on this host than those of mothers that preferred the novel host type. There was, however, no equivalent effect of host preference on body mass in the novel host. Below we discuss our findings in more detail.

Weak maternal effects. We used an original and a novel host type to examine whether mothers affect their offspring's phenotype in a host-

specific way. Previous studies have shown that environmental stress can

either increase or decrease the magnitude of maternal effects (Charmantier

and Garant, 2005; Rowiński and Rogell, 2017). Maternal effects variation is

sometimes not evident in resource rich conditions as differences among

mothers are reduced compared to a poor environment where differences

are exacerbated (e.g. Parichy and Kaplan (1992)). If both host types in our

study are resource rich (especially when few eggs are laid per bean) this

could explain the negligible maternal effects that we see here. Previous

quantitative genetic studies employing a similar approach to ours have

reported significant maternal effects in *C. maculatus*. This has been,

however, in other hosts (adzuki (*Vigna angularis*) in Fox (1993)) or when

testing the effect of bean removal after beetle emergence on adult traits

(Messina and Fry, 2003). The host species we used are both thought to be

highly favourable to *C. maculatus* (Messina, 2004a; Fox and Messina, 2018)

(discussed further below). This, in addition to the fact we prevented larval

competition with our experimental design, may have substantially reduced

the role of maternal effects in the current experiment.

Heritability was higher in the novel host type, but evolutionary potential

was similar in both hosts. Heritability of fitness-related traits is traditionally

assumed to be low, as additive genetic variation is expected to be depleted

by natural selection (Mousseau and Roff, 1987; Hill, 2010). Egg size, body

size, fecundity or lifespan were nevertheless previously shown to have

intermediate to high heritability in *C. maculatus* (h^2 from 0.27 to 0.74) (Fox,

1993; Messina, 1993; Fox *et al.*, 2003; Fox and Messina, 2018). Here, we

recorded considerable heritability in all life-history traits (except larval

survival), with the highest values occurring for daughter body mass and son

adult lifespan (both h^2 > 0.65). Heritability was notably higher in the novel

host environment (mung bean), with the one exception of adult lifespan of

daughters. Holloway *et al.* (1990) proposed that a novel environment should

yield higher additive genetic variance due to new genes being expressed

thereby uncovering cryptic genetic variation (Hoffmann and Merilä, 1999).

Indeed, Messina and Fry (2003) found that *C. maculatus* exhibited increased

additive genetic variance for longevity under a novel environment (absence

of host beans after emergence). Our results indicate, however, that it was

lower residual variance in the novel host (mung bean), rather than increased

additive genetic variation, that was responsible for higher heritability in the

novel environment. Residual phenotypic variation may entail non-additive

genetic, environmental variance and developmental noise (Rowiński and

Rogell, 2017) and the influence of at least some of these factors apparently

declined in the novel host type. More elaborated breeding design would be

required (see e.g. Tucić and Šešlija (2007), or Bilde *et al.* (2008)), however,

573 to examine their respective effects in *C. maculatus*. Interpretation of

574 heritability as the proportion of additive genetic versus total variance can

575 therefore become problematic (Hansen *et al.,* 2011) and evolvability offers

576 a more conclusive comparison of evolutionary potential (Houle, 1992;

577 Hansen *et al.*, 2011). Similar values of trait evolvability indicated that the

578 traits measured in our study possess similar evolutionary potential in the

579 two host types.

580  Genetic trade-offs are thought to prevent populations of parasites or

581 herbivores from becoming 'masters of all trades' and performing well on

582 multiple hosts (Joshi and Thompson, 1995; Agrawal *et al.*, 2010). In

583 phytophagous insects, performance trade-offs on different hosts are,

584 however, rarely reported (Ueno *et al.*, 2003; Scheirs *et al.*, 2005; Messina

585 and Durham, 2015). This includes studies of *C. maculatus* where offspring

586 from the same family usually perform similarly across hosts without

587 exhibiting a G×E interaction (Fox, 1993; Guntrip and Sibly, 1998; Messina,

588 2004a). We also found strong positive cross-environmental genetic

589 correlations indicating parallel reaction norms for offspring from different

590 families when reared on the two host types. Only the development duration

591 of sons showed a low genetic correlation between the two host types

592 indicating that it could evolve differently on cowpea and mung (Via and

593 Lande, 1985). It is, however, important to bear in mind that breeding design

studies cannot definitively conclude there are no genetic trade-offs in

adaptation to different hosts because they lack mechanistic explanation.

This can be obtained through analysis of long-term selected lines (Agrawal

et al., 2010) or data on the potential genes involved (Gompert and Messina,

2016).

Offspring performed better on the novel host type. We treated mung bean

as a novel host type. This was despite mung bean regularly being used to

rear other strains of *C. maculatus* (e.g. the Southern India strain (Fox, Bush,

et al., 2004)), and regarded with cowpea as a favourable host species for *C.*

maculatus (Fox and Messina, 2018). Our stock population was kept on

cowpea for at least 100 generations. In addition, cowpea strains are overall

more viable than those from mung bean (Messina, 2004a) and the cowpea

is generally considered to be the ancestral host for *C. maculatus* (Kébé *et al.*,

2017). Due to these reasons and the fact that a previous study showed that

long-term maintenance on a specific host yields decreased viability on other

hosts (Messina, 2004a), we predicted a lower suitability of the novel host

type (mung bean). Our finding that offspring performed better on the novel

host type (mung bean) was therefore unexpected. While offspring survival

was almost 90 % on cowpea, it was even higher on the novel host (93 % on

mung bean). There was no obvious cost to the offspring of rearing in the

novel host, as the duration of larval development was shorter on mung bean.

However, beetles that emerged from mung beans were larger and lived

longer as adults than those that emerged from cowpeas. It is not

straightforward, however, to tell which of the two host species is generally

better for *C. maculatus*, because it depends on other factors such as the level

of competition (Messina, 2004a; Paukku and Kotiaho, 2008; Fox and

Messina, 2018), temperature (Stillwell *et al.*, 2007) and the population's

evolutionary history (Messina, 2004a). In the present study, we eliminated

effects of larval competition by rearing one larva per bean. We therefore

removed the fitness advantage that cowpea may provide due to its larger

size when there is larval competition (Fox and Savalli, 1998). Surprisingly,

relative bean size within a given host type also negatively affected body mass

and lifespan as well as extending the developmental duration of daughters.

We cannot reject the hypothesis that the host effect recorded in our

experiment was confounded by egg laying order, as eggs were laid first on

mung bean (see Methods for the experimental justification). The lower

performance of offspring on cowpea might have been because, following

the preference trial, females were first provided with mung beans to lay

eggs, and only after that with cowpeas. There are, however, several reasons

to think that an effect of laying order in our experiment is unlikely. The

lifetime fecundity of *C. maculatus* is 60-80 eggs (Mitchell, 1975; Credland

and Wright, 1989; Wilson and Hill, 1989; Messina and Fry, 2003). The dams

in our experiment laid around 30 eggs per 24 h and probably still possessed

more than half of their lifetime fecundity after they had finishing laying in

our experiment. Crucially, past studies suggest there is no negative effect of

maternal age on offspring performance on the first day of egg laying (i.e. the

relevant time period in our study) (Wasserman and Asami, 1985; Fox, 1993;

Fox and Dingle, 1994; Iglesias-Carrasco *et al.*, 2018; Zhang *et al.*, in

preparation). We therefore think that the sequence of host types available

for egg laying is unlikely to explain the poorer offspring performance on

cowpea.

Host preference of *C. maculatus* increased daughter body mass and rearing

host affected host preference. An egg laying preference for a particular host

often predicts offspring performance on that host in phytophagous insects

(Gripenberg *et al.*, 2010). However, despite the critical role of oviposition

choice for larval success in *C. maculatus*, of the five offspring traits we

measured, the strength of dam preference for cowpea only correlated with

daughter body mass. Dams that showed a stronger preference for cowpea

produced heavier-than-average daughters on cowpea, but not when laying

on mung. This contributes to mother fitness through higher daughter

reproductive potential in cowpea as body size is a good proximate measure

for fecundity in *C. maculatus* (Messina and Fry, 2003). In mung bean,

however, daughters' body mass was not affected by their mothers' host

preference and the preference-performance relationship thus only holds for

daughter body mass in cowpea. The weak overall relationship between

mother's host preference for one host type and the performance of her

offspring on that host suggests that *C. maculatus* females might aim to

maximize their own fitness and not that of individual offspring (i.e. they

prioritise offspring number over offspring quality), perhaps through saving

searching time for the best host (Wood *et al.*, 2018).

Previous studies show that strains of *C. maculatus* maintained for > 40

generations on mung bean or cowpea exhibit a strong preference for their

recent host type (Messina, 2004b). Fox, Stillwell, *et al.* (2004) used different

host strains to demonstrate that switching a strain to another host for just a

single generation has no effect on host preference. We found that while

daughters reared on cowpea preferred cowpea over the novel host (mung

bean), those reared on mung beans showed no preference for either host

type. This suggests that there might be a G×E interaction affecting host egg

laying preferences, but tracking of more generations is necessary to test

whether the host preference evolves after the host switch (as in Fox,

Stillwell, *et al.* (2004), or Messina (2004b)).

Conclusion. Maternal effects were low for all measured traits in our study

for offspring reared on either a novel or a familiar host. A next step in the

exploration of whether M×E interactions affect estimates of genetic

variation, and of G×E interactions specifically, would be to conduct

experiments using less favourable novel hosts of *C. maculatus*, for example

adzuki bean (Fox, 1993) or lentil (Gompert and Messina, 2016). The host

species we used are both considered to be of high quality (Fox and Messina,

2018). By creating more challenging conditions for offspring, maternal

effects might emerge as an important host-specific factor that affects

variation in offspring performance (Fox *et al.*, 1997; Räsänen and Kruuk,

2007). Despite the fact that maternal effects were negligible in our study,

we still argue that it is important testing for M×E interactions. The M×E

effects have potential to bias measurements of G×E and the lack of research

on M×E interaction is an obstacle towards understanding their role in

adaptive processes.

Acknowledgements

We appreciate detailed comments by three anonymous referees that helped

us to improve the manuscript. We thank Timothée Bonnet and Loeske E. B.

Kruuk (RSB ANU) for very helpful discussions about phenotypic variation

partitioning and Bayesian mixed-effect models. We are thankful to Carolina

Dolan, Krish S. Sanghvi and Elroy Kwan Au (RSB ANU) for their help with the

experiment. MV and PJCC were both supported by 6-month Endeavour

fellowships from the Australian Government.

Author contribution

MV, PJCC, MLH and MDJ conceived and designed the study; MV, PJCC and

MIC collected data; MV analysed the data and drafted the manuscript; and

PJCC, MIC, MLH and MDJ revised it. All authors approved the final version of

the manuscript.

Conflict of interest

We declare that we do not have any conflict of interest.

Data archiving

Data with accompanying R code will be made available after manuscript

acceptance.

Supplementary Information

Table S1 and Annotated Registration.

References

Agrawal AA (2000). Host-range evolution: Adaptation and trade-offs in

fitness of mites on alternative hosts. *Ecology* **81**: 500–508.

Agrawal AA, Conner JK, Rasmann S (2010). Tradeoffs and negative

correlations in evolutionary ecology. In: Futuyma D, Levinton J, Eanes

W (eds) *Evolution since Darwin: the first 150 years*, Sinauer, pp. 243–

268.

Van Asch M, Julkunen-Tiito R, Visser ME (2010). Maternal effects in an

insect herbivore as a mechanism to adapt to host plant phenology.

Funct Ecol **24**: 1103–1109.

Berthouly A, Cassier A, Richner H (2008). Carotenoid-induced maternal

effects interact with ectoparasite burden and brood size to shape the

trade-off between growth and immunity in nestling great tits. *Funct*

Ecol **22**: 854–863.

Bilde T, Friberg U, Maklakov AA, Fry JD, Arnqvist G (2008). The genetic

architecture of fitness in a seed beetle: Assessing the potential for

indirect genetic benefits of female choice. *BMC Evol Biol* **8**.

Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, *et*

al. (2009). Generalized linear mixed models: a practical guide for

ecology and evolution. *Trends Ecol Evol* **24**: 127–35.

Cahenzli F, Erhardt A (2013). Transgenerational acclimatization in an

herbivore-host plant relationship. *Proc R Soc B Biol Sci* **280**.

Charmantier A, Garant D (2005). Environmental quality and evolutionary

potential: Lessons from wild populations. *Proc R Soc B Biol Sci* **272**:

1415–1425.

Credland PF, Wright AW (1989). Factors affecting female fecundity in the

cowpea seed beetle, *Callosobruchus maculatus* (Coleoptera:

Bruchidae). *J Stored Prod Res* **25**: 125–136.

Fox CW (1993). Maternal and genetic influences on egg size and larval

performance in a seed beetle. *Heredity* **73**: 509–517.

Fox CW, Bush ML, Roff DA, Wallin WG (2004). Evolutionary genetics of

lifespan and mortality rates in two populations of the seed beetle,

Callosobruchus maculatus. *Heredity* **92**: 170–181.

Fox CW, Bush ML, Wallin WG (2003). Maternal age affects offspring

lifespan of the seed beetle. *Funct Ecol* **17**: 811–820.

Fox CW, Czesak ME (2000). Evolutionary ecology of progeny size in

arthropods. *Annu Rev Entomol* **45**: 341–369.

Fox CW, Dingle H (1994). Dietary mediation of maternal age effects on

offspring performance in a seed beetle (Coleoptera: Bruchidae). *Funct*

Ecol **8**: 600–606.

Fox CW, Messina FJ (2018). Evolution of larval competitiveness and

associated life-history traits in response to host shifts in a seed beetle.

J Evol Biol **31**: 302–313.

Fox CW, Savalli UM (1998). Inheritance of environmental variation in body

size: Superparasitism of seeds affects progeny and grandprogeny body

size via a nongenetic maternal effect. *Evolution* **52**: 172.

Fox CW, Savalli UM (2000). Maternal effects mediate host expansion in a

seed-feeding beetle. *Ecology* **81**: 3–7.

Fox CW, Stillwell RC, Amarillo-Suárez AR, Czesak ME, Messina FJ (2004).

Genetic architecture of population differences in oviposition

behaviour of the seed beetle *Callosobruchus maculatus*. *J Evol Biol* **17**:

1141–1151.

Fox CW, Thakar MS, Mousseau TA (1997). Egg size plasticity in a seed

beetle: An adaptive maternal effect. *Am Nat* **149**: 149–163.

Galloway LF (2005). Maternal effects provide phenotypic adaptation to

local environmental conditions. *New Phytol* **166**: 93–100.

Gompert Z, Messina FJ (2016). Genomic evidence that resource-based

trade-offs limit host-range expansion in a seed beetle. *Evolution* **70**:

1249–1264.

Gripenberg S, Mayhew PJ, Parnell M, Roslin T (2010). A meta-analysis of

preference-performance relationships in phytophagous insects. *Ecol*

Lett **13**: 383–393.

Guntrip J, Sibly RM (1998). Phenotypic plasticity, genotype-by-environment

interaction and the analysis of generalism and specialization in

Callosobruchus maculatus. *Heredity* **81**: 198–204.

Guntrip J, Sibly RM, Holloway GJ (1997). The effect of novel environment

and sex on the additive genetic variation and covariation in and

between emergence body weight and development period in the

cowpea weevil, *Callosobruchus maculatus* (Coleoptera, Bruchidae).

Heredity **78**: 158–165.

Hadfield JD (2010). MCMC methods for multi-response generalized linear

mixed models: The MCMCglmm R package. *J Stat Softw* **33**.

Hansen TF, Pélabon C, Houle D (2011). Heritability is not evolvability. *Evol*

Biol **38**: 258–277.

Hill WG (2010). Understanding and using quantitative genetic variation.

Philos Trans R Soc B Biol Sci **365**: 73–85.

Hoffmann AA, Merilä J (1999). Heritable variation and evolution under

favourable and unfavourable conditions. *Trends Ecol Evol* **14**: 96–101.

Holloway GJ, Povey SR, Sibly RM (1990). The effect of new environment on

adapted genetic architecture. *Heredity* **64**: 323–330.

Houle D (1992). Comparing evolvability and variability of quantitative traits.

Genetics **130**: 195–204.

Iglesias-Carrasco M, Brookes S, Kruuk LEB, Head ML (2020). The effects of

competition on fitness depend on the sex of both competitors. *Ecol*

Evol **10**: 9808–9826.

Iglesias-Carrasco M, Jennions MD, Zajitschek SRK, Head ML (2018). Are

females in good condition better able to cope with costly males?

Behav Ecol **29**: 876–884.

Joshi A, Thompson JN (1995). Trade-offs and the evolution of host

specialization. *Evol Ecol* **9**: 82–92.

Kawecki TJ (1995). Expression of genetic and environmental variation for

life history characters on the usual and novel hosts in *Callosobruchus*

maculatus (Coleoptera: Bruchidae). *Heredity* **75**: 70–76.

Kébé K, Alvarez N, Tuda M, Arnqvist G, Fox CW, Sembène M, *et al.* (2017).

Global phylogeography of the insect pest *Callosobruchus maculatus*

(Coleoptera: Bruchinae) relates to the history of its main host, *Vigna*

unguiculata. *J Biogeogr* **44**: 2515–2526.

Kruuk LEB (2004). Estimating genetic parameters in natural populations

using the 'animal model'. *Philos Trans R Soc B Biol Sci* **359**: 873–890.

Kruuk LEB, Hadfield JD (2007). How to separate genetic and environmental

causes of similarity between relatives. *J Evol Biol* **20**: 1890–1903.

Leftwich PT, Nash WJ, Friend LA, Chapman T (2019). Contribution of

maternal effects to dietary selection in Mediterranean fruit flies.

Evolution **73**: 278–292.

Lynch M, Walsh B (1998). *Genetics and Analysis of Quantitative Traits*.

Sinauer.

McAdam AG, Garant D, Wilson AJ (2014) The effects of others' genes:

maternal and other indirect genetic effects. In: Charmantier A, Garant

D, Kruuk LEB (eds) *Quantitative Genetics in the Wild*. Oxford University

Press, pp. 81-104.

McGlothlin JW, Galloway LF (2014). The contribution of maternal effects to

selection response: An empirical test of competing models. *Evolution*

68: 549–558.

Messina FJ (1993). Heritability and 'evolvability' of fitness components in

*Callosobruchus maculatus. Heredity (***71**: 623–629.

Messina FJ (2004a). Predictable modification of body size and competitive

ability. *Evolution* **58**: 2788–2797.

Messina FJ (2004b). How labile are the egg-laying preferences of seed

beetles? *Ecol Entomol* **29**: 318–326.

Messina FJ, Durham SL (2015). Loss of adaptation following reversion

suggests trade-offs in host use by a seed beetle. *J Evol Biol* **28**: 1882–

1891.

Messina FJ, Fry JD (2003). Environment-dependent reversal of a life history

trade-off in the seed beetle *Callosobruchus maculatus*. *J Evol Biol* **16**:

501–509.

Messina FJ, Lish AM, Gompert Z (2018). Variable responses to novel hosts

by populations of the seed beetle *Callosobruchus maculatus*

(Coleoptera: Chrysomelidae: Bruchinae). *Environ Entomol* **47**: 1194–

1202.

Messina FJ, Morrey JL, Mendenhall M (2007). Why do host-deprived seed

beetles 'dump' their eggs? *Physiol Entomol* **32**: 259–267.

Messina FJ, Slade AF (1997). Inheritance of host-plant choice in the seed

beetle *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Ann*

Entomol Soc Am **90**: 848–855.

Mitchell R (1975). The evolution of oviposition tactics in the bean weevil,

Callosobruchus maculatus (F.). *Ecology* **56**: 696–702.

Moore MP, Whiteman HH, Martin RA (2019). A mother's legacy: the

strength of maternal effects in animal populations. *Ecol Lett* **22**: 1620–

1628.

Mousseau TA, Fox CW (1998). The adaptive significance of maternal

effects. *Trends Ecol Evol* **13**: 403–407.

Mousseau TA, Roff DA (1987). Natural selection and the heritability of

fitness components. *Heredity* **59**: 181–197.

Parichy DM, Kaplan RH (1992). Maternal effects on offspring growth and

development depend on environmental quality in the frog *Bombina*

orientalis. *Oecologia* **91**: 579–586.

Paukku S, Kotiaho JS (2008). Female oviposition decisions and their impact

on progeny life-history traits. *J Insect Behav* **21**: 505–520.

Plaistow SJ, Benton TG (2009). The influence of context-dependent

maternal effects on population dynamics: An experimental test. *Philos*

Trans R Soc B Biol Sci **364**: 1049–1058.

Price TN, Leonard A, Lancaster LT (2017). Warp-speed adaptation to novel

hosts after 300 generations of enforced dietary specialisation in the

seed beetle *Callosobruchus maculatus* (Coleoptera: Chrysomelidae:

Bruchinae). *Eur J Entomol* **114**: 257–266.

R Core Team (2020) R: A language and environment for statistical

computing. v. 4.0.0. R Foundation for Statistical Computing, Vienna,

Austria.

Räsänen K, Kruuk LEB (2007). Maternal effects and evolution at ecological

879   time-scales. *Funct Ecol* **21**: 408–421.

880   Roff DA, Wilson AJ (2014) Quantifying Genotype-by-Environment

881   interactions in laboratory systems. In: Hunt J, Hosken D (eds)

882   *Genotype-by-Environment Interactions and Sexual Selection*. John

883   Wiley and Sons, Ltd., pp. 101-136.

884   Rossiter M (1996). Incidence and consequences of inherited environmental

885   effects. *Annu Rev Ecol Syst* **27**: 451–476.

886   Rossiter M (1998) The role of environmental variation in parental effects

887   expression. In: Mousseau TA, Fox CW (eds) *Maternal Effects as*

888   *Addaptations*. Oxford University Press, pp. 112-134.

889   Rowiński PK, Rogell B (2017). Environmental stress correlates with

890   increases in both genetic and residual variances: A meta-analysis of

891   animal studies. *Evolution* **71**: 1339–1351.

892   Saltz JB, Bell AM, Flint J, Gomulkiewicz R, Hughes KA, Keagy J (2018). Why

893   does the magnitude of genotype-by-environment interaction vary?

894   *Ecol Evol* **8**: 6342–6353.

895   Scheirs J, Jordaens K, De Bruyn L (2005). Have genetic trade-offs in host use

896   been overlooked in arthropods? *Evol Ecol* **19**: 551–561.

Shama LNS, Strobel A, Mark FC, Wegner KM (2014). Transgenerational

plasticity in marine sticklebacks: Maternal effects mediate impacts of

a warming ocean. *Funct Ecol* **28**: 1482–1493.

Spiegelhalter DJ, Best NG, Carlin BP, Van Der Linde A (2002). Bayesian

measures of model complexity and fit. *J R Stat Soc Ser B Stat Methodol*

64: 583–616.

Stillwell RC, Wallin WG, Hitchcock LJ, Fox CW (2007). Phenotypic plasticity

in a complex world: Interactive effects of food and temperature on

fitness components of a seed beetle. *Oecologia* **153**: 309–321.

Tucić N, Šešlija D (2007). Genetic architecture of differences in oviposition

preference between ancestral and derived populations of the seed

beetle Acanthoscelides obtectus. *Heredity* **98**: 268–273.

Ueno H, Fujiyama N, Yao I, Sato Y, Katakura H (2003). Genetic architecture

for normal and novel host-plant use in two local populations of the

herbivorous ladybird beetle, *Epilachna pustulosa*. *J Evol Biol* **16**: 883–

895.

Vega-Trejo R, Head ML, Jennions MD, Kruuk LEB (2018). Maternal-by-

environment but not genotype-by-environment interactions in a fish

without parental care. *Heredity* **120**: 154–167.

Via S, Lande R (1985). Genotype-environment interaction and the evolution

of phenotypic plasticity. *Evolution* **39**: 505–522.

de Villemereuil P, Schielzeth H, Nakagawa S, Morrissey MB (2016). General

methods for evolutionary quantitative genetic inference from

generalized mixed models. *Genetics* **204**: 1281–1294.

Wasserman SS, Asami T (1985). The effect of maternal age upon fitness of

progeny in the southern cowpea weevil, *Callosobruchus maculatus*.

Oikos **45**: 191–196.

Wilson AJ, Coltman DW, Pemberton JM, Overall ADJ, Byrne KA, Kruuk LEB

(2005). Maternal genetic effects set the potential for evolution in a

free-living vertebrate population. *J Evol Biol* **18**: 405–414.

Wilson K, Hill L (1989). Factors affecting egg maturation in the bean weevil

Callosobruchus maculatus. *Physiol Entomol* **14**: 115–126.

Wilson AJ, Réale D, Clements MN, Morrissey MB, Postma E, Walling CA, *et*

al. (2010). An ecologist's guide to the animal model. *J Anim Ecol* **79**:

13–26.

Wolf JB, Wade MJ (2009). What are maternal effects (and what are they

not)? *Philos Trans R Soc London B* **364**: 1107–1115.

Wood CW, Wice EW, del Sol J, Paul S, Sanderson BJ, Brodie ED (2018).

Constraints imposed by a natural landscape override offspring fitness

effects to shape oviposition decisions in wild forked fungus beetles.

Am Nat **191**: 524–538.

TABLES

Table 1. Model comparison to estimate the best random effects structure for minimal animal MCMC models. The table shows the difference between Deviance Information Criterion (DIC) for each candidate model and the model with the lowest DIC (denoted as '-'). Model 'G' contained additive genetic variance only. The model with 'M' included also maternal effects variance. Models where variance was estimated separately '$_{sep}$' for each host type are 'G_{sep}' or 'M_{sep}'. Models 'G_{cov}' (or 'M_{cov}') estimated covariance '$_{cov}$' between the two host types in addition to the host-specific variance. When the host-specific interaction was fitted as a random effect (models '$_{sep}$' and '$_{cov}$'), we also estimate residual variance separately for each host type. Further details on model fitting are in the Methods. Please note that the low DIC for model 'G_{cov}' means that the genetic covariance between the host types is different from zero. It is, however, inconclusive about Genotype-by-Environment interaction as the covariance can still be highly positive (i.e. the rank order of genotypes does not change). So we also examined the cross-environmental genetic correlation (r_G) estimates from the model given in Table 2. DIC is not reliable for model comparison in non-Gaussian traits (Hadfield, 2010; Wilson *et al.*, 2010) and we thus did not perform model selection for larval survival and host egg laying preference of daughters and used model G_{cov} for inference.

Model	Random effects	Development duration		Body mass		Adult lifespan	
		daughters	sons	daughters	sons	daughters	sons
G	animal	137.6	225.4	82	78.8	70.4	125.6
G+M	animal + dam	151.8	233.3	136.3	115.9	79.4	153.2
G_{sep}	idh(HOST):animal	10.4	-	18.1	46.3	15.6	72.8
G_{cov}	us(HOST):animal	-	0.4	-	-	-	-
$G_{sep}+M_{sep}$	idh(HOST):animal + idh(HOST):dam	42.5	29.6	97.5	108.9	39.2	133.1
$G_{cov}+M_{sep}$	us(HOST):animal + idh(HOST):dam	23.7	33.5	31.3	22.1	16.3	17.6
$G_{cov}+M_{cov}$	us(HOST):animal + us(HOST):dam	30.1	31.3	53.3	63.6	17.2	34.6

954 **Table 2.** Phenotypic variance partitioning based on the best model from the model comparison. We used model 'G$_{cov}$' for all the traits (Table 1)

955 as it estimates cross-environmental genetic covariance and models with DIC difference <2 are usually considered equally supported. We

956 estimated host-specific additive genetic effects (V$_A$), proportion of additive genetic variance in total phenotypic variance (V$_P$) - heritability (h^2),

957 additive genetic variance standardized over trait mean - evolvability (CV$_A$), and cross-environmental genetic correlation (r$_G$). The estimates and

958 their credible intervals (in brackets) come from posterior distribution of sex-specific animal MCMC models.

Host type	Par.	Development duration		Body mass		Adult lifespan	
		daughters (N = 1 420)	sons (N = 1 384)	daughters (N = 1 431)	sons (N = 1 399)	daughters (N = 1 425)	sons (N = 1 389)
original	V$_A$	0.430 (0.105,0.678)	0.448 (0.208,0.876)	0.215 (0.147,0.317)	0.070 (0.034,0.114)	19.981 (11.423,26.851)	9.694 (5.578,14.841)
	V$_P$	1.768 (1.578,1.948)	2.167 (1.979,2.409)	0.501 (0.446,0.555)	0.310 (0.282,0.343)	39.267 (35.595,45.099)	28.298 (24.908,31.138)
	h^2	0.229 (0.084,0.387)	0.210 (0.082,0.374)	0.447 (0.327,0.615)	0.225 (0.122,0.358)	0.459 (0.301,0.624)	0.333 (0.218,0.505)
	CV$_A$	0.020 (0.012,0.027)	0.021 (0.014,0.030)	0.084 (0.066,0.097)	0.075 (0.058,0.100)	0.195 (0.148,0.226)	0.184 (0.146,0.234)
novel	V$_A$	0.349 (0.213,0.622)	0.527 (0.355,0.892)	0.250 (0.176,0.319)	0.103 (0.068,0.148)	16.664 (10.973,27.715)	16.304 (11.261,24.817)
	V$_P$	0.956 (0.853,1.083)	1.154 (1.028,1.339)	0.374 (0.340,0.426)	0.211 (0.186,0.236)	37.645 (33.595,43.241)	26.675 (23.763,31.265)
	h^2	0.365 (0.241,0.606)	0.495 (0.348,0.708)	0.649 (0.501,0.786)	0.513 (0.355,0.669)	0.507 (0.334,0.684)	0.627 (0.480,0.854)
	CV$_A$	0.020 (0.016,0.026)	0.026 (0.020,0.031)	0.077 (0.065,0.087)	0.082 (0.067,0.099)	0.168 (0.131,0.204)	0.208 (0.176,0.260)
	r$_G$	0.706 (0.277,0.981)	0.147 (-0.279,0.533)	0.901 (0.670,0.995)	0.996 (0.750,1.000)	0.790 (0.615,0.982)	0.967 (0.777,0.999)

Table 3. The effect of host type, other experimental variables and dam host preference on three offspring life-history traits (fixed-effects

estimates from the full animal models). The intercept always corresponds to the trait average on the original host type and the 'novel HOST

effect' is the effect of the novel host type (i.e. difference between the two means). The same applies to interaction terms, where the

interaction gives the estimate for the difference between the effect on the novel and the original host type. We embolden any credible

intervals (in brackets) that do not overlap zero. The estimates and their credible intervals come from posterior distribution of sex-specific

animal MCMC models.

Term	Development duration [days]		Body mass [mg]		Adult lifespan [days]	
	daughters (N = 1 420)	sons (N = 1 384)	daughters (N = 1 431)	sons (N = 1 399)	daughters (N = 1 425)	sons (N = 1 389)
Intercept (original HOST)	31.899 (31.713,32.103)	31.442 (31.187,31.607)	5.784 (5.660,5.904)	3.466 (3.414,3.593)	23.62 (22.432,24.739)	16.298 (15.338,17.259)
novel HOST effect	**-1.285 (-1.388,-1.115)**	**-1.148 (-1.296,-0.935)**	**0.728 (0.650,0.792)**	**0.349 (0.303,0.408)**	**3.338 (2.595,3.975)**	**2.355 (1.792,2.885)**
bean mass (original)	**0.145 (0.035,0.229)**	0.105 (-0.017,0.189)	**-0.083 (-0.130,-0.032)**	**-0.039 (-0.089,-0.011)**	**-0.499 (-0.930,-0.111)**	**-0.365 (-0.742,-0.038)**
dam mating order	0.060 (-0.002,0.136)	0.054 (-0.013,0.134)	0.007 (-0.037,0.052)	0.006 (-0.017,0.047)	-0.306 (-0.766,0.069)	0.210 (-0.116,0.587)
day mated (original)	**0.326 (0.206,0.454)**	**0.122 (0.028,0.291)**	0.058 (-0.006,0.149)	**0.074 (0.037,0.139)**	0.443 (-0.113,1.273)	**0.797 (0.199,1.262)**
dam preference for cowpea (original)	-0.073 (-0.180,0.031)	-0.025 (-0.174,0.068)	**0.086 (0.027,0.144)**	0.000 (-0.042,0.046)	0.219 (-0.366,0.733)	0.128 (-0.267,0.572)
Interaction: bean mass (novel)	-0.092 (-0.196,0.032)	0.031 (-0.131,0.138)	0.047 (-0.022,0.102)	0.040 (-0.011,0.091)	0.456 (-0.052,1.124)	-0.060 (-0.463,0.560)
Interaction: day mated (novel)	-0.017 (-0.141,0.142)	0.112 (-0.026,0.326)	-0.057 (-0.127,0.020)	-0.018 (-0.072,0.035)	-0.625 (-1.272,0.117)	-0.360 (-0.910,0.237)
Interaction: dam preference (novel)	0.051 (-0.056,0.206)	0.051 (-0.084,0.228)	**-0.081 (-0.146,-0.015)**	0.008 (-0.038,0.063)	-0.369 (-0.916,0.402)	-0.109 (-0.590,0.470)

Table 4. Phenotypic variance partitioning (A) and full model inference (B) based on G_{cov}

model for the non-Gaussian traits – larval survival and host egg laying preference. The

estimates for the variance parameters (A) are back-transformed to original data scale. The

effect of host type, the additional experimental variables and dam host preference (B) for

larval survival is on logit scale and for host preference on log scale. We embolden any

credible intervals (in brackets) that do not overlap zero. The intercept always corresponds to

the trait average on the original host type and the 'novel HOST effect' is the effect of the

novel host type (i.e. difference between the two means). The same applies to interaction

terms, where the interaction gives the estimate for the difference between the effect on the

novel and the original host type. The estimates and their credible intervals come from

posterior distribution of animal MCMC models.

A)

Host type	Par.	Larval survival	Host preference
		juveniles (N = 3 146)	daughters (N = 665)
original	V_A	0.006	<0.001
	V_P	0.094	0.671
	h^2	0.060	0.001
	CV_A	0.084	0.033
novel	V_A	0.003	0.001
	V_P	0.065	0.522
	h^2	0.040	0.003
	CV_A	0.055	0.071
	r_G	0.286* (-0.469,0.985)*	-0.046* (-0.905,0.948)*

979 * these estimates are on the transformed (latent) scale

981   B)

Term	Larval survival	Host preference
	juveniles (N = 3 146)	daughters (N = 665)
Intercept (original HOST)	2.868 (2.341,3.441)	-0.409 (-0.528,-0.338)
novel HOST effect	0.512 (-0.094,1.447)	**-0.247 (-0.324,-0.180)**
bean mass (original)	-0.161 (-0.353,0.050)	0.010 (-0.046,0.049)
dam mating order	-0.021 (-0.206,0.124)	-0.001 (-0.027,0.047)
day mated (original)	0.095 (-0.153,0.350)	-0.053 (-0.094,0.004)
dam preference for cowpea (original)	-0.042 (-0.258,0.183)	0.014 (-0.041,0.057)
Interaction: bean mass (novel)	-0.035 (-0.376,0.221)	0.017 (-0.075,0.076)
Interaction: day mated (novel)	-0.122 (-0.501,0.240)	-0.016 (-0.066,0.083)
Interaction: dam preference (novel)	0.242 (-0.106,0.549)	-0.037 (-0.096,0.049)

 Figure caption

Figure 1. Illustration of how maternal effects might differ between environments (host beans

in our study). Panel A) shows that ignoring the potential for environment-specific maternal

effects, as indicated by the colour of points representing maternal effects in the original (black

points) and the novel host (white points), could be misleading. Panel B) represents a scenario

where variance due to maternal effects (V_M) differs between the environments, i.e. the first

type of Maternal-by-Environment interaction (M×E). Panel C) illustrates the other type of M×E

where the rank of maternal effects differs between the hosts. The evidence for the shift in

ranking comes from the low cross-environmental correlation in maternal effects (r_M) despite

the maternal effects variance (V_M) being comparable between the hosts. Panel D) represents

a combination of B) and C).